

Cyclopropanone Hemiacetals

JACQUES SALAÜN

Laboratoire des Carbocycles, Université de Paris-Sud, 91405 Orsay Cedex, France

Received March 16, 1983 (Revised Manuscript Received June 8, 1983)

Contents

I. Introduction	619
II. Preparation of Cyclopropanone Hemiacetals	619
III. Chemical Properties	620
1. Reactions with Metal Ions	620
2. Reactions with Molecular Oxygen and Peroxides	621
3. Reactions with Acids and Bases	621
4. Solvolytic Reactions	622
5. Reactions with Nucleophiles	622
IV. Synthetic Applications	623
1. Ring Expansion into β -Lactams and 2-Pyrroline via 1-Aminocyclopropanols	623
2. Ring Expansion into Cyclobutanones via 1-Vinylcyclopropanols	625
3. Ring Expansion into Cyclopentanones via 1-(Trimethylsiloxy)-1-vinylcyclopropanes	625
V. Miscellaneous	628
1. Cyclopropanone Thiohemiacetal	628
2. [2 + 2] Cycloaddition of Cyclopropanone Diethyl Acetal	629
3. 1-Ethoxy-1-(trimethylsiloxy)cyclopropane as Homoenolate Anion Precursors	629
VI. Biological Activities of Cyclopropanone Precursors	629
VII. References	630

I. Introduction

Cyclopropanone chemistry has received considerable attention in recent years.^{1,2} Nevertheless, the chemistry of this unusually reactive class of ketones has previously found limited use in synthesis mainly because of the difficulties encountered in the preparation and handling of such strained systems. A first significant step in studying the chemistry of cyclopropanones has resulted from the discovery of derivatives capable of yielding the parent ketone or equivalent species in the reaction medium; among them, the cyclopropanone hemiacetal provides incontestably a convenient and storable source of the three-membered ring.² A second decisive step has recently been passed over with the discovery of a new preparation of this synthon, which now readily available, becomes a participant of choice in a number of useful chemical transformations including particularly cyclopropanol, methylenecyclopropane, cyclobutanone, β -lactam, γ -butyrolactone, cyclopentanone, and pyrroline derivative formation.

Two previous reviews concerning this challenging field, one by Turro in 1969¹ and one by Wasserman² in 1974² deal with the chemistry of cyclopropanone. The aim of this article is to review, up to now, the peculiar aspects of the cyclopropanone hemiacetal chemistry, and their recent applications in synthesis.




Jacques Salaün was born in Ollioules, France, and was graduated engineer from the Ecole Supérieure de Chimie de Caen in 1962. He obtained the Doctor-Engineer and Doctorat-ès-Sciences Physiques degrees from the University of Caen in 1965 and 1967, respectively, on research undertaken with Professor J. M. Conia on the stereochemistry and ring contraction of cyclobutanones. He did postdoctoral work on the [10]annulene system with Professor S. Masamune at Edmonton, Canada (1968), and in the field of vinyl cations with Professor M. Hanack at Saarbrücken (1974) and Tübingen (1976), Germany. He started his career at the Centre National de la Recherche Scientifique and became in 1975 Maître de Recherche in the Laboratoire des Carbocycles (associated to C.N.R.S.) at the University of Paris-Sud (Orsay). His main research interests include carbocation chemistry and the preparation and synthetic applications of small ring compounds.

II. Preparation of Cyclopropanone Hemiacetals

Due to its high reactivity cyclopropanone in the presence of alcohol immediately forms a hemiacetal sufficiently stable to permit isolation.¹⁻⁶


So, by adding an ethereal solution of diazomethane containing methanol to ketene, von Lipp has unexpectedly obtained the first cyclopropanone hemiacetal, i.e., the 1-methoxycyclopropanol (eq 1).³ Therefore,



the various methods of preparation of cyclopropanone^{1,2} lead to cyclopropanone hemiacetals either stepwise by adding alcohol to the preformed cyclopropanone^{4,5} or directly by synthesizing the three-membered ketone in the presence of methanol³ or ethanol.⁶ Thus, by using the most straightforward formation of cyclopropanone, ring-substituted cyclopropanone hemiacetals have been prepared either from alkyl (or aryl) diazo compounds and ketene or, alternatively from diazomethane and alkylketenes.


For instance, hemiacetals 1-3 have been obtained by reaction of dimethyl-,⁷ silyl-, and germylketenes⁸ with diazomethane and methanol, respectively; while, hemiacetals 4 (Ar = C₆H₅, *p*-ClC₆H₄, *p*-CH₃C₆H₄; R = H, CH₃) have been obtained by passing ketene through a solution of aryldiazomethane in CFC₃ or CCl₄ con-

taining an excess of methanol.⁹




However, these procedures are laborious and quite hazardous, particularly when carried out on a large scale.

To date, the cyclopropanone ethyl hemiacetal can be readily obtained in good yields by simple methanolysis¹⁰ of the 1-ethoxy-1-(trimethylsilyloxy)cyclopropane, product of the acyloin-type cyclization of commercial ethyl 3-chloropropanoate by sodium in refluxing ether in the presence of trimethylsilyl chloride¹⁵ (eq 2). The



preparation of 1-ethoxy-2-methyl-1-(trimethylsilyloxy)cyclopropane leading to 2-methylcyclopropanone hemiacetal has recently been reported following this procedure.¹²


An alternate pathway to these 1-alkoxy 1-silyloxy cyclopropanes, precursors of cyclopropanone hemiacetals, is provided by the addition of carbenes, generated from alkylidene iodide and diethylzinc, to the trimethylsilyl enol ethers of carboxylic esters (eq 3).¹³




$\text{R}_1 = \text{H}, \text{CH}_3, \text{C}_6\text{H}_{11}, \text{C}_6\text{H}_5; \text{R}_2 = \text{C}_6\text{H}_5, \text{CH}_3; \text{R}_1, \text{R}_2 = (\text{CH}_2)_4$

Similarly, it had been previously reported that the addition of the Simmons-Smith reagent ($\text{CH}_2\text{I}_2 + \text{Cu-Zn}$ couple) to 1-ethoxycyclopropyl acetate (or benzoate) provided 1-ethoxycyclopropyl acetate,¹⁴ which upon reaction with methanol yields the cyclopropanone methyl hemiacetal along with methyl acetate and methyl propionate.¹⁵


Furthermore, 1-(dimethylamino)cyclopropanol,¹⁶ prepared by addition of a tenfold excess of dimethylamine to cyclopropanone¹⁷ undergoes methanolysis to 1-methoxycyclopropanol (eq 4).¹⁸




Other pathways to the cyclopropanone hemiacetal consist of photochemical decarbonylation of tetramethylcyclobutanedione in ethanol¹⁹ or methanol²⁰ giving the corresponding tetramethylcyclopropanone hemiacetals and isobutyrate (eq 5) and electroreduc-



tion of highly alkylated α, α' -dihalo ketones²¹ in acetonitrile containing a twofold excess of methanol²² leading with good yields to the corresponding cyclopropanone hemiacetal (eq 6).



Although the reduction of the dimethylmalonic acid acyl of 1,3-dibromoacetone gave a low yield of cyclopropanone acyl, it has been shown that the reduction of this acyl with methanol is efficiently catalyzed upon addition of a trace amount of $\text{BF}_3 \cdot \text{OEt}_2$ complex to induce the immediate formation of cyclopropanone methyl hemiacetal (eq 7).²²




III. Chemical Properties

1. Reactions with Metal Ions


Like cyclopropanols,^{23,24} cyclopropanone hemiacetals are usually susceptible to oxidative cleavage at room temperature by metal ions with low oxidation potentials, such as $\text{Cu}^{\text{II}}, \text{Fe}^{\text{II}}, \text{Ce}^{\text{II}}$.^{24,25}

These one-electron oxidations lead to α cleavage of the highly unstable cyclopropyloxy radical intermediate into propionyl radical, which was detected by ESR,²⁴ (eq 8). The product composition depends on the metal




ion, its ligands, and on the concentration and addition rate of the reactants.²⁴ On the other hand, cyclopropanone dimethyl acetal fails to react under the same condition suggesting that the hydroxyl group of the hemiacetal is essential for the oxidation and is the first site of attack by the electrophilic metallic ion.²⁶

Arylcyclopropanone hemiacetals undergo oxidation by a variety of oxidizing agents. Although the common key intermediate is the ring-opened phenyl-stabilized β -propionate radical, the product distribution depends on the nature of the reactants (eq 9).²⁷




Because β -propionate radicals are not readily available from other sources, the oxidation of cyclopropanone hemiacetal in the presence of activated olefins constitutes a means of chain lengthening with a three-carbon unit (eq 10).²⁸




2. Reactions with Molecular Oxygen and Peroxides

While unsubstituted cyclopropanone hemiacetal reacts very sluggishly with oxygen, 2,2-dimethylcyclopropanone hemiacetal absorbs 1 equiv of molecular oxygen at room temperature and undergoes oxidative cleavage to give a β -hydroperoxy ester (eq 11).²⁹




A similar hydroperoxy ester has been isolated in 97% yield from the oxidation of tetramethylcyclopropanone hemiacetal.³⁰

Interestingly, the arylcyclopropanone hemiacetals when treated with oxygen lead mainly to peroxy lactones (eq 12).²⁷ The absorption of oxygen is strongly




accelerated in the presence of a catalytic amount of copper-amine complex; so a solution of phenylcyclopropanone methyl hemiacetal in methanol containing 2.5% Cu(II) nitrate-pyridine gives rise to about 80% uptake of oxygen within 30 min at room temperature, whereas the uncatalyzed reaction takes several hours.

The reaction mixture consists mainly of methyl benzoylacetate (eq 13).²⁷




Addition of a hydroperoxide to a solution of 2,2-dimethylcyclopropanone hemiacetal in CDCl₃ at room temperature, displays, almost instantaneously, negative peaks in the NMR spectrum.²⁹ This CIDNP effect can generally be initiated by any hydroperoxide or other source of alkoxy radicals.

In this respect, di-*tert*-butyl peroxide (TBPO) at 60 °C appears to be the most convenient source of *tert*-butoxy radicals.²⁹ The enhanced reactivity of the hydroxyl group is caused by stabilization of the developing oxygen radical by overlap with the orbitals of the neighboring C₁-C₂ bond, thus facilitating ring opening (eq 14).




The oxidation of phenylcyclopropanone hemiacetal by *tert*-butoxy radicals gives rise to a phenyl-stabilized radical from cyclopropyl ring opening that displays a CIDNP effect in agreement with simulated NMR spectra. This benzylic radical then undergoes either dimerization or disproportionation reactions, as shown by the observed absorption/emission (A/E) multiplet effect (eq 15).³¹




3. Reactions with Acids and Bases

The cyclopropanone methyl hemiacetal is smoothly converted into methyl propionate in the presence of acid (eq 16).^{32,33}




The ring cleavage is believed to proceed as in the case of cyclopropanols,^{32,33} via protonation of the cyclopropane C₁-C₂ bond.³⁴

On the other hand, 1-methoxy-2-phenylcyclopropanol reacts quantitatively with acids to yield phenylacetones (eq 17).³⁵ In the same way, cyclopropanone methyl




hemiacetal (or its hydrate) undergoes C₂-C₃ cyclopropyl bond cleavage in FSO₃H-liquid SO₂.³⁶ However, 1,1-diethoxy-2-phenylcyclopropane (2-phenylcyclopropanone ethyl acetal) is resistant to the action of 33% aqueous sulfuric acid.³⁷ Evidently, the presence of a hydroxyl group has a strong effect on the formation of the Woodward-Hoffmann type C₂-C₃ bond fission,^{38,39} but, as the hydroxyl group is less basic than the methoxy group, protonation of the hydroxyl group, followed by elimination of water and simultaneous ring opening appears to be unlikely.

The protonation of the methoxyl group of the hemiacetal and methanol elimination simultaneously with proton abstraction from the hydroxyl group giving rise to arylcyclopropanone as key intermediate, followed by O-protonation of this ketone with concerted C₂-C₃ bond rupture into 2-hydroxyallylic cation is reported as shown in eq 18.³⁵





The formation of the intermediate 2-arylcyclopropanone seems to be supported by the occurrence of a relatively rapid alkoxy group exchange (vide infra, the reaction of cyclopropanone hemiacetal with alcohols). Thus, on heating at 40 °C in perdeuterated methanol in the presence of sulfuric acid, 66% of methoxy group is exchanged within 1.5 h, whereas in the lack of acid, 50% of deuterated methyl hemiketal is obtained on heating at 40 °C for 45 h (eq 19).³⁵



Cyclopropanone hemiacetals, similarly to cyclopropanols, isomerize under basic conditions by C₁-C₂ ring fission into propionic esters;^{1,25} the ring opening is controlled by the formation of the most stable intermediate carbanion⁴⁰ and by steric factors.⁴¹

The aryl group of 2-arylcyclopropanone hemiacetals, which isomerize at room temperature with methanolic sodium methoxide or with weaker bases like tertiary amines, has a strong accelerating effect on the ring opening of the intermediate anion. The rearrangement products are exclusively 3-arylpropionic esters, in full agreement with the products obtained in the Favorskii reaction⁴² of α -aryl- α' -chloro ketones (eq 20).³⁵




4. Solvolytic Reactions


Nucleophilic substitution at cyclopropyl carbon is generally difficult and often takes place with considerable amounts of ring-opened product.⁴³ Likewise, upon solvolytic conditions, cyclopropyl derivatives usually undergo concerted ionization and disrotatory ring opening into allyl cation.³⁹ However, such a ring opening can be prohibited by steric⁴⁴ and conjugate interactions^{10a,45-48} or when an electron-donating substituent (RS,⁴⁹ ArS,⁵⁰ R₂N,^{18,51,52} RO⁵⁰ ... is attached to the three-membered ring in the α -position to the leaving group.

As a matter of fact, 1-chloro-1-methoxycyclopropane is prepared from cyclopropanone methyl hemiacetal via 1-methoxy-1-(tosyloxy)cyclopropane or via 1-methoxy-1-(trimethylsiloxy)cyclopropane and substitution by

chlorine of the trimethylsiloxy group with thionyl chloride (eq 21).⁴⁷




Then, the chlorine atom can be replaced quantitatively by methoxy, either on heating in pure methanol or in the presence of silver tetrafluoroborate at -30 °C. It is assumed that the chlorine substitution proceeds via a stabilized methoxy-substituted cyclopropyl cation (eq 22).⁴⁷



This reaction is similar to the solvolysis of 1-cyclopropyl-1-tosyloxy,^{45a} or 1-cyclopropyl-1-chlorocyclopropanes,^{45b} or 1-(*p*-anisyl)-1-chlorocyclopropane,⁴⁸ or 1-ethynyl-1-(tosyloxy)cyclopropanes,^{10a,46} which proceed mainly without ring rupture.


Thus, for instance, the solvolysis of the tosylates of 1-ethynylcyclopropanols, prepared from cyclopropanone ethyl hemiacetal (vide infra), proceeds via a stabilized mesomeric cyclopropyl-cyclopropylidenevinyl cation (eq 23).^{10a,46} However, although calculations indicate



that electron-releasing substituents may render 1-substituted cyclopropyl cations more stable than their 2-substituted allyl counterparts,⁵³ the theoretical expectations of the 2-substituted allyl cation ring closure into cyclopropyl cation have not been obtained experimentally, in spite of several attempts (eq 23).⁵⁴


5. Reactions with Nucleophiles

On standing at 25 °C for 1 week with methanol 65% of cyclopropanone ethyl hemiacetal is converted into its methyl hemiacetal; the conversion is completed within 2 weeks (eq 24).¹⁵



Reciprocally, excess of ethanol transforms cyclopropanone methyl hemiacetal into the corresponding ethyl hemiacetal. These changes are pictured, as shown in eq 24 in terms of a cyclopropane-cyclopropanone hemiacetal equilibrium.¹⁵


Cyclopropanone ethyl hemiacetal reacts with aniline at 25 °C to yield 1-(phenylamino)cyclopropanol and 1,1-dianilino-cyclopropane, (eq 25).



With excess aniline, the diaminocyclopropane is produced, almost quantitatively.^{2,15} Besides reactions with alcohols and amines, the cyclopropanone hemiacetal undergoes nucleophilic attack by various other reagents.


First of all, the reactions of the hemiacetal with carbanionic reagents such as vinyl (eq 26)^{6,55} and acetylenic (eq 27)^{6,10a,56} Grignard reagents leading to 1-vinyl and 1-alkynylcyclopropanols, respectively, have been reported.

Two equivalents of the Grignard reagent are involved in eq 27^{10a,56} due to the equilibrium, cyclopropanone hemiacetal = cyclopropanone + ethanol. On the other hand, the hemiacetal does not undergo nucleophilic reactions with organolithium reagents such as lithium




cyanide,^{57b} ethynyllithium,^{57b} or aryllithium.⁵⁸

A simple solution to this difficulty was proposed by Brown and Rao;⁵⁸ thus, upon treatment with an equimolar amount of methylmagnesium iodide, the cyclopropanone ethyl hemiacetal is converted into a species, likely the magnesium salt derivative, which is then able to react with aryllithium to give the expected 1-arylcyclopropanol (eq 28).⁵⁸



So, with its covalent O-Mg bond, the cyclopropanone hemiacetal magnesium salt readily breaks down into cyclopropanone, which then reacts with nucleophiles; whereas, the intermediate lithium salt, obtained in the reaction of the hemiacetal with organolithium reagents, has a steady ionic O-Li bond and appears unreactive (eq 29).




As shown in Scheme I, this cyclopropanone ethyl hemiacetal magnesium salt (5) reacts with lithium aluminum hydride to give cyclopropanol (6) in 75% yield (whereas the reaction of LiAlH₄ with the hemiacetal itself leads to cyclopropanol in 45% yield);^{57b} it reacts with lithium cyanide to yield cyclopropanone cyanohydrin (7),^{59a} a precursor of cyclobutanone^{59b} and of cleonine,^{59c} in 75% yield; it reacts with organometallic reagents (Mg, Li, Al) to lead to 1-substituted cyclopropanols 8, 9, and 10 but undergoes ring opening with the Reformatsky reagent BrZnCH₂CO₂Et; it reacts with alkylidenephosphoranes⁵⁷ offering an alternate pathway to alkylidenecyclopropanes (11), usually prepared from


the Wittig reaction of cyclopropylidenephosphorane with carbonyl compounds,⁶⁰ it reacts smoothly with phosphonate carbanion to give ethyl cyclopropylideneacetate (12); but it is not reactive with nitrogen and sulfur ylides, such as diazomethane and dimethylsulfonium and dimethylloxosulfonium methyllide. All these readily available cyclopropanol or methylenecyclopropane derivatives provide convenient keys to enter the field of small ring compounds.⁵⁷

The addition of arylidenetriphenylphosphoranes to the magnesium salt depends on the nature of the substituent of the phenyl ring; thus, an electron-withdrawing para substituent (e.g., X = NO₂) prevents the Wittig reaction (eq 30).⁵⁷


Although it has been corroborated that no reaction occurs between cyclopropanone hemiacetal itself and



phosphoranes under these conditions (i.e., in ether at reflux for ~40 h), benzyl 2-bromocyclopropylideneacetate is prepared from the hemiacetal and the suitable phosphorane in refluxing benzene and in the presence of benzoic acid (eq 31).⁶¹



Then, cyclization with a 4-mercaptoazetidinone (for the preparation of β-lactam from cyclopropanone hemiacetal vide infra) leads to spirocyclopropane norpenicillanic acid derivatives, endowed with antibiotic activities (eq 32).⁶¹




Like cyclopropanone hemiacetal magnesium salt, 1-acetoxycyclopropanol reacts smoothly with a variety of nucleophiles including CN⁻, N₃⁻, R₂N⁻, RO⁻, and RS⁻ to give the corresponding cyclopropanone adducts.¹⁶ Furthermore, 1-ethoxycyclopropyl acetate, reacts with lithium aluminum hydride or Grignard reagents to lead to cyclopropanol and 1-alkylcyclopropanols, respectively.^{14,15}


IV. Synthetic Applications

1. Ring Expansion Into β-Lactams and 2-Pyrrolidine via 1-Aminocyclopropanols

The importance of β-lactams in the penicillins,⁶² cephalosporins,⁶³ and thienamycin⁶⁴ and the recent discovery of antibiotic activity among monocyclic β-lactams⁶⁵ such as, for example, norcardicins 13 or the β-lactamase inhibitor clavulanic acid (14)⁶⁷ have recently


SCHEME I. Reactions of the Cyclopropanone Hemiacetal Magnesium Salt with Some Nucleophilic Reagents⁵⁷


intensified research toward the synthesis of this system.^{67,68}




Among the different procedures that have been developed for incorporating a 2-azetidinone unit,⁶⁹ the ring expansion of cyclopropanol amines provides a simple and convenient route to these attractive small-ring compounds.⁶⁸⁻⁷¹


Previously, 1-aminocyclopropanols were directly obtained by adding amines, in the place of methanol (vide supra, eq 1) to the product of reaction of diazomethane with ketene, at -78°C (eq 33).^{70,72}



These carbinol amines may be converted into *N*-chloro derivatives by using *tert*-hypochlorite; then, treatment of the *N*-halo derivative with silver ion in acetonitrile leads to β -lactam (eq 34).⁷⁰




Alternatively, such β -lactams are available from the reaction of cyclopropanone hemiacetal with sodium azide in buffered acetone (eq 35).⁸




More recently, like cyclopropanone hemiacetal itself (vide supra, eq 2) 1-aminocyclopropanols are readily available following the Rühlmann procedure, from the

reductive cyclization of the piperidine of 3-chloropropionic acid by sodium in the presence of trimethylchlorosilane and conversion of the intermediate silyl ether into the 1-piperidinocyclopropanol by methanolic tetrabutylammonium fluoride (eq 36).⁷³




The hydroxyl group of 1-(alkylamino)cyclopropanols is easily displaced by all common nucleophiles including amines, alcohols, thiols, hydrogen cyanide⁷⁴ and hydrazoic acid,¹⁸ C-H acidic compounds, which can react in Mannich-type condensations,⁷⁵ Grignard reagents or indole, *N*-methylpyrrole, silyl enol ethers, and species containing active methylene groups in the presence of titanium tetrachloride.⁷³


The reaction does not proceed by direct substitution (S_N2) but by the intermediary of a *N,N*-dialkylcyclopropaneiminium ion (eq 37).




Thus, for example, upon treatment with KCN in acetic acid a nitrile derivative is obtained in good yields, which then can add cyclopropyllithium in ether at -78°C (eq 38).



On heating in xylene, the imine undergoes vinylcyclopropane-cyclopentene ring enlargement to the enamine, followed by isomerization into the cyclic imine tautomer (eq 39).



Although prolonged heating of the pyrroline does not lead to further ring expansion, a second cyclopropylimine rearrangement takes place on heating in the presence of anhydrous hydrobromic acid at 140°C for 10 min, giving after hydrolysis the pyrrolizidinone (eq 40).⁷⁴




This sequence opens a route to pyrrolizidines that is of obvious interest in natural product synthesis.

2. Ring Expansion into Cyclobutanones via 1-Vinylcyclopropanols


Four-carbon annelation reactions⁷⁶ have been used in recent years as a key step in the construction of complex organic molecules and many synthetically useful transformations of cyclobutanones have been described.⁷⁷

Besides the initial preparation based on the addition of diazomethane to ketenes,³ the main recent approaches to cyclobutanones include the cycloaddition of dichloroketene to active olefins and reductive halogen removal,⁷⁸ the epoxidation of alkylidenecyclopropanes (vide supra) into oxaspiropentanes, which then undergo lithium-induced ring enlargement,⁷⁹⁻⁸¹ the condensation of cyclopropyl sulfur ylides with aldehydes and ketones leading to intermediate oxaspiropentanes,⁷⁷ the dithiane or methyl methylthiomethyl sulfoxide cyclization of 1,3-dihaloalkanes,⁸² and the ring enlargement of 1-vinylcyclopropanols.^{6,55,81b,83}


So, vinylcyclopropanols, readily available from the cyclopropanone hemiacetal and vinylmagnesium bromides (vide supra, eq 26), undergo ring expansion into cyclobutanones with a variety of electrophilic reagents; with hydrobromide, perbenzoic acid, and *tert*-butyl hypochlorite, 2-alkyl-, 2-hydroxymethyl-, and 2-chloroalkylcyclobutanones are obtained, respectively.⁶ The rearrangement takes place most probably through the intermediate cyclopropylcarbinyl cation (eq 41).



With trioxymethylene and dibenzylamine hydrochloride in refluxing ethanol 1-vinylcyclopropanol undergoes a Mannich-type reaction to yield the corresponding 2-(2-(dibenzylamino)ethyl)cyclobutanone (eq 42).⁵⁵




When 2-methyl-1-vinylcyclopropanol is treated with dry hydrogen bromide in methylene chloride at 0 °C for 5 min a mixture of *trans*- and *cis*-2,3-dimethylcyclobutanones (ratio 3:1) is formed (eq 43).⁵⁵ This




result is consistent with the preferred migration of the

more highly substituted carbon atom, recently observed in the peracid oxidation of methylenecyclopropanes.⁸⁴

On simple heating at 100 °C in liquid phase or in a gas chromatograph, 1-cyclopentyl- and 1-cyclohexenylcyclopropanols are quantitatively converted into the corresponding spiroketones (eq 44).^{81b,83}



The stereochemistry and mechanism of such a thermal ring enlargement have been determined by the examination of the product of rearrangement of a perdeuteriocyclopropanol. Although thermally allowed, a concerted process⁸⁹ is ruled out by the occurrence of an anti configuration for the deuterium atom on the carbon 5 of the spiroketone, implying an intramolecular stereoscopic *cis* addition of this deuterium from O-D to the double bond (eq 45).⁸³




Such cyclobutanones appear to be very useful intermediates for the creation of a wide range of structural units. Thus, for instance, these cyclic ketones undergo unusually facile Baeyer-Villiger oxidation, even with basic hydrogen peroxide, to lead to γ -butyrolactones,⁷⁷ which are known to cyclize to cyclopentenones upon acid treatment.⁸⁵ They can undergo ring cleavage by nucleophiles,^{77,86} or one-carbon ring expansion reactions with diazomethane to offer an alternative three-carbon annelation method.⁸⁷

3. Ring Expansion into Cyclopentanones via 1-(Trimethylsiloxy)-1-vinylcyclopropanes

The synthesis of cyclopentanoid compounds is still a subject of present interest because of the discovery of a growing number of naturally occurring substances of biological importance that contain the five-membered ring moiety.⁸⁸ Besides the methodology based essentially upon the conjugate addition of cuprates to cyclopentenones,⁸⁹ the recent approaches include intramolecular ring closure of acyclic precursors,⁹⁰ [3 + 2]⁹¹ and [4 + 1]⁹² cycloaddition reactions, ring contraction,⁹³ or ring enlargement of cyclic precursors.^{77,87,94-97}


Among them, the thermal vinylcyclopropane-cyclopentene rearrangement⁹⁸ of 1-siloxy-1-vinylcyclopropanes into cyclopentanone silyl enol ethers that are then able to undergo either further regiospecific alkylation into 2,3-disubstituted cyclopentanones⁹⁴⁻⁹⁶ or dehydrosilylation into cyclopentenones^{99,100} constitutes an efficient three-carbon annelation process (eq 46).



The silylated vinylcyclopropanols can be prepared from the base-induced ring opening of the oxasipropentanes obtained in the condensation of diphenylsulfonium cyclopropylide with carbonyl compounds,^{77,99} from the silver Simmons-Smith cyclopropanation of the silyl enol ether of α -ethylenic ketones,⁹⁵ from 1,2-disiloxycyclobutene,¹⁰⁰ or from the cyclopropanone ethyl hemiacetal.⁹⁷

Effectively, upon treatment with 1 equiv of methylmagnesium bromide and 1 equiv of acetylenic magnesium bromide or lithium, the hemiacetal is converted, in good yields, into propargylic cyclopropanols (vide supra, Scheme I).

Lithium aluminum hydride reduction of acetylenic cyclopropanols in refluxing tetrahydrofuran for 1 h leads exclusively to the (*E*)-1-vinylcyclopropanols;⁹⁷ while, reduction with dicyclopentadienyltitanium hydride, prepared from isobutylmagnesium halides and a catalytic amount of $(\eta^5\text{-C}_5\text{H}_5)_2\text{TiCl}_2$ ¹⁰¹ offers exclusively the (*Z*)-1-vinylcyclopropanols,¹⁰² (eq 47). Then,




the cyclopropanols are O-silylated by action of trimethylsilyl chloride and triethylamine in the presence of a catalytic amount (5%) of dimethyl sulfoxide.¹⁰³

An advantage of this procedure over the previous methods^{77,95} resides in the use of cheap and readily available starting materials. In this way, the (*Z*)- and (*E*)-1-(trimethylsiloxy)-1-vinylcyclopropanes reported in Table I are obtained in good yields.


While 1-vinylcyclopropanols in general, undergo on heating at 100 °C quantitative $\text{C}_3 \rightarrow \text{C}_4$ ring enlargement into cyclobutanones (vide supra, eq 44), on the other hand, silylated (*Z*)- and (*E*)-1-vinylcyclopropanols undergo quantitative regiospecific thermal $\text{C}_3 \rightarrow \text{C}_5$ ring enlargement into 1-siloxycyclopentenes either on heating in the liquid phase in sealed tube at 300 °C for 30 min^{95,97,102} or by passing through a conditioned hot tube at 330 °C with a contact time of $<4 \text{ s}$ ⁹⁴ or by flash thermolysis at 600 °C for 10 ms.^{94b,97,102}

Thus, for instance the 1-cyclopentenylcyclopropanol can be rearranged either into the spiro[3.4]octan-1-one⁸³ or, after O-silylation, into the regiospecific enol silyl ether of bicyclo[3.3.0]octan-1-one^{94,95} (eq 48).




The effect of the trimethylsiloxy group on the thermal vinylcyclopropane \rightarrow cyclopentene rearrangement that requires for 15 49.6 kcal/mol,^{104,105} appears highly dependent on the position of this substituent on the system. Thus, placed on the cyclopropane ring as in 16 this substituent facilitates the rearrangement by about 5 kcal/mol¹⁰⁶ whereas, placed on the double bond of 17 the siloxy group hampers the rearrangement by 3 kcal/mol.^{96,106}

This effect is responsible for the specificity of the




rearrangement. As a matter of fact, when two vinylcyclopropane moieties can be a priori involved in the rearrangement, the driving substituent effect of the siloxy group leads exclusively to the 3-cyclopropylcyclopentanone silyl enol ether (eq 49).⁹⁷

On the other hand, trimethylsilyl substituents that are known to destabilize the α -radical¹⁰⁷ have an op-




posite effect. Thus, it has been observed that pyrolysis at 570 °C of 1-(1-(trimethylsilyl)cyclopropyl)-1-cyclopropylethylene leads exclusively to 1-(1-trimethylsilyl)cyclopropyl)cyclopentene;¹⁰⁸ contrary to the siloxy group, the silyl substituent exerts a rate-retarding effect on the rupture of the cyclopropane ring (eq 50).



A similar exclusive participation of the less substituted cyclopropane ring has also been reported in the thermal rearrangement of 1-(1-piperidinocyclopropyl)-1-cyclopropylketimine (vide supra, eq 39).⁷⁴

The conjugation of the vinylic bond of the vinylcyclopropane system with unsaturated substituents lowers the activation energy required for the thermal rearrangement,¹⁰⁵ thus, for example, the activation energy for the rearrangement of the 1-(trimethylsiloxy)-1-(4-trimethylsilyl)-1-buten-3-ynyl)cyclopropane is estimated to be less than 40 kcal/mol indicating additional substituent effects of the siloxy group and of the conjugated triple bond for the ring enlargement, which occurs on heating at 226 °C only (eq 51).⁹⁷



The thermal $\text{C}_3 \rightarrow \text{C}_5$ ring enlargement occurs indifferently either from the (*Z*) or from the (*E*)-1-(trimethylsiloxy)-1-vinylcyclopropanes, under the same conditions (e.g., flash thermolysis at 600 °C, eq 52).¹⁰²






TABLE I. Cyclopentanones from Cyclopropanone Hemiacetal via 1-Siloxy-1-vinylcyclopropanes⁹⁷

	$R_1-C\equiv C-M$						
	R_1	M	R_1	R_1	R_1	R_1	R_2
a	CH ₃	MgBr	CH ₃	CH ₃	CH ₃	CH ₃	CH ₃
b	Ph	MgBr	Ph	Ph	Ph	Ph	CH ₃
c	<i>p</i> -CH ₃ C ₆ H ₄	MgBr	<i>p</i> -CH ₃ C ₆ H ₄	<i>p</i> -CH ₃ C ₆ H ₄	<i>p</i> -CH ₃ C ₆ H ₄	<i>p</i> -CH ₃ C ₆ H ₄	CH ₃
d	<i>c</i> -C ₃ H ₅	MgBr	<i>c</i> -C ₃ H ₅	<i>c</i> -C ₃ H ₅	<i>c</i> -C ₃ H ₅	<i>c</i> -C ₃ H ₅	CH ₃
e	CH ₂ =CH	MgBr	CH ₂ =CH	CH ₂ =CH	CH ₂ =CH	CH ₂ =CH	CH ₂ =CH-CH ₂
f	Me ₃ SiC≡C	Li	Me ₃ SiC≡C	Me ₃ SiC≡C	Me ₃ SiC≡C	Me ₃ SiC≡C	CH ₂ =CH-CH ₂

SCHEME II. Total Synthesis of (±)-11-Deoxyprostaglandin E₂ Methyl Ester from the Cyclopropanone Ethyl Hemiacetal^{2,97}


^a (a) IMgCH_3 , $\text{Me}_3\text{SiC}\equiv\text{C}-\text{C}\equiv\text{ClLi}$,¹²² 67%; (b) DHP, PPTS, HCCl_3 , 100%; (c) KF, $2\text{H}_2\text{O}$, DMF, 91%; (d) *n*-BuLi, THF, $\text{CH}_3(\text{CH}_2)_4\text{CHO}$, 70%; (e) EtOH, PPTS, 55 °C, 95%; (f) LiAlH_4 , THF, 65 °C, 87%; (g) ClSiMe_3 , NEt_3 , Me_2SO , 83%; (h) Flash vacuum thermolysis at 600 °C, 100%; (i) NH_2Li , NH_3 , methyl *cis*-7-bromo-5-heptenoate, 44.5%.

However, although the thermal $\text{C}_3 \rightarrow \text{C}_7$ ring enlargement of butadienylcyclopropanes into cycloheptadiene derivatives is well-known,^{109,110} the thermolysis of the (*E*)-1-butadienyl-1-(trimethylsiloxy)cyclopropane leads exclusively to the 3-vinylcyclopentanone silyl enol ether, (eq 53).⁹⁷




Rate acceleration of 10^{10} – 10^{17} times the rate of the Oxy-Cope rearrangement are induced by potassium alkoxide;¹¹¹ this oxyanionic substituent effect provides, among others, a useful stereocontrolled synthetic entry to prostaglandins.¹¹²

On the other hand, it has been reported that the lithium salts of 2-vinyl-1-cyclopropanols undergo the vinylcyclopropane-cyclopentene rearrangement at room temperature (eq 54).¹¹³



But unfortunately, such a dramatic rate enhancement does not occur with the potassium salts of the 1-vinylcyclopropanols and of the enols of cyclopropyl ketones, as shown in eq 55.¹¹⁴ Metal-promoted vi-




nylcyclopropane \rightarrow cyclopentene rearrangements using stoichiometric amounts of rhodium ($(\text{C}_2\text{H}_4)_2\text{Rh}(\text{acac})$)¹¹⁵ have also been attempted, however with limited success.¹¹⁶

Acid or basic hydrolyses of the thermolysis products, i.e., the 1-siloxycyclopentenones, unmask the carbonyl groups and provide the corresponding cyclopentanones.^{77,95,97} Treatment with methyl lithium¹¹⁷ or with lithium amide in ammonia¹¹⁸ allows the generation of the lithium enolates derived from the 1-siloxycyclopentenones, which can then be regioselectively alkylated to introduce further alkyl groups leading to 2,3-disubstituted cyclopentanones, which encompass a broad class of important, naturally occurring substances.^{88,100,119}

Thus, upon treatment with methyl lithium and methyl iodide or allyl bromide, the α,β -disubstituted cyclopentanones reported in the Table I, are obtained in 54–66% yields.⁹⁷

Furthermore, the O-silylated cyclopentanone enol ethers can be directly alkylated in the presence of Lewis acids.¹²⁰ For example, phenylthioalkylation of silyl enol ethers using α -chloroalkyl phenyl sulfides in the presence of TiCl_4 or ZnBr_2 leads after reductive Raney nickel desulfurization or oxidative sodium metaperiodate desulfurization to α -alkylated or α -alkylenated cyclopentanones, respectively, in high yields (eq 56).¹²¹




Alkylation of enol silyl ethers is also effectively promoted by fluoride ion.¹²²


β -Acetylenic ketones are valued synthetic precursors but owing to the inability of organocuprates to transfer alkynyl groups¹²³ the usual 1,4-addition cannot be employed to effect the alkylation of cyclopentanones. Although it has been reported recently that dialkyl-

alkynylalanes are capable of selective alkylation of cyclopentenones in the presence of catalyst prepared from $\text{Ni}(\text{acac})_2$ and $i\text{-Bu}_2\text{AlH}$,¹²⁴ a very convenient pathway to these challenging β -alkynylcyclopentanones is provided by the thermal rearrangement of the readily available 1-(trimethylsiloxy)-1-(4-(trimethylsilyl)-1-buten-3-ynyl)cyclopropane into the 3-(silylethynyl)cyclopentanone silyl enol ether (vide supra, Table I, entry f).⁹⁷

α -Methylene carbonyl compounds are also useful synthetic intermediates and the number of known natural substances containing the α -methylene ketone moiety is growing rapidly.¹²⁵ Dimethylmethyleneammonium iodide¹²⁶ appears sufficiently electrophilic to react with silylated enol ethers in a Mannich reaction to provide β -amino ketones, which lead, after addition of methyl iodide and base-induced elimination, to regiospecific α,β -unsaturated ketones.¹²⁷ Upon similar treatment, the 2-methylene-3-(silylethynyl)cyclopentanone is obtained in 35% overall yield from the corresponding silylated enol ether (eq 57).⁹⁷



Similar regiospecificity is observed when the silylated enol ethers are first cleaved with methyl lithium, and the lithium enolate trapped by formal at -78°C to give a (hydroxymethyl)cyclopentanone, which then undergoes dehydration by means of methanesulfonyl chloride and a tertiary amine, to give the same α -methylene-cyclopentanone in 40% overall yield (eq 58).⁹⁷



Such lithium enolates are also trapped by dimethyl(methylene)ammonium trifluoroacetate.¹²⁸ An alternate method of α -methylenation relies on regiospecific phenylthiomethylation¹²¹ followed by oxidative removal of sulfur.^{121,129} (vide supra, eq 56).

α -Methylene-cyclopentanones have been used with success as key intermediates in the synthesis of prostaglandins,¹³⁰ so, the 3-((trimethylsilyl)ethynyl)-2-methylene-cyclopentanone obtained in eq 57 and 58 from the cyclopropanone hemiacetal (vide supra) really constitutes a crucial intermediate in the synthesis of these challenging compounds.⁹⁷ However, in order to

illustrate the efficiency of the cyclopropanone hemiacetal, as synthon toward complex molecules the total convergent synthesis of the (\pm)-11-deoxyprostaglandin E_2 methyl ester is reported in Scheme II.


So, upon treatment with 1 equiv of methylmagnesium bromide (vide supra) and 1 equiv of (trimethylsilyl)-butadiynyl lithium,¹³¹ the cyclopropanone ethyl hemiacetal is transformed into the 1-((trimethylsilyl)butadiynyl)cyclopropanol. Addition of dihydropyran in the presence of a catalytic amount (10 mol %) of PPTS,¹³² desilylation by potassium fluoride in DMF,¹³³ metalation with 1 equiv of $n\text{-BuLi}$ and condensation with hexanal lead, after elimination of the protecting THP group,¹³² to a dipropargylic diol. Then, lithium aluminum hydride reduction of the two triple bonds followed by double silylation in the presence of dimethyl sulfoxide¹⁰³ give the silylated (E,E)-1-butadienylcyclopropanol derivative. Flash thermolysis at 600°C provides quantitative ring enlargement into the expected 1-(trimethylsiloxy)-3-(3-(trimethylsiloxy)-1-octenyl)-cyclopentene. This prostaglandin precursor,¹¹⁸ is finally alkylated by means of lithium amide in liquid ammonia and a fourfold excess of methyl (Z)-7-bromo-5-heptenoate to yield a mixture of (\pm)-11-desoxyprostaglandin E_2 and (\pm)-11-desoxy-15-epiprostaglandin E_2 methyl esters, readily isolable by liquid chromatography, Scheme II.⁹⁷

V. Miscellaneous


1. Cyclopropanone Thiohemiacetal

It has been shown that 1-aminocyclopropanols undergo substitution of the hydroxyl group by all common nucleophiles, via the intermediary of a cyclopropane-dialkyliminium ion (vide supra, eq 37). Likewise, the formation and reactivity of 1-alkylthiocyclopropanols have been recently reported.


Thus, the cyclopropanone thiohemiacetals are formed upon simple addition of alkane- or arene-thiols to cyclopropanone (eq 59).¹³⁴



They are transformed into 3-chlorothiopropanoate, upon oxidation with FeCl_3 , in a reaction similar to the one-electron oxidation of cyclopropanone hemiacetal (vide supra) (eq 60).¹³⁴




The hydroxy group of the thiohemiacetal can be replaced by halogen, upon treatment with HBr or HCl , via a relatively stable (alkylthio)cyclopropyl cation (eq 61).¹³⁴




Like 1-chlorocyclopropyl methyl ether (vide supra,

eq 22⁴⁷) 1-chlorocyclopropyl sulfides are methanolized via a sulfur-stabilized cyclopropyl cation without any rupture of the three-membered ring (eq 62).¹³⁵ Such




1-halocyclopropyl sulfides can be transformed into 1-substituted cyclopropyl sulfides with a variety of nucleophiles including formate, iodide, azide, hydrosulfide, furan, hydride¹³⁴

1-Bromocyclopropyl sulfide undergoes oxidation with hydrogen peroxide into sulfoxide and sulfone (eq 63).¹³⁴




It reacts with sodium methanethiolate in protic or aprotic solvents to yield cyclopropyl sulfides. The reduction proceeds via the 1-(methylthio)cyclopropyl cation; then, attack by thiolate on the sulfur forms a sulfonium cyclopropylide, which by intramolecular hydrogen shift, gives the reduction product (eq 64).¹³⁶



2. [2 + 2] Cycloaddition of Cyclopropanone Diethyl Acetal

Cyclopropanones undergo [4 + 3] cycloaddition with conjugated dienes such as furan and cyclopentadiene at 0 °C or *N*-methylpyrrole at -78 °C to yield seven-membered ring adducts.¹³⁷ On the other hand, cyclopropanone ethyl hemiacetal does not add cyclopentadiene or furan in attempted similar cycloadditions carried out from 0 to 37 °C.¹³⁸ Reaction between 2,3-dichloro-5,6-dicyano-1,4-benzoquinone and cyclopropanone methyl hemiacetal yields thermally unstable cycloadducts.¹³⁹

However, cyclopropanone acetals (or thioacetals) and tetracyanoethylene give, in a thermal [$\sigma 2 + \pi 2$] cycloaddition, 2,2,3,3-tetracyanocyclopentanone acetals (or thioacetals). With respect to the parent compound methyl substituents on the cyclopropane ring retard the cycloaddition reaction, whereas a phenyl group, (see eq 65) accelerates the reaction.¹⁴⁰




3. 1-Ethoxy-1-(trimethylsiloxy)cyclopropane as Homoenoate Anion Precursors^{12,141}


The concept of homoenoate anion has become the

major subject of an unceasing number of recent papers.¹⁴² In this respect 1-ethoxy-1-(trimethylsiloxy)cyclopropane, the new source of cyclopropanone ethyl hemiacetal (vide supra, eq 2), can work well as a homoenoate anion precursor, i.e., the β -anion of ethyl propionate.

Thus, it reacts with titanium tetrachloride to provide, in high yield, the isolable dimeric titanium homoenoate as deep purple crystals (eq 66).^{12,141}



This homoenoate undergoes addition of a wide range of saturated aliphatic and aromatic aldehydes to yield γ -hydroxy esters, usually isolated as the derived γ -lactones (eq 67).^{12,141} Other common Lewis acids (i.e.,




Hg(OAc)₂, SnCl₄, Bu₃SnOTf) besides TiCl₄ are totally ineffective or give different products.

VI. Biological Activities of Cyclopropanone Precursors


Both naturally occurring and synthetic cyclopropanol compounds are irreversible inhibitors of particular target enzymes.¹⁴³ The inactivation derives from the ability of the cyclopropanone precursors, i.e., 1-substituted cyclopropanols, to undergo nucleophilic addition of the enzyme.

Thus, for example, the toxic mushroom constituent coprine,¹⁴⁴ produced by the *Coprinus atramentarius* mushroom, undergoes hydrolysis of the amide bond by a glutaminase enzyme to provide the 1-aminocyclopropanol, which then forms a hemithioacetal (vide supra, eq 37) upon nucleophilic addition of the active cysteinyl SH group of an aldehyde dehydrogenase to yield inactive aldehyde dehydrogenase molecules (eq 68).¹⁴⁵ The activity, denoted as "antabuse activity",



ascribed to the inhibition of the enzyme aldehyde dehydrogenase is responsible for acetaldehyde intoxication after alcohol consumption.

Some cyclopropylamines are antidepressant inhibitors of monoamine oxidase. Thus, the tranlylcypamine 18 is oxidized by the amine oxidase into cyclopropylimine, which then is able to capture an active site of the enzyme to form inactive cyclopropanone hemithioaminal adducts.¹⁴⁶ Similarly, the cytochrome mono-



oxygenase P₄₅₀ isozyme in mammalian liver probably oxygenates also the benzylcyclopropylamine 19 into a cyclopropanone derivative.¹⁴⁷ On the other hand, the methylenecyclopropane amino acid 20 or hypoglycin A, a toxic metabolite of the Jamaica ackee fruit¹⁴⁸ is oxidized into an acyl-CoA thiolester that blocks reversible the isovaleryl-CoA dehydrogenase.¹⁴³ The methylenecyclopropane 21 is a suicide substrate for the flavoenzyme desaturase, butyryl-CoA dehydrogenase.¹⁴⁹

Finally, the 1-alkynylcyclopropanols prepared from the cyclopropanone ethyl hemiacetal (vide supra, eq 33^{10a} and Scheme I⁵⁷ present antibacterial activities and are inhibitors of the PG synthetase (indomethacin).¹⁵⁰

Other interesting biological activities of these compounds can be expected, so it appears worthwhile to develop efficient routes to cyclopropanone-derived compounds.

VII. References

- (1) Turro, N. J. *Acc. Chem. Res.* **1969**, *2*, 25.
- (2) Wasserman, H. H.; Clark, G. M.; Turley, P. C. *Top. Curr. Chem.* **1974**, *73*.
- (3) von Lipp, P.; Koster, R. *Liebigs Ann. Chem.* **1932**, *499*, 1; *Chem. Ber.* **1931**, *64*, 2823.
- (4) Turro, N. J.; Hammond, W. B. *J. Am. Chem. Soc.* **1966**, *88*, 3672; *Tetrahedron* **1968**, *24*, 6029.
- (5) Schaafsma, S. E.; Steinberg, H.; de Boer, T. J. *Recl. Trav. Chim. Pays-Bas* **1966**, *85*, 70.
- (6) Wasserman, H. H.; Cochoy, R. E.; Baird, M. S. *J. Am. Chem. Soc.* **1969**, *91*, 2375.
- (7) Turro, N. J.; Hammond, W. B. *Tetrahedron* **1968**, *24*, 6017.
- (8) Zaitseva, G. S.; Bogdanova, G. S.; Bankov, Y. I.; Lutsenko, I. F. *J. Organometal. Chem.* **1976**, *21*, C21. Bogdanova, G. S.; Zaitseva, G. S.; Lutsenko, I. F. *Zh. Obsch. Khim.* **1979**, *49*, 2784.
- (9) Bakker, B. H.; Steinberg, H.; de Boer, T. J. *Recl. Trav. Chim. Pays-Bas* **1975**, *94*, 50.
- (10) (a) Salaün, J. *J. Org. Chem.* **1976**, *41*, 1237. (b) Salaün, J.; Marguerite, J. *Org. Synth.*, in press.
- (11) Rühlmann, K. *Synthesis* **1971**, 236.
- (12) Kuwajima, I.; Nakamura, E.; Shimada, J. **1982**, ICOS IV, Tokyo Symposium.
- (13) Rousseau, G.; Sloughi, N.; *Tetrahedron Lett.* **1983**, *24*, 1251.
- (14) Wasserman, H. H.; Clagett, D. C. *Tetrahedron Lett.* **1964**, 341.
- (15) Wasserman, H. H.; Clagett, D. C. *J. Am. Chem. Soc.* **1966**, *88*, 5368.
- (16) Van Tilborg, W. J. M.; Steinberg, H.; de Boer, T. J. *Recl. Trav. Chim. Pays-Bas* **1974**, *93*, 287.
- (17) Van Tilborg, W. J. M.; Steinberg, H.; de Boer, T. J. *Synth. Commun.* **1973**, *3*, 189.
- (18) Van Tilborg, W. J. M.; Steinberg, H.; de Boer, T. J. *Recl. Trav. Chim.-Pays Bas* **1974**, *93*, 290.
- (19) Richey, H. G., Jr.; Richey, J. M.; Clagett, D. C. *J. Am. Chem. Soc.* **1964**, *86*, 3906.
- (20) Turro, N. J.; Leermakers, P. A.; Wilson, H. R.; Neckers, D. C.; Byers, G. W.; Vesley, G. F. *J. Am. Chem. Soc.* **1965**, *87*, 2613.
- (21) Fry, A. F.; Scoggins, R. *Tetrahedron Lett.* **1972**, 4079.
- (22) van Tilborg, W. J. M.; Plomp, R.; de Ruiter, R.; Smit, C. J.; *Recl. Trav. Chim. Pays-Bas* **1980**, *99*, 206.
- (23) Depuy, C. H.; Arney, W. C.; Gilbson, D. H. *J. Am. Chem. Soc.* **1968**, *90*, 1830.
- (24) Schaafsma, S. E.; Steinberg, H.; de Boer, T. J. *Recl. Trav. Chim. Pays-Bas* **1966**, *85*, 73.
- (25) Schaafsma, S. E.; Molenaar, E. J. F.; Steinberg, H.; de Boer, T. J. *Recl. Trav. Chim. Pays-Bas* **1968**, *87*, 1301.
- (26) Bakker, B. H. Dissertation, Amsterdam, 1973, Chapter 4.
- (27) Bakker, B. H.; Steinberg, H.; de Boer, T. J. *Recl. Trav. Chim. Pays-Bas* **1976**, *95*, 274.
- (28) Schaafsma, S. E.; Jorritsma, R.; Steinberg, H.; de Boer, T. J. *Tetrahedron Lett.* **1973**, 827.
- (29) Bakker, B. H.; Schilder, G. J. A.; Reints Bok, T.; Steinberg, H.; de Boer, T. J. *Tetrahedron* **1973**, *29*, 93.
- (30) Gibson, D. H.; Depuy, C. H. *Tetrahedron Lett.* **1969**, 2203.
- (31) Bakker, B. H.; Reints Bok, T.; Steinberg, H.; de Boer, T. J. *Recl. Trav. Chim. Pays-Bas* **1977**, *96*, 31.
- (32) Depuy, C. H.; Breitbeil, F. W.; Debrun, K. R. *J. Am. Chem. Soc.* **1966**, *88*, 3347.
- (33) Wharton, P. S.; Bair, T. I. *J. Org. Chem.* **1966**, *31*, 2480.
- (34) Turro, N. J.; Hammond, W. B. *J. Am. Chem. Soc.* **1967**, *89*, 1028.
- (35) Bakker, B. H.; van Ramesdonk, H. J.; Steinberg, H.; de Boer, T. J. *Recl. Trav. Chim. Pays-Bas* **1975**, *94*, 64.
- (36) de Boer, T. J. *Angew. Chem., Int. Ed. Engl.* **1972**, *11*, 321.
- (37) Dull, M. F.; Abend, P. G. *J. Am. Chem. Soc.* **1959**, *81*, 2588.
- (38) Woodward, R. B.; Hoffmann, R. *J. Am. Chem. Soc.* **1965**, *87*, 395.
- (39) Hoffmann, R.; Woodward, R. B. *Angew. Chem., Int. Ed. Engl.* **1969**, *8*, 556.
- (40) Cram, D. J. "Fundamentals of Carbanion Chemistry"; Academic Press: New York, 1965; p 243.
- (41) Rappe, C.; Knutsson, L.; Turro, W. J.; Gagosian, R. B. *J. Am. Chem. Soc.* **1970**, *92*, 2032.
- (42) Kende, A. *Org. React.* (N.Y) **1960**, *11*, 261.
- (43) Depuy, C. H. *Acc. Chem. Res.* **1968**, *1*, 33. An intramolecular S_N2 type substitution on cyclopropane with inversion of configuration has been reported: Seebach, D.; Damman, R.; Lindner, H. J.; Kitschke, B. *Helv. Chim. Acta* **1979**, *62*, 1143.
- (44) Berson, J. A.; Olin, S. S. *J. Am. Chem. Soc.* **1970**, *92*, 1087.
- (45) (a) Howell, B. A.; Jewett, J. G. *J. Am. Chem. Soc.* **1971**, *93*, 798. (b) Landgrebe, J. A.; Becker, L. W. *Ibid.* **1968**, *90*, 395.
- (46) Martin, R. A.; Landgrebe, J. A. *J. Org. Chem.* **1972**, *37*, 1996.
- (47) Salaün, J. *J. Org. Chem.* **1977**, *42*, 28.
- (48) van der Vecht, J. R.; Steinberg, H.; de Boer, T. J. *Recl. Trav. Chim. Pays-Bas* **1977**, *96*, 313.
- (49) van der Vecht, J. R.; Dirks, R. J.; Steinberg, H.; de Boer, T. J. *Recl. Trav. Chim. Pays-Bas* **1977**, *96*, 309.
- (50) Braun, M.; Seebach, D. *Chem. Ber.* **1976**, *109*, 669.
- (51) Schollkopf, U.; Ruban, E.; Toune, P.; Riedel, K. *Tetrahedron Lett.* **1970**, 5077.
- (52) Wasserman, H. H.; Baird, M. S. *Tetrahedron Lett.* **1971**, 3721.
- (53) Jongejan, E.; van Tilborg, W. J. M.; Dusseau, C. H. V.; Steinberg, H.; de Boer, T. J. *Tetrahedron Lett.* **1972**, 2359.
- (54) Radom, L.; Hariharan, P. C.; Pople, J. A.; Schleyer, P. v. R. *J. Am. Chem. Soc.* **1973**, *95*, 6531. Radom, L.; Pople, J. A.; Schleyer, P. v. R. *Ibid.* **1973**, *95*, 8193.
- (55) Salaün, J. *J. Org. Chem.* **1978**, *43*, 2809.
- (56) Wasserman, H. H.; Hearn, M. J.; Cochoy, R. E. *J. Org. Chem.* **1980**, *45*, 2874.
- (57) Salaün, J. *J. Org. Chem.* **1977**, *42*, 28.
- (58) (a) Salaün, J.; Fadel, A. *Tetrahedron Lett.* **1979**, 4375. (b) Salaün, J.; Bennani, F.; Compain, J. C.; Fadel, A.; Ollivier, J. *J. Org. Chem.* **1980**, *45*, 4129.
- (59) Brown, H. C.; Rao, C. G. *J. Org. Chem.* **1978**, *43*, 3602.
- (60) (a) van Tilborg, W. J. M.; Schaafsma, S. E.; Steinberg, H.; de Boer, T. J. *Recl. Trav. Chim. Pays-Bas* **1967**, *86*, 419. (b) Stork, G.; Depey, J. C.; D'Angelo, J. *Tetrahedron Lett.* **1975**, 389. (c) Cleonine is an amino(1-hydroxycyclopropyl)-acetic acid contained in cleomycin: Kato, K. *Tetrahedron Lett.* **1980**, *21*, 4925.
- (61) Sisido, K.; Utimoto, K. *Tetrahedron Lett.* **1966**, 3267. Utimoto, K.; Tamura, M.; Sisido, K. *Tetrahedron* **1973**, *29*, 1169.
- (62) Schweizer, E. E.; Berninger, C. J.; Thompson, J. G. *J. Org. Chem.* **1968**, *33*, 336.
- (63) Osborne, N. F. U.K. Patent 2049660, 1981; *Chem. Abstr.* **1981**, *95*, 62190g; *J. Chem. Soc., Perkin Trans 1* **1982**, 1435.
- (64) Clarke, H. T.; Johnson, J. R.; Robinson, R. "Chemistry of Penicillin"; Princeton University Press: Princeton, NJ, 1949.
- (65) Flynn, E. H. "Cephalosporins and Penicillins"; Academic Press: New York, 1972.
- (66) Johnston, D. B. R.; Schmitt, S. M.; Bouffard, F. A.; Chris-

- tensen, B. G. *J. Am. Chem. Soc.* **1978**, *100*, 313 and references cited therein.
- (65) Bose, A. K.; Manhas, M. S.; Kapur, J. C.; Sharma, S. D.; Amin, S. G. *J. Med. Chem.* **1974**, *17*, 541. Mukerjee, A. K.; Singh, A. K. *Tetrahedron* **1978**, *34*, 1731. Isaacs, N. S.; Summan, G.; Reading, C. *J. Antibiot.* **1982**, *35*, 589.
- (66) Hashimoto, M.; Komoni, T.; Kamiya, T. *J. Am. Chem. Soc.* **1976**, *98*, 3023.
- (67) Howarth, T. J.; Brown, A. G.; King, T. J. *J. Chem. Soc., Chem. Commun.* **1976**, 266.
- (68) Wasserman, H. H.; Lipshutz, B. H.; Tremper, A. W.; Wu, J. S. *J. Org. Chem.* **1981**, *46*, 2991 and references cited therein.
- (69) Mukerjee, A. K.; Srivastava, R. C. *Synthesis* **1973**, 387. Isaacs, N. S. *Chem. Rev.* **1976**, *76*, 181.
- (70) Wasserman, H. H.; Adickes, H. W.; Espejo de Ochoa, O. *J. Am. Chem. Soc.* **1971**, *93*, 5586. Wasserman, H. H. *Angew. Chem., Int. Ed. Engl.* **1972**, *11*, 332.
- (71) Wasserman, H. H.; Glazer, E. A.; Hearn, M. J. *Tetrahedron Lett.* **1973**, 4855.
- (72) Wasserman, H. H.; Glazer, E. A. *J. Org. Chem.* **1975**, *40*, 1505.
- (73) Wasserman, H. H.; Dion, R. P. *Tetrahedron Lett.* **1982**, *23*, 785.
- (74) Wasserman, H. H.; Dion, R. P. *Tetrahedron Lett.* **1982**, *23*, 1413.
- (75) van Tilborg, W. J. M.; Dooyeward, G.; Steinberg, H.; de Boer, T. J. *Tetrahedron Lett.* **1972**, 1677.
- (76) Jung, M. E. *Tetrahedron* **1976**, *32*, 3 and references cited therein.
- (77) Trost, B. M. *Top. Curr. Chem.* **1973**, *41*, 1. Trost, B. M. *Acc. Chem. Res.* **1974**, *7*, 85. Trost, B. M. *Pure Appl. Chem.* **1975**, *43*, 565.
- (78) Brady, W. T. *Tetrahedron* **1981**, *37* (Suppl. 17), 2949 and references cited therein.
- (79) Crandall, J. K.; Paulson, D. R. *J. Org. Chem.* **1968**, *33*, 991; *Tetrahedron* **1969**, 2751.
- (80) Wiserman, J. R.; Chan, H. F. *J. Am. Chem. Soc.* **1970**, *92*, 4749.
- (81) (a) Salaün, J.; Conia, J. M. *J. Chem. Soc., Chem. Commun.* **1971**, 1957. (b) Salaün, J.; Garnier, B.; Conia, J. M. *Tetrahedron*, **1974**, *30*, 1413. (c) Salaün, J.; Champion, J.; Conia, J. M. *Org. Synth.* **1977**, *57*, 36.
- (82) Corey, E. J.; Seebach, D. *Angew. Chem., Int. Ed. Engl.* **1965**, *4*, 1075, 1077. Seebach, D.; Jones, N. R.; Corey, E. J. *J. Org. Chem.* **1968**, *33*, 300. Seebach, D.; Beck, A. K. *Org. Synth.* **1971**, *51*, 76. Ogura, K.; Yamashita, M.; Suzuki, M.; Tsuchihashi, G. *Tetrahedron Lett.* **1974**, 3653.
- (83) Salaün, J.; Conia, J. M. *Tetrahedron Lett.* **1972**, 2849.
- (84) Crandall, J. K.; Conover, W. W. *J. Org. Chem.* **1978**, *43*, 3533.
- (85) Ansell, M. F.; Palmer, M. H. *Q. Rev. Chem. Soc.* **1964**, *18*, 211. Eaton, P. E.; Carlson, G. R.; Lee, J. T. *J. Org. Chem.* **1973**, *38*, 4071.
- (86) Goldstein, S.; Vannes, P.; Houge, C.; Frisque-Hesbain, A. M.; Wiaux-Zarmar, C.; Ghosez, L.; Germain, G.; Declercq, J. R.; van Meerssche, M.; Arrieta, J. M. *J. Am. Chem. Soc.* **1981**, *103*, 4616.
- (87) Greene, A. E.; Depres, J. P. *J. Am. Chem. Soc.* **1979**, *101*, 4003.
- (88) Mikołajczyk, M.; Grzejszczak, S.; Lyzwa, P. *Tetrahedron Lett.* **1982**, *23*, 2237 and references cited therein.
- (89) Posner, G. M.; Sterling, J. L.; Whitten, C. E.; Lentz, C. M.; Burnelle, D. J. *J. Am. Chem. Soc.* **1975**, *97*, 107; Mitra, A. "The Synthesis of Prostaglandins"; Wiley Interscience: New York, 1977.
- (90) Ellison, R. A. *Synthesis* **1973**, 397. Ho, T. L. *Synth. Commun.* **1974**, *4*, 265; Trost, B. M.; Vladuchik, W. C. *J. Org. Chem.* **1979**, *44*, 148.
- (91) Noyori, R.; Yokoyama, K.; Makino, S.; Hayakawa, Y. *J. Am. Chem. Soc.* **1972**, *94*, 1772; **1973**, *95*, 2722. Hayakawa, Y.; Yokoama, K.; Noyori, R. *Ibid.* **1979**, *101*, 6429 and references cited therein.
- (92) Corey, E. J.; Walinsky, S. W. *J. Am. Chem. Soc.* **1972**, *94*, 8932.
- (93) Buchi, G.; Egger, B. *J. Org. Chem.* **1971**, *36*, 2021.
- (94) (a) Trost, B. M.; Bogdanowicz, M. J. *J. Am. Chem. Soc.* **1973**, *95*, 289, 5311. (b) Trost, B. M.; Nishimura, Y.; Yamamoto, K.; Elvain, S. S. *Ibid.* **1979**, *101*, 1328.
- (95) Girard, C.; Amice, P.; Barnier, J. P.; Conia, J. M. *Tetrahedron Lett.* **1974**, 3329. Conia, J. M. *Pure Appl. Chem.* **1975**, *43*, 317. Girard, C.; Conia, J. M. *J. Chem. Res., Synop.* **1978**, 182.
- (96) Monti, S. A.; Cowherd, F. G.; Mc Aninch, T. W. *J. Org. Chem.* **1975**, *40*, 858 and references cited therein.
- (97) Salaün, J.; Ollivier, J. *Nouv. J. Chim.* **1981**, *5*, 587.
- (98) von Doering, W. E.; Roth, W. R. *Angew. Chem., Int. Ed. Engl.* **1963**, *2*, 115. Willcott, M. R.; Cargle, V. H. *J. Am. Chem. Soc.* **1967**, *89*, 723; **1969**, *91*, 4310. Skattebol, L. *Tetrahedron* **1967**, *23*, 1107.
- (99) Trost, B. M.; Kurozumi, S. *Tetrahedron Lett.* **1974**, 1929.
- (100) Salaün, J.; Almirantis, Y. *Tetrahedron* **1983**, *39*, 2421.
- (101) Sato, F.; Ishikawa, H.; Sato, M. *Tetrahedron Lett.* **1980**, *21*, 365; **1981**, *22*, 85. Sato, F.; Ishikawa, H.; Wanatabe, H.; Miyake, T.; Sato, M. *J. Chem. Soc., Chem. Commun.* **1981**, 718.
- (102) Salaün, J.; Ollivier, J., unpublished results. Ollivier, J. *Thesis, Orsay*, 1982.
- (103) Visser, R. G.; Bos, H. J. T.; Brandsma, L. *Recl. Trav. Chim. Pays-Bas* **1980**, *99*, 70.
- (104) Flowers, M. C.; Frey, H. M. *J. Chem. Soc.* **1961**, 3547. Wellington, C. A. *J. Phys. Chem.* **1962**, *66*, 1671.
- (105) Willcott, M. R.; Cargill, R. L.; Sears, A. B. *Prog. Phys. Org. Chem.* **1972**, *9*, 25.
- (106) Trost, B. M.; Scudder, P. H. *J. Org. Chem.* **1981**, *46*, 506.
- (107) Sommer, L. H.; Dorfman, F.; Goldberg, G. M.; Whitmore, F. G. *J. Am. Chem. Soc.* **1946**, *68*, 488.
- (108) Paquette, L. A.; Wells, G. J.; Horn, K. A.; Tu Hsin Yan. *Tetrahedron Lett.* **1982**, *23*, 263.
- (109) Grimme, W.; Doering, W. E. *Chem. Ber.* **1973**, *106*, 1765. Heger, K.; Grimme, W. *Angew. Chem.* **1976**, *88*, 62.
- (110) Schiess, P.; Funschilling, P. *Helv. Chim. Acta* **1976**, *59*, 1756. Okamura, W. H.; Snider, W. H.; Katz, T. J. *Tetrahedron Lett.* **1968**, 3367. Paquette, L. A.; Halusta, R. J. *J. Org. Chem.* **1970**, *35*, 132. Eberbach, W.; Konig, G.; Trostmann, V. *Tetrahedron Lett.* **1979**, 4649.
- (111) Evans, D. A.; Golob, A. M. *J. Am. Chem. Soc.* **1975**, *97*, 4765. Evans, D. A.; Baillargeon, D. J.; Nelson, J. V. *Ibid.* **1978**, *100*, 2242. Evans, D. A.; Nelson, J. V. *Ibid.* **1980**, *102*, 774.
- (112) Paquette, L. A.; Crouse, G. D.; Sharrow, A. K. *J. Am. Chem. Soc.* **1980**, *102*, 3972.
- (113) Danheiser, R. L.; Martinez-Davila, C.; Morin, J. M., Jr. *J. Org. Chem.* **1980**, *45*, 1340.
- (114) Salaün, J.; Almirantis, Y., unpublished results.
- (115) Murakami, M.; Nishida, S. *Chem. Lett.* **1979**, 927. Aris, V.; Brown, J. M.; Coneely, J. A.; Golding, B. T.; Williamson, R. H. *J. Chem. Soc., Perkin Trans 2* **1975**, *4*; **1979**, 962. Solomon, R. G.; Kachinski, J. L. C. *J. Am. Chem. Soc.* **1977**, *99*, 1043.
- (116) Hudlicky, T.; Koszyk, F. J.; Kutchan, T. M.; Sheth, J. P. *J. Org. Chem.* **1980**, *45*, 5020. Hudlicky, T.; Koszyk, F. J.; Dochwat, D. M.; Cantrell, G. I. *J. Org. Chem.* **1981**, *46*, 2911.
- (117) Stork, G.; Hudrlik, P. F. *J. Am. Chem. Soc.* **1968**, *90*, 4462, 4464. House, H. O.; Czuba, L. J.; Gall, M.; Olmstead, H. D. *J. Org. Chem.* **1969**, *34*, 2324. House, H. O.; Gall, M.; Olmstead, H. D. *Ibid.* **1971**, *36*, 2364. House, H. O. "Modern Synthetic Reactions", 2nd ed.; Benjamin: Menlo Park, CA, 1972; p 568.
- (118) Patterson, J. W., Jr.; Fried, J. H. *J. Org. Chem.* **1974**, *39*, 2506.
- (119) Liotta, D.; Barnum, C. S.; Saindame, M. *J. Org. Chem.* **1978**, *46*, 4301 and references cited therein.
- (120) For a recent review on Lewis acid induced α -alkylation of carbonyl compounds, see: Reetz, M. T. *Angew. Chem., Int. Ed., Engl.* **1982**, *21*, 96. See also: Colvin, E. "Silicon in Organic Synthesis"; Butterworths: London, 1981; p 221 and references cited therein.
- (121) Paterson, I. *Tetrahedron Lett.* **1979**, 1519. Paterson, I.; Fleming, I. *Ibid.* **1979**, 993, 995, 2179.
- (122) Noyori, R.; Yokoyama, K.; Sakata, J.; Kuwajima, I.; Nakamura, E.; Shimizu, M. *J. Am. Chem. Soc.* **1977**, *99*, 1265. Noyori, R.; Nishida, I.; Sakata, J.; Nishizawa, M. *Ibid.* **1980**, *102*, 1223. Noyori, R.; Nishida, I.; Sakata, J. *Tetrahedron Lett.* **1980**, *21*, 2085. Olofson, R. A.; Cuomo, J. *Ibid.* **1980**, 819.
- (123) House, H. O.; Fischer, W. F. *J. Org. Chem.* **1969**, *34*, 3615.
- (124) Hansen, R. T.; Carr, D. B.; Schwartz, J. *J. Am. Chem. Soc.* **1978**, *100*, 2244. Schwartz, J.; Carr, D. B.; Hansen, R. T.; Dayrit, F. M. *J. Org. Chem.* **1980**, *45*, 3053.
- (125) Roberts, J. L.; Borromeo, P. S.; Poulter, C. D. *Tetrahedron Lett.* **1977**, *19*, 1621 and references cited therein.
- (126) Schreiber, J.; Maag, H.; Nashimoto, N.; Eschenmoser, A. *Angew. Chem., Int. Ed. Engl.* **1971**, *10*, 330.
- (127) Danishefsky, S.; Kitahawa, T.; Mc Kee, R.; Schuda, P. F. *J. Am. Chem. Soc.* **1976**, *98*, 6715. Danishefsky, S.; Prisbilla, M.; Lipisko, B. *Tetrahedron Lett.* **1980**, *21*, 805. Danishefsky, S.; Guingant, A.; Prisbilla, M. *Ibid.* **1980**, *21*, 2033.
- (128) Holy, N. L.; Wang, Y. F. *J. Am. Chem. Soc.* **1977**, *99*, 944.
- (129) Reich, H. J.; Renga, J. M. *J. Chem. Soc., Chem. Commun.* **1974**, 135. Trost, B. M.; Salzmann, T. N. *J. Am. Chem. Soc.* **1973**, *95*, 5321.
- (130) Stork, G.; Isobe, M. *J. Am. Chem. Soc.* **1975**, *97*, 4745, 6260.
- (131) Holmes, A. B.; Jemmings-White, C. L. D.; Schulthess, A. H. *J. Chem. Soc., Chem. Commun.* **1979**, 840.
- (132) Miyashita, N.; Yoshikoshi, A.; Grieco, P. A. *J. Org. Chem.* **1977**, *42*, 3772.
- (133) Corey, E. J.; Fleet, G. W. J.; Kato, M. *Tetrahedron Lett.* **1973**, 3963.
- (134) Jorritsma, R.; Steinberg, H.; de Boer, T. *J. Recl. Trav. Chim. Pays-Bas* **1981**, *100*, 184.
- (135) Jorritsma, R.; Steinberg, H.; de Boer, T. *J. Recl. Trav. Chim. Pays-Bas* **1981**, *100*, 94.
- (136) Jorritsma, R.; Steinberg, H.; de Boer, T. *J. Recl. Trav. Chim. Pays-Bas* **1982**, *101*, 288.

- (137) Fort, A. W. *J. Am. Chem. Soc.* **1962**, *84*, 4979. O'Brien, J. P.; Rachlin, A. I.; Treitel, S. *J. Med. Chem.* **1969**, *12*, 1112. Edelson, S. S.; Turro, N. J. *J. Am. Chem. Soc.* **1970**, *92*, 2770. Turro, N. J.; Edelson, S. S. *J. Am. Chem. Soc.* **1968**, *90*, 4499.
- (138) Salaün, J., unpublished results.
- (139) Bakker, B. H. Thesis, Amsterdam, 1973.
- (140) Noordstrand, A. A. P.; Steinberg, H.; de Boer, T. J. *Tetrahedron Lett.* **1975**, *30*, 2611. Wiering, R. G.; Steinberg, H. *J. Org. Chem.* **1981**, *46*, 1663. Wiering, P. G.; Verhoeven, J. W.; Steinberg, H. *J. Am. Chem. Soc.* **1981**, *103*, 7675. Wiering, P. G.; Steinberg, H. *Recl. Trav. Chim. Pays-Bas* **1982**, *101*, 267.
- (141) Nakamara, E.; Kuwajuma, I. *J. Am. Chem. Soc.* **1977**, *99*, 7360.
- (142) For a recent review see: Werstiuk, N. H. *Tetrahedron* **1983**, *39*, 205.
- (143) Walsh, C. *Tetrahedron* **1982**, *38*, 892, and references cited therein.
- (144) Lindberg, P.; Bergman, R.; Wickberg, B. *J. Chem. Soc., Chem. Commun.* **1975**, 946.
- (145) Wiseman, J.; Tayrien, G.; Abeles, R. *Biochemistry* **1980**, *19*, 4222.
- (146) Paech, C.; Salach, J.; Singer, T. *J. Biol. Chem.* **1980**, *255*, 2700.
- (147) Hanzlik, R.; Kishore, V.; Tullman, R. *J. Med. Chem.* **1979**, *22*, 759.
- (148) Hassall, C.; Reyle, K. *Biochem. J.* **1955**, *60*, 334.
- (149) Ghisla, S.; Wenz, A.; Thorpe, C. "Enzyme Inhibitors"; Brodbeck, V., Ed.; Verlag: Weinheim 1980; p 43.
- (150) Salaün, J.; Bruneau, P., unpublished results.