

Chemical Reviews

Volume 85, Number 5 October 1985

Synthetic Methods and Structure–Reactivity Relationships in Electron-Rich Sulfur–Nitrogen Rings and Cages

TRISTRAM CHIVERS

Department of Chemistry, The University of Calgary, Calgary, Alberta, Canada T2N 1N4

Received November 20, 1984 (Revised Manuscript Received March 21, 1985)

Contents

I. Introduction	341	IV. Spectroscopic Studies	352
II. General Synthetic Methods	342	A. Photoelectron Spectra	352
A. The Reaction of Ammonia or Ammonium Salts with Sulfur Halides and Related Cyclocondensation Reactions	343	B. UV–Visible and MCD Spectra	353
B. Preparation of S–N Heterocycles from S_4N_4	343	C. ^{15}N NMR Spectra	354
1. Reactions with Halogens or Other Oxidizing Agents	343	D. ESR Spectra	355
2. Reactions with Nucleophiles or Reducing Agents	344	E. IR and Raman Spectra	355
3. Reactions with Metal Halides or Organometallic Reagents	344	V. Reactions	356
4. Reactions with Acetylenes	344	A. Adduct Formation	356
C. Syntheses from Cyclothiazyl Halides	344	1. With Lewis Acids	356
1. $S_3N_2Cl^+Cl^-$	344	2. With Olefins	357
2. $(NSCl)_3$	345	B. Redox Reactions	357
D. Syntheses from Si–N or Sn–N Reagents	345	1. Oxidation	357
E. Syntheses from Metallocyclothiazenes	345	2. Reduction	358
F. Syntheses from S–N Cations	346	C. Ring Contraction	358
G. Syntheses from S–N Anions	346	1. Reactions with Reducing Agents or Nucleophiles	358
H. Syntheses from S–N Oxides	346	2. Reactions with Electrophiles	358
III. Molecular and Electronic Structures	346	3. Thermolysis	359
A. Monocyclic Systems	346	4. Elimination of NSX	359
1. Four-Membered Rings	347	D. Ring Expansion	359
2. Five-Membered Rings	348	1. Dimerization	359
3. Six-Membered Rings	348	2. Insertion Reactions	359
4. Seven-Membered Rings	349	3. Oxidation	360
5. Eight-Membered Rings	350	E. Chain Formation	360
6. Ten-Membered Rings	350	1. Polymerization	360
7. Summary and Conclusions	351	2. With Nucleophiles	360
B. Bicyclic and Tricyclic S–N Rings and Cages	351	3. Thermolysis	360
1. The Nature of S–S Interactions	351	VI. Summary and Conclusions	360
2. Tetrasulfur Tetranitride, S_4N_4	351	VII. Acknowledgments	361
3. Bicyclic Compounds, $1,5-E(NSN)_2E$	352	VIII. References and Notes	361
4. Disubstituted Derivatives, $1,5-X_2S_4N_4$	352		
5. Tetrasulfur Pentanitride Anion and Cation, $S_4N_5^{\pm}$, and Pentasulfur Hexanitride, S_5N_6	352		
6. Derivatives of Tetrasulfur Pentanitride, S_4N_5X	352		

I. Introduction

During the last 15 years the study of sulfur–nitrogen (S–N) compounds has progressed from relative obscurity to a position in the vanguard of main-group chemistry. In part, this rise to prominence originated with the discovery of the unusual properties of the conducting polymer $(SN)_x$.¹ Concurrently, the variety of chemical transformations and structures found for



Tristram Chivers, a native of Bath, England, received his B.Sc., Ph.D., and D.Sc. degrees all from the University of Durham, England. He did postdoctoral work during the years 1964–1969 at the Universities of Cincinnati, Sussex, and British Columbia. He joined the faculty at the University of Calgary in 1969, was Head of the Chemistry Department (1977–1982), and now continues as Professor of Chemistry. His research interests are in the general area of main-group chemistry, particularly phosphorus and sulfur compounds, with emphasis on synthetic, structural, and electrochemical studies.

sulfur–nitrogen heterocycles have attracted both experimental and theoretical interest. In a recent book² Heal stated "...the reaction chemistry of these substances (S–N rings) deserves to rank with that of boranes for novelty and interest." Investigations of the chemistry of boron hydrides continue to play a leading role in the development of our understanding of electron-deficient compounds and their influence has been extended to an appreciation of the structures of other cluster systems and organometallic complexes.³ The purpose of the present review is to demonstrate that studies of cyclothiazenes have similarly provided a framework for comprehending the behavior of π -electron-rich inorganic systems. This point will be emphasized by comparing their structures and properties with those of π -electron-precise rings, e.g., cyclophosphazenes and aromatic hydrocarbons. [In an electron-rich S–N ring the number of π -electrons exceeds the number of atomic centers in the ring. It

follows that the number of π -electrons is equal to the number of atomic centers in an electron-precise ring. Strictly speaking, this definition cannot be applied to cages since the distinction between σ - and π -electrons is not clear-cut for nonplanar molecules.]

Of the numerous reviews devoted to various aspects of sulfur–nitrogen chemistry, those on cyclic S–N compounds by Banister⁴ and Roesky⁵ are the most comprehensive. The latter covers the literature to the end of 1977 and is organized according to (a) ring size and (b) coordination number at sulfur and nitrogen. The wide variety of sulfur–nitrogen compounds can be attributed to the range of oxidation states (0–6) and coordination numbers (2–4) available to sulfur in combination with nitrogen (or other sulfur atoms).⁵ In addition nitrogen may be two-coordinate (unsaturated) or three-coordinate (saturated). This article will focus on π -electron-rich rings, i.e., those containing two-coordinate sulfur and two-coordinate nitrogen, and related bicyclic and cage molecules.

The survey begins with a discussion of general synthetic approaches to both binary S–N rings and heterocyclothiazenes in which one or more sulfur atoms have been replaced by a heteroatom, e.g., C, Si, Sn, Pb, P, As, Se, or a transition metal (plus associated ligands). There follows a detailed account of the molecular and electronic structures of important S–N heterocycles, a knowledge of which is necessary for understanding their spectroscopic properties and chemical reactions, which are described in subsequent sections.

II. General Synthetic Methods

To a casual observer the methods used for the preparation of S–N heterocycles are often obscure and even the expert can find the unpredictable nature of such reactions perplexing. The aim of this section is to provide a coherent account of synthetic strategies in terms of the typical reaction types exemplified in Table I. This overview begins with a description of the preparation of S_4N_4 and cyclothiazyl halides from ammonia or ammonium halides and sulfur halides and related condensation reactions. This is followed by

TABLE I. Some Examples of Synthetic Routes to S–N Heterocycles

1.	cyclocondensation	$NH_3 + SCl_2 \xrightarrow{Cl_2} S_4N_4$ $NH_4Cl + S_2Cl_2 + S_8 \rightarrow S_3N_2Cl^+Cl^-$ $Ni(S_2N_2H)_2 + S_2Cl_2 \xrightarrow{Et_3N} S_4N_2$ $RC(NH_2)NH + SCl_2 \rightarrow RC(NSN)_2CR$
2.	oxidation	$S_4N_4 + S_2O_6F_2 \rightarrow [S_4N_4^{2+}][FSO_3^-]_2$ $S_3N_3^- \xrightarrow{O_2} S_3N_3O^- \xrightarrow{O_2} S_3N_3O_2^-$
3.	reduction	$S_4N_3^+Cl^- \xrightarrow{KI} S_4N_4$ $(NSCl)_3 \xrightarrow{Fe} S_4N_4$
4.	cycloaddition	$NS_2^+ + RC \equiv CR' \rightarrow RR'C_2NS_2^+$ $2/3(NSCl)_3 + Me_2NCN \rightarrow (Me_2NCN)(NSCl)_2$ $NWC1_3 + S_4N_4 \rightarrow Cl_3WS_2N_3$
5.	metathesis	$(NSCl)_3 + Me_3SiNSNSiMe \rightarrow S_4N_5Cl$ $2S_3N^- + CuCl \rightarrow Cu(S_3N)_2^-$
6.	ring closure	
7.	ring expansion	$S_4N_4 + NS^+ \rightarrow S_5N_5^+$ $S_3N_2Cl^+Cl^- + R_2PN(SiMe_3)_2 \rightarrow (R_2PN)(SN)_2$ $(Ph_2PN)_2(NSCl) + Ph_3Sb \rightarrow (Ph_2PN)_4(SN)_2$
8.	ring contraction	$Cl_2S_4N_4 + Cl_2 \xrightarrow{-NSCl} (NSCl)_3$ $S_4N_4 + N_3^- \rightarrow S_3N_3^-$ $S_4N_4 + Ph_3E \rightarrow Ph_3E=N-S_3N_3 \quad (E = P, As)$




Figure 1. Preparation of S-N heterocycles from ammonia or ammonium salts and sulfur halides: (i) reflux, (ii) CCl₄ or CH₂Cl₂, (iii) Cl₂, (iv) Fe, Hg or Ph₃Sb, (v) S₈/CS₂, (vi) S₂Cl₂, (vii) KI.

details of the wide variety of S-N heterocycles that can be obtained by treating S₄N₄ with various reagents. Syntheses based on the use of the cyclothiazyl halides, S₃N₂Cl₂ and (NSCl)₃, are described next and this section concludes with an account of preparative routes which employ Si-N or Sn-N reagents, metallocyclothiazenes, S-N cations, S-N anions, and S-N oxides. For details of the various methods available for the preparation of *specific* ring systems the reader is referred to a recent book.⁶



In this section S-N rings will be represented by molecular formulas. Their molecular structures are discussed in section III and schematic drawings of the structures are depicted in Figures 8 and 16.

A. The Reaction of Ammonia or Ammonium Salts with Sulfur Halides and Related Cyclocondensation Reactions

The cyclocondensation reaction of ammonia (or ammonium salts) with sulfur halides is the most important route to S-N heterocycles since it provides an easy preparation of several key starting materials (Figure 1).⁷ When conducted in carbon tetrachloride⁸ or methylene dichloride⁹ this reaction is the standard synthesis of S₄N₄, which, in turn, is used for the preparation of numerous sulfur-nitrogen ring systems (see section IIB). The reaction of ammonia with S₂Br₂ has been adapted for the synthesis of small quantities of ³³S-labeled S₄N₄.¹⁰ The rapid addition of S₂Cl₂ to *aqueous* ammonia at 0–5 °C provides a quick and convenient source of S₄N₂.^{11,12}

S₃N₂Cl⁺Cl⁻, another important reagent in S-N chemistry (see section IIC1), is formed as an intermediate in the synthesis of S₄N₄. It is best prepared, however, by the treatment of S₂Cl₂ with sulfur and ammonium chloride at 150–160 °C;¹³ urea can be used instead of NH₄Cl.¹⁴ Chlorination of S₃N₂Cl⁺Cl⁻ with Cl₂¹³ or, more conveniently, SO₂Cl₂ produces (NSCl)₃¹⁵ which is reduced to S₄N₄ by Fe,¹⁶ Hg,¹⁷ or Ph₃Sb.¹⁸ This reaction sequence (Figure 1) has been used to produce ¹⁵N-labeled S₄N₄.¹⁸ Alternatively, treatment of S₃N₂Cl⁺Cl⁻ with S₂Cl₂ produces S₄N₃⁺Cl⁻, another intermediate in the S₄N₄ synthesis, which is reduced to S₄N₄ by potassium iodide.^{18–20} By contrast the reduction of (NSCl)₃ with sodium azide or Me₃SiN₃ produces (SN)_x.²¹

The related cyclocondensation reactions between amidines and SCl₂, in the presence of an HCl acceptor, provides a source of eight-membered heterocyclothiazenes.²²




The cage anion S₄N₅O⁻ is obtained on hydrolysis of the mixture formed by the reaction of thionyl chloride with liquid ammonia.²³

B. Preparation of S-N Heterocycles from S₄N₄

The molecule S₄N₄ exhibits very versatile chemical behavior and provides a source of many S-N heterocycles. For example, the thermolysis of S₄N₄ at ca. 300 °C using a Ag-wool catalyst generates S₂N₂,²⁴ the precursor of the polymer (SN)_x.¹ The syntheses of other S-N rings from S₄N₄ can be classified as follows: reactions of S₄N₄ with (a) halogens or other oxidizing agents, (b) nucleophiles or reducing agents, (c) metal halides or organometallic reagents, and (d) acetylenes.

1. Reactions with Halogens or Other Oxidizing Agents

The controlled halogenation of S₄N₄ with Cl₂ at –60 °C,²⁵ AgF₂,^{25b} or F₂ diluted with an inert gas²⁶ produces 1,5-S₄N₄X₂ (X = Cl, F). The reaction of S₄N₄ with



liquid bromine or ICl yields the salts S₄N₃⁺X₃⁻ (X₃ = Br₃, ICl₂).²⁷ In CS₂ solution, the solvent participates in the reaction to give CS₃N₂Br⁺Br₃⁻.²⁸ By contrast, Br₂ or ICl *vapor* react with solid S₄N₄ to give highly conducting polymers of the type (SNBr_{0.4})_x.^{29,30}


The oxidation of S₄N₄ leads to several different cyclic sulfur-nitrogen cations depending on the reaction conditions. The use of excess Lewis acid favors oxidation over simple adduct formation. For example, S₄N₄²⁺ is obtained by treatment of S₄N₄ with excess SbCl₅, AsF₅, SbF₅, or HSO₃F or by oxidizing S₄N₄·AlCl₃ with chlorine in the presence of a stoichiometric amount of AlCl₃.³¹ S₄N₄ is also oxidized quantitatively to (S₄N₄²⁺)(SO₃F⁻)₂ by peroxydisulfuryl difluoride S₂O₆F₂.³²



The dimeric cation S₆N₄²⁺ is prepared by the oxidation of S₄N₄ with (S₈²⁺)(AsF₆⁻)₂,³³ CF₃SO₃H,³⁴ or FeCl₃.³⁵ Although the cation radical S₃N₂⁺ has frequently been detected by ESR spectroscopy (section IVD) in solutions of S₄N₄ with various oxidizing agents, e.g., AsF₅,³³ (CF₃SO₂)₂O,³⁴ or AlCl₃,³⁷ it has only been obtained in the solid state as the AsF₆⁻ salt prepared from S₄N₄ and (Te₆⁴⁺)(AsF₆⁻)₄.³³

Ring contraction also occurs when S₄N₄ is treated with electrophiles such as acyl halides or acid anhydrides.^{38–40} The products are derivatives of the five-membered S₃N₂ ring. In the case of (CF₃)₃COCl, however, oxidative addition of (CF₃)₃CO groups to give N₄S₄[OC(CF₃)₃]₄ takes place.^{41a} The eight-membered ring is also retained when S₄N₄ reacts with the radical (CF₃)₂NO· to give di- and tetrasubstituted derivatives.^{41b,c}

The treatment of S₄N₄ with AsF₅ in the presence of the stoichiometric amount of elemental sulfur and a *trace* of bromine is a convenient source of the linear NS₂⁺ cation,⁴² a useful reagent in cycloaddition reactions (section IIF).






Figure 2. Preparation of S-N heterocycles from S_4N_4 and nucleophilic reagents: (i) $R_4N^+N_3^-$, (ii) Ph_3E ($E = P, As$) in C_6H_6 , (iii) Ph_3P in CH_3CN , (iv) R_2NH , (v) Ph_2PH , (vi) Ph_2PCL .




Figure 3. Preparation of metallocyclothiazenes from S_4N_4 : (i) $Co_2(CO)_8/NO$, (ii) $(Ph_3P)_2PtC_5H_4$, (iii) $CpCo(CO)_2$ or $CpCoCOD$, (iv) VCl_4 , (v) $(PhCN)_2PtCl_2$, (vi) $Cp_2Ti(CO)_2$.

2. Reactions with Nucleophiles or Reducing Agents

The reactions of S_4N_4 with nucleophiles produce a variety of S-N rings and cages (Figure 2). From the structures of the products it appears that the initial nucleophilic attack causes ring opening to give a linear intermediate which cyclizes and, in some cases, reacts further to give bicyclic or cage molecules.

For example, the degradation of S_4N_4 by anionic nucleophiles, e.g., N_3^- , S^{2-} , or CN^- , produces the six-membered ring, $S_3N_3^-$, and the cage anion, $S_4N_5^-$.⁴³ The latter ion is also formed on treatment of S_4N_4 with secondary amines.⁴⁴ Derivatives of a six-membered ring, $Ph_3E=N-S_3N_3$ ($E = P$ or As), are obtained when S_4N_4 reacts with Ph_3P ⁴⁵ (or, preferably, $Ph_3PNSiMe_3$)⁴⁶ in benzene or Ph_3As ^{47,48} (or Ph_3AsNH)⁴⁸ in CH_2Cl_2 . In a polar solvent, e.g., acetonitrile, the reaction with Ph_3P also produces $1,5-(Ph_3PN)_2S_4N_4$ and $[(Ph_3PN)_3S^+][S_4N_5^-]$.⁴⁹

Cyclophosphthiazenes of the type $(R_2PN)_n(SN)_2$ can be prepared by the reaction of S_4N_4 with R_2PPR_2 ($R = Me, Ph$)^{50,51} or Ph_2PH ^{51,52} in toluene at reflux. These heterocyclothiazenes include a six-membered ring ($n = 1$) and two structural isomers of an eight-membered ring ($n = 2$). A six-membered ring, $(Ph_2PN)_2(NSCl)$, is obtained in high yield when S_4N_4 and Ph_2PCL are heated at reflux in acetonitrile.⁵³

3. Reactions with Metal Halides or Organometallic Reagents

Although S_4N_4 typically forms 1:1 adducts with Lewis acids (section VA1), the interaction with metal halides or organometallic reagents frequently gives rise to ring opening (Figure 3). For example, the reaction of S_4N_4 with $CuCl_2$ ⁵⁴ or aluminum trihalides^{55,56} results in the formation of adducts of S_2N_2 . In the latter case $NS_2^+AlCl_4^-$ is also produced.⁵⁷

It is well-known from earlier work that S_4N_4 reacts with transition-metal halides in polar solvents to give





Figure 4. Preparation of S-N heterocycles from $S_3N_2Cl^+Cl^-$: (i) heat, (ii) $RN(SiMe_3)_2$, (iii) $(R_f)_2PN(SiMe_3)_2$, $R_f = CF_3, C_2F_5$, (iv) $Me_2NC(NH)NH_2 \cdot HCl$, (v) $SO_2(NH_2)_2$, (vi) $HCOOH$.

complexes of the chelating ligands S_3N^- and $S_2N_2H^-$ and some improvements in the preparative and purification procedures have been described.⁵⁸ In the case of $PdCl_2$ this reaction also produces $Pd_2(S_3N)_2(S_3N_2)$ in which the S_3N_2 ligand bridges two Pd atoms in a symmetric, bidentate fashion.⁵⁹



$$M = Ni, Co, Pd, Pt$$

Recently a variety of other metallocyclothiazenes have been obtained by treatment of S_4N_4 with organometallic reagents or metal halide complexes although the yields are low in many cases. As indicated in Figure 3, these include examples of the five-membered rings MS_2N_2 ($M = Co$,^{60a} Pt ,^{60b} Pb ,^{60c}) and MS_3N ($M = Co$),⁶¹ the six-membered ring MS_2N_3 ($M = V$,^{62a,b} Mo ,^{62c-e} W ^{62c-e}), and the eight-membered rings MS_4N_3 ($M = Pt$)⁶³ and MS_3N_4 ($M = Ti$).⁶⁴

4. Reactions with Acetylenes

Although the reactions of S_4N_4 with acetylenes yield 1,2,5-dithiazoles as the major products,^{65a} several novel C-N-S heterocycles have been characterized as minor products.^{65b,c} These include the seven-membered rings 1,3,5,2,4-trithiadiazepine and 1,3,5,2,4,6-trithiatiazepine (section IIIA4).^{65b} The former can be viewed as the result of cycloaddition of the acetylene to the S_3N_2 moiety, while the formation of the latter requires cleavage of the acetylene $C \equiv C$ bond.

C. Syntheses from Cyclothiazyl Halides

1. $S_3N_2Cl^+Cl^-$

The easily obtained cyclothiazyl halide $S_3N_2Cl^+Cl^-$ (section IIA) is useful for preparing other S-N ring systems via condensation reactions. For example, it can be converted to S_3N_2O by formic acid,¹⁴ and the reaction with $SO_2(NH_2)_2$ produces $S_4N_4O_2$ (Figure 4).¹⁴ Heating at ca. 90 °C or treatment with SO_2 transforms yellow $S_3N_2Cl_2$ into dark green $(S_3N_2Cl)_2$, which contains the $S_6N_4^{2+}$ cation.⁶⁶ Metathetical reactions with Si-N reagents result in the replacement of chlorine atoms by imino substituents to give covalent derivatives of the S_3N_2 ring,^{14,67,68} but with $R_2PN(SiMe_3)_2$ ($R = CF_3, C_2F_5$) the six-membered ring $[R_2PN][SN]_2$ is formed.⁶⁹ The condensation reaction with dimethylguanidine hydrochloride, $Me_2NC(NH)NH_2 \cdot HCl$, also produces a six-membered ring, $(Me_2NCN)(NSCl)_2$, in low yield.⁷⁰




Figure 5. Preparation of S-N heterocycles from $(\text{NSCl})_3$: (i) RCN or $\text{RCH}=\text{NN}=\text{CHR}$, (ii) $\text{Me}_3\text{SiNSNSiMe}_3$, (iii) $\text{S}_4\text{N}_4/\text{FeCl}_3$ in SOCl_2 , (iv) Me_2NCN .


2. $(\text{NSCl})_3$

The potential of $(\text{NSCl})_3$ as a reagent for the synthesis of other sulfur-nitrogen heterocycles is limited by the ease with which the trimeric ring is cleaved to the monomer.^{17,71} This dissociation is manifested by a reversible color change from pale yellow to mint green for solutions of $(\text{NSCl})_3$ in CCl_4 ,^{25a} CH_3CN ,⁷² or THF ⁷² above ca. 55 °C and has been exploited in the *in situ* generation of NS^+ salts, which undergo an insertion reaction with S_4N_4 to give the S_5N_5^+ cation (Figure 5).^{73,74} In SO_2 solution $(\text{NSCl})_3$ forms a 1:1 adduct with SbCl_5 , formulated as $[(\text{ClSN})_2(\text{SN})^+][\text{SbCl}_6^-]$,⁷⁵ but the addition of more SbCl_5 produces $\text{S}_4\text{N}_4^{2+}$.⁷⁶ Heterocyclic rings containing carbon, RCN_2S_2^+ , are obtained when $(\text{NSCl})_3$ is heated at reflux with azines, $\text{RCH}=\text{NN}=\text{CHR}$,⁷⁷ or nitriles, RCN ($\text{R} = t\text{-Bu, Ph, CCl}_3$).⁷⁸ In the case of CF_3CN , the six-membered ring $(\text{CF}_3\text{CN})(\text{NSCl})_2$ is also isolated.⁷⁹ With Me_2NCN , however, both six-membered and eight-membered rings, $(\text{Me}_2\text{NCN})(\text{NSCl})_2$ and 1,3- $(\text{Me}_2\text{NCN})_2(\text{NSCl})_2$, can be prepared in good yields via a cycloaddition reaction of Me_2NCN with NSCl units generated from $(\text{NSCl})_3$ in CCl_4 at 60 °C.⁸⁰

Several reactions of $(\text{NSCl})_3$ with nucleophilic or reducing reagents result in ring expansion to give S-N cages. For example, the chemical reduction of $(\text{NSCl})_3$ to S_4N_4 is included in Figure 1.¹⁶⁻¹⁸ The cathodic reduction of $(\text{NSCl})_3$ in CH_2Cl_2 yields S_4N_4 at room temperature and $\text{S}_5\text{N}_5\text{Cl}$ at -20 °C with current efficiencies of 90% and 49%, respectively.⁸¹ The metathetical reaction of $(\text{NSCl})_3$ with $\text{Me}_3\text{SiNSNSiMe}_3$ provides an excellent source of the bicyclic S_4N_5^+ cation,⁸² while the related anion S_4N_5^- is present in liquid ammonia solutions of $(\text{NSCl})_3$.⁸³ Surprisingly, the treatment of $(\text{NSCl})_3$ with $(\text{Me}_3\text{Sn})_3\text{N}$ produces S_5N_6 .⁸⁴

D. Syntheses from Si-N or Sn-N Reagents

The use of $(\text{Me}_3\text{Sn})_3\text{N}$ instead of ammonia allows kinetic control of the reaction with S_2Cl_2 and results in the formation of a five-membered metathiazene ring.⁸⁵



The reagent $\text{Me}_3\text{SiN}=\text{S}=\text{NSiMe}_3$ is extremely useful for the preparation of cyclic S-N compounds (Figure 6). A number of these reactions, e.g., with RAsCl_2 ($\text{R} = \text{Me}$,⁸⁶ Ph ,^{87a} $t\text{-Bu}$ ^{87b}), MeSiCl_3 ,^{88,89} $\text{Cl}_2\text{S}_4\text{N}_4$,⁸⁴ $(\text{NSCl})_3$,⁸² and $(\text{Ph}_2\text{PN})(\text{NSCl})_2$ ⁹⁰ involve metathesis and proceed in a predictable manner to give S-N rings or






Figure 6. Preparation of S-N heterocycles from $\text{Me}_3\text{SiNSNSiMe}_3$: (i) RAsCl_2 , (ii) 1,5- $\text{Cl}_2\text{S}_4\text{N}_4$, (iii) MeOH , (iv) RCOCl , (v) PCl_5 , (vi) PF_5 .


cages by the introduction of $\text{N}=\text{S}=\text{N}$ bridges. By contrast, treatment of $\text{Me}_3\text{SiN}=\text{S}=\text{NSiMe}_3$ with PCl_5 ⁹¹ or PF_5 ⁹² unexpectedly gives a six-membered ring and a bicyclic compound, respectively, (Figure 6). Furthermore, the reaction with MeAsCl_2 produces a five-membered ring MeAsS_2N_2 when the arsenic reagent is in excess.⁸⁶

Some other unpredictable reactions of $\text{Me}_3\text{SiN}=\text{S}=\text{NSiMe}_3$ which are useful for the synthesis of S-N rings or cages include (a) the production of S_4N_5^- upon methanolysis,^{83,93} (b) the reactions with acid chlorides³⁹ or anhydrides⁹⁴ to give derivatives of the S_3N_2 ring, and (c) the formation of $\text{S}_4\text{N}_4\text{O}_2$ and $\text{S}_5\text{N}_5^+\text{S}_3\text{N}_3\text{O}_4^-$ on treatment with FSO_2NSO .⁹⁵

An interesting development of the application of silylated sulfur diimides in the synthesis of S-N heterocycles involves the reaction of ArNSNSiMe_3 with SCL_2 .⁹⁶ The metathetical elimination of Me_3SiCl is followed by an intramolecular condensation involving electrophilic attack of the sulfenyl chloride at the *o*-position of the aromatic ring to give inorganic analogues of naphthalene^{96a} and phenazulene rings.^{96b}




Ring closure to give benzo-1,3,2-dithiazolium chloride also occurs when benzene-1,2-disulfenyl chloride is treated with trimethylsilyl azide.⁹⁷






E. Syntheses from Metallocthiazenes

There are several reports of the use of metallocthiazenes in the preparation of S-N heterocycles. For example, the five-membered rings $\text{S}_3\text{N}_2\text{O}$ and $\text{S}_2\text{N}_2\text{CO}$ are conveniently prepared by the metathetical reactions of $\text{Me}_2\text{SnS}_2\text{N}_2$ with SOF_2 ⁹⁸ or COF_2 ,⁹⁹ respectively.



In a similar approach the readily prepared complex $\text{Ni}(\text{S}_2\text{N}_2\text{H})_2$ (section IIB3) takes part in combined metathesis-condensation reactions to give five- and six-membered S-N rings.¹⁰⁰



Further development of the synthetic applications of metallocyclothiazenes can be anticipated in view of the recent discovery of new ring systems (Figure 3).

F. Syntheses from S-N Cations

The cations SN^+ and S_2N^+ represent potential building blocks in the construction of S-N heterocycles. For example, the insertion reaction of NS^+ with S_4N_4 is described in section IIC2 and this type of reaction could probably be extended to other S-N heterocycles. The easily obtained NS_2^+ cation⁴² undergoes cycloaddition reactions with acetonitrile or alkynes in SO_2 to give almost quantitative yields of the 1,3,2,4-dithiazolium and 1,3,2-dithiazolium cations, respectively (section IIIA2).¹⁰¹ A more complex reaction occurs between NS_2^+ and S_4N_4 to give the nitrogen-bridged $(\text{S}_3\text{N}_2)_2\text{N}^+$ cation.¹⁰²


The preparation of S_4N_4 by the reduction of $\text{S}_4\text{N}_3^+\text{Cl}^-$ with KI is mentioned in section IIA.¹⁸⁻²⁰ Highly explosive, covalent derivatives of the S_4N_5^+ cation are obtained by treating $\text{S}_4\text{N}_5\text{Cl}$ with silylated sulfidimides.¹⁰³

G. Syntheses from S-N Anions

The acyclic anions S_3N^- ¹⁰⁴ and S_4N^- ¹⁰⁵ are potential sources of metallocyclothiazenes via metathetical reactions with metal halides. This procedure works well for S_3N^- with Cu(I) or Ag(I) halides,¹⁰⁶ but with other transition-metal halides, e.g., Ni(II) or Co(II), $\text{MS}_2\text{N}_2\text{H}$ complexes are also formed and, in other cases, redox reactions occur.^{104b}

The reactions of the cyclic S_3N_3^- ¹⁰⁷ or cage S_4N_5^- ¹⁰⁸ anions with electrophiles provide a rich source of other S-N heterocycles, but, in most instances, alternative synthetic routes are preferable. For example, the reactions of S_4N_5^- with Br_2 or I_2 provided the first synthesis of S_5N_6 ,¹⁰⁸ which is best prepared, however, from $\text{Cl}_2\text{S}_4\text{N}_4$ and $\text{Me}_3\text{SiNSNSiMe}_3$ (Figure 6).⁸³

Oxidation of S_3N_3^- with electrophiles frequently produces S_4N_4 (and $(\text{SN})_x$),¹⁰⁷ but controlled addition of molecular oxygen gives oxyanions in which the six-membered ring is preserved.¹⁰⁹



H. Syntheses from S-N Oxides

The reactions of $\text{S}_4\text{N}_4\text{O}_2$ with nucleophiles have been investigated in detail (Figure 7). The methoxide ion attacks the sulfur atom opposite to the SO_2 group to give the $\text{MeOS}_4\text{N}_4\text{O}_2^-$ anion, in which the eight-membered ring remains intact.^{89,110} Other nucleophiles promote opening and, subsequently, contraction of the ring. For example, the six-membered rings $\text{S}_3\text{N}_3\text{O}_2^-$ and $\text{O}_2\text{S}_3\text{N}_3(\text{MMe}_3)$ are formed on treatment of $\text{S}_4\text{N}_4\text{O}_2$ with azide ion^{109b,111a} or $(\text{Me}_3\text{M})_3\text{N}$ ($\text{M} = \text{Si}, \text{Sn}$),^{111b} respectively. With Ph_3As a derivative of the five-membered S_3N_2 ring is obtained.¹¹²





Figure 7. Reactions of $\text{S}_4\text{N}_4\text{O}_2$ with nucleophiles (i) NaOMe , (ii) $(\text{Me}_3\text{Sn})_3\text{N}$ or Me_3SnN_3 , (iii) $(\text{Me}_3\text{M})_3\text{N}$ ($\text{M} = \text{Si}, \text{Sn}$), (iv) N_3^- , (v) Ph_3As , (vi) Ph_3As , (vii) in acetone for a few days.

The reactions of $\text{S}_4\text{N}_4\text{O}_2$ with electrophiles have been less well studied but the reagent SO_3 produces a variety of sulfur-nitrogen oxides including $(\text{S}_3\text{N}_3\text{O}_4)_2\text{S}$,^{113a} $[\text{S}_6\text{N}_4^{2+}][\text{S}_3\text{N}_3\text{O}_4^-]_2$,^{113b} and the six-membered ring $\text{S}_3\text{N}_2\text{O}_5$.^{113b}

Titanium tetrachloride promotes the elimination of SO_2 from $\text{S}(\text{NSO})_2$ to give an adduct of S_2N_2 .¹¹⁴



The cycloaddition reaction between the electrochemically generated NSO^- ion and S_2N_2 (from S_4N_4) has been reported recently.¹¹⁵



III. Molecular and Electronic Structures

A. Monocyclic Systems

In 1972 Banister proposed that planar S-N heterocycles belong to a class of "electron-rich aromatics", which conform to the well-known Hückel ($4n + 2$)- π -electron rule.¹¹⁶ On the reasonable assumption that each sulfur contributes two and each nitrogen one electron to the π system, S_2N_2 (6π), S_4N_3^+ (10π), and S_5N_5^+ (14π) were cited as examples in support of this contention. However, this simple counting procedure belies the fact that the surplus of electrons is accommodated in π^* (or $n\pi$) orbitals in these electron-rich systems and does not predict, for example, that the π bond orders in S-N heterocycles are significantly lower than those in the corresponding aromatic hydrocarbons.¹¹⁷⁻¹²⁰

The electronic structures of planar S-N rings have been discussed at the simple Hückel level (HMO) by Gimarc and Trinajstić,¹¹⁷ at the extended Hückel level (EHMO) by Gleiter¹¹⁸ and, more recently, at the ab initio level by Laidlaw and Trsic.¹¹⁹ The S-N rings typically accommodate more π electrons than their neutral hydrocarbon analogues. Perhaps the most extreme example is S_3N_3^- ($10\pi e^-$); cf. C_6H_6 ($6\pi e^-$). In considering the " π -electron richness" of the S_3N_3^- ion it should be noted that all electron ab initio MO calculations show that the lowest empty orbitals in benzene are the degenerate π^* pair so that $\text{C}_6\text{H}_6^{4-}$ would also be a 10π -electron system. The stability of S_3N_3^- relative to $\text{C}_6\text{H}_6^{4-}$ can be attributed, in part, to its lower ionic charge and the greater ability of nitrogen to accommodate excess negative charge. However, because of its weaker net π bonding, planar S_3N_3^- should be less stable than neutral C_6H_6 . Moreover, the S-N ring has




Figure 8. Schematic representation of the structures of monocyclic binary S-N rings and heterocyclothiazenes. Details of X-ray structures can be found in the following references: 1,²⁴ 2 (R = CCl₃,^{121a} Cl^{121b}), 3,¹⁰¹ 4,¹⁰¹ 5,¹²² 6,^{40,123} 7,¹²⁴ 8,³³ 9,¹²⁵ 10,¹²⁶ 11 (R = Ph,⁵⁰ Me₃SiNH¹²⁷), 12,^{109b} 13,¹²⁸ 14,^{12,129} 15 (X = O⁻,¹⁰⁹ NPPH₃,⁴⁵ NAsPh₃⁴⁷), 16,¹³⁰ 17,^{65b} 18,^{65b} 19,^{96b} 20,³¹ 21,¹³¹ 22,²² 23 (R = Ph,⁵¹ CF₃⁶⁹), 24,^{96a} 25.¹³²

no exocyclic hydrogen substituents which would tend to stabilize lone-pair distributions. It would appear that the larger separation of nuclei, $d(\text{S-N}) \sim 1.6$ vs. $d(\text{C-C}) \sim 1.4$ Å, reduces the lone-pair repulsions so that the advantages of the π network of planar S_3N_3^- are retained. The electronic structures of other S-N rings will involve additional factors and the balance between a π -electron-rich planar structure and other ring conformations is often delicate. Nevertheless the principles controlling the stability of S_3N_3^- appear to be fundamental to all S-N rings.

Although *ab initio* MO calculations are to be preferred because of their less arbitrary input, HMO and EHMO calculations can give a useful indication of the π -electronic structures of π -electron-rich S-N rings. It is important, however, to recognize their limitations. For example, the choice of Coulomb parameters α_{N} and α_{S} is particularly difficult. The assumption that nitrogen is more electronegative than sulfur is appropriate for the frontier π orbitals but is not likely correct for the deeper π orbitals.¹⁴⁸ This is reflected, for example, in the polarization toward sulfur calculated for the π orbitals of S_3N_3^- .^{128b} Even though the net charge on nitrogen is negative, and is largely due to the π elec-

trons, it should be emphasized that the largest π population is on the sulfur centers. This is simply a reflection of the fact that each sulfur formally contributes two π electrons to the π system whereas nitrogen donates only one. Analysis of charge redistribution in S-N anions or cations is somewhat arbitrary. However, if one presumes that the ionic charge is initially distributed uniformly over all centers then, in accord with the usual concept of electronegativity, nitrogen normally accumulates charge at the expense of sulfur in these systems.

In the ensuing discussion the electronic structures of the S-N heterocycles illustrated in Figure 8 will be described according to ring size. The level of the theoretical approach for each ring system is indicated and, unless specified otherwise, the rings can be assumed to be planar.

1. Four-Membered Rings

The S_2N_2 molecule, 1, is π -isoelectronic with the pseudoaromatic cyclobuta-1,3-diene dianion, $\text{C}_4\text{H}_4^{2-}$, in the valence shell.¹³³ The upper occupied levels are not degenerate in 1, however, due to the lower symmetry (D_{2h}) of S_2N_2 compared to $\text{C}_4\text{H}_4^{2-}$ (D_{4h}). The electronic




Figure 9. Qualitative π MO scheme for the S_3N_2 ring.^{118,137}

structure has been discussed at various theoretical levels.¹³⁴ Ab initio MO calculations^{134a,136} indicate that the HOMO is a nonbonding π MO located on sulfur, which is of slightly higher energy than the nonbonding nitrogen-based MO, and this conclusion is supported by UV-photoelectron spectral data (section IVA).¹³⁴⁻¹³⁶

Ab initio valence-bond calculations, with sulfur 3d orbitals omitted, indicate that the primary Lewis valence-bond structure of S_2N_2 is a spin-paired diradical (or "long-bond") structure,^{134b} in agreement with an INDO VB study.^{134c}

2. Five-Membered Rings

Although the neutral S_3N_2 ring (8π) is unknown it provides a convenient reference point for the discussion of the electronic structure of five-membered rings. Unlike its hydrocarbon analogue $C_5H_5^-$ (D_{5h}), there are no degenerate levels in the π system of S_3N_2 (C_{2v} , Figure 9). However, MNDO calculations indicate only a small energy difference between $2a_2$ and $3b_1$, so that a triplet ground state seems probable.^{118,137} The formation of a stable cyclic dianion $S_3N_2^{2-}$ (10π) is unlikely since both π^* levels would be fully occupied. On the other hand, the dication $S_3N_2^{2+}$ (6π) should be stable but has not been structurally characterized, although salts have been prepared.¹³⁸ The structures of several π -isoelectronic heterocyclothiazenes, 2-4, have been determined by X-ray crystallography.^{101,121}

The radical cation $S_3N_2^+$, 8, is produced in solution by treatment of S_4N_4 with various oxidizing agents, but has been obtained in the solid state only as the AsF_6^- salt (section IIB1).³³ The dimer $S_6N_4^{2+}$, in which two S_3N_2 rings are connected by two long S-S bonds (ca. 3.0 Å), is present in other salts.^{34,35} The related cation





Figure 10. Schematic representation of the occupied MO's of 5 or 6.¹³⁷

$Se_4S_2N_4^{2+}$ has a similar structure.³⁶ On the basis of semiempirical (EHMO) calculations, the bonding in the dimer has been described in terms of an electron-rich ($6e^-$) four-center bond using the dimerization of $H_2S_2^+$ as a model.^{118,137} Recent ab initio SCF calculations of charge densities in $S_6N_4^{2+}$ show, however, that the long S-S bonds have the characteristics of closed shell rather than shared interactions.¹³⁹

A dimer with a similar structure to $S_6N_4^{2+}$, except that the two rings adopt a cis rather than a trans conformation, is formed by the 7- π -electron radical obtained by the one-electron reduction of 2.¹²⁵

The five-membered rings containing one three-coordinate sulfur atom $S_3N_2Cl^+$, 5, and S_3N_2X , 6 ($X = O, NR$), are related to 8. The structure of 5 can be viewed formally as an association of $S_3N_2^+$ with Cl. The linear combination of the $p\pi$ orbitals of the four remaining centers provides a set of butadiene-like orbitals of which the lowest three are occupied (6π , Figure 10). The occupation of ψ_3 is expected to strengthen S-N bond a and weaken slightly bonds b and c in agreement with structural data.¹²² Similar considerations apply to the neutral molecules 6 which are perceived as the union of $S_3N_2^+$ with O^- or RN^- .^{118,137}

3. Six-Membered Rings

The $S_3N_3^-$ ion, 13, is an interesting example of an electron-rich six-membered ring. It has been confirmed as a 10- π -electron system by a variety of ab initio MO calculations^{128b,140} and by spectroscopic measurements (section IVB). The planar $S_3N_3^-$ ring, with approximately equal S-N bond distances,¹²⁸ has a π manifold reminiscent of benzene with the important difference that the degenerate π^* levels ($2e''$) are fully occupied in 13 (Figure 11). Consequently the π bond order in $S_3N_3^-$ is lower than that in benzene^{117,128b} and the framework bonds are considerably weaker (section VE3).

The replacement of a sulfur atom in $S_3N_3^-$ by another heteroatom to give, for example, $R_2PS_2N_3$, 11, or the introduction of one (15) or two (12) substituents on one sulfur lowers the symmetry of the ring to C_{2v} or C_s . As a result there are no degenerate π levels for 11, which has eight endocyclic π electrons according to ab initio MO calculations (Figure 11).⁵⁰ The experimental geometry, in which the phosphorus atom is tilted out of the NSNSN plane by 0.28 Å, is slightly lower in energy (ca. 15 kcal mol⁻¹) than a planar PS_2N_3 ring. Calculations of the distribution of π charge density suggest that the internal salt model $R_2P^+S_2N_3^-$ is a useful approximate description of 11. The electronic structure of the $S_3N_3O_2^-$ ion, 12, can be described in a similar way, i.e., a combination of a sulfone (SO_2) group with an 8- π -electron pentadienyl-like NSNSN⁻ fragment.^{109b} Consequently the rings in 11 and 12 retain many of the




Figure 11. π MO scheme for $S_3N_3^-$ and PS_2N_3 showing the frontier orbitals of the PS_2N_3 ring.^{50,141}




Figure 12. Composition of the occupied π MO's and the LUMO of the $P_2SN_3^+$ ring.¹⁴¹

structural and electronic features of the thiazene unit and the frontier orbitals are sulfur-based (Figure 11).⁵⁰

In contrast, the substitution of S^+ for R_2P in $R_6P_3N_3$ to give the π -electron-precise $P_2SN_3^+$ ring results in more profound adjustments to the electronic structure of the ring since the energy of the π -type orbital on sulfur is *intermediate* between the energies of the two groups of π MO's of the phosphazene (NPNPN) unit. Ab initio MO calculations¹⁴¹ suggest that the π system in $P_2SN_3^+$ can be analyzed in terms of the π systems of the fragments, PNP^+ and NSN . As shown in Figure 12, the lowest π MO, $2b_1$, is bonding throughout NSN while the next, $2a_2$, is primarily a nitrogen-centered lone-pair orbital. The HOMO, $3b_1$, is localized on the PNP unit [$d(P-N) = 1.58 \text{ \AA}$]. The π overlap populations indicate stronger P–N π bonding in this unit than in the P–N bonds connecting the PNP and NSN units consistent with the longer distances found for these P–N bonds [$d(P-N) = 1.66 \text{ \AA}$].¹⁴¹

The "fragment approach" has also been used to rationalize the molecular and electronic structures of S_4N_2 , **14**,¹² which have been investigated by CNDO/2,¹⁴² EHMO,¹⁴³ MNDO,¹² and ab initio¹² (with configuration interaction)¹⁴⁴ methods. The CNDO/2 calculations predict a planar geometry,¹⁴² but all other theoretical approaches favor the half-chair conformation over the planar ring by 6–14 kcal mol⁻¹. Indeed the molecular




Figure 13. Correlation diagram for the π orbitals of SSS , NSN , and planar S_4N_2 .¹²

structure proposed by Palmer et al.,^{144a} on the basis of ab initio MO calculations, bears a remarkable resemblance to the observed geometry in which the dihedral angle is 55° ,^{12,129} cf. calculated value of 57° .^{144a} The S_4N_2 ring can be viewed as the result of the interaction of two pseudo-allyl fragments, SSS ($6\pi e^-$) and NSN ($4\pi e^-$).¹² As indicated in Figure 13 the bonding contribution of $2b_1$ to the S–S bonds is largely offset by the occupation of $3b_1$ which is antibonding in the SSS fragment. Similarly, the occupancy of both $1a_2$ and $2a_2$ effects a partial "cancellation" of bonding and antibonding contributions in the S–N connecting bonds. In contrast, for the NSN fragment only the bonding MO, $1b_1$, is occupied and one may therefore expect significant net π bonding in the NSN unit. This picture is consistent with the short S–N bonds observed for this unit ($d(S-N) = 1.56 \text{ \AA}$) relative to the S–N connecting bonds ($d(S-N) = 1.68 \text{ \AA}$).^{12,129}

4. Seven-Membered Rings

The well-known cation $S_4N_3^+$, **16**, a rare representative of a delocalized 10- π -electron seven-membered ring, is isoelectronic with the hypothetical [7]annulenide trianion $C_7H_7^{3-}$.^{145,146} Ab initio MO calculations indicate that the two highest occupied π MO's, $2a_2$ and $3b_2$, are nearly degenerate and they are antibonding and bonding, respectively, with respect to the S–S unit (Figure 14).¹⁴⁶ Consequently, the π bonding across this unit is weaker than in the thiazene segment of the ring. In fact, the S–S bond lengths (2.07–2.09 \AA) are typical for single bonds. The pseudodegeneracy of the higher π levels of $S_4N_3^+$ is reminiscent of the electronic structure of the hypothetical *planar* $S_3N_3^+$ ion ($8\pi e^-$, Figure 11),^{128b} but it is a stable configuration since both levels are fully occupied in $S_4N_3^+$. In a *half-chair* conforma-




Figure 14. The two highest occupied ($2a_2$ and $3b_2$) and the two lowest unoccupied ($3a_2$ and $4b_2$) π MO's and (a) UV-visible and (b) MCD spectra of $S_4N_3^+$.^{145,146}

tion, however, the e'' (HOMO) orbitals of $S_3N_3^+$ are split into nondegenerate levels and EHMO calculations indicate that this conformation is preferred over a planar ring by 0.43 eV.¹⁴³

Two additional examples of 10- π -electron seven-membered cyclothiazenes, **17** and **18**, have been structurally characterized recently,^{65b} but details of their electronic structures are not available. The related ring system, **19**, is unknown, but EHMO calculations of π charge densities suggest that it is a component of a CSN heterocycle based on a phenazulene ring system (eq 8).^{96b}

5. Eight-Membered Rings

The $S_4N_4^{2+}$ cation is planar with equal S-N bond lengths (D_{4h}) (ca. 1.55 Å) except in the case of $[S_4N_4^{2+}][SbCl_6^-]_2$ in which the cation is slightly puckered towards a boat conformation.^{31b} CNDO and EHMO calculations indicate that the planar conformation is stable but easily deformable.³² This conclusion is supported by MNDO¹⁴⁷ and ab initio MO calculations,¹⁴⁸ which also demonstrate that $S_4N_4^{2+}$ is a fully delocalized 10- π -electron system. The latter calculations show that the strong π network in $S_4N_4^{2+}$ is more than sufficient to overcome the loss of S-S σ -bonding in the $S_4N_4^{2+}$ cage. The π manifold for $S_4N_4^{2+}$ is similar to that of the isoelectronic $C_8H_8^{2-}$ ion except that the degeneracy of the middle levels is split into two primarily nonbonding orbitals, $1b_{1u}$ and $1b_{2u}$, on sulfur and nitrogen, respectively (Figure 15). The former is higher in energy (cf. S_2N_2). The π^* levels in $S_4N_4^{2+}$ are unoccupied.

The recent claim of the isolation of $[S_4N_4^+][FeCl_4^-]$,¹⁴⁹ in which the cation would be a 11- π -electron ring, should be viewed with circumspection since the structural parameters and other physical data reported for this salt do not eliminate the possibility that the cation is $S_4N_4H^+$ (section VA1).¹⁵⁰

In $S_4N_4O_2$, **21**, the sulfone sulfur atom lies 1.52 Å and the two adjacent nitrogen atoms are 0.41 Å above the mean plane of the five-atom S_3N_2 unit which is coplanar to within 0.01 Å.¹³¹ The metallocyclothiazene $Cp_2TiS_3N_4$ is structurally related to **21**, but less puckered and the variations in S-N bond lengths suggest a somewhat different electronic structure in these two




Figure 15. π MO's and upper occupied σ MO's of $S_4N_4^{2+}$.¹⁴⁸

rings.⁶⁴ In **21** the structural data indicate it consists of two N=S=N units linked by a sulfur atom and an SO_2 group, but this molecule has also been described as a delocalized 10- π -electron S_3N_4 system linked by an SO_2 group.

The geometry of eight-membered heterocyclothiazenes, $1,5\text{-E}(\text{NSN})_2\text{E}$, is markedly dependent on the effective electronegativity of the heteroatoms E. When E = CPh, **22**, the heterocyclic ring is planar and the C-N and S-N bond lengths are consistent with a delocalized 10- π -electron system.^{22a} This view of the electronic structure of the tetrazocine ring, $S_2N_4C_2$, has been confirmed by ab initio^{22b,152,153} and MNDO^{22b,154} MO calculations. When E = CNMe₂ or PR₂ (R = Me, Ph), however, the molecule adopts a folded (butterfly) structure with a cross-ring S-S interaction of 2.4–2.5 Å (section IIIB2). By contrast the formation of transannular S-S bonding contacts is not geometrically favorable for the 1,3-isomers, **23**. These ring systems consist of a planar S_2N_3 unit with P atoms on the opposite (R = Ph)⁵¹ or the same (R = CF₃)⁶⁹ side of the plane.

The 10- π -electron monocation, $RCS_3N_4^+$ (R = CF₃), a hybrid of **20** and **22**, has a planar structure.¹⁵⁵

6. Ten-Membered Rings

$S_5N_5^+$ is an almost planar 14- π -electron ten-membered ring. Both heart-shaped, **25a**,^{132d} and azulene-like, **25b**,^{132a,b,c} conformations of the ring have been reported for different salts. It is possible that the heart-shaped structure is a consequence of the superposition of two azulene-like rings, i.e., structural disorder.¹⁵⁶ However, MNDO calculations,^{157a} in contrast to earlier EHMO results,^{157b} have shown that **25a** and **25b** have almost equal energies. Consequently, **25a** could be the most stable isomer in certain salts.

Although it is not a monocyclic system, benzo-1,3-dithia-2,4-diazene, **24**, has an essentially planar structure and can be considered as an S-N analogue of naphthalene with two additional electrons, i.e., 12- π -electrons.^{96a}

7. Summary and Conclusions

MO calculations have shown that the energies of the π manifolds of S-N rings are lower and the size of the HOMO-LUMO gaps is smaller for S-N rings compared to their hydrocarbon or cyclophosphazene counterparts. The excess of electrons is accommodated in $n\pi$ or π^* (antibonding) orbitals, which are primarily sulfur based. Various consequences of the occupation of these π^* levels and the small HOMO-LUMO gap, e.g., intense colors, redox activity, cycloaddition reactions, and susceptibility to nucleophilic degradation, are discussed in sections IV and V.

B. Bicyclic and Tricyclic S-N Rings and Cages

1. The Nature of S-S Interactions

There is a strong tendency, especially among larger S-N rings (eight atoms or more), to reduce the number of antibonding electrons in the π manifold by the formation of long transannular S-S bonds (2.4–2.7 Å). This gives rise to bicyclic (folded), tricyclic, or cage structures (Figure 16). The nature of the S-S interactions in these unusual compounds has been a central issue in attempts to understand their structures and bonding. Theoretical approaches have included the empirical procedure of Banister,¹⁶⁷ semiempirical (primarily EHMO and MNDO) and ab initio MO calculations, and the use of "increased-valence" structures.¹⁶⁸

Banister has rationalized the cage framework of S_4N_4 on the assumption that all six edges of the almost regular S_4 tetrahedron represent S-S bonds,¹⁶⁷ and this conceptually simple view can be extended to explain the structures of $S_4N_5^+$ and $S_4N_5^-$. In contrast, an EHMO analysis of various conformations of S_4N_4 by Gleiter requires the formation of only two S-S σ -bonds in the cage structure¹¹⁸ and ab initio MO calculations confirm that only the unbridged pairs of sulfur atoms of the S_4 tetrahedron in an S-N cage take part in a bonding interaction.^{82b} The overlap populations indicate that these S-S bonds are weak^{82b} and the calculated charge densities are characteristic of closed shell rather than shared interactions.¹³⁹

Although there are as yet no simple, universal bonding rules for rationalizing the structures of S-N cages, MO calculations on individual molecules do provide some helpful insight as shown by the following examples.

2. Tetrasulfur Tetranitride, S_4N_4

Tetrasulfur tetranitride, **26**, has been the subject of many theoretical discussions at both the semiempirical and ab initio levels in recent years.^{82b,120b,134a,139} The unusual structure of S_4N_4 , in which the sulfur atoms are three coordinate and the nitrogen atoms are two coordinate, is the inverse of that found for the isoelectronic cages, α - P_4S_4 and As_4S_4 . The preference for the more electronegative atoms to occupy the lowest coordination sites in these molecules has been explained in terms of




Figure 16. Schematic representation of bicyclic, tricyclic, and cage S-N and P-S-N molecules and ions. Details of the X-ray structural investigations can be found in the following references: **26**,¹⁵⁸ **27** (E = CNMe₂,^{22a} PMe₂,¹⁵⁹ PPh₂,⁵¹), **28**,¹⁶⁰ **29**,^{53a} **30** (X = Ph₃PN,⁴⁹ Cl,¹⁶¹ NMe₂,¹⁶²), **31**,^{108,163} **32**,^{128c,164} **33**,^{82,165} **34**,²³ **35**.¹⁶⁶

the rule of topological charge stabilization.¹⁶⁹ EHMO calculations for a 44-electron X_8 unit with D_{4d} geometry indicate that these sites are expected to carry the negative charge and, hence, will be favored by the more electronegative atoms.

Gleiter has presented an intuitively appealing explanation of the cage structure based on an EHMO analysis.^{118,170} A planar (D_{4h}) S_4N_4 molecule would be a 12- π -electron system with an open-shell configuration (Figure 15), unstable with respect to Jahn-Teller distortion. Correlation of the MO's of the hypothetical planar molecule with those of the experimental (D_{2d}) conformation shows that the orbital degeneracy of the ground state is lost in the latter and four of the previously π^* electrons are accommodated in S-S σ bonds. The possibility of four-center bonding between all four sulfur atoms has been suggested on the basis of electron-density measurements,¹⁵⁸ but calculated charge densities indicate that these S-S bonds involve closed-shell interactions.¹³⁹

Ab initio calculations identify the HOMO of **26** as a nitrogen-based lone-pair orbital and not the cross-ring S-S bond, while the LUMO is the S-S (σ^*) orbital




Figure 17. The LUMO's of S_4N_4 .

(Figure 17).^{82b,134a,171} This conclusion is supported by recent measurements of the X-ray fluorescence spectra of S_4N_4 .¹⁷²

3. Bicyclic Compounds, 1,5-E(NSN)₂E

The structural alternatives for molecules of the type 1,5-E(NSN)₂E, planar, **22**, or folded, **27**, can be understood by comparing their electronic structures with that of the unperturbed 10- π -electron system in $S_4N_4^{2+}$, **20**. Heteroatom incorporation transforms the non-bonding π HOMO of **20** into an antibonding orbital localized over the NSN linkages.^{22b,152-154} Strong electron-donor substituents on the heteroatom increase the electron density on the ring in **27** and, hence, weaken the S-N bonds. Puckering of the molecule alleviates this weakness by redistributing the orbital into a transannular S-S bonding interaction. Other related examples of cross-ring S-S bonds include the cyclophosphathiazenes **28**¹⁶⁰ and **29**.^{53a}

4. Disubstituted Derivatives, 1,5-X₂S₄N₄

The effect of adding π -donor substituents to the 1,5-sulfur atoms of S_4N_4 is similar to that of heteroatom substitution, i.e., a folded structure, **30**, is formed. MNDO calculations of the heats of formation of planar and puckered [$d(S-S) = 2.6 \text{ \AA}$] forms of **30** (X = Cl, F, NH₂) predict a higher stability for the latter conformation.¹⁵⁴ In the solid state these derivatives (X = Ph₃PN,⁴⁹ Cl,¹⁶¹ NMe₂)¹⁶² all exhibit an exo, endo configuration of the substituents as depicted by structure **30**. By contrast, the ¹⁹F NMR spectra of solutions of **30** (X = F¹⁷³ or ON(CF₃)₂)¹⁷⁴ indicate equivalent substituents (section IVC).

5. Tetrasulfur Pentanitride Anion and Cation, S₄N₅[±], and Pentasulfur Hexanitride, S₅N₆

The tetrasulfur pentanitride ions **32** and **33** can be considered as disubstituted derivatives of S_4N_4 in which the substituents (a bridging N atom) are in endo, endo positions. In the anion **32** there is one cross-ring S-S bond [$d(S-S) = 2.66^{128c}$ or 2.71 \AA^{164}], whereas in the corresponding cation **33** these sulfur atoms are separated by a distance (ca. $4.0 \text{ \AA}^{82,165}$) which exceeds the sum of the van der Waals radii for sulfur (3.7 \AA). These ions, and related S-N cages, e.g., S₅N₆, **31**, have been examined by the EHMO method by using a model in which the NSN units containing the unbridged sulfur atoms were treated as two pseudo-allylic 5- π -electron fragments.¹⁷⁵ If the bridging atom or group is able to withdraw π^* -electron density from the S_4N_4 cage, e.g., N⁺, the S-S bonding interaction will be weakened and a large S-S separation is expected. By contrast, if the

bridge provides a high-lying filled orbital, e.g., N⁻ or NSN, a stable structure with a short S-S distance is predicted.

Ab initio MO calculations provide further insight into the bonding in S-N cages. For example, the calculated overlap populations indicate that the introduction of an N⁻ bridge weakens the S-N bonds in the S_4N_4 cage,^{82b} consistent with the observed thermal instability of the $S_4N_5^-$ anion (section VC3).^{105,176} The X-ray structural parameters have prompted the suggestion that S₅N₆, **31**, can be viewed as NSN and S_4N_4 units linked through weak S-N bonds. The S-N connecting bonds are long, ca. 1.70 \AA , compared to those in the NSN bridge, ca. 1.54 \AA .^{108,163} The latter unit appears to have some "in molecule diradical" character,¹⁷⁷ which may account for the facile (and, in the solid state, explosive) thermal decomposition of S₅N₆ to S_4N_4 and sulfur.¹⁰⁸

6. Derivatives of Tetrasulfur Pentanitride, S₄N₅X

Two structural isomers are possible for S-N cages of the type S_4N_5X . In the first, exemplified by $S_4N_5O^-$, **34**, the exocyclic substituent is attached to a sulfur atom which is bonded to three nitrogens, whereas in the other isomer the sulfur atom bearing the substituent is connected to only two nitrogens, e.g., S_4N_5F , **35**. In the latter case, there is no cross-ring S-S bond and the exocyclic substituent may adopt a syn or anti conformation with respect to the S_4N_5 cage. The anti conformation has been established by X-ray crystallography for S_4N_5F , although MNDO calculations suggest there is very little energy difference between the two conformers.¹⁶⁶ However, the calculated barrier to inversion at the three-coordinate sulfur atom, which is necessary for the anti \rightarrow syn isomerization, is substantial ($46.7 \text{ kcal mol}^{-1}$).¹⁶⁶

IV. Spectroscopic Studies

In this section special emphasis will be placed on spectroscopic studies, e.g., photoelectron, UV-visible, MCD, and ESR spectra, which provide experimental data to corroborate calculations of the electronic structure of electron-rich S-N rings and cages described in section III.

A. Photoelectron Spectra

Photoelectron spectra (PES) supply information about the binding energies of either inner-core (X-ray PES) or valence-level (UV-PES) electrons. The data obtained provide experimental support for MO calculations of electronic energy levels and atomic charges in various molecules. UV-PES are most readily obtained for reasonably volatile compounds and studies of S-N heterocycles have included S₂N₂,^{134-136,178} derivatives of the S₃N₂ ring,^{179,180} S₄N₂,¹⁴⁴ S₄N₄,^{134,178,179,181} S₄N₄O₂,¹⁷⁹ and the eight-membered rings RC(NSN)₂CR (R = Ph, Me₂N;^{22b,152} R = *t*-Bu^{22b}).

The UV-PES for S₂N₂ show that the two upper π levels are nearly degenerate with the π_S orbital (primarily nonbonding electrons located on the two sulfur atoms) being of slightly higher energy ($\Delta E \sim 0.5 \text{ eV}$) than π_N .¹³⁴ The binding energies for S ($2p_{3/2}$) and N (1s) obtained from the X-ray PES data are consistent with charge transfer of ca. 0.19 e^- from sulfur to nitro-

gen, somewhat smaller than the value obtained (0.40 e⁻) from MNDO calculations.¹⁷⁸

Several groups have reported the UV^{134,181} and X-ray^{134,180,181} PES of S₄N₄ and good agreement between experimental ionization potentials and Hartree-Fock-Slater eigenvalues has been obtained.¹⁸² The X-ray PES data indicate a slightly greater S → N charge transfer in S₄N₄ compared to S₂N₂ consistent with MO calculations by the MNDO¹⁷⁸ or SCF-Xα¹⁸³ methods. The UV and X-ray PES data are also compatible with a modest S-S bonding interaction in S₄N₄.^{178,181}

The UV-PES of *t*-BuC(NSN)₂CBu-*t*, **27** (E = *t*-BuC), shows four well-resolved peaks with ionization potentials less than 11 eV and good agreement with calculated orbital energies for the highest five MO's of the model compound, **27** (E = CH),^{22b} was obtained. The experimental (UV-PES) and calculated values (ab initio MO calculations with configuration interaction) for the first nine ionization potentials of S₄N₂, **14** were also found to be in reasonable agreement.^{144b}

The X-ray PES of S₄N₄O₂ exhibits distinct binding energy values for three types of sulfur atoms and two pairs of unequivalent nitrogen atoms consistent with the known structure **21**.¹⁸⁰ Similarly, the X-ray PES of S₃N₂ derivatives show three sulfur (2p) and two nitrogen (1s) binding energies as expected for **6**.¹⁷⁹

B. UV-Visible and MCD Spectra

In contrast to unsaturated organic ring systems or cyclophosphazenes, electron-rich S-N heterocycles usually exhibit intense colors ranging from yellow, **13**, through orange, **16**, **20**, and **25**, and red, **14**, to blue, **11** and **24**. The colors are due to low energy ($\pi^* \rightarrow \pi^*$ or $n \rightarrow \pi^*$) transitions. Laidlaw and Trsic have had considerable success in correlating the experimental values (from electronic spectra) with excitation energies calculated by the transition-state method (Table II).^{182,184} Unambiguous assignments of UV-visible absorption bands to the appropriate electronic transitions can usually be made by comparing the calculated transition moments for the various alternatives.

In several instances the technique of magnetic circular dichroism (MCD) spectroscopy has been used to corroborate assignments based on UV-visible spectroscopic data. For example the assignment of the intense 360-nm band for S₃N₃⁻ to a π^* (2e'') → π^* (2a₂'') (HOMO → LUMO) excitation^{128b} has been confirmed by the measurement of the MCD spectrum of S₃N₃⁻.¹⁸⁶ The interpretation relies on the application of the Platt perimeter model,¹⁸⁷ as developed by Michl.¹⁸⁸ The S₃N₃⁻ ion is an unusual example of a (4N + 2)-electron *n*-center (*N* = 2, *n* = 2*N* + 2) π -electron perimeter, which will have two π -holes. As such, the 2e'' → 2a₂'' transition, which is doubly degenerate (Figure 11), is predicted to generate a negative *A* term in the MCD spectrum, as a result of the splitting of the excited state by the magnetic field. The MCD spectrum of S₃N₃⁻ does indeed display the characteristic shape of a negative *A* term.¹⁸⁶

The MCD spectrum of S₄N₃⁺, **16**, indicates that each of the two electronic absorption bands at 340 and 263 nm consists of two pairs of nearly degenerate transitions.¹⁴⁵ This conclusion has recently been confirmed by Hartree-Fock-Slater calculations of the electronic transitions (Figure 14 and Table II).¹⁴⁶ The UV-visible

TABLE II. Experimental and Calculated Electronic Transition Energies for S-N Rings and Cages

S-N heterocycle	λ_{\max} , eV			ref
	expt	calcd	assignment	
S ₄ N ₂	2.72	2.48	7a'' → 11a'	12, 144c
	3.29	3.16	10a'' → 11a'	
S ₃ N ₃ ⁻	3.44	3.60 ^a	2e'' → 2a ₂ ''	128b, 182
Ph ₃ P=N-S ₃ N ₃	2.59		<i>e</i>	109b, 189
	3.76			
Ph ₃ As=N-S ₃ N ₃	2.54		<i>e</i>	109b, 189
	3.69			
OS ₃ N ₃ ⁻	2.44		<i>e</i>	109b
	3.65			
(R ₂ PN)(SN) ₂	2.13-2.28 ^b	2.21 ^c	2a ₂ → 4b ₁	50
	4.59 ^d	4.43 ^c	3b ₁ → 4b ₁	
O ₂ S ₃ N ₃ ⁻	2.21		<i>e</i>	109b
	3.42			
S ₄ N ₃ ⁺	3.53	3.64	2a ₂ → 3a ₂	145, 146
	3.78	3.73	3b ₂ → 3a ₂	
	4.64	4.40	2a ₂ → 4b ₂	
	4.96	4.48	3b ₂ → 4b ₂	
S ₄ N ₄ ²⁺	3.58	3.71	1b _{1u} → 2e _g	148
	4.13	3.75	4e _u → 2e _g	
	4.73	4.73	1b _{2u} → 2e _g	
	4.43	4.08	2b _{1u} → 2b _{2g}	
<i>t</i> -BuC(NSN) ₂ Bu- <i>t</i>	3.40	3.36	2b _{1u} → 2b _{2g}	22b
	4.43	4.08	1a _u → 2b _{2g}	
S ₄ N ₄ H ⁺	5.30	5.15	2b _{1u} → 2b _{3g}	229
	2.62		<i>e</i>	
S ₅ N ₅ ⁺	3.82			185
	2.91		<i>e</i>	
S ₄ N ₅ ⁻	4.06		<i>e</i>	105b, 182
	3.49	3.55 ^a		
S ₅ N ₆	3.31	3.02	5a ₂ → 7b ₂	108b, 177
	4.96		11a ₁ → 7b ₂	

^a Obtained by adding 0.50 eV to the one-electron energy differences.¹⁸² ^b Exact value depends on the nature of R. ^c R = H. ^d R = Me. ^e See text.

spectrum of S₄N₄²⁺, a classic example of an eight-atom 10- π -electron perimeter, shows peaks at 346 and 262 nm attributed to $n\pi_S \rightarrow \pi^*$ and $n\pi_N \rightarrow \pi^*$ transitions, respectively, which should give rise to a pair of positive *A* terms in the MCD spectrum.^{148,188} The electronic spectrum of S₅N₅⁺ has not been assigned unambiguously, but it has been suggested that one of the bands (327 or 426 nm) is due to an $n \rightarrow \pi^*$ transition.¹⁵⁷

Monosubstituted derivatives of the type S₃N₃X, **15** (where X = O⁻, Ph₃PN, Ph₃AsN) exhibit remarkably similar electronic spectra (Table II) and molecular structures.^{120b} The result of the addition of a single exocyclic substituent on the π -electronic structure of the S₃N₃ ring is the loss of degeneracy for the HOMO's (2e''). Consequently, the strong absorptions at 480-510 and 330-340 nm have been assigned to the HOMO (π^*) → LUMO (π^*) and HOMO-1 (π^*) → LUMO (π^*) transitions, respectively.^{109b} This assignment is supported by the MCD spectra of Ph₃E=N-S₃N₃ (E = P, As).¹⁸⁹

The similarity between the electronic structures of (R₂PN)(SN)₂, **11**, and S₃N₃O₂⁻, **12**, referred to in section IIIA3, is also reflected in their UV-visible spectra. Both these heterocycles have an intense purple color due to a visible absorption band at ca. 560 nm attributed to the HOMO (π^*) → LUMO (π^*) transition.^{50,109b}

The pale yellow color of 1,5-(R₂PN)₂(SN)₂, **27** (E = Ph₂P, Me₂P) is in contrast to the dark orange color ($\lambda_{\max} \sim 460$ nm) of the structural isomers 1,3-(R₂PN)₂(SN)₂, **23**. In the absence of a cross-ring S-S bond, the latter isomers are π -electron-rich (10- π electrons) and the strong visible absorption band has been tentatively

TABLE III. ^{15}N NMR Parameters of Unsaturated S-N Heterocycles and Related Acyclic Systems

compound	chemical shift ^a	$^2J_{\text{NN}}$, ^b Hz	ref
S_4N_4	123.2		18
$1,5\text{-Cl}_2\text{S}_2\text{N}_4^{\text{c}}$	~172 (2 N), 182 (2 N)		120b
$1,5\text{-(Ph}_3\text{P=N)}_2\text{S}_4\text{N}_4$	153.6 (2 N), 159.8 (2 N)		18
$\text{S}_4\text{N}_5\text{Cl}$	39.3 (1 N), 225.1 (4 N)		18
S_4N_5^-	53.3 (1 N), 138.7 (4 N)		18
$\text{S}_4\text{N}_5\text{O}^-$	104.0 (1 N), 129.0 (2 N), 147.9 (2 N)		109b
S_3N_3^-	148.4		18
$\text{S}_3\text{N}_3\text{O}^-$	225.4 (2 N), 288.3 (1 N)		109b
$\text{S}_3\text{N}_3\text{-N=PPh}_3$	203.5 (2 N), 282.5 (1 N)		18
$\text{S}_3\text{N}_3\text{-N=AsPh}_3$	209.6 (2 N, d), 282.6 (1 N, tr)	2.2	18
$\text{S}_3\text{N}_3\text{O}_2^-$	215.2 (2 N), 353.3 (1 N)		109b
$(\text{Ph}_2\text{PN})(\text{SN})_2$	116.2 (2 N, d), 336.2 (1 N, tr)	2.8	50b
$(\text{Me}_2\text{PN})(\text{SN})_2$	123.2 (2 N, d), 338.9 (1 N, tr)	3.0	50b
$[(\text{PhO})_2\text{PN}](\text{SN})_2$	135.8 (2 N, d), 407.4 (1 N, tr)	2.7	50b
S_4N_2	266.3		18
$\text{S}_3\text{N}_2\text{O}^{\text{d}}$	255.8 (1 N), 326.0 (1 N)		195
S_4N_3^+	366.4 (1 N, tr), 376.1 (2 N, d)		18,
		194	
$[\text{SSNSS}]^-$	244.2		196
$\text{Ph}_3\text{P=N}_a\text{SN}_b\text{SS}$	86.9 (N_a), 472.0 (N_b)		18
$[\text{SSNS}]^-$	616.7		196

^a δ (in ppm) downfield of $\text{NH}_3(\text{l})$ at 25 °C (d = doublet; tr = triplet).
^b Coupling constants are only available for highly enriched (99%) samples; coupling to exocyclic substituents, e.g., ^{31}P , is not included in this data.
^c Measured at -80 °C in CH_2Cl_2 .
^d Chemical shifts were quoted relative to aqueous NH_4NO_3 , for which the shift is 376.25 ppm downfield of $\text{NH}_3(\text{l})$ at 25 °C.

ascribed to a HOMO (π^*) \rightarrow LUMO (π^*) transition.⁵¹

The UV-visible spectrum of S_4N_4 has been measured in different solvents over a range of temperatures¹⁹⁰ and the earlier assignments have been revised.¹⁸¹

C. ^{15}N NMR Spectra

The ^{14}N NMR chemical shifts of several S-N heterocycles have been reported, but the lines are usually very broad due to the quadrupole moment of ^{14}N ($I = 1$; 99.6%) and low symmetry around the nitrogen nuclei, e.g., S_4N_2 ($\Delta h_{1/2} = 110$ Hz),¹⁹¹ S_4N_4 ($\Delta h_{1/2} \sim 600$ Hz),¹⁹² and S_4N_5^- ($\Delta h_{1/2} = 1600$ Hz).⁹³ By contrast, the ^{15}N nucleus ($I = 1/2$) gives rise to narrow signals and is, in principle, a more useful probe for studying the structures of S-N rings and cages. It is usually necessary, however, to prepare ^{15}N -enriched samples due to the low natural abundance of ^{15}N (0.37%) and poor solubility of many S-N heterocycles. In view of the expense involved, this procedure is only warranted in carefully selected cases, e.g., when ^{15}N - ^{15}N coupling patterns can provide crucial structural information.

The first application of ^{15}N NMR spectroscopy in inorganic chemistry¹⁹³ involved the S-N heterocycle, S_4N_3^+ . In this classic example, reported by Logan and Jolly in 1965,¹⁹⁴ the ^{15}N NMR spectrum exhibits the doublet-triplet pattern expected for structure 16. More recently the ^{15}N NMR spectra of a variety of S-N rings and cages have been reported (Table III). The ^{15}N chemical shifts span a wide range of values, 40–400 ppm, which is even more pronounced for related acyclic systems. For example, the chemical shift difference for the two inequivalent nitrogen atoms in Ph_3PNSNSS is ca. 385 ppm¹⁸ and for the catena anions SSNSS^- and SSNS^- the δ values are 244 and 617 ppm, respectively.¹⁹⁶ In certain unsaturated nitrogen derivatives, e.g., nitrosyl (XN=O) or azo (XN=N_x) compounds, there is a




Figure 18. ^{15}N NMR spectrum of $\text{Ph}_2\text{PS}_2^*\text{N}_3$ (*N = 99% ^{15}N).⁵⁰




Figure 19. ^{15}N NMR spectrum of $\text{Ph}_3\text{As=N-S}_3\text{N}_3$ (99% enriched in ^{15}N).¹⁸

correlation between the nitrogen NMR chemical shifts, δ , and the wavelengths, λ , of the lowest electronic transition ($n \rightarrow \pi^*$).¹⁹³ This correlation does not apply to electron-rich S-N systems because the intense visible absorption bands are either magnetic dipole forbidden (e.g., $\pi^* \rightarrow \pi^*$) or the charge circulation is closer to sulfur than to nitrogen, i.e., the LUMO's are based mainly on sulfur.¹⁹³ Consequently, the λ values are not relevant to nitrogen shielding.

A few applications of ^{15}N NMR spectroscopy to structural problems in S-N heterocycles have been reported. The ^{15}N NMR spectra of the compounds $\text{R}_2\text{PS}_2\text{N}_3$ all show a characteristic pattern of a downfield doublet of triplets and an upfield doublet of doublets (Figure 18) consistent with the six-membered ring structure, 11, established by X-ray crystallography for $\text{R} = \text{Ph}$. Thus the NMR data confirm that the other derivatives ($\text{R} = \text{Me}$, OPh), which are thermally unstable oils, also have cyclic structures. The coupling constant, $^2J_{^{15}\text{N}-^{15}\text{N}}$, in these neutral compounds is significantly smaller (2–3 Hz) than the corresponding value (8.5 Hz) for the cation, S_4N_3^+ .

Derivatives of the type XS_3N_3 , 15, exhibit similar coupling constants (ca. 2 Hz) and characteristic chemical shifts of ca. 200–220 and 280–290 ppm for the nonequivalent endocyclic nitrogen atoms (Figure 19). This information afforded the first indication that the anion $\text{S}_3\text{N}_3\text{O}^-$ was among the products of the oxidation

of $S_3N_3^-$ by molecular oxygen (section IIG).^{109,120b}

¹⁵N NMR spectroscopy would be the technique of choice for the study of fluxional behavior in S-N cages. For example, a series of degenerate intramolecular rearrangements can be constructed leading to equivalence of all five nitrogen atoms in $S_4N_5^-$, **32**.¹⁷⁵ No experimental evidence in support of stereochemical nonrigidity in S-N cages has been presented, however, probably because the activation energy necessary for such a process is sufficient to disrupt the cage, e.g., by loss of N_2S (section VC3).¹⁰⁵

Three structural isomers, with different orientations of the exocyclic substituents, are possible for 1,5- $X_2S_4N_2$ derivatives, **30**. In the solid state an exo, endo configuration of substituents is usually found (section IIIB4).^{49,161,162} This isomer should exhibit two distinct ¹⁵N NMR signals, corresponding to two inequivalent pairs of nitrogen atoms, and this pattern is found for $X = NPPH_3$ ¹⁸ and $X = Cl$ at low temperatures.^{120b} However, the ¹⁹F NMR spectra of **30** ($X = F$ ¹⁷³ or $ON-(CF_3)_2$ ¹⁷⁴) show a single fluorine resonance indicative of either an endo,endo or, more likely, and exo,exo conformation. It is unlikely that exo,endo \rightarrow exo,exo isomerization occurs rapidly at ambient temperatures since the necessary inversion at sulfur would require 47 kcal mol⁻¹ for $X = F$, according to MNDO calculations.¹⁵⁴

No examples of the application of ³³S NMR spectroscopy in S-N chemistry are known due to the inherent difficulties associated with the relatively large quadrupole moment of ³³S ($I = 3/2$; 0.76%) and the low symmetry around the sulfur nuclei in most S-N heterocycles. It is likely, however, that reasonably narrow signals and characteristic chemical shifts could be obtained for sulfur in an approximately tetrahedral environment, e.g., SN_2O_2 in **12** or spirocyclic SN_4 in **29** (cf. SO_4^{2-}).¹⁹⁷

D. ESR Spectra


The frequent and rapid color changes observed in certain reactions of S-N heterocycles suggest the involvement of free radicals. The best characterized is the radical cation $S_3N_2^+$, **8**, which usually dimerizes in the solid state (section IIIA2).³³ This radical is easily produced in solution when S_4N_4 is oxidized, e.g., by $(CF_3SO_2)_2O$ ³⁴ or AsF_5 ,³⁶ or on treatment of other S-N compounds, e.g., $R_{4-n}Si(NSN)_nSiR_{4-n}$ ($n = 1-3$) or $R_3SiNSNSNSNSiR_3$ with $AlCl_3$ in CH_2Cl_2 .³⁷ The ESR spectrum of $S_3N_2^+$ shows a characteristic 1:2:3:2:1 quintet consistent with coupling of the unpaired electron with two equivalent ¹⁴N nuclei ($I = 1$).^{33,34,36,37} The spectrum of this radical enriched in ¹⁵N (99%) and ³³S (59.2%) also confirms the presence of two equivalent sulfurs and a third, unique sulfur nucleus.¹⁹⁸ Further analysis of the ESR data indicates that the unpaired electron is accommodated in a π orbital which is antibonding with respect to the S-S bond ($2a_2$ in Figure 9).^{137,198}

The related 7- π -electron radical 5-methyl-1,3,2,4-dithiazol-2-yl, prepared in solution by reduction of the cation **3** with silver powder or electrochemically, is also long lived and does not dimerize even at $-100^\circ C$.¹⁰¹ The ESR spectrum consists of three equally intense lines (attributed to coupling with one of the inequivalent nitrogen nuclei) each of which consists of a

1:4:7:7:4:1 sextet apparently due to exactly equal coupling to the other nitrogen nucleus and the three equivalent protons of the methyl group. The isomeric radical $PhCN_2S_2$, 4-phenyl-1,2,3,5-dithiazolyl, prepared by the reduction of the cation **2** with metals, is probably present in solid samples of the dimer since the ESR spectrum exhibits a broad signal ($g \sim 2.00$) characteristic of a single unpaired electron.¹²⁵ The ESR spectrum of the six-membered, seven π -electron radical $Ph_2C_2N_3S$, **10**, ($g = 2.0059$) consists of seven lines with equal hyperfine coupling constants to all three nitrogen nuclei.¹²⁶ This radical dimerizes via a long S-S bond in the solid state.

The thermally unstable radical anion $S_4N_4^-$ is formed when S_4N_4 is reduced by potassium metal or electrochemically in ether solvents.^{199,200} This radical gives rise to a nine-line ESR spectrum, consistent with the presence of four equivalent nitrogen nuclei ($a_N = 1.185$ G,²⁰⁰ 1.17 ± 0.2 G²⁰¹), and has a half-life of ca. 8 s at $-30^\circ C$. The decay of $S_4N_4^-$ obeys a first-order rate law ($E_A = 11.1 \pm 0.5$ kcal mol⁻¹) and probably occurs via intramolecular bond rupture²⁰¹ to give the $S_3N_3^-$ ion.⁴⁴

Extensive ESR investigations by Sutcliffe and co-workers,²⁰²⁻²⁰⁷ using isotopic enrichment, have established that the long-lived radicals 1,3,2-dithiazol-2-yl, **36** and 1,3,2-dithiazolidin-2-yl, **37**, are formed in the reactions of S_4N_2 , **14**, with strained olefins or alkynes, respectively. The large g factors found for **37** suggest



partial delocalization over the SNS unit but the substituents, R and R', have little effect on the hyperfine coupling constants.²⁰⁷ The benzo derivative of **37** exhibits similar ESR parameters.⁹⁷ The ESR spectrum of the isomeric radical, **38** (³³S-enriched), prepared from *o*-substituted anilines with sulfur and Br_2 or Cl_2 , indicates that two nonequivalent sulfur atoms are present.²⁰⁸

E. IR and Raman Spectra

There are only a few examples of a complete vibrational analysis of an electron-rich S-N heterocycle. The IR and Raman spectra of S_2N_2 , **1**, have been interpreted on the basis of D_{2h} geometry.²⁰⁹⁻²¹¹ The earlier assignments²¹² of IR-active vibrations have been confirmed,²¹¹ but there is disagreement over the numerical values of the harmonic force constants.²⁰⁹⁻²¹¹

The technique of ¹⁵N-enrichment has been used in several cases to distinguish S-N from S-S^{12,213} or anion vibrations.²¹⁴ The IR and Raman spectra of $S_4^{14}N_4$ and $S_4^{15}N_4$ have been assigned in accordance with the D_{2d} structure, **26**.²¹³ Force constants were calculated and good agreement between observed and calculated wave numbers was obtained.^{213,215} A strong Raman band at 218 cm⁻¹ for solid S_4N_4 was assigned to the symmetric stretching vibration of the weak cross-ring S-S bonds. Indeed Raman spectroscopy is a useful probe for detecting transannular S-S interactions in bicyclic or cage S-N molecules or ions.¹⁵⁹ The strongly Raman active

TABLE IV. S-S Bond Lengths and Raman Stretching Frequencies in S-N Bicyclic and Cage Compounds

S-N heterocycle	<i>d</i> (S-S), Å	ν (S-S), cm ⁻¹	ref
(Ph ₂ PN) ₄ (SN) ₂	2.38	245	160a
S ₅ N ₆	2.43	269	108
1,5-(Ph ₃ P=N) ₂ S ₄ N ₄	2.45	259	49, 216
1,5-Cl ₂ S ₄ N ₄	2.48	260	161, 216
Ph ₂ P(NSN) ₂ PPh ₂	2.53	268	51
Me ₂ P(NSN) ₂ PMe ₂	2.55	250	159
S ₄ N ₄ ⁺	2.60	218	158, 213
S ₄ N ₅ O ⁻	2.63	222	23, 213
S ₄ N ₅ ⁻	2.66, 2.71	186	128c, 164, 216

vibrations occur at frequencies in the range 180–300 cm⁻¹ for S-S bond lengths of 2.4–2.7 Å (Table IV). These bonds are substantially longer than a S-S single bond (ca. 2.06 Å) for which vibrational frequencies of 400–420 cm⁻¹ are observed.²¹⁷

The vibrational spectra of S₄N₄²⁺ cation as its FSO₃⁻ or AlCl₄⁻ salts are consistent with a flexible planar structure with *D*_{4h} symmetry (see section IIIA5).^{32,214} A normal coordinate analysis has been carried out for S₄N₄²⁺ and gave a value of 6.56 mdyn Å⁻¹ for the S-N valence force constant.²¹⁴ The vibrational spectra of S₄¹⁴N₂ and S₄¹⁵N₂ suggest a structural change (increase in symmetry) in going from the solid state to solutions.¹²

The IR and Raman spectra of Na⁺, K⁺, and R₄N⁺ salts of the S₃N₃⁻ ion, **13**, have been assigned according to the *D*_{3h} symmetry found by X-ray crystallography for the *n*-Bu₄N⁺ salt.^{128b} For the Cs⁺ salt, however, additional weak bands, indicative of a lowering of the symmetry of the anion, were observed. A comparison of the IR spectra of various S₃N₂ derivatives, **6**, has led to the identification of four characteristic vibrations of the S₃N₂ ring at ca. 990, 920, 740, and 660 (±20) cm⁻¹.⁴⁰

V. Reactions

The knowledge of electronic structures and, in particular, the composition and energies of frontier orbitals, developed in section III, is helpful in understanding many aspects of the reactivity of electron-rich S-N rings and cages. In this section we shall examine the rich diversity of their chemical behavior in the context of that information. The discussion is organized according to the following unifying themes, which portray the fate of S-N heterocycles in various reactions: (a) adduct formation, (b) redox reactions, (c) ring contraction, (d) ring expansion, and (e) chain formation.

A. Adduct Formation

1. With Lewis Acids

Electron-rich S-N heterocycles behave as weak Lewis bases with nitrogen as the coordination site in the formation of adducts with Lewis acids. For example, disulfur dinitride forms numerous complexes of the type S₂N₂·L (L = BCl₃,²¹⁸ SbCl₅,^{219,220} SnCl₄,¹¹⁴ BeCl₂,¹¹⁴ VCl₄,^{62a} [(Ph₃P)₂(CO)₂RuCl]⁺,¹¹⁴ and TiCl₄¹¹⁴) and S₂N₂·2L (L = SbCl₅,^{219,220} AlCl₃,⁵⁵ AlBr₃,^{56,114} CuCl₂⁵⁴). The latter adducts all have similar structures, **39**, in which a planar S₂N₂ ligand is coordinated through nitrogen atoms to the Lewis acid. The S-N bond distances are not significantly different from those in free S₂N₂ and the bond angles at N and S are ca. 95 and 85°, respectively.^{55,220} The displacement of S₂N₂ from its complex with AlCl₃ by S₄N₄ at 80 °C in high vacuum





Figure 20. Composition and relative energies of the π MO's of S₄N₄H⁺.²²⁹

has been proposed as the method of choice for the preparation of S₂N₂.²²¹



S₄N₄ reacts with a wide variety of Lewis acids to give adducts claimed to include 2:1, 1:1, 1:2, and 1:4 stoichiometries.²²² Adducts of the type S₄N₄·L are the most common and have been structurally characterized by X-ray crystallography for L = BF₃,²²³ AsF₅,²²⁴ SbCl₅,²²⁵ SO₃,^{226a} FSO₂NCO,^{226b} TaCl₅,²²⁷ and FeCl₃.²²⁸ All these adducts have a similar structure, **40**, in which one nitrogen atom is coordinated to the Lewis acid and the S₄N₄ ring is flattened into a puckered boat with no cross-ring interactions. A similar structure is adopted by the S₄N₄ rings in *trans*-SnCl₄·2S₄N₄^{60c} and the protonated cation S₄N₄H⁺.²²⁹



The formation of the N-H σ bond primarily involves the nitrogen-based σ lone-pair orbital (4e) of S₄N₄. The protonation of S₄N₄ lowers the symmetry and lifts the orbital degeneracy of the 12- π -electron ground state (Figure 20). MNDO MO calculations also reveal that there is a polarization of π charge towards the protonated nitrogen atom. The six occupied π MO's of S₄N₄H⁺ consist of three bonding (1b₁, 2b₁, 1a₂), two nonbonding (3b₁, 2a₂), and one antibonding (4b₁), which largely offsets the bonding contribution of 2b₁ (Figure

20).²²⁹ The net result of the localization of π^* electrons in the skeletal bonds to the protonated nitrogen is a strengthening of the SNSNSNS fragment at the expense of the S-N(H) bonds. Indeed a product of ring contraction, $S_3N_2NH_2^+$, is also formed in the protonation of S_4N_4 (section VC2). In contrast to the other 1:1 adducts, the complexes $CuX \cdot S_4N_4$ ($X = Cl, Br$) have polymeric structures, 41, in which two S_4N_4 ligands bridge two copper atoms via 1,3-N,N coordination sites.²³⁰

In addition to simple adduct formation, the interaction of S_4N_4 with excess Lewis acid can lead to oxidation to give S-N cations (section IIB1) or cleavage of the eight-membered S_4N_4 ring to give S_2N_2 adducts. The latter process occurs spontaneously with $CuCl_2$,⁵⁴ Al_2X_6 ($X = Cl$,⁵⁵ Br ⁵⁶), VCl_4 ,⁶² and, on heating, with $S_4N_4 \cdot TiCl_4$ and $S_4N_4 \cdot 2BeCl_2$.¹¹⁴ Infusible solids of composition $S_2N_2 \cdot SnX_2$ ($X = Cl, Br, OMe, NMe_2$), which may have polymeric structures, are obtained from S_4N_4 and tin(II) salts.²³¹

The sulfur-nitrogen oxides S_3N_2O , 6 ($X = O$), and $S_4N_4O_2$, 21, form adducts in which coordination to either exocyclic oxygen atoms or ring nitrogen atoms is observed. For example, coordination occurs through oxygen in $S_3N_2O \cdot L$ ($L = AsF_5, SbF_5$)¹⁹⁵ and $2S_3N_2O \cdot L$ ($L = SnCl_4, TiCl_4$),¹⁹⁵ whereas in complexes with silver(I), $[Ag(S_3N_2O)_2]^+ AsF_6^-$ ²³² and $[Ag(S_4N_4O_2)_4]^+ AsF_6^-$,²³³ the nitrogen atoms are bound to the metal. In the latter case all the coordinated nitrogen atoms are adjacent to the sulfur(VI) center.

Coordination occurs via oxygen in adducts of S_2N_2CO , 7, with Lewis acids, e.g., $[Zn(S_2N_2CO)_6][AsF_6]_2$.²³⁴


2. With Olefins

A number of electron-rich S-N heterocycles undergo reversible cycloaddition reactions with olefins, e.g., $S_4N_4 \cdot 2L$ ($L =$ norbornene,²³⁵ norbornadiene (NBD),²³⁵ or *trans*-cyclooctene²³⁶). At first there was disagreement over the regiochemistry of the addition, but it has now been established by X-ray crystallography that one C=C bond in each diene molecule adds across an SNS unit in $S_4N_4 \cdot 2NBD$.^{237,238} The preference for the 1,3-S,S mode of addition can be rationalized from a consideration of the LUMO of S_4N_4 (Figure 17), which is antibonding with respect to opposite pairs of sulfur atoms. Consequently, the HOMO of the olefin will add concertedly across nearest neighbor pairs of sulfur atoms.²³⁹

An S,S mode of addition is also observed for the 1:1 adducts of NBD with $(R_2PN)(SN)_2$, 11,^{50b,69,90} 1,3- $(R_2PN)_2(SN)_2$, 23 ($R = Ph$),⁵¹ and $Ph_3P=NS_3N_3$, 15 ($X = Ph_3PN$).²⁴⁰ This regiochemistry is readily rationalized from a consideration of the composition of the frontier orbitals of 11 (Figure 11).^{50b} Both the HOMO and LUMO are primarily sulfur based and of the correct symmetry to overlap with the LUMO and HOMO, respectively, of an olefin. The formation of the NBD adduct of 11 occurs very rapidly and is of practical value in characterizing and storing unstable derivatives, e.g., $R = Me$,^{50b} F ,⁹⁰ CF_3 ,⁶⁹ since the free heterocycle can be regenerated by mild heating of the adduct. The rates of addition of NBD to other S-N heterocycles differ significantly, that for 15 ($X = Ph_3PN$) being several orders of magnitude slower than that for 11 ($R = Ph$). Comparisons of the energies of the interacting frontier orbitals in these cycloadditions have led Oakley et al.

to suggest that the HOMO (olefin)-LUMO (SN heterocycle) interaction controls the kinetics of these reactions.²⁴⁰

The interaction of NBD with S_4N_2 , 14, results in rupture of the cyclic skeleton of 14 and produces 42,²⁴¹ presumably via insertion of the olefin into an S-S bond, 43²⁴² and the radical 36 (section IVD).




B. Redox Reactions

Electron-rich S-N heterocycles are easily oxidized. This process usually occurs without change in the size of ring and, since it involves the removal of π^* electrons, results in a strengthening of the π framework. The same S-N heterocycles are also good *electron acceptors* since the acceptor orbitals (LUMO's) are low lying relative to those in aromatic hydrocarbons or cyclophosphazenes (section III). The LUMO's are π^* orbitals, antibonding with respect to S-N linkages. Consequently, the addition of electrons will promote ring opening to give chains which often recyclize to smaller rings. In this section redox reactions which occur with retention of ring size will be discussed and those which take place with contraction or expansion of the ring are considered in sections VC and VD.


1. Oxidation

The weak intra- or intermolecular S-S bonds found in S-N bicyclic, cage, or dimeric molecules are easily cleaved by halogens. The oxidative addition of Cl_2 to S_4N_4 must be conducted at low temperatures to avoid the decomposition of 1,5- $S_4N_4Cl_2$ (30 ($X = Cl$), section IIB1). By contrast, the bifunctional derivatives, 44,



obtained from the addition of halogens across the S-S bond of heterocyclothiazenes, 27 ($E = Ph_2P$,²⁴³ $CNMe_2$)^{80,244} have much higher thermal stability.

Dimers such as $(PhCN_2S_2)_2$ (section IIIA2) or $S_6N_4O_4$, 45,²⁴⁵ also react readily with halogens. In the former case the products are salts of the cation 2,²⁴⁶ while 45 reacts with Cl_2 to give $S_3N_2O_2Cl_2$, 46, which decomposes at $-30^\circ C$ to $S_3N_3O_2Cl$, 47.²⁴⁷






Figure 21. $\Delta E = \Delta[\epsilon(\pi \text{ LUMO}) - \epsilon(\pi \text{ LOMO})]$ vs. $E_{1/2}$ for $(\text{Ph}_2\text{PN})_2(\text{SN})^+$ (\square), $(\text{Ph}_2\text{PN})(\text{SN})_2$ (Δ), S_3N_3^- (\circ), and $(\text{Ph}_2\text{PN})_3$ (\bullet).¹⁴¹

The six-membered rings, **11** and **12**, are readily oxidized by a source of Cl_2 , e.g., SO_2Cl_2 or PhICl_2 , to give $(\text{Ph}_2\text{PN})(\text{NSCl})_2$ ⁹⁰ and $(\text{NSO}_2)(\text{NSCl})_2^-$,²⁴⁸ respectively.

Surprisingly, only a few examples of the oxidation of neutral S–N heterocycles to the corresponding cations have been reported, e.g., $\text{S}_4\text{N}_4 \rightarrow \text{S}_4\text{N}_4^{2+}$ (eq 3).

2. Reduction

Fukui et al. have pointed out that the energetically low-lying LUMO of S_4N_4 implies *electron-deficient* properties.²⁴⁹ This assertion is substantiated by the facile electrochemical reduction of S_4N_4 at -0.93 V (vs. $\text{Ag}/0.1$ M AgClO_4) to the anion radical S_4N_4^- , which takes on an additional electron at low temperatures to give the unstable dianion $\text{S}_4\text{N}_4^{2-}$.²⁵⁰ In the presence of a proton source the product is $\text{S}_4\text{N}_4\text{H}_4$.²⁵¹

The reduction of the cations **2–4** to the corresponding radicals, e.g., **9**, can be achieved with silver powder, sodium dithionite or via a reversible electrochemical reduction (cyclic voltammetry).¹⁰¹ The radical **10** is produced by the reduction of the corresponding cation, as a chloride salt, with triphenylantimony.¹²⁶

The electrochemical reduction potentials of the series of related six-membered rings, P_3N_3 , P_2SN_3^+ , PS_2N_3 , and S_3N_3^- , correlate well with the energies of the LUMO's adjusted for the effect of charge (Figure 21).¹⁴¹ Both of the electron-rich rings, **11** ($\text{R} = \text{Ph}$) and **13**, are better electron acceptors than the electron-precise P_3N_3 ring as a consequence of the lowering of the LUMO energy by the substitution of the more electronegative sulfur atoms for phosphorus in the PSN rings (section IIIA3).

C. Ring Contraction

Ring contraction is a common feature of the reactions of electron-rich S–N heterocycles. It can occur under the influence of nucleophilic or reducing agents, electrophiles, or on thermolysis.

1. Reactions with Reducing Agents or Nucleophiles

The outcome of nucleophilic degradation is frequently similar to that observed for the reduction of S–N rings. This parallel behavior can be rationalized since both processes will populate the π^* (S–N antibonding) LUMO's and, hence, foster ring-opening reactions. The anion radical S_4N_4^- , formed by the electrochemical reduction of S_4N_4 , undergoes an intramolecular bond rupture with an activation energy of 11.1


± 0.5 kcal mol⁻¹²⁰¹ to give S_3N_3^- , **13**.^{44,251} The nucleophilic degradation of S_4N_4 by anionic nucleophiles also produces salts of **13** (section IIB2). Similarly, both the electrochemical reduction¹¹⁵ and the nucleophilic cleavage of $\text{S}_4\text{N}_4\text{O}_2$, **21**, yield the ring-contracted anion $\text{S}_3\text{N}_3\text{O}_2^-$, **12**.^{109b,111a}

The polarity of the S–N bond, $\text{S}^{\delta+}-\text{N}^{\delta-}$, is expected to attract the initial nucleophilic attack to the sulfur atom, e.g., the formation of $\text{MeOS}_4\text{N}_4\text{O}_2^-$ from **21** and methoxide ion (Figure 7). However, subsequent rearrangement often gives products in which the nucleophile is bonded to nitrogen. For example, derivatives of the type $\text{Ph}_3\text{E}=\text{N}-\text{S}_3\text{N}_3$ ($\text{E} = \text{P}, \text{As}$) are formed from the reactions of S_4N_4 with Ph_3E ($\text{E} = \text{P}, \text{As}$).^{45,47,48} Other phosphorus(III) reagents, e.g., Ph_2PX , stimulate ring-opening reactions of S_4N_4 and, if X is a good leaving group, the phosphorus atom is incorporated into the ring-contracted product, e.g., **11** and **23** ($\text{X} = \text{H}$)^{51,52} or $(\text{Ph}_2\text{PN})_2(\text{NSCl})$.⁵³

2. Reactions with Electrophiles


Ring-contraction reactions of electron-rich S–N heterocycles under the influence of electrophiles commonly involve the formation of the five-membered ring S_3N_2^+ , **8**, or its derivatives $\text{S}_3\text{N}_2\text{X}$, **6**. The radical cation is obtained when S_4N_4 is treated with oxidizing agents, e.g., $(\text{CF}_3\text{SO}_2)_2\text{O}$,³⁴ AsF_5 ,³³ or AlCl_3 ,³⁷ while **6** is produced from S_4N_4 and acyl halides or acid anhydrides ($\text{X} = \text{RC}(\text{O})\text{N}=\text{O}$)^{38–40} and on protonation ($\text{X} = \text{NH}_2^+$).²⁵² The same products are obtained from the reactions of S_3N_3^- , **13**, with electrophiles, e.g., CCl_3COCl ^{43c} or $\text{HBF}_4 \cdot \text{Et}_2\text{O}$.²⁵²

The reasons for the instability of the six-membered ring, **48**, relative to **6** ($\text{X} = \text{NH}_2^+$) have been discussed by Oakley et al.²⁵² in the light of MNDO MO calcula-



tions on the model compound $\text{S}_3\text{N}_3\text{H}$ (**48**, $\text{R} = \text{H}$). The formation of the N–H σ bond probably involves the nitrogen-based σ lone-pair orbital (4e) of S_3N_3^- ^{128b} and results in extensive polarization of the π^* electron density into the skeletal bonds to the protonated nitrogen (cf. $\text{S}_4\text{N}_4\text{H}^+$, section VA1). Consequently the π bond order in those S–N bonds is reduced to zero, while that in the remaining S–N linkages is increased to ca. 0.3 (cf. 0.11 in S_3N_3^-).²⁵² The localized weakening of the cyclic structure results in the observed ring contraction.

The reaction of S_4N_4 with an excess of SO_3 produces the six-membered ring $\text{S}_3\text{N}_2\text{O}_5$, **49**,²⁵³ and $(\text{S}_6\text{N}_4^{2+})(\text{S}_3\text{N}_3\text{O}_4^-)_2$,^{113b} in which the anion **50** is iso-electronic with **49**, and has a remarkably similar structure.²⁵⁴



The ring contraction of S_4N_4 to S_4N_3^+ , **16**, occurs with certain electrophiles, e.g., liquid Br_2 or ICl ,²⁷ or S_2Cl_2 .²


3. Thermolysis

Electron-rich S-N heterocycles are thermodynamically unstable with respect to the loss of N₂ and their structural frameworks are weaker than those of cyclophosphazenes due, in part, to the partial occupation of π* MO's. In addition there is often a low kinetic barrier to decomposition and caution must be exercised to avoid explosions. Under suitably controlled conditions, however, thermal decomposition can be used to prepare other S-N compounds. The classic example is the generation of S₂N₂ by thermolysis of S₄N₄ at ca. 300 °C using a silver-wool catalyst.²⁴ More recently, other cyclothiazenes, e.g., Ph₃As=N-S₃N₃⁴⁸ and S₄N₃⁺Cl⁻,²⁵⁵ have been used to produce S₂N₂ under milder conditions.

The thermally initiated reductive elimination of an unstable NSN fragment²⁵⁶ converts bicyclic to monocyclic S-N rings, e.g., S₄N₅⁻ → S₃N₃⁻.¹⁰⁵ Kinetic measurements of this process for **45** (E = Me₂P) gave an activation energy of 24.5 ± 1.4 kcal mol⁻¹ for the release of NSN.⁹⁰ The thermolysis of **45** (E = F₂P) has been used to prepare the unstable six-membered ring **11** (E = F₂P) as an NBD adduct.⁹⁰ The reverse of this process, the cycloaddition of an NSN moiety is comparable to the cycloaddition of olefins to an S-N ring in terms of frontier-orbital considerations since the NSN unit and an olefin are isolobal.⁹⁰

4. Elimination of NSX

The best known example of ring contraction via loss of NSCl occurs in the synthesis of (NSCl)₃ by chlorination of S₄N₄. The intermediate product 1,5-Cl₂S₄N₄ (**30**, X = Cl) undergoes an eight- → six-membered ring contraction, with elimination of NSCl either prior to or immediately after the addition of the second mole of Cl₂.²⁵⁷ Although S₄N₄F₄ is thermally more stable than S₄N₄Cl₄,^{258a} it undergoes ring contraction with elimination of NSF on treatment with AsF₅ (eq 18).^{258b} The six-membered ring (NSF)₃ is the major product of the fluorination of S₄N₄ under certain conditions (section IIB1).²⁵⁹



The chlorination of 1,3-(Ph₂PN)₂(SN)₂, **23** (R = Ph), produces the six-membered ring (Ph₂PN)₂(NSCl), presumably via loss of NSCl from 1,3-(Ph₂PN)₂(NSCl)₂.²⁴³

D. Ring Expansion

Ring expansion reactions of electron-rich S-N rings, although less common than ring contraction, have been observed in several instances. Insertion reactions or the dimerization of reactive monomer units are the most prevalent examples of this type of behavior.

1. Dimerization

Considerable interest has focussed on the mechanism of the polymerization of S₂N₂ to (SN)_x (section VE1), but S₂N₂ is also known to dimerize to S₄N₄ in organic solvents in the presence of nucleophiles.^{260,261} Fukui et al.²⁶² have explained the structural reorganization involved in this process in terms of the HOMO-LUMO




Figure 22. Orbital interactions in the dimerization of S₂N₂ to S₄N₄.²⁶²

interactions of two S₂N₂ molecules oriented so that the sulfur atoms of one molecule interact with the nitrogen atoms of the other (Figure 22). In this way two new S-N σ bonds, S₁-N₃ and N₂-S₄ are formed and the S-N π bonds, S₁-N₂ and N₃-S₄, are weakened. In addition, an S₂-S₃ interaction develops. Deformation of this "stacking model" to give the S₄N₄ cage can be achieved by twisting about the S₂-S₃ fulcrum with formation of an S₁-S₄ bond.²⁶²

Reduction of (Ph₂PN)₂(NSCl) with triphenylantimony yields the twelve-membered ring, **28**.¹⁶⁰ The ease with which this ring-opening dimerization occurs has been attributed to the strongly S-N antibonding character of the LUMO (π*) of the P₂SN₃⁺ ring (Figure 12).¹⁴¹ In contrast, reduction of (PhCN)₂(NSCl) produces the radical **10**, which dimerizes without cleavage of the ring.¹²⁶

Certain derivatives of **11** undergo a spontaneous six- to eight-membered ring expansion under ambient conditions. In the two known examples, however, different isomers of the eight-membered ring were isolated. The decomposition of **11** (R = Me) produced the 1,5-isomer **27** (E = Me₂P)¹⁵⁹ whereas the 1,3-isomer **23** was obtained from **11** (R = CF₃).⁶⁹ It is possible that these products are formed by the dimerization of an (R₂PN)(SN) moiety generated by spontaneous cleavage of the ring in **11**.


2. Insertion Reactions

The preparation of S₅N₅⁺ salts, **25**, from NS⁺ and S₄N₄ (sections IIC2 and IIF) is the best known example of a ring expansion reaction in S-N chemistry. In a recent variation of this approach, S₅N₅⁺Cl⁻ was prepared by the direct reaction of S₄N₄ with (NSCl)₃ in formic acid.²⁶³ Ring-expansion reactions also occur in the preparation of S₄N₄ from S₃N₂Cl⁺, **5**, via S₄N₃⁺ (Figure 1), but the mechanism of these transformations is uncertain.²

The reactions of Si-N reagents with cyclothiazyl halides can bring about ring expansion. For example, the six-membered rings **11** (R = CF₃, C₂F₅) may be prepared from S₃N₂Cl⁺Cl⁻ and R₂PN(SiMe₃)₂.⁶⁹ Six- → eight-membered ring expansion occurs when (Ph₂PN)₂(NSCl) is treated with Me₃SiNSO²⁶⁵ or Me₃SiNSNSiMe₃,^{53a,264} to give **23** (R = Ph) or **29**, respectively. The latter transformation can be viewed as a double insertion of SN groups from the sulfur diimide (NSN) reagent into separate P₂SN₃ rings with the common sulfur atom becoming the spirocyclic center.

A six- → eight-membered ring expansion also takes place in the reaction of (Me₂NCN)(NSCl)₂ with Me₃SiNSO to give 1,5-Me₂NC(NSN)₂S₂Cl, **51**, a hybrid of **27** (E = Me₂NC) and **30** (X = Cl) with a similar

folded structure and an endo chlorine atom.⁸⁰



3. Oxidation

The oxidation of anions, $S_3N_3^-$ or $S_4N_5^-$ (**32**) with halogens results in substantial structural reorganizations to give S_4N_4 ¹⁰⁷ and S_5N_6 (**31**),¹⁰⁸ respectively. The same products are obtained from the electrochemical oxidation of $S_3N_3^-$ and $S_4N_5^-$.^{250a} Interestingly, the reduction of $(NSCl)_3$ with metals^{16,17} or azides²¹ also produces S_4N_4 , in addition to $(SN)_x$.²¹ It is possible that a common intermediate, the $S_3N_3\cdot$ radical, is involved in these redox reactions of the trithiazyl ring.

In contrast to the ring contraction reaction observed when **23** ($R = Ph$) is oxidized by Cl_2 (section VC4), an eight- \rightarrow twelve-membered ring expansion to give $(Ph_2PN)_4(SN)_2^{2+}$, the dication of **28**, occurs on treatment of **23** with Br_2 or Cl_2 .

E. Chain Formation

The formation of open-chain intermediates is implicated in a number of ring-contraction reactions described in section VC. In this section the production of stable catena molecules and ions from S-N rings will be discussed. Ring \rightarrow chain transformations can occur spontaneously or under the influence of nucleophiles or electrophiles, heat, or light.

1. Polymerization



The spontaneous topochemical polymerization of S_2N_2 to $(SN)_x$ at 0 °C,²⁴ the most prominent example of a polymerization reaction in S-N chemistry, has been described in detail by Labes et al.¹ On the basis of EHMO calculations, this process is both thermally and photochemically allowed,²⁶⁶ and the photo-induced polymerization occurs at -65 °C in tetrahydrofuran.²⁶⁷ INDO-type calculations²⁶⁸ for S_2N_2 and several deformed structures of $(SN)_2$, as well as the partially polymerized units, $(SN)_4$ and $(SN)_6$, show that the triplet biradical nature (section III1) emerges in deformed structures of $(SN)_2$ in accord with the ESR signal ($g = 2.005$) observed at the initial stage of polymerization.²⁴ No triplet character is predicted for $(SN)_4$ or $(SN)_6$ consistent with the fact that the system gradually becomes diamagnetic as polymerization proceeds.

The direct polymerization of S_4N_4 can be induced by a radiofrequency discharge.²⁶⁹ The reactions of powdered S_4N_4 with bromine or the interhalogens, ICl or IBr , yield highly conducting polymers or approximate composition $(SNBr_{0.4})_x$ and $[(SN)(IX)_{0.4}]_x$, respectively.^{29,30} The polymerization of S-N rings to $(SN)_x$ can also be promoted electrochemically. The electrochemical reduction of salts of the $S_5N_5^+$ cation, **25**, in SO_2 or CH_2Cl_2 at low temperatures produces microcrystalline $(SN)_x$.^{270,271}

2. With Nucleophiles

Nucleophilic attack ruptures the cage of S_4N_4 to give open-chain intermediates which, in many cases, cyclize

to give ring-contracted products (Figure 2 and section VC1). Some nucleophiles, however, produce stable S-N chains. The cleavage of S_4N_4 by Si-N or Sn-N reagents is the most widely studied example of this behavior (eq 19 and 20).²⁷²⁻²⁷⁸ At elevated temperatures the latter



reaction proceeds further to give the five-membered ring $(Me_2SnN_2S_2)_2$ (cf. eq 6). The reaction of S_4N_4 with the primary amine, $PhCH_2NH_2$, produces an open-chain tetrasulfide, $PhCHNSSSSNCHPh$.²⁷⁹


Treatment of S_4N_4 with aryl Grignard reagents yields small amounts of $ArSNSNSAr$, in addition to the major product, $ArSSAr$.^{280,281} The reaction of S_4N_4 with diazo compounds, e.g., Ph_2CN_2 , gives the catenated derivative, $Ph_2CNSNSNSNCPh_2$,²⁸² a member of a class of compounds used as electrophotographic pigments.²⁸³

The primary product of the reaction of $S_4N_4O_2$ with Ph_3As is a derivative of the S_3N_2 ring, which decomposes in acetone with the elimination of S_4N_4 and sulfur to give the 11-atom chain, $Ph_3AsNSO_2NSNSNSNSO_2NAsPh_3$ (Figure 7).²⁸⁴

The reaction of $S_3N_2Cl^+Cl^-$, **5**, with amines can also lead to chains. For example, when **5** is treated with Me_3SiNMe_2 in the molar ratio 1:3 the product is $Me_3SiNSNSNMe_2$. The reaction of **5** with $C_6F_5N(SnMe_3)_2$ gives $C_6F_5NSNSNSNC_6F_5$.²⁸⁵

3. Thermolysis

Thermally induced ring contraction reactions were described in section VC3. In a few cases, the thermal breakdown of electron-rich S-N rings produces stable catenated species. For example, the six-membered rings $S_3N_3^-$, **13**, and $Ph_3P=N-S_3N_3$, **15** ($X = Ph_3P=N$), undergo ring opening in boiling acetonitrile to give $SSNSS^-$ ^{105,176} and $Ph_3P=NSNSS$,⁴⁸ respectively. Both the anion and the neutral molecule have a catenated cis-trans structure, **52**, with short (ca. 1.9 Å) polar, terminal S-S bonds. It is pertinent to note that ab



initio MO calculations on the uniform deformation of the $S_3N_3^-$ ion suggest the decomposition channel which gives $NSN + SNS^-$.²⁸⁶

The S_7N^- ion, obtained by the deprotonation²⁸⁷ or electrochemical reduction²⁸⁸ of S_7NH , decomposes above ca. -50 °C to give **52** ($X = S^-$).

VI. Summary and Conclusions

Many aspects of the molecular structures, chemical behavior, and spectroscopic properties of electron-rich S-N heterocycles derive from the fact that the HOMO and LUMO are both normally π^* or $n\pi$ orbitals. The result of the partial occupation of π^* orbitals is a weakened π framework which is readily deformed or even disrupted under the influence of heat or upon adduct formation. The intense colors of many S-N

rings are also a manifestation of their electron richness and can be attributed to $\pi^* \rightarrow \pi^*$ (or $n\pi \rightarrow \pi^*$) electronic transitions.

Electron-rich S-N heterocycles are easily oxidized. They are also good electron acceptors and this property can be attributed to the low energy of their acceptor orbitals (LUMOs) relative to those of the corresponding aromatic hydrocarbons or cyclophosphazenes. The LUMO's are usually strongly antibonding with respect to the S-N bonds and, consequently the usual result of addition of an electron or nucleophilic attack is ring opening to give catenated species or ring-contracted products.

Although the possibilities for the preparation of novel binary sulfur-nitrogen heterocycles appear to be limited, there are already indications that the scope for the incorporation of heteroatoms such as carbon, phosphorus, or transition metals into these ring systems is considerable. These heterocyclothiazenes may prove to be suitable precursors for linear polymers, related to $(SN)_x$, but containing carbon or other heteroatoms as well as thiazyl units. In addition to an alternative synthetic approach, electrochemical techniques should provide valuable information on the redox properties of new compounds.

The characteristically small HOMO-LUMO energy gaps found for electron-rich sulfur-nitrogen rings suggest photochemical activity. To date, however, there has been no major effort to study the photochemistry of these compounds. Mechanistic studies of the ubiquitous ring-contraction and ring-expansion reactions also represent a virgin field. Finally, an evaluation of the effect of the amalgamation of unsaturated sulfur-nitrogen rings with transition metals on their electronic structures and chemical reactivity represents an interesting challenge.

VII. Acknowledgments

I thank Professor W. G. Laidlaw for many helpful discussions particularly with regard to section III. The financial support of the Natural Sciences and Engineering Research Council of Canada and the award of a Killam Resident Fellowship by the University of Calgary are also gratefully acknowledged.

VIII. References and Notes

- (1) Labes, M. M.; Love, P.; Nichols, L. F. *Chem. Rev.* **1979**, *79*, 1.
- (2) Heal, H. G. "The Inorganic Heterocyclic Chemistry of Sulfur, Nitrogen and Phosphorus"; Academic Press: London, 1980.
- (3) Wade, K. *Adv. Inorg. Chem. Radiochem.* **1976**, *18*, 1.
- (4) Banister, A. J. *MTP Int. Rev. of Sci.: Inorg. Chem., Ser. Two* **1975**, *3*, 41.
- (5) (a) Roesky, H. W. *Adv. Inorg. Chem. Radiochem.* **1979**, *22*, 239. (b) Roesky, H. W. *Angew. Chem., Int. Ed. Engl.* **1979**, *18*, 91.
- (6) Chivers, T. In "The Chemistry of Inorganic Ring Systems"; Haiduc, I., Sowerby, D. B., Eds.; Wiley-Interscience: London, in press.
- (7) Banister, A. J.; Smith, N. R. M. *J. Chem. Educ.* **1982**, *59*, 1058.
- (8) Villena-Blanco, M.; Jolly, W. L. *Inorg. Synth.* **1967**, *9*, 98. Banister, A. J.; *Inorg. Synth.* **1977**, *17*, 197.
- (9) MacLean, G. K.; Passmore, J.; Rao, M. N. S., private communication.
- (10) Apter, S.; Carruthers, M.; Sutcliffe, L. H. *Inorg. Chim. Acta* **1978**, *31*, L455.
- (11) Niinisto, L.; Laitinen, R. *Inorg. Nucl. Chem. Lett.* **1976**, *12*, 191.
- (12) Chivers, T.; Coddling, P. W.; Laidlaw, W. G.; Liblong, S. W.; Oakley, R. T.; Trsic, M. J. *Am. Chem. Soc.* **1983**, *105*, 1186.
- (13) Jolly, W. L.; Maguire, K. D. *Inorg. Synth.* **1967**, *9*, 102.
- (14) Roesky, H. W.; Schaper, W.; Petersen, O.; Müller, T. *Chem. Ber.* **1977**, *110*, 2695.
- (15) Alange, G. G.; Banister, A. J.; Bell, B. *J. Chem. Soc., Dalton Trans.* **1972**, 2399.
- (16) (a) Banister, A. J.; Fielder, A. J.; Hey, R. G.; Smith, N. R. M. *J. Chem. Soc., Dalton Trans.* **1980**, 1457. (b) Banister, A. J.; Fielder, A. J.; Hauptman, Z. V.; Smith, N. R. M. U.S. Patent 4268491; *Chem. Abstr.* **1981**, *95*, 83144k.
- (17) Berthet, M. P.; Vincent, H.; Monteil, Y. *Z. Naturforsch., B: Anorg. Chem., Org. Chem.* **1980**, *35B*, 329.
- (18) Chivers, T.; Oakley, R. T.; Scherer, O. J.; Wolmershäuser, G. *Inorg. Chem.* **1981**, *20*, 914. Chivers, T.; Oakley, R. T. unpublished results.
- (19) Peake, S. C.; Downs, A. J. *J. Chem. Soc., Dalton Trans.* **1974**, 859.
- (20) Vincent, H.; Monteil, Y.; Berthet, M. P. *J. Inorg. Nucl. Chem.* **1980**, *42*, 5.
- (21) (a) Passmore, J.; Rao, M. N. S. *J. Chem. Soc., Chem. Commun.* **1980**, 1268. (b) Kennett, F. A.; MacLean, G. K.; Passmore, J.; Rao, M. N. S. *J. Chem. Soc., Dalton Trans.* **1982**, 851.
- (22) (a) Ernest, I.; Holick, W.; Rihs, G.; Schomburg, D.; Shoham, G.; Wenkert, D.; Woodward, R. B. *J. Am. Chem. Soc.* **1981**, *103*, 1540. (b) Gleiter, R.; Bartetzko, R.; Cremer, D. *J. Am. Chem. Soc.* **1984**, *106*, 3437.
- (23) (a) Steudel, R.; Luger, P.; Bradaczek, H. *Angew. Chem., Int. Ed. Engl.* **1973**, *12*, 316. (b) Luger, P.; Bradaczek, H.; Steudel, R. *Chem. Ber.* **1976**, *109*, 3441.
- (24) (a) Mikulski, C. M.; Russo, P. J.; Saran, M. S.; MacDiarmid, A. G.; Garito, A. F.; Heeger, A. J. *J. Am. Chem. Soc.* **1975**, *97*, 6358. (b) Cohen, M. J.; Garito, A. F.; Heeger, A. J.; MacDiarmid, A. G.; Mikulski, C. M.; Saran, M. S.; Kleppinger, J. *J. Am. Chem. Soc.* **1976**, *98*, 3844.
- (25) (a) Vincent, H.; Monteil, Y. *Synth. React. Inorg. Met.-Org. Chem.* **1978**, *8*, 51. (b) Zborilova, L.; Gebauer, P. *Z. Anorg. Allg. Chem.* **1979**, *448*, 5. (c) Zborilova, L.; Gebauer, P. *Z. Chem.* **1979**, *19*, 32.
- (26) Ruppert, I. *J. Fluorine Chem.* **1982**, *20*, 241.
- (27) Wolmershäuser, G.; Street, G. B. *Inorg. Chem.* **1978**, *17*, 2685.
- (28) (a) Wolmershäuser, G.; Street, G. B.; Smith, R. D. *Inorg. Chem.* **1979**, *18*, 383. (b) Wolmershäuser, G.; Kruger, C.; Tsay, Y.-H. *Chem. Ber.* **1982**, *115*, 1126.
- (29) Street, G. B.; Bingham, R. L.; Crowley, J. I.; Kuyper, J. *J. Chem. Soc., Chem. Commun.* **1977**, 464.
- (30) (a) Akhtar, M.; Chiang, C. K.; Heeger, A. J.; MacDiarmid, A. G. *J. Chem. Soc., Chem. Commun.* **1977**, 846. (b) Akhtar, M.; Chiang, C. K.; Heeger, A. J.; Milliken, J.; MacDiarmid, A. G. *Inorg. Chem.* **1978**, *17*, 1539.
- (31) (a) Gillespie, R. J.; Slim, D. R.; Tyrer, J. D. *J. Chem. Soc., Chem. Commun.* **1977**, 253. (b) Gillespie, R. J.; Kent, J. P.; Sawyer, J. F.; Slim, D. R.; Tyrer, J. D. *Inorg. Chem.* **1981**, *20*, 3799.
- (32) Sharma, R. D.; Aubke, F.; Paddock, N. L. *Can. J. Chem.* **1981**, *59*, 3157.
- (33) (a) Gillespie, R. J.; Kent, J. P.; Sawyer, J. F. *Inorg. Chem.* **1981**, *20*, 3784. (b) Gillespie, R. J.; Ireland, P. R.; Vekris, J. E. *Can. J. Chem.* **1975**, *53*, 3147.
- (34) (a) Roesky, H. W.; Hamza, A. *Angew. Chem., Int. Ed. Engl.* **1976**, *15*, 226. (b) Krebs, B.; Henkel, G.; Pohl, S.; Roesky, H. W. *Chem. Ber.* **1980**, *113*, 226.
- (35) Thewalt, U.; Burger, M. *Z. Naturforsch., B: Anorg. Chem., Org. Chem.* **1981**, *36B*, 293.
- (36) Gillespie, R. J.; Kent, J. P.; Sawyer, J. F. *Inorg. Chem.* **1981**, *20*, 4053.
- (37) Giordan, J.; Bock, H.; Eiser, M.; Roesky, H. W. *Phosphorus Sulfur* **1982**, *13*, 19.
- (38) (a) Roesky, H. W.; Grimm, L. F. *Angew. Chem., Int. Ed. Engl.* **1972**, *11*, 642. (b) Roesky, H. W.; Janssen, E. *Chem. Ber.* **1975**, *108*, 2531.
- (39) Lidy, W.; Sundermeyer, W.; Verbeek, W. *Z. Anorg. Allg. Chem.* **1974**, *406*, 228.
- (40) Steudel, R.; Rose, F.; Reinhardt, R.; Bradaczek, H. *Z. Naturforsch., B: Anorg. Chem., Org. Chem.* **1977**, *32B*, 488.
- (41) (a) Gard, G. L.; Shreeve, J. M. *J. Am. Chem. Soc.* **1982**, *104*, 5566. (b) Emel'us, H. J.; Forder, R. A.; Poulet, R. J.; Sheldrick, G. M. *J. Chem. Soc. D* **1970**, 1483. (c) Mews, R. J. *Fluorine Chem.* **1981**, *18*, 155.
- (42) Banister, A. J.; Hey, R. G.; MacLean, G. K.; Passmore, J. *Inorg. Chem.* **1982**, *21*, 1679.
- (43) (a) Bojes, J.; Boorman, P. M.; Chivers, T. *Inorg. Nucl. Chem. Lett.* **1976**, *12*, 551. (b) Bojes, J.; Chivers, T. *J. Chem. Soc., Chem. Commun.* **1977**, 453. (c) Bojes, J.; Chivers, T. *Inorg. Chem.* **1978**, *17*, 318.
- (44) Bojes, J.; Chivers, T.; Drummond, I.; MacLean, G. K. *Inorg. Chem.* **1978**, *17*, 3668.
- (45) (a) Holt, E. M.; Holt, S. L. *Chem. Commun.* **1970**, 1704. (b) Holt, S. L. *J. Chem. Soc., Dalton Trans.* **1974**, 1990.

- (46) Ruppert, I.; Bastian, V.; Appel, R. *Chem. Ber.* **1974**, *107*, 3426.
- (47) Holt, E. M.; Holt, S. L.; Watson, K. J. *J. Chem. Soc., Dalton Trans.* **1977**, 514.
- (48) Chivers, T.; Cordes, A. W.; Oakley, R. T.; Swepston, P. N. *Inorg. Chem.* **1981**, *20*, 2376.
- (49) (a) Bojes, J.; Chivers, T.; MacLean, G.; Oakley, R. T.; Cordes, A. W. *Can. J. Chem.* **1979**, *57*, 3171. (b) Bojes, J.; Chivers, T.; Cordes, A. W.; MacLean, G.; Oakley, R. T. *Inorg. Chem.* **1981**, *20*, 16.
- (50) (a) Burford, N.; Chivers, T.; Oakley, R. T.; Cordes, A. W.; Swepston, P. N. *J. Chem. Soc., Chem. Commun.* **1980**, 1204. (b) Burford, N.; Chivers, T.; Cordes, A. W.; Laidlaw, W. G.; Noble, M. C.; Oakley, R. T.; Swepston, P. N. *J. Am. Chem. Soc.* **1982**, *104*, 1282.
- (51) Burford, N.; Chivers, T.; Richardson, J. F. *Inorg. Chem.* **1983**, *22*, 1482.
- (52) Burford, N.; Chivers, T.; Rao, M. N. S.; Richardson, J. F. *Adv. Chem. Ser.* **1983**, *232*, 81.
- (53) (a) Chivers, T.; Rao, M. N. S.; Richardson, J. F. *J. Chem. Soc., Chem. Commun.* **1982**, 982. (b) Chivers, T.; Rao, M. N. S. *Inorg. Chem.* **1984**, *23*, 3605.
- (54) Thewalt, U.; Müller, B. Z. *Anorg. Allg. Chem.* **1980**, *462*, 214.
- (55) Thewalt, U.; Burger, M. *Angew. Chem., Int. Ed. Engl.* **1982**, *21*, 634.
- (56) Thewalt, U.; Holl, K. Z. *Naturforsch.* **1984**, *39b*, 145.
- (57) Thewalt, U.; Berhalter, K.; Müller, P. *Acta Crystallogr., Sect. B: Struct. Crystallogr. Cryst. Chem.* **1982**, *B38*, 1280.
- (58) Woollins, J. D.; Grinter, R.; Johnson, M. K.; Thomson, A. J. *J. Chem. Soc., Dalton Trans.* **1980**, 1910 and references cited therein.
- (59) Thewalt, U. Z. *Naturforsch.* **1982**, *37b*, 276.
- (60) (a) Edelmann, F. J. *Organomet. Chem.* **1982**, *228*, C47. (b) Drews, R.; Behrens, U.; Edelmann, F., private communication. (c) Martan, H.; Weiss, J. Z. *Anorg. Allg. Chem.* **1984**, *514*, 107.
- (61) Herberhold, M.; Haumaier, L.; Schubert, U. *Inorg. Chim. Acta* **1981**, *49*, 21.
- (62) (a) Roesky, H. W.; Anhaus, J.; Schmidt, H. G.; Sheldrick, G. M.; Noltemeyer, M. *J. Chem. Soc., Dalton Trans.* **1983**, 1207. (b) Hanich, J.; Krestel, M.; Müller, U.; Dehnicke, K.; Rehder, D. Z. *Naturforsch., B: Anorg. Chem., Org. Chem.* **1984**, *39B*, 1686. (c) Anhaus, J.; Siddiqi, Z. A.; Schimkowiak, J.; Roesky, H. W. Z. *Naturforsch.* **1984**, *39b*, 1722. (d) Kynast, U.; Conrad, E.; Müller, U.; Dehnicke, K. Z. *Naturforsch., B: Anorg. Chem., Org. Chem.* **1984**, *39B*, 1680. (e) Anhaus, J.; Jones, P. G.; Noltemeyer, M.; Pinkert, W.; Roesky, H. W.; Sheldrick, G. M. *Inorg. Chim. Acta* **1985**, *97*, L7.
- (63) Endres, H.; Galantai, E. *Angew. Chem., Int. Ed. Engl.* **1980**, *19*, 653.
- (64) Marcellus, C. G.; Oakley, R. T.; Cordes, A. W.; Pennington, W. T. *J. Chem. Soc., Chem. Commun.* **1983**, 1451.
- (65) (a) Tashiro, M.; Mataka, S. *Heterocycles* **1976**, *4*, 1243. (b) Mataka, S.; Takahashi, K.; Yamada, Y.; Tashiro, M. *J. Heterocycl. Chem.* **1979**, *16*, 1009. (c) Daley, S. T. A. K.; Rees, C. W.; Williams, D. J. *J. Chem. Soc., Chem. Commun.* **1984**, 55. (d) Daley, S. T. A. K.; Rees, C. W.; Williams, D. J. *J. Chem. Soc., Chem. Commun.* **1984**, 57.
- (66) Small, R. W. H.; Banister, A. J.; Hauptman, Z. V. *J. Chem. Soc., Dalton Trans.* **1984**, 1377.
- (67) Roesky, H. W.; Janssen, E. *Chem.-Ztg.* **1974**, *98*, 260.
- (68) Montenarh, M.; Appel, R. Z. *Naturforsch., B: Anorg. Chem., Org. Chem.* **1976**, *31B*, 902.
- (69) Roesky, H. W.; Lucas, J.; Noltemeyer, M.; Sheldrick, G. M. *Chem. Ber.* **1984**, *117*, 1583.
- (70) Roesky, H. W.; Schäfer, P.; Noltemeyer, M.; Sheldrick, G. M. Z. *Naturforsch. B: Anorg. Chem., Org. Chem.* **1983**, *38B*, 347.
- (71) Patton, L.; Jolly, W. L. *Inorg. Chem.* **1970**, *9*, 1079.
- (72) Greenhough, T. J.; Kolthammer, B. W. S.; Legzdins, P.; Trotter, J. *Inorg. Chem.* **1979**, *18*, 3548.
- (73) (a) Banister, A. J.; Dainty, P. J. *J. Chem. Soc., Dalton Trans.* **1972**, 2658. (b) Banister, A. J.; Clarke, H. G. *J. Chem. Soc., Dalton Trans.* **1972**, 2661. (c) Banister, A. J.; Clarke, H. G. *Inorg. Synth.* **1977**, *17*, 188.
- (74) Mews, R. *Angew. Chem., Int. Ed. Engl.* **1976**, *15*, 691.
- (75) Mews, R.; Wagner, D. L.; Glemser, O. Z. *Anorg. Allg. Chem.* **1975**, *412*, 148.
- (76) Gillespie, R. J.; Sawyer, J. F.; Slim, D. R.; Tyrer, J. D. *Inorg. Chem.* **1982**, *21*, 1296.
- (77) Roesky, H. W.; Müller, T. *Chem. Ber.* **1978**, *111*, 2969.
- (78) (a) Alange, G. G.; Banister, A. J.; Bell, B.; Millen, P. W. J. *Chem. Soc., Perkin Trans. 1* **1979**, 1192. (c) Banister, A. J.; Smith, N. R. M.; Hey, R. G. *J. Chem. Soc., Perkin Trans. 1* **1983**, 1181.
- (79) Höfs, H.-U.; Hartmann, G.; Mews, R.; Sheldrick, G. M. Z. *Naturforsch., B: Anorg. Chem., Org. Chem.* **1984**, *39B*, 1389.
- (80) Chivers, T.; Richardson, J. F.; Smith, N. R. M. *Mol. Cryst. Liq. Cryst.* **1985**, *125*, 447.
- (81) Fritz, H. P.; Bruchhaus, R. *Chem. Ber.* **1984**, *117*, 1920.
- (82) (a) Chivers, T.; Fielding, L. *J. Chem. Soc., Chem. Commun.* **1978**, 212. (b) Chivers, T.; Fielding, L.; Laidlaw, W. G.; Trsic, M. *Inorg. Chem.* **1979**, *18*, 3379.
- (83) Scherer, O. J.; Wolmershäuser, G. *Chem. Ber.* **1977**, *110*, 3241.
- (84) (a) Roesky, H. W.; Rao, M. N. S.; Nakajima, T.; Sheldrick, W. S. *Chem. Ber.* **1979**, *112*, 3531. (b) Sheldrick, W. S.; Rao, M. N. S.; Roesky, H. W. *Inorg. Chem.* **1980**, *19*, 538.
- (85) (a) Roesky, H. W.; Wiezer, H. *Angew. Chem., Int. Ed. Engl.* **1973**, *12*, 674. (b) Roesky, H. W.; Wiezer, H. *Chem. Ber.* **1974**, *107*, 3186. (c) Roesky, H. W. Z. *Naturforsch., B: Anorg. Chem., Org. Chem.* **1976**, *31B*, 680.
- (86) Scherer, O. J.; Wies, R. *Angew. Chem., Int. Ed. Engl.* **1971**, *10*, 812.
- (87) (a) Alcock, N. W.; Holt, E. M.; Kuyper, J.; Mayerle, J. J.; Street, G. B. *Inorg. Chem.* **1979**, *18*, 2237. (b) Gieren, A.; Betz, H.; Hübner, T.; Lamm, V.; Herberhold, M.; Guldner, K. Z. *Anorg. Allg. Chem.* **1984**, *513*, 160. The anion NSN^{2-} has been used instead of $\text{Me}_3\text{SiNSNSiMe}_3$ for the preparation of the *t*-Bu derivative.
- (88) Roesky, H. W.; Wiezer, H. *Angew. Chem., Int. Ed. Engl.* **1974**, *13*, 146.
- (89) Roesky, H. W.; Witt, M.; Krebs, B.; Henkel, G.; Korte, H.-J. *Chem. Ber.* **1981**, *114*, 201.
- (90) Burford, N.; Chivers, T.; Oakley, R. T.; Oswald, T. *Can. J. Chem.* **1984**, *62*, 712.
- (91) Pohl, S.; Petersen, O.; Roesky, H. W. *Chem. Ber.* **1979**, *112*, 1545.
- (92) (a) Appel, R.; Ruppert, I.; Milker, R.; Bastian, V. *Chem. Ber.* **1974**, *107*, 380. (b) Weiss, J.; Ruppert, I.; Appel, R. Z. *Anorg. Chem. Allg. Chem.* **1974**, *406*, 329.
- (93) Scherer, O. J.; Wolmershäuser, G. *Angew. Chem., Int. Ed. Engl.* **1972**, *11*, 642.
- (94) (a) Roesky, H. W.; Grimm, L. F. *Angew. Chem., Int. Ed. Engl.* **1972**, *11*, 642. (b) Steudel, R.; Rose, F.; Reinhardt, R.; Bradaczek, H. Z. *Naturforsch., B: Anorg. Chem., Org. Chem.* **1977**, *32B*, 488.
- (95) Roesky, H. W.; Böwing, W. G.; Rayment, I.; Shearer, H. M. M. *J. Chem. Soc., Chem. Commun.* **1975**, 735.
- (96) (a) Koenig, H.; Oakley, R. T. *J. Chem. Soc., Chem. Commun.* **1983**, 73. (b) Codding, P. W.; Koenig, H.; Oakley, R. T. *Can. J. Chem.* **1983**, *61*, 1562.
- (97) Wolmershäuser, G.; Schnauber, M.; Wilhelm, T. *J. Chem. Soc., Chem. Commun.* **1984**, 573.
- (98) (a) Roesky, H. W.; Wiezer, H. *Angew. Chem., Int. Ed. Engl.* **1975**, *14*, 258. (b) Roesky, H. W.; Holtschneider, G.; Wiezer, H.; Krebs, B. *Chem. Ber.* **1976**, *109*, 1358.
- (99) Roesky, H. W.; Wehner, E. *Angew. Chem., Int. Ed. Engl.* **1975**, *14*, 498.
- (100) Heitmann, A.; Edelmann, F. Z. *Naturforsch., B: Anorg. Chem., Org. Chem.* **1983**, *38B*, 521.
- (101) MacLean, G. K.; Passmore, J.; Schriver, M. J.; White, P. S.; Bethell, D.; Pilkington, R. S.; Sutcliffe, L. H. *J. Chem. Soc., Chem. Commun.* **1983**, 807.
- (102) MacLean, G. K.; Passmore, J.; White, P. S. *J. Chem. Soc., Dalton Trans.* **1984**, 211.
- (103) Roesky, H. W.; Graf, C.; Rao, M. N. S. *Chem. Ber.* **1980**, *113*, 3815.
- (104) (a) Bojes, J.; Chivers, T. *J. Chem. Soc., Chem. Commun.* **1980**, 1023. (b) Bojes, J.; Chivers, T.; Laidlaw, W. G.; Trsic, M. *J. Am. Chem. Soc.* **1982**, *104*, 4837.
- (105) (a) Chivers, T.; Oakley, R. T. *J. Chem. Soc., Chem. Commun.* **1979**, 752. (b) Chivers, T.; Laidlaw, W. G.; Oakley, R. T.; Trsic, M. *J. Am. Chem. Soc.* **1980**, *102*, 5773.
- (106) Bojes, J.; Chivers, T.; Codding, P. W. *J. Chem. Soc., Chem. Commun.* **1981**, 1171.
- (107) Chivers, T.; Rao, M. N. S. *Can. J. Chem.* **1983**, *61*, 1957.
- (108) (a) Chivers, T.; Proctor, J. *J. Chem. Soc., Chem. Commun.* **1978**, 642. (b) Chivers, T.; Proctor, J. *Can. J. Chem.* **1979**, *57*, 1286.
- (109) (a) Chivers, T.; Oakley, R. T.; Cordes, A. W.; Pennington, W. T. *J. Chem. Soc., Chem. Commun.* **1981**, 1214. (b) Chivers, T.; Cordes, A. W.; Oakley, R. T.; Pennington, W. T. *Inorg. Chem.* **1983**, *22*, 2429.
- (110) Roesky, H. W.; Witt, M.; Krebs, B.; Korte, H. J. *Angew. Chem., Int. Ed. Engl.* **1979**, *18*, 415.
- (111) (a) Witt, M.; Roesky, H. W. Z. *Anorg. Allg. Chem.* **1984**, *515*, 51. (b) Roesky, H. W.; Witt, M.; Diehl, M.; Bats, J. W.; Fuess, H. *Chem. Ber.* **1979**, *112*, 1372.
- (112) Roesky, H. W.; Witt, M.; Clegg, W.; Isenberg, W.; Noltemeyer, M.; Sheldrick, G. M. *Angew. Chem., Int. Ed. Engl.* **1980**, *19*, 943.
- (113) (a) Roesky, H. W.; Clegg, W.; Schimkowiak, J.; Schmidt, M.; Witt, M.; Sheldrick, G. M. *J. Chem. Soc., Dalton Trans.* **1982**, 2117. (b) Roesky, H. W.; Witt, M.; Schimkowiak, J.; Schmidt, M.; Noltemeyer, M.; Sheldrick, G. M. *Angew. Chem., Int. Ed. Engl.* **1982**, *21*, 538.
- (114) Roesky, H. W.; Anhaus, J.; Sheldrick, W. S. *Inorg. Chem.* **1984**, *23*, 75.
- (115) Chivers, T.; Hojo, M. *Inorg. Chem.* **1984**, *23*, 4088.
- (116) Banister, A. J. *Nature (London)* **1972**, *237*, 92.

- (117) Gimarc, B. M.; Trinajstić, N. *Pure Appl. Chem.* **1980**, *52*, 1443.
- (118) Gleiter, R. *Angew. Chem., Int. Ed. Engl.* **1981**, *20*, 444.
- (119) Laidlaw, W. G.; Trsic, M. "Proceeding of the Applied Quantum Chemistry Symposium"; Smith, V. H., Ed.; D. Reidel: Dordrecht, Netherlands, in press.
- (120) (a) Chivers, T. *Acc. Chem. Res.* **1984**, *17*, 2336. (b) Chivers, T.; Oakley, R. T. *Top. Curr. Chem.* **1982**, *102*, 117.
- (121) (a) Andreasen, O.; Hazell, A. C.; Hazell, R. G. *Acta Crystallogr. Sect. B: Struct. Crystallogr. Cryst. Chem.* **1977**, *B33*, 1109. (b) Höfs, H. U.; Mews, R.; Clegg, W.; Noltemeyer, M.; Schmidt, M.; Sheldrick, G. M. *Chem. Ber.* **1983**, *116*, 416.
- (122) (a) Shearer, H. M. M.; Banister, A. J.; Halfpenny, J.; Whitehead, G. *Polyhedron* **1983**, *2*, 149. (b) Isenberg, W.; Homsy, N. K.; Anhaus, J.; Roesky, H. W.; Sheldrick, G. M. *Z. Naturforsch., B: Anorg. Chem., Org. Chem.* **1983**, *38B*, 808.
- (123) (a) Roesky, H. W.; Holtschneider, G.; Wiezer, H.; Krebs, B. *Chem. Ber.* **1976**, *109*, 1358. (b) Roesky, H. W.; Witt, M.; Clegg, W.; Isenberg, W.; Noltemeyer, M.; Sheldrick, G. M. *Angew. Chem., Int. Ed. Engl.* **1980**, *19*, 943. (c) Rayment, I.; Shearer, H. M. M.; Roesky, H. W. *J. Chem. Soc., Dalton Trans.* **1982**, 883.
- (124) Roesky, H. W.; Wehner, E.; Zehnder, E.-J.; Deiseroth, H.-J.; Simon, A. *Chem. Ber.* **1978**, *111*, 1670.
- (125) Vegas, A.; Pérez-Salazar, A.; Banister, A. J.; Hey, R. G. *J. Chem. Soc., Dalton Trans.* **1980**, 1812.
- (126) Cordes, A. W.; Hayes, P. J.; Josephy, P. D.; Koenig, H.; Oakley, R. T.; Pennington, W. T. *J. Chem. Soc., Chem. Commun.* **1984**, 1021.
- (127) (a) Appel, R.; Halstenberg, M. *Angew. Chem., Int. Ed. Engl.* **1976**, *15*, 695. (b) Weiss, J. *Acta Crystallogr. Sect. B: Struct. Crystallogr. Cryst. Chem.* **1977**, *B33*, 2272.
- (128) (a) Bojes, J.; Chivers, T. *J. Chem. Soc., Chem. Commun.* **1978**, 391. (b) Bojes, J.; Chivers, T.; Laidlaw, W. G.; Trsic, M. *J. Am. Chem. Soc.* **1979**, *101*, 4517. (c) Martan, H.; Weiss, J. *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.* **1983**, *C39*, 959.
- (129) (a) Chivers, T.; Coddling, P. W.; Oakley, R. T. *J. Chem. Soc., Chem. Commun.* **1981**, 584. (b) Small, R. W. H.; Banister, A. J.; Hauptman, Z. V. *J. Chem. Soc., Dalton Trans.* **1981**, 2188.
- (130) (a) Moss, G.; Guru Row, T. N.; Coppens, P. *Inorg. Chem.* **1980**, *19*, 2396. (b) Mayerle, J. J.; Wolmershäuser, G.; Street, G. B. *Inorg. Chem.* **1979**, *18*, 1161. (c) Ziegler, M. L.; Weidenhammer, K. *Z. Anorg. Allg. Chem.* **1977**, *434*, 152 and references cited therein.
- (131) Jones, P. G.; Pinkert, W.; Sheldrick, G. M. *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.* **1983**, *C39*, 827.
- (132) (a) Gillespie, R. J.; Sawyer, J. F.; Slim, D. R.; Tyrer, J. D. *Inorg. Chem.* **1982**, *21*, 1296. (b) Banister, A. J.; Durrant, J. A.; Rayment, I.; Shearer, H. M. M. *J. Chem. Soc., Dalton Trans.* **1976**, 928. (c) Roesky, H. W.; Böwing, W. G.; Rayment, I.; Shearer, H. M. M. *J. Chem. Soc., Chem. Commun.* **1975**, 735. (d) Hazell, A. C.; Hazell, R. G. *Acta. Chem. Scand.* **1972**, *26*, 1987.
- (133) McKennis, J. S.; Brenner, L.; Schweiger, J. R.; Pettit, R. *J. Chem. Soc., Chem. Commun.* **1972**, 365.
- (134) (a) Findlay, R. H.; Palmer, M. H.; Downs, A. J.; Egddell, R. G.; Evans, R. *Inorg. Chem.* **1980**, *19*, 1307, and references cited therein. (b) Skrzeneck, F. L.; Harcourt, R. D. *J. Am. Chem. Soc.* **1984**, *106*, 3934. (c) Fujimoto, H.; Yokoyama, T. *Bull. Chem. Soc. Jpn.* **1980**, *53*, 800.
- (135) (a) Frost, D. C.; Le Geyt, M. R.; Paddock, N. L.; Westwood, N. P. C. *J. Chem. Soc., Chem. Commun.* **1977**, 217. (b) Sharma, J.; Downs, D. S.; Iqbal, Z.; Owens, F. J. *J. Chem. Phys.* **1977**, *67*, 3045.
- (136) Palmer, M. H. *Z. Naturforsch., A: Phys., Phys. Chem., Kosmophys.* **1984**, *39A*, 102.
- (137) Gleiter, R.; Bartetzko, R.; Hofmann, P. *Z. Naturforsch., B: Anorg. Chem., Org. Chem.* **1980**, *35B*, 1166.
- (138) Roesky, H. W.; Dietl, M. *Chem. Ber.* **1973**, *106*, 3101.
- (139) Tang, T. H.; Bader, R. F. W. *Abstr. Pap. Int. Chem. Congr. Pacific Basin Soc.* **1984**, 05E50.
- (140) (a) Chivers, T.; Laidlaw, W. G.; Oakley, R. T.; Trsic, M. *Inorg. Chim. Acta* **1981**, *53*, L189. (b) Smith, V. H., Jr.; Sabin, J. R.; Broclawik, E.; Mrozek, J. *Inorg. Chim. Acta* **1983**, *77*, L101. (c) Nguyen, M.-T.; Ha, T.-K. *J. Mol. Struct.* **1983**, *105*, 129.
- (141) Burford, N.; Chivers, T.; Hojo, M.; Laidlaw, W. G.; Richardson, J. F. *Inorg. Chem.* **1985**, *24*, 709. In the general formula $P_2S_3-N_3$ the two exocyclic substituents on each phosphorus atom have been omitted.
- (142) Adkins, R. R.; Turner, A. G. *J. Am. Chem. Soc.* **1978**, *100*, 1383.
- (143) Zhu, J.-K.; Gimarc, B. M. *Inorg. Chem.* **1983**, *22*, 1996.
- (144) (a) Palmer, M. H.; Wheeler, J. R.; Findlay, R. H.; Westwood, N. P. C.; Lau, W. M. *J. Mol. Struct.* **1981**, *86*, 193. (b) Palmer, M. H.; Lau, W. M.; Westwood, N. P. C. *Z. Naturforsch., A: Phys., Phys. Chem., Kosmophys.* **1982**, *37a*, 1061. (c) Palmer, M. H. *Z. Naturforsch., A: Phys., Phys. Chem., Kosmophys.* **1983**, *38A*, 74.
- (145) Waluk, J. W.; Michl, J. *Inorg. Chem.* **1982**, *21*, 556.
- (146) Trsic, M.; Laidlaw, W. G. *Inorg. Chem.* **1984**, *23*, 1981.
- (147) Turner, A. G. *Inorg. Chim. Acta* **1982**, *65*, L201.
- (148) Trsic, M.; Laidlaw, W. G.; Oakley, R. T. *Can. J. Chem.* **1982**, *60*, 2281.
- (149) Müller, U.; Conradi, E.; Demant, U.; Dehnicke, K. *Angew. Chem., Int. Ed. Engl.* **1984**, *23*, 237.
- (150) Cordes, A. W.; Marcellus, C. G.; Noble, M. C.; Oakley, R. T.; Pennington, W. T. *J. Am. Chem. Soc.* **1983**, *105*, 6008.
- (151) Roesky, H. W.; Witt, M.; Krebs, B.; Korte, H.-J. *Angew. Chem., Int. Ed. Engl.* **1979**, *18*, 415.
- (152) Boutique, J. P.; Riga, J.; Verbist, J. J.; Delhalle, J.; Fripiat, J. G.; Haddon, R. C.; Kaplan, M. L. *J. Am. Chem. Soc.* **1984**, *106*, 312.
- (153) Millefiori, S.; Millefiori, A.; Granozzi, G. *Inorg. Chim. Acta* **1984**, *90*, L55.
- (154) Oakley, R. T. *Can. J. Chem.* **1984**, *62*, 2763.
- (155) Höfs, H.-U.; Hartmann, G.; Mews, R.; Sheldrick, G. M. *Angew. Chem., Int. Ed. Engl.* **1984**, *23*, 988.
- (156) Banister, A. J.; Durrant, J. A. *J. Chem. Res. Miniprint* **1978**, 1912. Banister, A. J.; Durrant, J. A. *J. Chem. Res. Miniprint* **1978**, 1931.
- (157) (a) Bartetzko, R.; Gleiter, R. *Z. Naturforsch., B: Anorg. Chem., Org. Chem.* **1981**, *36B*, 956. (b) Bartetzko, R.; Gleiter, R. *Inorg. Chem.* **1978**, *17*, 995.
- (158) Delucia, M. L.; Coppens, P. *Inorg. Chem.* **1978**, *17*, 2336.
- (159) Burford, N.; Chivers, T.; Coddling, P. W.; Oakley, R. T. *Inorg. Chem.* **1982**, *21*, 982.
- (160) (a) Chivers, T.; Rao, M. N. S.; Richardson, J. F. *J. Chem. Soc., Chem. Commun.* **1983**, 186. (b) Burford, N.; Chivers, T.; Rao, M. N. S.; Richardson, J. F. *Inorg. Chem.* **1984**, *23*, 1946.
- (161) Zak, Z. *Acta Crystallogr., Sect. B: Struct. Sci.* **1981**, *B37*, 23.
- (162) (a) Roesky, H. W.; Rao, M. N. S.; Graf, C.; Gieren, A.; Hädicke, E. *Angew. Chem., Int. Ed. Engl.* **1981**, *20*, 592. (b) Roesky, H. W.; Pelz, C.; Gieren, A.; Hädicke, E. *Z. Naturforsch., B: Anorg. Chem., Org. Chem.* **1981**, *36B*, 1437.
- (163) Roesky, H. W.; Rao, M. N. S.; Nakajima, T.; Sheldrick, W. S. *Chem. Ber.* **1979**, *112*, 3531.
- (164) Flues, W.; Scherer, O. J.; Weiss, J.; Wolmershäuser, G. *Angew. Chem., Int. Ed. Engl.* **1976**, *15*, 379.
- (165) (a) Isenberg, W.; Mews, R. *Z. Naturforsch., B: Anorg. Chem., Org. Chem.* **1982**, *37B*, 1388.
- (166) Isenberg, W.; Mews, R.; Sheldrick, G. M.; Bartetzko, R.; Gleiter, R. *Z. Naturforsch., B: Anorg. Chem., Org. Chem.* **1983**, *38B*, 1563.
- (167) Banister, A. J. *Nature (London), Phys. Sci.* **1972**, *239*, 69.
- (168) (a) Harcourt, R. D.; Hügel, R. M. *J. Inorg. Nucl. Chem.* **1981**, *43*, 239. (b) Harcourt, R. D. *J. Inorg. Nucl. Chem.* **1977**, *39*, 237.
- (169) (a) Burdett, J. K. *Acc. Chem. Res.* **1982**, *15*, 34. (b) Gimarc, B. M. *J. Am. Chem. Soc.* **1983**, *105*, 1979.
- (170) Gleiter, R. *J. Chem. Soc.* **1970**, 3174.
- (171) Millefiori, S.; Millefiori, S. *J. Chem. Res., Synop.* **1980**, 244.
- (172) Dolenko, G. N.; Zibarev, A. V.; Pykhov, V. Z. *Russ. J. Inorg. Chem. (Engl. Transl.)* **1981**, *26*, 1476.
- (173) Zhorilova, L.; Gebauer, P. *Z. Anorg. Allg. Chem.* **1979**, *448*, 5.
- (174) Mews, R. *J. Fluorine Chem.* **1981**, *18*, 155.
- (175) Bartetzko, R.; Gleiter, R. *Chem. Ber.* **1980**, *113*, 1138.
- (176) Bojes, J.; Chivers, T.; Oakley, R. T. *Inorg. Synth.*, submitted for publication.
- (177) Trsic, M.; Wagstaff, K.; Laidlaw, W. G. *Int. J. Quantum Chem.* **1982**, *22*, 903.
- (178) Brant, P.; Weber, D. C.; Ewing, C. T.; Carter, F. L.; Hashmall, J. A. *Inorg. Chem.* **1980**, *19*, 2829.
- (179) Andreocci, M. V.; Bossa, M.; Di Castro, V.; Furlani, C.; Maltogno, G.; Roesky, H. W. *Gazz. Chim. Ital.* **1979**, *109*, 1.
- (180) Andreocci, M. V.; Bossa, M.; Di Castro, V.; Furlani, C.; Maltogno, G.; Roesky, H. W. *Z. Phys. Chem. (Munich)* **1979**, *118*, 137.
- (181) Salaneck, W. R.; Lin, J. W.-p.; Paton, A.; Duke, C. B.; Ceasar, G. B. *Phys. Rev. B: Solid State* **1976**, *13*, 4517.
- (182) Trsic, M.; Laidlaw, W. G. *Int. J. Quantum Chem., Quantum Chem. Symp.* **1983**, *17*, 367.
- (183) Salahub, D. R.; Messmer, R. P. *J. Chem. Phys.* **1976**, *64*, 2039.
- (184) Ziegler, T.; Rauk, A.; Baerends, E. J. *Theor. Chim. Acta* **1977**, *43*, 261.
- (185) (a) Zahradnik, R.; Banister, A. J.; Clarke, H. G. *Collect. Czech. Chem. Commun.* **1973**, *38*, 998. (b) Banister, A. J.; Clarke, H. G. *Inorg. Synth.* **1977**, *17*, 188.
- (186) Waluk, J. W.; Michl, J. *Inorg. Chem.* **1981**, *20*, 963.
- (187) Platt, J. R. *J. Chem. Phys.* **1949**, *17*, 484.
- (188) Michl, J. *J. Am. Chem. Soc.* **1978**, *100*, 6801.
- (189) Waluk, J. W.; Chivers, T.; Oakley, R. T.; Michl, J. *Inorg. Chem.* **1982**, *21*, 832.
- (190) Braterman, P. S. *J. Chem. Soc.* **1965**, 2297.
- (191) Nelson, J.; Heal, H. G. *J. Chem. Soc. A* **1971**, 136.

- (192) Mason, J. *J. Chem. Soc. A* **1969**, 1567.
 (193) Mason, J. *Chem. Rev.* **1981**, *81*, 205.
 (194) Logan, N.; Jolly, W. L. *Inorg. Chem.* **1965**, *4*, 1508.
 (195) Roesky, H. W.; Kuhn, M.; Bats, J. W. *Chem. Ber.* **1982**, *115*, 3025.
 (196) Bojes, J.; Chivers, T.; Laidlaw, W. G.; Trsic, M. *J. Am. Chem. Soc.* **1982**, *104*, 4837.
 (197) Wasylshen, R.; Connor, C.; Friedrich, J. *Can. J. Chem.* **1984**, *62*, 981.
 (198) Fairhurst, S. A.; Preston, K. F.; Sutcliffe, L. H. *Can. J. Chem.* **1984**, *62*, 1124.
 (199) Chapman, D.; Massey, A. G. *Trans. Faraday Soc.* **1962**, *58*, 1291.
 (200) Meinzer, R. A.; Pratt, D. W.; Myers, R. J. *J. Am. Chem. Soc.* **1969**, *91*, 6623.
 (201) Williford, J. D.; Van Reet, R. E.; Eastman, M. P.; Prater, K. B. *J. Electrochem. Soc.* **1973**, *120*, 1498.
 (202) Brinkman, M. R.; Sutcliffe, L. H. *J. Magn. Reson.* **1977**, *28*, 263.
 (203) Fairhurst, S. A.; McIlwaine, W. R.; Sutcliffe, L. H. *J. Magn. Reson.* **1979**, *35*, 121.
 (204) Rolfe, S.; Griller, D.; Ingold, K. U.; Sutcliffe, L. H. *J. Org. Chem.* **1979**, *44*, 3515.
 (205) Fairhurst, S. A.; Sutcliffe, L. H. *J. Chem. Soc., Faraday Trans. 1* **1980**, *76*, 1490.
 (206) Fairhurst, S. A.; Pilkington, R. S.; Sutcliffe, L. H. *J. Chem. Soc., Faraday Trans. 1* **1983**, *79*, 439.
 (207) Fairhurst, S. A.; Pilkington, R. S.; Sutcliffe, L. H. *J. Chem. Soc., Faraday Trans. 1* **1983**, *79*, 925.
 (208) Mayer, R.; Domshe, G.; Bleisch, S.; Barth, A.; Stasko, A. *Z. Chem.* **1981**, *21*, 264.
 (209) Bragin, J.; Evans, M. V. *J. Chem. Phys.* **1969**, *51*, 268.
 (210) Iqbal, Z.; Downs, D. S. *Solid State Commun.* **1976**, *20*, 1147.
 (211) Karpfen, A.; Schuster, P.; Petkov, J.; Lischka, H. *J. Chem. Phys.* **1977**, *66*, 5167.
 (212) Warn, J. R. W.; Chapman, D. *Spectrochim. Acta.* **1966**, *22*, 1371.
 (213) Steudel, R. *Z. Naturforsch., A: Phys., Phys. Chem., Kosmophys.* **1981**, *36A*, 850.
 (214) Kent, J. P. Ph.D. Thesis, McMaster University, Hamilton, Ontario, Canada, 1984.
 (215) Turowski, A.; Appel, R.; Sawodny, W.; Molt, K. *J. Mol. Struct.* **1978**, *48*, 313.
 (216) Chivers, T.; Lau, C.; Oakley, R. T., unpublished results.
 (217) Steudel, R. *Angew. Chem., Int. Ed. Engl.* **1975**, *14*, 655.
 (218) Patton, R. L.; Jolly, W. L. *Inorg. Chem.* **1969**, *8*, 1392.
 (219) Patton, R. L.; Jolly, W. L. *Inorg. Chem.* **1969**, *8*, 1389.
 (220) Patton, R. L.; Raymond, K. N. *Inorg. Chem.* **1969**, *8*, 2426.
 (221) Roesky, H. W.; Anhaus, J. *Chem. Ber.* **1982**, *115*, 3682.
 (222) Alange, G. G.; Banister, A. J. *J. Inorg. Nucl. Chem.* **1978**, *40*, 203 and references cited therein.
 (223) Drew, M. G. B.; Templeton, D. H.; Zalkin, A. *Inorg. Chem.* **1967**, *6*, 1607.
 (224) Gillespie, R. J.; Kent, J. P.; Sawyer, J. F. *Acta Crystallogr. Sect. B: Struct. Sci.* **1980**, *B36*, 655.
 (225) Neubauer, D.; Weiss, J. *Z. Anorg. Allg. Chem.* **1960**, *303*, 28.
 (226) (a) Gieren, A.; Dederer, B.; Roesky, H. W.; Amin, N.; Petersen, O. *Z. Anorg. Allg. Chem.* **1978**, *440*, 119. (b) Gieren, A.; Hahn, C.; Dederer, B.; Roesky, H. W.; Amin, N. *Z. Anorg. Allg. Chem.* **1978**, *447*, 179.
 (227) Thewalt, U.; Albrecht, G. *Z. Naturforsch., B: Anorg. Chem., Org. Chem.* **1982**, *37B*, 1098.
 (228) Thewalt, U. *Z. Naturforsch., B: Anorg. Chem., Org. Chem.* **1980**, *35B*, 855.
 (229) Cordes, A. W.; Marcellus, C. G.; Noble, M. C.; Oakley, R. T.; Pennington, W. T. *J. Am. Chem. Soc.* **1983**, *105*, 6008.
 (230) (a) Thewalt, U. *Angew. Chem., Int. Ed. Engl.* **1976**, *15*, 365. (b) Thewalt, U.; Müller, B. *Z. Naturforsch. B: Anorg. Chem., Org. Chem.* **1982**, *37B*, 828.
 (231) Das, M. K.; Bibber, J. W.; Zuckerman, J. J. *Inorg. Chem.* **1982**, *21*, 2864.
 (232) Roesky, H. W.; Thomas, M.; Schimkowiak, J.; Schmidt, M.; Noltemeyer, M.; Sheldrick, G. M. *J. Chem. Soc., Chem. Commun.* **1982**, 790.
 (233) Roesky, H. W.; Thomas, M.; Schmidt, H. G.; Clegg, W.; Noltemeyer, M.; Sheldrick, G. M. *J. Chem. Soc., Dalton Trans.* **1983**, 405.
 (234) Roesky, H. W.; Thomas, M.; Noltemeyer, M.; Sheldrick, G. M. *Angew. Chem., Int. Ed. Engl.* **1982**, *21*, 858.
 (235) Brinkman, M. R.; Allen, C. W. *J. Am. Chem. Soc.* **1972**, *94*, 1550.
 (236) Mock, W. L.; Mehrota, I. *J. Chem. Soc., Chem. Commun.* **1976**, 123.
 (237) Griffin, A. M.; Sheldrick, G. M. *Acta Crystallogr., Sect. B: Struct. Crystallogr. Cryst. Chem.* **1975**, *B31*, 895.
 (238) Ertl, G.; Weiss, J. *Z. Anorg. Allg. Chem.* **1976**, *420*, 155.
 (239) Yamabe, T.; Tanaka, K.; Tachibana, A.; Fukui, K.; Kato, H. *J. Phys. Chem.* **1979**, *83*, 767.
 (240) Liblong, S. W.; Oakley, R. T.; Cordes, A. W.; Noble, M. C. *Can. J. Chem.* **1983**, *61*, 2062.
 (241) Koenig, H.; Oakley, R. T.; Cordes, A. W.; Noble, M. C. *Can. J. Chem.* **1983**, *61*, 1185.
 (242) Cordes, A. W.; Koenig, H.; Noble, M. C.; Oakley, R. T. *Inorg. Chem.* **1983**, *22*, 3375.
 (243) Burford, N.; Chivers, T.; Rao, M. N. S.; Richardson, J. F. *Inorg. Chem.* **1984**, *23*, 1946.
 (244) Chivers, T.; Richardson, J. F.; Smith, N. R. M., submitted for publication in *Inorg. Chem.*
 (245) Roesky, H. W.; Emmert, R.; Gries, T. *Chem. Ber.* **1984**, *117*, 404.
 (246) Banister, A. J.; Smith, N. R. M.; Hey, R. G. *J. Chem. Soc., Perkin Trans. 1* **1983**, 1181.
 (247) Roesky, H. W.; Emmert, R.; Noltemeyer, M.; Sheldrick, G. M. *Z. Naturforsch., B: Anorg. Chem., Org. Chem.* **1984**, *39B*, 701.
 (248) Oakley, R. T.; Shevalier, M.; Cordes, A. W.; Pennington, W. L. *Abstr. Pap.—Am. Chem. Soc.* **1982**, *184th*, INOR 86.
 (249) Tanaka, K.; Yamabe, T.; Tachibana, A.; Kato, H.; Fukui, K. *J. Phys. Chem.* **1978**, *82*, 2121.
 (250) (a) Chivers, T.; Hojo, M. *Inorg. Chem.* **1984**, *23*, 1526. (b) Tweh, J. W.; Turner, A. G. *Inorg. Chim. Acta* **1981**, *48*, 173.
 (251) Hojo, M. *Bull. Chem. Soc. Jpn.* **1980**, *53*, 2856.
 (252) Marcellus, C. G.; Oakley, R. T.; Cordes, A. W.; Pennington, W. T. *Can. J. Chem.* **1984**, *62*, 1822.
 (253) Rodek, E.; Amin, N.; Roesky, H. W. *Z. Anorg. Allg. Chem.* **1979**, *457*, 123.
 (254) Roesky, H. W.; Bowing, W. G.; Rayment, I.; Shearer, H. M. *M. J. Chem. Soc., Chem. Commun.* **1975**, 735.
 (255) Banister, A. J.; Hauptman, Z. V. *J. Chem. Soc., Dalton Trans.* **1980**, 731.
 (256) Laidlaw, W. G.; Trsic, M. *Inorg. Chem.* **1981**, *20*, 1792.
 (257) (a) Nelson, J.; Heal, H. G. *Inorg. Nucl. Chem. Lett.* **1970**, *6*, 429. (b) Vincent, H.; Monteil, Y. *Synth. React. Inorg. Met.-Org. Chem.* **1978**, *8*, 51.
 (258) (a) Glemser, O.; Mews, R. *Angew. Chem., Int. Ed. Engl.* **1979**, *18*, 91. (b) Mews, R.; Wagner, D. L.; Glemser, O. *Z. Anorg. Allg. Chem.* **1975**, *412*, 148.
 (259) Maraschin, N. J.; Lagow, R. J. *J. Am. Chem. Soc.* **1972**, *94*, 8601.
 (260) Heal, H. G. *Adv. Inorg. Chem. Radiochem.* **1972**, *15*, 375.
 (261) Emeléus, H. J. *Endeavour* **1973**, *32*, 76.
 (262) Tanaka, K.; Yamabe, T.; Noda, K.; Fukui, K.; Kato, H. *J. Phys. Chem.* **1978**, *82*, 1453.
 (263) Zborilova, L.; Gebauer, P. *Z. Anorg. Allg. Chem.* **1981**, *483*, 44.
 (264) Chivers, T.; Rao, M. N. S.; Richardson, J. F. *Inorg. Chem.*, in press.
 (265) Chivers, T.; Lensink, C., unpublished results.
 (266) Burdett, J. K. *J. Am. Chem. Soc.* **1980**, *102*, 5458.
 (267) Love, P.; Kao, H. I.; Myer, G. H.; Labes, M. M. *J. Chem. Soc., Chem. Commun.* **1978**, 301.
 (268) Yamabe, T.; Tanaka, K.; Fukui, K.; Kato, H. *J. Phys. Chem.* **1977**, *81*, 727.
 (269) Witt, M. W. R.; Bailey, W. I.; Lagow, R. J. *J. Am. Chem. Soc.* **1983**, *105*, 1668.
 (270) Banister, A. J.; Hauptman, Z. V.; Kendrick, A. G. *J. Chem. Soc., Chem. Commun.* **1983**, 1017.
 (271) Fritz, H. P.; Bruchhaus, Z. *Naturforsch., B: Anorg. Chem., Org. Chem.* **1983**, *38B*, 1375.
 (272) Roesky, H. W. In "Sulfur in Organic and Inorganic Chemistry"; Senning, A., Ed.; Marcel Dekker: New York, 1982; Vol. 4, p 15.
 (273) Roesky, H. W.; Dietl, M. *Angew. Chem., Int. Ed. Engl.* **1973**, *12*, 424.
 (274) Roesky, H. W.; Wiezer, H. *Angew. Chem., Int. Ed. Engl.* **1973**, *12*, 674.
 (275) Roesky, H. W.; Wiezer, H. *Chem. Ber.* **1974**, *107*, 3186.
 (276) Roesky, H. W. *Z. Naturforsch. B: Anorg. Chem., Org. Chem.* **1976**, *31B*, 680.
 (277) Ruppert, I.; Bastian, V.; Appel, R. *Chem. Ber.* **1974**, *107*, 3186.
 (278) Ruppert, I.; Bastian, V.; Appel, R. *Chem. Ber.* **1974**, *108*, 2329.
 (279) (a) Sasaki, Y.; Olsen, F. P. *Can. J. Chem.* **1971**, *49*, 271. (b) Barrick, J. C.; Calvo, C.; Olsen, F. P. *Chem. Commun.* **1971**, 1043.
 (280) Banister, A. J.; House, J. R. *J. Inorg. Nucl. Chem.* **1971**, *33*, 4057.
 (281) Mataka, S.; Takahashi, K.; Yamamoto, H.; Tashiro, M. *J. Chem. Soc., Perkin Trans. 1* **1980**, 2417.
 (282) (a) Holt, E. M.; Holt, S. L. *J. Chem. Soc., Chem. Commun.* **1973**, 36. (b) Holt, E. M.; Holt, S. L.; Watson, K. J. *J. Chem. Soc., Dalton Trans.* **1974**, 1347.
 (283) Grushkin, B. U.S. Patents 3 615 409, 3 616 393, 1971; *Chem. Abstr.* **1972**, *76*, 40262, 40283.
 (284) Witt, M.; Roesky, H. W.; Noltemeyer, M.; Clegg, W.; Schmidt, M.; Sheldrick, G. M. *Angew. Chem., Int. Ed. Engl.* **1981**, *20*, 974.

- (285) Roesky, H. W.; Ambrosius, K. *Isr. J. Chem.* **1978**, *17*, 132.
- (286) Bénard, M.; Laidlaw, W. G.; Paldus, J. *Can. J. Chem.*, in press.
- (287) (a) Chivers, T.; Drummond, I. *J. Chem. Soc., Chem. Commun.* **1973**, 734. (b) Chivers, T.; Drummond, I. *Inorg. Chem.* **1974**, *13*, 1222. (c) Bojes, J.; Chivers, T.; Drummond, I. *Inorg. Synth.* **1978**, *18*, 203.
- (288) Chivers, T.; Hojo, M. *Inorg. Chem.* **1984**, *23*, 2738.