

Bioactive Metabolites of Symbiotic Marine Microorganisms

Jun'ichi Kobayashi* and Masami Ishibashi

Faculty of Pharmaceutical Sciences, Hokkaido University, Sapporo 060, Japan

Received November 12, 1992 (Revised Manuscript Received February 5, 1993)

Contents

I. Introduction	1753
II. Bacteria	1754
A. Bacteria Isolated from Sponges	1754
B. Others	1755
III. Fungi	1755
A. Fungi Isolated from Marine Animals	1755
B. Fungi Isolated from Marine Plants	1756
IV. Dinoflagellates	1756
A. Symbiotic Dinoflagellates of Flatworms	1756
B. Others	1763
V. Other Microalgae	1763
VI. Marine Natural Products Assumed To Be of Microbial Origins	1763
VII. Conclusion	1767
VIII. References	1767

I. Introduction

Marine microorganisms are of considerable current interest as a new promising source of bioactive substances. Recently marine microorganisms have proven to produce a variety of chemically interesting and biologically significant secondary metabolites and some of them are expected to serve as lead compounds for drug development or pharmacological tools for basic studies of life sciences.¹ Recent studies have also suggested that some bioactive compounds isolated from marine invertebrates such as sponges, coelenterates, mollusks, or protochordates^{2,3} are truly originating from symbiotic microorganisms (*e.g.* bacteria, fungi, blue-green algae, dinoflagellates, or haptophytes).

It was a striking topic when the real origin of tetrodotoxin⁴ was demonstrated to be marine bacteria isolated from a variety of marine organisms such as a red alga,⁵ a xanthid crab,⁶ or the skin of a pufferfish.⁷ Saxitoxin,⁴ a well-known paralytic shellfish poison, is produced by dinoflagellates of the genus *Gonyaulax* and the blue-green alga *Aphanizomenon flos-aquae*.⁸ A Gram-positive bacteria (Aalxll strain),⁹ which is associated with the Japanese ivory shell *Babylonia japonica*, was shown to be responsible for the production of surugatoxins, the causative toxins of a food poisoning outbreak in 1965 at Suruga Bay.¹⁰ Okadaic acid, a potent inhibitor of protein phosphatases first isolated from a sponge *Halichondria okadai*,¹¹ is produced by a dinoflagellate *Prorocentrum lima*.¹²

Since it is quite impossible to collect unlimited amounts of marine animals or plants, isolation and cultivation of symbiotic microorganisms are of great importance for sufficient and constant supply of bioactive compounds. This review describes the recent studies on bioactive substances isolated from symbiotic



Jun'ichi Kobayashi was born in 1949 at Hirosaki, Japan. He completed his B.S., in 1973, and M.S. degrees, in 1975, from Hokkaido University with Yoshihisa Mizuno on the studies of nucleic acid synthesis. In 1975 he joined Mitsubishi-Kasei Institute of Life Sciences where he worked on the synthesis and conformational analyses of bioactive peptides. After receipt of his Ph.D. from Hokkaido University in 1979, he initiated his research program on marine natural products chemistry. He did postdoctoral studies at University of Illinois with K. L. Rinehart from 1982 to 1984. In 1989 he was appointed as a full professor at Hokkaido University, Faculty of Pharmaceutical Sciences, where he still continues his research career. His main research interest is the search for bioactive substances from marine organisms and their application to the basic research of life sciences as well as the development of new drugs.



Masami Ishibashi was born in 1957 in Saga, Japan. He received his B.S. in 1980 and his Ph.D. in 1985 under the direction of Takeyoshi Takahashi from the University of Tokyo. He then spent four years as a postdoctoral associate at University of Hawaii with R. E. Moore, Mitsubishi-Kasei Institute of Life Sciences with Jun'ichi Kobayashi, and Kitasato Institute with Satoshi Omura. He joined the faculty of Toho University in 1989 and moved to Hokkaido University in 1990, where he is currently an associate professor of pharmacognosy. His major research interests are concerned with natural products chemistry.

microorganisms associated with marine animals or plants. In sections II and III bioactive natural products isolated from symbiotic marine bacteria and fungi are surveyed, while sections IV and V mainly deal with our


recent work on bioactive substances from symbiotic marine dinoflagellates and haptophytes.

II. Bacteria

A. Bacteria Isolated from Sponges


Bioactive metabolites of marine sponges have been extensively studied by marine natural products chemists. The unique metabolites of sponges are, however, often very minor constituents. It is often found that structurally similar compounds are obtained from sponges of different species and sometimes sponges of the same species contained quite different metabolites. These facts as well as the structural characteristics of the compounds strongly suggested that microorganisms living in or on sponges are responsible for the production of many bioactive compounds. Recently some of secondary metabolites which had been isolated from extracts of sponges were obtained from the culture media of marine bacteria isolated from sponges.

Cardellina and co-workers examined the microorganisms growing on or in sponges from Bermudian waters. From a specimen of the sponge *Tedania ignis* a bright orange-pigmented, Gram-positive bacterium was isolated and identified as a member of the genus *Micrococcus*. Culturing of the *Micrococcus* sp. was successfully carried out and from the culture media three diketopiperazines (1–3) were isolated by chromatographic techniques, including centrifugal counter-current chromatography.¹³ Compounds 1–3 proved to be identical to those previously isolated from extracts of the sponge *Tedania ignis*.¹⁴ From the fermentation




culture extracts of the same *Micrococcus* sp. four benzothiazoles (4–7) were obtained.¹⁵ These benzothiazoles were previously known as volatile constituents of cranberries, an aroma constituent of tea leaves, a fermentation product of yeast, or synthesized compounds.

Elyakov and co-workers reported that a *Vibrio* sp. of bacteria isolated from the sponge *Dysidea* sp. collected at Eastern Samoa biosynthesized brominated diphenyl ethers on the basis of GC-MS analyses using 3,5-dibromo-2-(3',5'-dibromo-2'-methoxyphenoxy)phenol (8) as a standard. Compound 8 was previously obtained from extracts of the sponge *Dysidea* sp.¹⁶




From a Japanese sponge *Halichondria okadai* a bacterium of a member of the genus *Alteromonas* was isolated and cultured in the laboratory. From the cultured mycelium of this bacterium we have isolated




a unique macrocyclic lactam, alteramide A (9), pos-



sessing cytotoxic activity [IC₅₀ (50% inhibitory concentration) values against P388, L1210, and KB cells, 0.1, 1.7, and 5.0 μg/mL, respectively].¹⁷ The structure of compound 9 was elucidated by extensive analyses of the spectral data (e.g. ¹H-¹H COSY, HMQC, HMBC, NOESY, and ROESY) as well as the chemical degradations into small fragments. Treatment of 9 with ozone followed by NaBH₄ reduction and acetylation afforded the bicyclo[3.3.0]octane derivative 10, while β-hydroxyornithine (11) was obtained from 9 through ozonolysis and oxidative workup with H₂O₂. The




relative stereochemistry of the bicyclo[3.3.0]octane unit was established from NOESY data obtained for 10 and the absolute configurations of C-23 and C-25 positions of 9 were defined through chiral HPLC examination of 11. Compound 9 possesses two diene groups which were vulnerable to an intramolecular photochemical [4 + 4] cycloaddition to give a hexacyclic compound 12. It should be noted that the structure of alteramide A (9) appears to be biogenetically related to those of an antibiotic ikarugamycin (13) produced by a terrestrial *Streptomyces* sp.¹⁸ and an antifungal and cytotoxic metabolite discoderamide (14) isolated from a Caribbean marine sponge *Discodermia dissoluta*.¹⁹ Since compound 9 was obtained from a bacterium isolated from a marine sponge, discoderamide (14) is strongly suggested to be of microbial origin.




B. Others

A marine-derived actinomycete of the genus *Streptomyces* was isolated from the surface of an unidentified gorgonian of the genus *Pacifigorgia* by Fenical and co-workers.²⁰ From the culture broth of the *Streptomyces* sp. two closely related novel compounds, octalactins A (15) and B (16), were isolated and their structures,



including relative stereochemistries, were firmly established, with 15 being defined by the single-crystal X-ray crystallographic analysis. These metabolites contain unusual saturated eight-membered lactone functionalities and the lactone ring has a boat-chair conformation with a cis lactone in the solid state, which was consistent with the result of the molecular mechanics calculation. Octalactin A (15) exhibited strong cytotoxic activity toward B-16-F10 murine melanoma and HCT-116 human colon tumor cell lines with the IC_{50} values of 0.0072 and 0.5 $\mu\text{g/mL}$, respectively, while quite surprisingly octalactin B (16) was completely inactive in the cytotoxicity assay.


Embryos of the shrimp *Palaemon macrodactylus* are known to resist infection by the pathogenic fungus *Lagenidium callinectes*. Fenical and co-workers found that a penicillin-sensitive bacterial strain (*Alteromonas* sp.) which was consistently isolated from healthy embryos effectively inhibited the growth of the fungus *L. callinectes* in vitro. This bacterial strain was shown to produce and release an antifungal compound, which was identified as 2,3-indolinedione (17, known as isatin).²¹ Bacteria-free embryos which were exposed to the fungus quickly died, whereas embryos re-inoculated with the bacteria or treated only with 2,3-indolinedione (17) thrived. Thus it was clearly revealed that the commensal *Alteromonas* sp. bacteria protect shrimp embryos from fungal infection by producing and liberating the antifungal metabolite 17.



III. Fungi


A. Fungi Isolated from Marine Animals

Chemical studies on the bioactive metabolites of fungal isolates from the marine environments have been limited to date. Marine fungi, however, can be expected as a potential source of new bioactive substances from the following studies. Researchers of the Sankyo group have isolated the novel platelet-activating factor (PAF) antagonists, phomactins A (18),²² B (19), B₁ (20), and B₂ (21),²³ from the cultured broth of a marine deuteromycetes, *Phoma* sp., which lived in the shell of a crab *Chionoecetes opilio* collected off the coast of Fukui prefecture, Japan. Phomactins are unique diterpenoids



and their structures, including the absolute configurations, were established by the X-ray analyses performed on a mono-*p*-bromobenzoyl derivative of phomactin A (18) as well as a diketone derived from phomactin B (19). Phomactin B₁ (20) and B₂ (21) were chemically correlated with phomactin B (19). Compounds 18–21 inhibited PAF-induced platelet aggregation (IC_{50} 1.0 $\times 10^{-5}$ M, 1.7 $\times 10^{-5}$ M, 0.98 $\times 10^{-5}$ M, and 0.16 $\times 10^{-5}$ M, respectively) and binding of PAF to its receptors (IC_{50} 0.23 $\times 10^{-5}$ M, $>10^{-5}$ M, 2.0 $\times 10^{-5}$ M, and 0.54 $\times 10^{-5}$ M, respectively). Phomactin A (18), however, exhibited no effect on adenosine diphosphate-, arachidonic acid-, and collagen-induced platelet aggregation. Phomactin A (18) was thus considered to be a new type of specific PAF antagonist.


Two lipophilic tripeptides, fellutamides A (22) and B (23),²⁴ have been isolated by us from the cultured mycelium of a fungus *Penicillium fellutanum* Biourge which was isolated from the gastrointestinal of the marine fish *Apogon endekataenia* Bleeker, collected off Manazuru beach, Kanagawa prefecture, Japan. The



structures of fellutamides were elucidated by applying several types of two-dimensional NMR techniques in combination with FAB MS/MS data. The stereochemistries of amino acid residues were determined by chiral HPLC and GC analyses and the absolute configuration of 3-hydroxydodecanoic acid obtained through hydrolysis was assigned by comparison of the optical rotation with those of standard samples. Fellutamides A (22) and B (23) were potently cytotoxic against murine leukemia P388 (IC_{50} 0.2 and 0.1 $\mu\text{g/mL}$, respectively) and L1210 (IC_{50} 0.8 and 0.7 $\mu\text{g/mL}$, respectively) cells and human epidermoid carcinoma KB (IC_{50} 0.5 and 0.7 $\mu\text{g/mL}$, respectively) cells in vitro. The fungus *Penicillium fellutanum* is not a marine-specific one, since the same species have been isolated from terrestrial sources. These peptides 22 and 23 which are structurally related to the leupeptines²⁵ exhibited stimulating activity on nerve growth factor (NGF) synthesis.²⁶

Numata and co-workers have isolated three quinazoline derivatives, fumiquinazolines A (24), B (25), and


C (26),²⁷ from the cultured mycelium of a strain of *Aspergillus fumigatus* which existed in the gastrointestinal tract of the saltwater fish *Pseudolabrus japonicus*.



Their structures were based on spectral and chemical evidences as well as X-ray diffraction analysis of 26. The absolute stereostructure of 26 was revealed by the production of L-(+)-alanine through its acid hydrolysis. Compounds 24–26 exhibited moderate cytotoxic activity against the cultured P388 lymphocytic leukemia cells.

B. Fungi Isolated from Marine Plants

A polyketide metabolite, obionin A (27),²⁸ has been isolated by Gloer and co-workers from the liquid culture of the marine fungus *Leptosphaeria obiones* (Crouan et Crouan) Saccoro, a halotolerant ascomycete originally isolated from the salt marsh grass *Spartina alterniflora* in the coastal marshlands of Sapelo Island, GA. The



culture of *L. obiones* originally exhibited potent brine shrimp toxicity. Obionin A (27), however, did not account for that activity. Since an ichthyotoxic hydroquinone metabolite was reported from a brown alga which oxidizes in air to the less active *o*-quinone,²⁹ a similar phenomenon might have occurred for the extract of the *L. obiones* culture. Obionin A (27) showed inhibitory activity against binding of the dopamine D-1 selective ligand ³H-SCH 23390 to bovine corpus striatum membrane with an IC₅₀ value of 2.5 μg/mL.

Gloer and co-workers also reported isolation of two isomeric δ-lactones, helicascalides A (28) and B (29),³⁰ from the Hawaiian mangrove ascomycete *Helicascus kanaloanus* Kohlmeyer. Their structures were based primarily on NMR studies and the absolute configuration was deduced from CD spectral data. Secondary metabolites with these structural features have never been obtained from terrestrial fungi. Marine fungi, therefore, may have potential capabilities to produce different types of compounds from those of terrestrial sources.





Figure 1. Host: flatworm *Amphiscolops magniviridis* (5 mm in length).




Figure 2. Symbiont: dinoflagellate *Amphidinium* sp. (Y-26, 20 μm in length).

IV. Dinoflagellates

A. Symbiotic Dinoflagellates of Flatworms

During the course of our studies on the search for bioactive substances from Okinawan marine organisms,^{31–33} we initiated a project on bioactive natural products of symbiotic marine microalgae about seven years ago. With help from previous studies by Yamasu who systematically collected symbiotic microalgae from marine invertebrates of Okinawan coastal waters,^{34,35} large-scale cultures of the microalgae were carried out in our laboratory. Figures 1 and 2 represent the host (flatworm) and the symbiont (dinoflagellate), respectively.


The first microalga we utilized for research was a dinoflagellate belonging to the genus *Amphidinium* (strain number Y-5, 25 μm in length and 20 μm in width) isolated from the inner tissue of the host, a flatworm of the genus *Amphiscolops* Graff, 1905 (500 μm in length and 220 μm in width, green color), which was living on algae or sea weeds such as *Enteromorpha* and *Jania* spp. and collected at Chatan beach, Okinawa Island.³⁶ The unialgal cultures of *Amphidinium* sp. (Y-5) were grown in 3-L glass bottles containing 2 L of sea water medium enriched with Provasoli's ES (Erd-Schreiber) supplement (Table I).^{37,38} Cultures were incubated statically at 25 °C in an apparatus where illumination from a fluorescent light source was supplied in a cycle of 16 h of light and 8 h of darkness. After 2 weeks the culture was harvested by suction of the supernatant media with an aspirator, followed by centrifugation, to yield harvested cells ranging from 0.3 to 0.5 g/L of culture. Harvested cells were extracted with methanol/toluene (3:1) followed by partitioning between toluene and water. The toluene-soluble fraction was subjected

Table I. Contents of Provasoli's ES supplement^a

distilled H ₂ O	100 mL
NaNO ₃	350 mg
sodium glycerophosphate	50 mg
Fe (as EDTA; ^b 1:1 mol)	2.5 mg
PII solution of metals ^c	25 mL
vitamin B ₁₂	10 μg
vitamin B ₁	0.5 mg
biotin	5 μg
TRIS ^d	500 mg
pH	7.8

^a This was filtered with sterile membrane filter and added to sterile sea water to make a 1% solution. ^b Ethylenediaminetetraacetic acid. ^c The solution consists of Fe (as Cl⁻, 1 mg), B (H₃BO₃, 20 mg), Mn (as Cl⁻, 4 mg), Zn (as Cl⁻, 500 μg), Co (as Cl⁻, 100 μg), Na₂-EDTA (100 mg), and distilled H₂O (100 mL). ^d Tris(hydroxymethyl)aminomethane.



Scheme I. Isolation Procedure for Amphidinolides A-D (30-33)








to repeated silica gel flash chromatography followed by reversed-phase HPLC resulting in isolation of four cytotoxic macrolides amphidinolides A (30),³⁹ B (31),⁴⁰ C (32),⁴¹ and D (33)⁴² (Scheme I).

Three other species of dinoflagellates of the genus *Amphidinium* (strain number Y-5', Y-25, and Y-26) were also investigated. Their host animals were also Okinawan flatworms [Y-5', *Amphiscolops* sp. (different from the host of Y-5); Y-25, *Amphiscolops breviviridis*; Y-26, *Amphiscolops magniviridis*]. From the extracts of the cultured cells of these strains of *Amphidinium* spp. four other cytotoxic macrolides amphidinolides E (34),⁴³ F (35),⁴⁴ G (36),⁴⁵ and H (37)⁴⁵ were isolated. The isolated yields and cytotoxicity data of amphidinolides A-H (30-37) are presented in Table II. In addition to potent cytotoxic activity, amphidinolides B (31) and C (32) were shown to activate rabbit skeletal muscle actomyosin ATPase activity.⁴⁶ The chemical structures of these macrolides (30-37) were elucidated mainly on the basis of extensive spectroscopic studies including several types of two-dimensional NMR experiments⁴⁷ (e.g. ¹H-¹H COSY, ¹H-¹³C COSY, HMBC, HMQC, and NOESY spectra).

Amphidinolide A (30), C₃₁H₄₆O₇, was suggested to possess an α,β,γ,δ-dienoate chromophore from its UV absorption spectrum (λ_{max} 265 nm). ¹H and ¹³C NMR data revealed the presence of one epoxide and three



31
33 C-21 epimer



37

Table II. Isolated Yields and Cytotoxicity Data of Amphidinolides

compounds	isolated yields ($\times 10^{-4}\%$) ^a strain				cytotoxicity (IC ₅₀ , $\mu\text{g/mL}$)	
	Y-5	Y-5'	Y-25	Y-26	L1210	KB
amphidinolide A	20	4	– ^b	–	2.0	5.7
amphidinolide B	10	–	–	0.8	0.00014	0.0042
amphidinolide C	15	–	–	0.3	0.0058	0.0046
amphidinolide D	4	–	–	–	0.019	0.08
amphidinolide E	–	3	–	–	2.0	10
amphidinolide F	–	–	–	0.1	1.5	3.2
amphidinolide G	–	–	20	–	0.0054	0.0059
amphidinolide H	–	–	17	–	0.00048	0.00052


^a On the basis of wet weight of the harvested cells. ^b “–” denotes “not isolated”.

exo-methylene groups. Detailed analysis of the ¹H–¹H COSY spectrum allowed assignment of all proton signals and clearly established the proton connectivities from H-2 to H₃-25, leading to a 20-membered macrolide structure. It was ambiguous whether a secondary methyl group was located at C-22 or C-23 due to overlapping of the signals for protons on C-22–C-24. It was, however, placed on C-22 because the ¹³C chemical shifts of the C-23–C-25 signals suggested the presence of an *n*-propyl group instead of an ethyl group.⁴⁸

Amphidinolide B (31), C₃₂H₅₀O₈, possessed an α -methyl- α,β -unsaturated ester moiety, which was shown by the UV absorption (λ_{max} 222 nm) as well as the ¹³C NMR chemical shifts. ¹H and ¹³C NMR data also suggested the presence of one *exo*-methylene, one epoxide, and one isolated ketone group. The ¹H–¹H COSY spectrum indicated the proton connectivities of three fragments (C-1–C-15, C-17–C-19, and C-21–C-26). Since the ¹H NMR chemical shifts for α positions to an sp³ quaternary carbon (C-16) and ketone group (C-20) could be discriminated and NOE were observed between the methyl protons on C-15 and C-16, these three fragments were shown to be connected through C-16 and C-20, leading to a total structure consisting of a 26-membered macrocyclic lactone ring.

Amphidinolide C (32), C₄₁H₆₂O₁₀, exhibited the UV absorption maximum at λ_{max} 240 nm, implying the presence of a diene chromophore. Interpretation of the ¹H and ¹³C NMR data suggested the presence of two tetrahydrofuran rings, two exomethylenes, and two isolated ketone groups. Extensive two-dimensional NMR experiments were carried out on the tetraacetate (38) prepared from 32. Three partial structures (C-2 ~ C-14, C-16 ~ C-17, and C-19 ~ C-34) were elucidated by analysis of the ¹H–¹H COSY and double relayed coherence transfer (RCT2) spectra. All protonated carbons were clearly assigned by ¹H–¹³C COSY via one-bond couplings. Geometries of the double bonds were determined from NOE data. The connection of three partial structures through carbonyl groups (C-1, C-15, and C-18) was established by analyzing the ¹H–¹³C long-range (two- and three-bond) couplings detected in the HMBC spectrum to give rise to a complete structure. The diene moiety (C-36, 9, 10, and 11) of 38 was slowly oxidized by air to afford a [4 + 2] cycloaddition product (39), the spectral data of which provided additional proof for the structure of amphidinolide C (32).

Amphidinolide E (34), C₃₀H₄₄O₆, showed a UV absorption maximum at λ_{max} 230 nm, corresponding to




39

a diene chromophore. The presence of two *exo*-methylenes and one tetrahydrofuran ring was inferred by analyzing the ¹H and ¹³C NMR data. ¹H–¹H COSY data afforded information on the proton connectivities for three partial structures (C-2–C-14, C-15–C-19, and C-20–C-27). Geometries of the four disubstituted double bonds were deduced on the basis of the ¹H–¹H coupling constants determined by a *J*-resolved two-dimensional NMR experiment. The partial structures were revealed to be linked to each other by ¹H chemical shift data together with the relayed correlations observed through an RCT-COSY spectrum to construct a 19-membered macrocyclic lactone ring with an alkyl side chain.

Spectral investigations of amphidinolides D (33), G (36), and H (37), whose molecular formulas are all C₃₂H₅₀O₈, revealed that they are structurally related closely to amphidinolide B (31): amphidinolides B (31) and D (33) are stereoisomers at C-21; amphidinolides G (36) and H (37) are regioisomers of the lactone-terminal positions (C-25 or C-26); amphidinolides B (31) and H (37) are only different in the position of one hydroxyl group (C-16 or C-26). Amphidinolide F (35), C₃₅H₅₂O₉, proved to be analogous to amphidinolide C (32). The structural difference between amphidinolides F (35) and C (32) was found in the length of the alkyl side chain, the former possibly being a biogenetic precursor of the latter.

Structural features of the amphidinolides are unique. Amphidinolides show a variation in the size of the macrocyclic lactone rings [19-membered, amphidinolide E (34); 20-membered, amphidinolide A (30); 26-membered, amphidinolides B (31), D (33), and H (37); 25-membered, amphidinolides C (32) and F (35); 27-membered, amphidinolide G (36)]. It should be noted that cytotoxic activity of amphidinolide B (31) and its related compounds (33, 36, and 37) are extremely strong (Table II). Amphidinolide D (33), an epimer of amphidinolide B (31) at the C-21 position, was, however, about 100 times less cytotoxic than the latter. During the isolation process of the extract of the strain Y-5, 1 mol of MeOH was added on the C-8/C-9 epoxide of amphidinolide B (31) to generate compound 40,⁴² the



40

IC₅₀ value of which was 0.081 $\mu\text{g/mL}$ against L1210 cell, being considerably weaker than that of 31 (1/600). These results implied that the stereochemistry at C-21 and the presence of an epoxide at C-8/C-9 position are quite important for the cytotoxic activity of these compounds, presumably due to the significant change of the






Figure 3. Phase-sensitive NOESY spectrum of Amphidinolide A (30) (500 MHz, C_6D_6).

molecular conformation. The cytotoxicities of amphidinolides C (32) and F (35) were also very different, the former being about 250 times as strong as the latter. The length of the side chain may significantly affect the potency of cytotoxic activity. It seems to be also noteworthy that amphidinolides A (30), B (31), C (32), and E (34) are quite unlike one another in molecular constitution and substitution patterns although they were obtained from the same species of dinoflagellate *Amphidinium* sp. (Y-5 or Y-5'). Their biogenetical relationships may raise an interesting subject.

A very important question remaining undefined is the stereochemistry of amphidinolides. All attempts to obtain crystals of the amphidinolides suitable for X-ray analysis have been unsuccessful. Efforts to elucidate the stereochemistry of amphidinolide A (30), however, were made by analyzing the phase-sensitive NOESY spectra of this 20-membered macrolide with nine chiral centers (Figure 3). Interpretation of the relative stereochemistry of conformationally mobile molecules by spectral data alone is still not easy and very few reports have ever been published on this subject.⁴⁹ The NOE data, however, may provide some suggestive information for the relative configurations and molecular conformations of such molecules.⁵⁰ We have proposed a possible stereostructure of amphidinolide A as 30a containing the relative configurations of the nine chiral centers of 30 and most sufficiently



satisfying the NOESY data obtained in several solvent systems.⁵¹ NOESY experiments on amphidinolide C (32) were also carried out (Figure 4) and the relative stereochemistry around the two tetrahydrofuran ring moieties were proposed as 32a and 32b. For unam-



biguous assignment of the stereochemistry of amphidinolides the following studies are required: (a) further large-scale culturing of the dinoflagellates *Amphidin-*





Figure 4. Phase-sensitive NOESY spectrum of Amphidinolide C (32) (500 MHz, C_6D_6).

Scheme II. Outline of the Synthesis of a Potential C-10-C-19 Fragment (41) of Amphidinolide A (30)



ium spp. to obtain an adequate quantity of these macrolides; (b) degradation experiments (ozonolysis or periodate oxidation) of these macrolides to lead to several small fragments; (c) stereoselective synthesis of all possible stereoisomers of the small fragments in optically active forms. These challenging problems are currently under investigation by us. Williard and co-workers independently described synthetic efforts on amphidinolide A (30) in order to make an unambiguous assignment of its stereochemistry.⁵² A potential C-10-C-19 fragment (41) of amphidinolide A (30) was prepared by a stereospecific route (Scheme II). It may be possible to prepare alternative optically pure stereoisomers of the diene 41 by utilizing similar reaction

sequences starting with (+)- or (-)-tartaric acid (42) and (+)- or (-)-methyl 3-hydroxy-2-methylpropionate (43), each of which is commercially available.

The quantity of these macrolides in the extracts of the cultured cells varied a little during the course of time. Extraction of the harvested cells was carried out repeatedly after about 300–400 g of cells (wet weight) were accumulated to give different batches of extracts of the same strain of the alga. From a batch of the extract of *Amphidinium* sp., strain Y-5, which was relatively rich in amphidinolides B (31) and D (33), we once isolated another component of interest, designated as Y5-726, which is most likely a macrolide related to the amphidinolides. Y5-726 was separated from the





Figure 5. ^1H NMR spectrum of Y5-726 (500 MHz, CDCl_3).

mixture of **31** and **33** by reversed-phase HPLC using 87% methanol as eluent. It was isolated in 0.0005% yield (wet weight) as colorless amorphous solid, $[\alpha]_{\text{D}}^{26} +4.5^\circ$ (c 1, CHCl_3); UV (MeOH) 238 nm (ϵ 29 000); IR (film) 3400, 2930, 1730, 1440, and 1050 cm^{-1} . The molecular weight of Y5-726 was suggested to be 726 by the FABMS data [glycerol matrix m/z 727 ($\text{M} + \text{H}$) $^+$ and 709 ($\text{M} + \text{H} - \text{H}_2\text{O}$) $^+$; diethanolamine (DEA) matrix m/z 832 ($\text{M} + \text{DEA} + \text{H}$) $^+$]. The ^1H NMR spectrum of Y5-726 (Figure 5) showed signals due to three secondary methyls [δ 1.00 (3 H, d, $J = 6.5$ Hz), 1.08 (3 H, d, $J = 7$ Hz), 1.67 (3 H, d, $J = 6.5$ Hz)], two vinyl methyls [δ 1.77 (3 H, s) and 1.88 (3 H, s)], two possible epoxide-bearing methine protons [δ 2.92 (1 H, dd, $J = 4.5$ and 2 Hz) and 3.14 (1 H, dd, $J = 5.5$ and 2 Hz)], eight oxymethine protons [δ 3.77 (2 H, m), 3.86 (1 H, br s), 3.95 (1 H, dd, $J = 12$ and 4.5 Hz), 4.07 (1 H, m), 4.17 (1 H, m), 4.23 (1 H, dd, $J = 13$ and 7 Hz), and 4.30 (1 H, m)], two possible *exo*-methylene protons [δ 4.95 (2 H, s)], and eight olefinic and one possible lactone-terminal protons [δ 5.36 (1 H, dd, $J = 15$ and 9 Hz), 5.45 (2 H, m), 5.54 (1 H, dd, $J = 9.5$ and 5.5 Hz), 5.59 (1 H, m), 5.76 (1 H, dd, $J = 15.5$ and 6.5 Hz), 5.84 (1 H, d, $J = 11$ Hz), 6.21 (1 H, d, $J = 15.5$ Hz), and 6.31 (1 H, dd, $J = 15$ and 11 Hz)]. The incomplete ^1H - ^1H

COSY spectrum of Y5-726 was recorded (Figure 6) and suggested the presence of partial structures of **44a-c**.



Further structural studies of Y5-726 were unfortunately precluded since the compound decomposed extensively during storage as a CDCl_3 solution in a refrigerator, probably due to contact with traces of hydrochloric acid in the CDCl_3 solvent. Unfortunately, Y5-726 was not isolated from the extracts of subsequent *Amphidinium* sp. cultures.






Figure 6. ^1H - ^1H COSY spectrum of Y5-726 (500 MHz, CDCl_3).

Nakamura and co-workers isolated the novel vasoconstrictive substances, named zooxanthellatoxins A and B,^{53,54} from a cultured zooxanthella (typical symbiotic dinoflagellate) of the genus *Symbiodinium*, which was isolated from an Okinawan flatworm *Amphiscolops* sp. Zooxanthellatoxins A and B (negative FABMS, m/z 2873 and 2835, respectively) proved to be sulfate esters of highly unsaturated polyols and induce the contractile response of the rabbit aorta at 7×10^{-7} M each. This contractile response was not affected by tetrodotoxin (a sodium channel blocker), histamine antagonists, or serotonin antagonists, while it was markedly inhibited by calcium antagonists such as verapamil or magnesium chloride. Thus this vasoconstrictive activity was suggested to result from an increase of the calcium permeability across the membrane. The structure of zooxanthellatoxin A was investigated through the periodate oxidation to give several fragments, the structures of which were elucidated (e.g. 45a-d) by extensive spectral analyses.




B. Others

A symbiotic dinoflagellate of the genus *Symbiodinium* was isolated from the inside of gill cells of the Okinawan bivalve *Fragum* sp. The procedure of culturing this dinoflagellate in the laboratory followed that of the dinoflagellate *Amphidinium* spp. described above. From the toluene-soluble fraction of the extract of the harvested cells a sphingosine derivative, symbioramide (46),⁵⁵ was isolated by using silica gel column



chromatographies. The structure of symbioramide (46) was elucidated on the basis of spectral and chemical means. The ¹H and ¹³C NMR data suggested that compound 46 belongs to ceramides. Acid hydrolysis of 46 afforded methyl 2-hydroxyoctadec-3(*E*)-enoate and 2(*S*)-amino-3(*R*)-hydroxyoctadecan-1-ol. The absolute configurations at C-2 and C-3 positions of the sphingosine part of 46 were thus determined. Symbioramide (46) exhibited weak cytotoxic activity against L1210 cells in vitro with an IC₅₀ value of 9.5 μg/mL. Additionally, compound 46 was found to be a sarcoplasmic reticulum (SR) Ca²⁺-ATPase activator. The Ca²⁺-ATPase in SR membrane plays an key role in muscle relaxation by energizing Ca²⁺ pumping from the cytoplasm into the lumen of SR.⁵⁶ Symbioramide (46) at 10⁻⁴ M activated SR Ca²⁺-ATPase activity by 30%. This is the first example of SR Ca²⁺-ATPase activator of marine origin. Symbioramide (46) may serve as a valuable chemical tool for studying the regulatory mechanisms of SR Ca²⁺-pumping systems. The α-hydroxy-β,γ-dehydro fatty acid contained in symbioramide (46) is rare from natural sources. Previously ceramides of α-hydroxy fatty acids were obtained from the sponge *Dysidea etheria*.⁵⁷ The ceramides isolated from the sponge may be of microbial origin since sponges are known to possess symbiotic microorganisms. The absolute configuration of the C-2 position of 2-hydroxyoctadec-3(*E*)-enoic acid moiety of symbioramide (46), which we left unassigned, was established as *R* on the basis of the total synthesis of symbioramide (46) achieved by Hino and co-workers.⁵⁸

Soft corals are known to possess symbiotic microalgae. We previously isolated a diterpene, 16-deoxysarcophine (47),⁵⁹ with potent Ca antagonistic activity from the




Okinawan soft coral *Sarcophyton* sp. and established the structure by X-ray analysis. This was the first example of a Ca antagonist from marine sources and of a nonalkaloid compound with such activity. Symbiotic microalgae of the soft coral *Sarcophyton* sp. were examined and a dinoflagellate belonging to the genus *Symbiodinium* was isolated from the polyps of the soft


coral. 16-Deoxysarcophine (47) was detected by HPLC analysis of the extracts of the dinoflagellate *Symbiodinium* sp.

V. Other Microalgae

The Haptophyceae are microscopic, unicellular algae, which are widely distributed in the oceans and often constitute a major proportion of marine phytoplankton. We encountered a haptophyte of the genus *Hymenomonas*, which was isolated from an unidentified cylindrical stony coral collected at Sesoko Island, Okinawa. The alga was mass cultured by the same procedure as that of the *Amphidinium* spp. of dinoflagellates. The extract of the harvested cells was partitioned between toluene and water and the aqueous phase was further extracted with chloroform. The toluene-soluble fraction was subjected to separation using Sephadex LH-20 and silica gel column chromatographies to give a sterol sulfate, hymenosulfate (48),⁶⁰ with potent Ca²⁺-




releasing activity in SR.⁶¹ The structure of 48 was elucidated by conversion of 48 into known (24*R*)-23,24-dimethylcholesta-(22*E*)-5,22-dien-3β-ol through acid hydrolysis. The presence of a sulfate group was confirmed by ion chromatography of sulfate ions liberated by hydrolysis. Hymenosulfate (48) is the first sterol sulfate isolated from marine microalgae. The major components of the toluene-soluble fraction of the extract of this haptophyte were glycolipids, mono- and digalactosyldiacylglycerols (49 and 50, respectively), while the chloroform layer contained mainly octadecatetraenoic acid (51) along with a small amount of monogalactosylmonoacylglycerol (52). In the SR, the Ca²⁺-releasing activity of 48 was 10 times more potent than that of caffeine, a well-known Ca²⁺ releaser. The glycolipids 49, 50, and 52 exhibited⁶⁰ inhibition of Na⁺,K⁺-ATPase activity⁶² with an IC₅₀ value of 2 × 10⁻⁵ M each.




VI. Marine Natural Products Assumed To Be of Microbial Origins

Recently, many fused tetra- and pentacyclic aromatic alkaloids have been isolated from a variety of marine organisms (Chart I),⁴ e.g., from sponges, amphimedine

Chart I



(53, *Amphimedon* sp.),⁶³ petrosamine (54, *Petrosia* sp.),⁶⁴ and dercitin (55, *Dercitus* sp.);^{65,66} from tunicates, ascididemine (56, *Didemnum* sp.),⁶⁷ cystodytins (57 and 58, *Cystodytes* sp.),^{68,69} and segolines (59–61, *Eudistoma* sp.);^{70,71} and from a sea anemone, calliactine (62, *Calliactis* sp.).^{72,73} These alkaloids, obtained from such diverse sources, might be assumed to originate from symbiotic microorganisms, since they possess a common tetracyclic ring system (63, a pyrido[4,3,2-*mn*]acridine skeleton or benzo-3,6-diazaphenanthroline ring) and are closely related to one another. For example, segoline




63


B (60) and isosegoline A (61) appear to correspond to regioisomers which were assumed to be generated through intramolecular [4 + 2] cycloaddition reaction from cystodytin B (58).

From a compound tunicate *Clavelina* sp. collected at Wakaya Island, Fiji, a pyrroloiminoquinone alkaloid, wakayin (64),⁷⁴ was isolated and this alkaloid exhibited cytotoxic, antimicrobial, topoisomerase II inhibitory, and DNA-damaging activities. The structure of compound 64 appears to be closely related to those of discorhabdin C (65)^{75–77} and prianosin A (66),^{78–80} which were isolated from marine sponges *Latrunculia* sp. and *Prianos melanos*, respectively. Wakayin (64) may be biogenetically derived from two molecules of tryptophan, while compounds 65 and 66 seem to be


biosynthesized from tyrosine and tryptophan. Since similar biogenetic origins could be assumed for these compounds, these metabolites may be produced by symbiotic microorganisms.



64



65



66

Several isoquinoline quinone alkaloids, such as renierone (67),⁸¹ mimosamycin (68),⁸¹ renieramycin E (69),⁸² and renierol (70) (Chart II),⁸³ have been isolated from marine sponges of the genera *Reniera* and *Xestospongia*. From a tunicate *Ecteinascidia turbinata* similar tetrahydroisoquinoline alkaloids, ecteinacidins (e.g., 71),^{84,85} were isolated. These alkaloids were likely to be biosynthesized by symbiotic microorganisms, since mimosamycin (68) was first isolated from *Streptomyces lavendulae* no. 314.⁸⁶ and this terrestrial microorganism also yielded saframycin B (72),⁸⁷ which

Chart II





Chart III






is similar in structure to renieramycins and ecteinacidins.






From sponges of the genus *Mycale* and *Theonella*, several polyether metabolites such as mycalamide A (73),^{86,89} onnamide A (74),^{90,91} and theopederin A (75)⁹² were isolated. Their structures have striking resemblances to pederin (76) (Chart III),⁹³ a vesicatory principle from blister beetles of the genus *Paederus*. It seems quite interesting that polyethers obtained from marine sponges and terrestrial insects bear such strong structural similarities.

Swinholide A (77)⁹⁴⁻⁹⁸ and bistheonelide A (78)⁹⁹⁻¹⁰² are potent cytotoxic macrodiolides isolated from marine

sponges of the genus *Theonella*. From the egg masses of the nudibranch *Hexabranchnus sanguineus* oxazole-containing macrolides, ulapualide A (79)¹⁰³ and kabiramide C (80),^{104,105} were isolated. The latter was also obtained from a sponge *Halichondria* sp. which contained a related macrolide halichondramide (81)^{106,107} as the major constituent. These macrolides (77-81) were structurally related to scytophycin B (82)¹⁰⁸ and tolytoxin A (83)¹⁰⁹ which were isolated from the laboratory-cultured terrestrial blue-green algae *Scytonema pseudohofmanni* and *Tolypothrix conglutinata* var. *colorata*, respectively. Three contiguous oxazole rings of ulapualide A (79) and kabiramide C




Scheme III. A Hypothetical Biogenetic Pathway for Three Oxazole Systems in Ulapualide A and Kabiramide C



macrolides are strongly suggested to be blue-green algae (Scheme IV). Electron microscopic analysis demonstrated that a symbiotic blue-green alga inhabited the marine sponge *Theonella swinhoei* that contained swinholides.⁹⁷

There are a number of secondary metabolites isolated from marine sponges or tunicates bearing the structural characteristics which were suggestive that those metabolites were produced by symbiotic microorganisms. Among them, macrolides and peptides are typical instances. From an Okinawan tunicate *Eudistoma cf. rigida* four cytotoxic macrolides, iejimalides A–D (84–87),^{110,111} were isolated. Since the iejimalides possess



- 84 $R^1=H, R^2=H$
 85 $R^1=Me, R^2=H$
 86 $R^1=H, R^2=SO_3Na$
 87 $R^1=Me, R^2=SO_3Na$

(80) may be assumed as being biosynthesized through naturally-occurring Beckmann rearrangements (Scheme III) from a precursor of 22-membered lactone ring, this ring size being the same as that of scytophycin B (82) or tolytoxin A (83). Nudibranchs are known to feed on sponges, whereas sponges are symbiotically associated with blue-green algae. Thus the real origins of these

a common structural feature consisting of a 24-membered macrocyclic lactone ring, they are presumably of microbial origin. Sponges belonging to the genus *Theonella* proved to contain numerous interesting bioactive substances. In particular, many cyclic peptides or peptide lactones were reported to be obtained from *Theonella* sponges and their structural features, including unusual amino acids or D-amino acids, suggest that they are likely to be originating from symbiotic microorganisms. Theonellapeptolide Id (88),^{112–116} a peptide lactone possessing Na^+, K^+ -ATPase inhibitory

- (6) Noguchi, T.; Jeon, J.-K.; Arakawa, O.; Sugita, H.; Deguchi, Y.; Shida, Y.; Hashimoto, K. *J. Biochem.* 1986, 99, 311-314.
- (7) Yotsu, M.; Yamazaki, T.; Meguro, Y.; Endo, A.; Murata, M.; Naoki, H.; Yasumoto, T. *Toxicol.* 1987, 25, 225-228.
- (8) Shimizu, Y. *Progress in the Chemistry of Organic Natural Products*; Herz, W., Grisebach, H., Kirby, G. W., Tamm, C., Eds.; Springer-Verlag: Wien, 1984; Vol. 45, pp 235-264.
- (9) Kosuge, T.; Zenda, H.; Tsuji, K. *Yakugaku Zasshi* 1987, 107, 665-675.
- (10) Kosuge, T.; Tsuji, K.; Hirai, K.; Fukuyama, T. *Chem. Pharm. Bull.* 1985, 33, 3059-3061.
- (11) Tachibana, K.; Scheuer, P. J.; Tsukitani, Y.; Kikuchi, H.; van Engen, D.; Clardy, J.; Gopichand, Y.; Schmitz, F. J. *J. Am. Chem. Soc.* 1981, 103, 2469-2471.
- (12) Murakami, Y.; Oshima, Y.; Yasumoto, T. *Bull. Jpn. Soc. Sci. Fish.* 1982, 48, 69-72.
- (13) Stierle, A. A.; Cardellina, J. H., II; Singleton, F. L. *Experientia* 1988, 44, 1021.
- (14) Schmitz, F. J.; Vanderah, D. J.; Hollenbeak, K. H.; Enwall, C. E. L.; Gopichand, Y.; Sengupta, P. K.; Hossain, M. B.; van der Helm, C. *J. Org. Chem.* 1983, 48, 3941-3945.
- (15) Stierle, A. A.; Cardellina, J. H., II; Singleton, F. L. *Tetrahedron Lett.* 1991, 32, 4847-4848.
- (16) Elyakov, G. B.; Kuznetsova, T.; Mikhailov, V. V.; Maltsev, I. I.; Voinov, V. G.; Fedoreyev, S. A. *Experientia* 1991, 47, 632-633.
- (17) Shigemori, H.; Bae, M.-A.; Yazawa, K.; Sasaki, T.; Kobayashi, J. *J. Org. Chem.* 1992, 57, 4317-4320.
- (18) Ito, S.; Hirata, Y. *Bull. Chem. Soc. Jpn.* 1977, 50, 1813-1820.
- (19) Gunasekera, S. P.; Gunasekera, M.; McCarthy, P. *J. Org. Chem.* 1991, 56, 4830-4833.
- (20) Tapiolas, D. M.; Roman, M.; Fenical, W.; Stout, T. J.; Clardy, J. *J. Am. Chem. Soc.* 1991, 113, 4682-4683.
- (21) Gil-Turnes, M. S.; Hay, M. E.; Fenical, W. *Science* 1989, 246, 116-118.
- (22) Sugano, M.; Sato, A.; Iijima, Y.; Oshima, T.; Furuya, K.; Kuwano, H.; Hata, T.; Hanzawa, H. *J. Am. Chem. Soc.* 1991, 113, 5463-5464.
- (23) Sugano, M.; Sato, A.; Iijima, Y.; Kuwano, H.; Haruyama, H.; Yoda, K.; Hata, T.; Hanzawa, H.; Oshima, T.; Furuya, K. *Tennen Yuki Kagobutsu Toronkai Yoshishu* 1991, 33, 699-706.
- (24) Shigemori, H.; Wakuri, S.; Yazawa, K.; Nakamura, T.; Sasaki, T.; Kobayashi, J. *Tetrahedron* 1991, 47, 8529-8534.
- (25) Aoyagi, T.; Takeuchi, T.; Matsuzaki, A.; Kawamura, K.; Kondo, S.; Hamada, M.; Maeda, K.; Umezawa, H. *J. Antibiot.* 1969, 22, 283-286.
- (26) Yamaguchi, K.; Tsuji, T.; Wakuri, S.; Yazawa, K.; Kondoh, K.; Shigemori, H.; Kobayashi, J. *Biosci. Biotech. Biochem.* 1993, 57, 195-199.
- (27) Numata, A.; Takahashi, C.; Matsushita, T.; Miyamoto, T.; Kawai, K.; Usami, Y.; Matsumura, E.; Inoue, M.; Ohishi, H.; Shingu, T. *Tetrahedron Lett.* 1991, 33, 1621-1624.
- (28) Poch, G. K.; Gloer, J. B. *Tetrahedron Lett.* 1989, 30, 3483-3486.
- (29) Gerwick, W. H.; Fenical, W. *J. Org. Chem.* 1981, 46, 22-27.
- (30) Poch, G. K.; Gloer, J. B. *J. Nat. Prod.* 1989, 52, 257-260.
- (31) Kobayashi, J.; Nakamura, H.; Ohizumi, Y.; Kobayashi, M.; Hirata, Y. *Pharmaceuticals and the Sea*; Jefford, C. W., Rinehart, K. L., Shield, L. S., Eds.; Technomic Publishing Co.: Basel, 1988; pp 73-78.
- (32) Kobayashi, M.; Kobayashi, J.; Ohizumi, Y. *Bioorganic Marine Chemistry*; Scheuer, P. J. Ed.; Springer-Verlag: Heiderberg, 1989; Vol. 3, pp 71-84.
- (33) Kobayashi, J. *New J. Chem.* 1990, 14, 741-745.
- (34) Yamasu, T. *Iden* 1988, 42, 12-20.
- (35) Yamasu, T. *The Coral Reefs of Okinawa*; Nishidaira, M., Ed.; Okinawa Kankyo Kagaku Kensa Center: Urazoe, Okinawa, 1988; pp 123-145.
- (36) Yamasu, T.; Okazaki, A. *Galaxea* 1987, 6, 61-68.
- (37) Provasoli, L. *Culture and Collection of Algae*; Watanabe, A., Hattori, A., Eds.; Japanese Society of Plant Physiology: Tokyo, 1968; pp 63-75.
- (38) Iwasaki, H. *Sourui Kenkyuhou*; Nishizawa, K., Chihara, M., Eds.; Kyouritau-Shuppan: Tokyo, 1979; pp 281-293.
- (39) Kobayashi, J.; Ishibashi, M.; Nakamura, H.; Ohizumi, Y.; Yamasu, T.; Sasaki, T.; Hirata, Y. *Tetrahedron Lett.* 1986, 27, 5755-5758.
- (40) Ishibashi, M.; Ohizumi, Y.; Hamashima, M.; Nakamura, H.; Hirata, Y.; Sasaki, T.; Kobayashi, J. *J. Chem. Soc., Chem. Commun.* 1987, 1127-1129.
- (41) Kobayashi, J.; Ishibashi, M.; Wälchli, M. R.; Nakamura, H.; Hirata, Y.; Sasaki, T.; Ohizumi, Y. *J. Am. Chem. Soc.* 1988, 110, 490-494.
- (42) Kobayashi, J.; Ishibashi, M.; Nakamura, H.; Ohizumi, Y.; Yamasu, T.; Hirata, Y.; Sasaki, T.; Ohta, T.; Nozoe, S. *J. Nat. Prod.* 1989, 52, 1036-1041.
- (43) Kobayashi, J.; Ishibashi, M.; Murayama, T.; Takamatsu, M.; Iwamura, M.; Ohizumi, Y.; Sasaki, T. *J. Org. Chem.* 1990, 55, 3421-3423.
- (44) Kobayashi, J.; Tsuda, M.; Ishibashi, M.; Shigemori, H.; Yamasu, T.; Hirota, H.; Sasaki, T. *J. Antibiot.* 1991, 44, 1259-1261.
- (45) Kobayashi, J.; Shigemori, H.; Ishibashi, M.; Yamasu, T.; Hirota, H.; Sasaki, T. *J. Org. Chem.* 1991, 56, 5221-5224.
- (46) Kobayashi, M.; Kajiwara, A.; Takahashi, M.; Ohizumi, Y.; Shoji, N.; Takemoto, T. *J. Biol. Chem.* 1984, 259, 15007-15009.
- (47) Ernst, R. R. *Angew. Chem., Int. Ed. Engl.* 1992, 31, 805-823.
- (48) Capon, R. J.; Faulkner, D. J. *J. Org. Chem.* 1984, 49, 2506-2508.
- (49) Sowinski, P.; Gariboldi, P.; Pawlak, J. K.; Borowski, E. *J. Antibiot.* 1989, 42, 1639-1642.
- (50) Lancelin, J.-M.; Beau, J.-M. *J. Am. Chem. Soc.* 1990, 112, 4060-4061.
- (51) Kobayashi, J.; Ishibashi, M.; Hirota, H. *J. Nat. Prod.* 1991, 54, 1435-1439.
- (52) O'Connor, S. J.; Willard, P. G. *Tetrahedron Lett.* 1989, 30, 4637-4640.
- (53) Asari, T.; Nakamura, H.; Murai, A.; Kondo, T.; Yoshida, K.; Ohizumi, Y. *Tennen Yuki Kagobutsu Toronkai Yoshishu* 1992, 34, 95-102.
- (54) Nakamura, H.; Asari, T.; Ohizumi, Y.; Kobayashi, J.; Yamasu, T.; Murai, A. *Toxicol.* 1993, 31, 371-376.
- (55) Kobayashi, J.; Ishibashi, M.; Nakamura, H.; Hirata, Y.; Yamasu, T.; Sasaki, T.; Ohizumi, Y. *Experientia* 1988, 44, 800-802.
- (56) Kobayashi, M.; Shoji, N.; Ohizumi, Y. *Biochim. Biophys. Acta* 1987, 903, 96-102.
- (57) Grode, S. H.; Cardellina, J. H., II. *Lipids* 1983, 18, 889-893.
- (58) Nakagawa, M.; Yoshida, J.; Hino, T. *Chem. Lett.* 1990, 1407-1410.
- (59) Kobayashi, J.; Ohizumi, Y.; Nakamura, H.; Yamakado, T.; Matsuzaki, T.; Hirata, Y. *Experientia* 1983, 39, 67-69.
- (60) Kobayashi, J.; Ishibashi, M.; Nakamura, H.; Ohizumi, Y.; Hirata, Y. *J. Chem. Soc., Perkin Trans. 1* 1989, 101-103.
- (61) Seino, A.; Kobayashi, M.; Kobayashi, J.; Fang, Y.-L.; Ishibashi, M.; Nakamura, H.; Momose, K.; Ohizumi, Y. *J. Pharmacol. Exp. Ther.* 1991, 256, 861-867.
- (62) Ohizumi, Y.; Ishida, Y.; Shibata, S. *J. Pharmacol. Exp. Ther.* 1982, 221, 748-752.
- (63) Schmitz, F. J.; Agarwal, S. K.; Gunasekera, S. P.; Schmidt, P. G.; Shooley, J. N. *J. Am. Chem. Soc.* 1983, 105, 4835-4836.
- (64) Molinsky, T. F.; Fahy, E.; Faulkner, D. J.; Van Dwyne, G. D.; Clardy, J. *J. Org. Chem.* 1988, 53, 1340-1341.
- (65) Gunawardana, G. P.; Kohmoto, S.; Gunasekera, S. P.; McConnell, O. J.; Koehn, F. E. *J. Am. Chem. Soc.* 1988, 110, 4856-4858.
- (66) Gunawardana, G. P.; Koehn, F. E.; Lee, A. Y.; Clardy, J.; He, H.-Y.; Faulkner, D. J. *J. Org. Chem.* 1992, 57, 1523-1526.
- (67) Kobayashi, J.; Cheng, J.-F.; Nakamura, H.; Ohizumi, Y.; Hirata, Y.; Sasaki, T.; Ohta, T.; Nozoe, S. *Tetrahedron Lett.* 1988, 29, 1177-1180.
- (68) Kobayashi, J.; Cheng, J.-F.; Wälchli, M. R.; Nakamura, H.; Hirata, Y.; Sasaki, T.; Ohizumi, Y. *J. Org. Chem.* 1988, 53, 1800-1804.
- (69) Kobayashi, J.; Tsuda, M.; Tanabe, A.; Ishibashi, M.; Cheng, J.-F.; Yamamura, S.; Sasaki, T. *J. Nat. Prod.* 1991, 54, 1634-1638.
- (70) Rudi, A.; Benayahu, Y.; Goldberg, L.; Kashman, Y. *Tetrahedron Lett.* 1988, 29, 3861-3864.
- (71) Rudi, A.; Kashman, Y. *J. Org. Chem.* 1989, 54, 5331-5337.
- (72) Cimino, G.; Crispino, S.; De Rosa, S.; De Stefano, S.; Gavagnin, M.; Sodano, G. *Tetrahedron* 1987, 43, 4023-4030.
- (73) Schmitz, F. J.; DeGuzman, F. S.; Hossain, M. B.; van der Helm, D. *J. Org. Chem.* 1991, 56, 804-808.
- (74) Copp, B. R.; Ireland, C. M.; Barrows, L. R. *J. Org. Chem.* 1991, 56, 4596-4597.
- (75) Perry, N. B.; Blunt, J. W.; McCombs, J. D.; Munro, M. H. G. *J. Org. Chem.* 1986, 51, 5476-5478.
- (76) Perry, N. B.; Blunt, J. W.; McCombs, J. D.; Munro, M. H. G. *Tetrahedron* 1988, 53, 1727-1734.
- (77) Perry, N. B.; Blunt, J. W.; Munro, M. H. G.; Higa, T.; Sakai, R. *J. Org. Chem.* 1988, 53, 4127-4128.
- (78) Kobayashi, J.; Cheng, J.-F.; Ishibashi, M.; Nakamura, H.; Ohizumi, Y.; Hirata, Y.; Sasaki, T.; Lu, H.; Clardy, J. *Tetrahedron Lett.* 1987, 28, 4939-4942.
- (79) Cheng, J.-F.; Ohizumi, Y.; Wälchli, M. R.; Nakamura, H.; Hirata, Y.; Sasaki, T.; Kobayashi, J. *J. Org. Chem.* 1988, 53, 4621-4624.
- (80) Kobayashi, J.; Cheng, J.-F.; Yamamura, S.; Ishibashi, M. *Tetrahedron Lett.* 1991, 32, 1227-1228.
- (81) Frincke, J. M.; Faulkner, D. J. *J. Am. Chem. Soc.* 1982, 104, 265-269.
- (82) He, H.-Y.; Faulkner, D. J. *J. Org. Chem.* 1989, 54, 5822-5824.
- (83) McKee, T. C.; Ireland, C. M. *J. Nat. Prod.* 1987, 50, 754-756.
- (84) Wright, A. E.; Forleo, D. A.; Gunawardana, G. P.; Gunasekera, S. P.; Koehn, F. E.; McConnell, O. J. *J. Org. Chem.* 1990, 55, 4508-4512.
- (85) Rinehart, K. L.; Holt, T. G.; Fregeau, N. L.; Stroh, J. G.; Kiefer, P. A.; Sun, F.; Li, L. H.; Martin, D. G. *J. Org. Chem.* 1990, 55, 4512-4515.
- (86) Arai, T.; Yazawa, K.; Mikami, Y. *J. Antibiot.* 1976, 29, 398-407.
- (87) Arai, T.; Takahashi, K.; Kubo, A.; Nakahara, S.; Sato, S.; Aiba, K.; Tamura, C. *Tetrahedron Lett.* 1979, 2355-2358.
- (88) Perry, N. B.; Blunt, J. W.; Munro, M. H. G.; Pannell, L. K. *J. Am. Chem. Soc.* 1988, 110, 4850-4851.
- (89) Perry, N. B.; Blunt, J. W.; Munro, M. H. G.; Thompson, A. M. *J. Org. Chem.* 1990, 55, 223-227.
- (90) Sakemi, S.; Ichiba, T.; Kohmoto, S.; Saucy, G.; Higa, T. *J. Am. Chem. Soc.* 1988, 110, 4851-4853.
- (91) Matsunaga, S.; Fusetani, N.; Nakao, Y. *Tetrahedron* 1992, 48, 8369-8376.
- (92) Fusetani, N.; Sugawara, T.; Matsunaga, S. *J. Org. Chem.* 1992, 57, 3828-3832.

- (93) Furusaki, A.; Watanabe, T.; Matsumoto, T.; Yanagiya, M. *Tetrahedron Lett.* **1968**, 6301-6304.
- (94) Carmeli, S.; Kashman, Y. *Tetrahedron Lett.* **1985**, 26, 511-514.
- (95) Kitagawa, I.; Kobayashi, M.; Katori, T.; Yamashita, M.; Tanaka, J.; Doi, M.; Ishida, T. *J. Am. Chem. Soc.* **1990**, 112, 3710-3712.
- (96) Doi, M.; Ishida, T.; Kobayashi, M.; Kitagawa, I. *J. Org. Chem.* **1991**, 56, 3629-3632.
- (97) Kobayashi, M.; Tanaka, J.; Katori, T.; Matsuura, M.; Yamashita, M.; Kitagawa, I. *Chem. Pharm. Bull.* **1990**, 38, 2409-2418.
- (98) Tsukamoto, S.; Ishibashi, M.; Sasaki, T.; Kobayashi, J. *J. Chem. Soc., Perkin Trans. 1* **1991**, 3185-3188.
- (99) Sakai, R.; Higa, T.; Kashman, Y. *Chem. Lett.* **1986**, 1499-1502.
- (100) Kato, Y.; Fusetani, N.; Matsunaga, S.; Hashimoto, K.; Sakai, R.; Higa, T.; Kashman, Y. *Tetrahedron Lett.* **1987**, 28, 6225-6228.
- (101) Tanaka, J.; Higa, T.; Kobayashi, M.; Kitagawa, I. *Chem. Pharm. Bull.* **1990**, 38, 2967-2970.
- (102) Kobayashi, J.; Tsukamoto, S.; Tanabe, A.; Sasaki, T.; Ishibashi, M. *J. Chem. Soc., Perkin Trans. 1* **1991**, 2379-2383.
- (103) Roesener, J. A.; Scheuer, P. J. *J. Am. Chem. Soc.* **1986**, 108, 846-847.
- (104) Matsunaga, S.; Fusetani, N.; Hashimoto, K.; Koseki, K.; Noma, M. *J. Am. Chem. Soc.* **1986**, 108, 847-849.
- (105) Matsunaga, S.; Fusetani, N.; Hashimoto, K.; Koseki, K.; Noma, M.; Noguchi, H.; Sankawa, U. *J. Org. Chem.* **1989**, 54, 1360-1363.
- (106) Kernan, M. R.; Faulkner, D. J. *Tetrahedron Lett.* **1987**, 28, 2809-2812.
- (107) Kernan, M. R.; Molinski, T. F.; Faulkner, D. J. *J. Org. Chem.* **1988**, 53, 5014-5020.
- (108) Ishibashi, M.; Moore, R. E.; Patterson, G. M. L.; Xu, C.; Clardy, J. *J. Org. Chem.* **1986**, 51, 5300-5306.
- (109) Carmeli, S.; Moore, R. E.; Patterson, G. M. L. *J. Nat. Prod.* **1990**, 53, 1533-1542.
- (110) Kobayashi, J.; Cheng, J.-F.; Ohta, T.; Nakamura, H.; Nozoe, S.; Hirata, Y.; Ohizumi, Y.; Sasaki, T. *J. Org. Chem.* **1988**, 53, 6147-6150.
- (111) Kikuchi, Y.; Ishibashi, M.; Sasaki, T.; Kobayashi, J. *Tetrahedron Lett.* **1991**, 32, 797-798.
- (112) Kitagawa, I.; Kobayashi, M.; Lee, N. K.; Shibuya, H.; Kawata, Y.; Sakiyama, F. *Chem. Pharm. Bull.* **1986**, 34, 2664.
- (113) Nakamura, H.; Kobayashi, J.; Nakamura, Y.; Ohizumi, Y.; Kondo, T.; Hirata, Y. *Tetrahedron Lett.* **1986**, 27, 4319-4322.
- (114) Kitagawa, I.; Lee, N. K.; Kobayashi, M.; Shibuya, H. *Chem. Pharm. Bull.* **1987**, 35, 2129.
- (115) Kitagawa, I.; Lee, N. K.; Kobayashi, M.; Shibuya, H. *Tetrahedron* **1991**, 47, 2169-2180.
- (116) Kitagawa, I.; Lee, N. K.; Kobayashi, M.; Shibuya, H. *Chem. Pharm. Bull.* **1991**, 39, 1177-1184.
- (117) Kobayashi, J.; Sato, M.; Murayama, T.; Ishibashi, M.; Wälchli, M. R.; Kanai, M.; Shoji, J.; Ohizumi, Y. *J. Chem. Soc., Chem. Commun.* **1991**, 1050-1052.
- (118) Kobayashi, J.; Sato, M.; Ishibashi, M.; Shigemori, H.; Nakamura, T.; Ohizumi, Y. *J. Chem. Soc., Perkin Trans. 1* **1991**, 2609-2611.
- (119) Isono, F.; Inukai, M.; Takahashi, S.; Haneishi, T.; Kinoshita, T.; Kuwano, H. *J. Antibiot.* **1989**, 42, 667.
- (120) Kobayashi, J.; Itagaki, F.; Shigemori, H.; Ishibashi, M.; Takahashi, K.; Ogura, M.; Nagasawa, S.; Nakamura, T.; Hirota, H.; Ohta, T.; Nozoe, S. *J. Am. Chem. Soc.* **1991**, 113, 7812-7813.
- (121) Itagaki, F.; Shigemori, H.; Ishibashi, M.; Nakamura, T.; Sasaki, T.; Kobayashi, J. *J. Org. Chem.* **1992**, 57, 5540-5542.