

LETTER TO THE EDITOR

A Unified Nomenclature System for the Insect Olfactory Coreceptor

Leslie B. Vosshall¹ and Bill S. Hansson²¹Laboratory of Neurogenetics and Behavior, Howard Hughes Medical Institute, The Rockefeller University, New York, NY, USA and ²Department of Evolutionary Neuroethology, Max Planck Institute for Chemical Ecology, Hans-Knoell-Strasse 8, D-07745 Jena, Germany

Correspondence to be sent to: Leslie B. Vosshall, Laboratory of Neurogenetics and Behavior, Howard Hughes Medical Institute, The Rockefeller University, 1230 York Avenue, Box 63, New York, NY, 10065, USA. e-mail: leslie.vosshall@rockefeller.edu

Accepted February 8, 2011

We propose a revised nomenclature for the insect olfactory coreceptor variously named “*Or83b*,” “*Or1*,” “*Or2*,” and “*Or7*.” This gene appears to be functionally orthologous across all insects and serves as a chaperoning coreceptor for the odor- and pheromone-specific subunits of the insect olfactory receptor gene family (Krieger et al. 2003; Larsson et al. 2004; Pitts et al. 2004; Nakagawa et al. 2005; Benton et al. 2006). The insect odorant receptors (*Ors*) are atypical 7-transmembrane domain proteins that form ligand-gated ion channels by assembling a ligand-selective subunit with the olfactory coreceptor (Nakagawa et al. 2005; Sato et al. 2008; Wicher et al. 2008). The *Or* gene family appears to be an insect-specific adaptation (Robertson et al. 2003) as neither ligand-selective *Ors* nor the olfactory coreceptor have been identified in any noninsect genomes (Penalva-Arana et al. 2009).

Although the insect olfactory coreceptor appears to perform the same function in all insects, there is no consistency in nomenclature across species. The current nomenclature system with different names in different insect species is confusing, unnecessary, and negatively impacts our ability to communicate our work to nonspecialists. The confusion is compounded by ambiguity in gene names: *Or2* and *Or7* are not unique names but names that vary in function across species. For instance, *Or2* in *Bombyx mori* refers to the olfactory coreceptor, whereas in *Anopheles gambiae*, *Or2* refers to a 2-methylphenol receptor. *Or7* is the coreceptor in *Anopheles* and *Aedes* but is an odor-specific receptor in various Lepidopteran species.

To avoid further confusion in the literature, we propose a new and unified nomenclature that abandons all the old names (*Or83b*, *Or1*, *Or2*, and *Or7*) and uses a single new name based on the consensus function of this protein across all insect species: “*Orco*,” short for “olfactory receptor co-

receptor.” The insect species will be signified using the 4-letter convention adopted by FlyBase, in which the first letter of the genus and first 3 letters of the species are followed by *Orco*. A backslash separates the species abbreviation and *Orco* to enhance legibility and facilitate computational data mining.

Some examples of the new gene names with 4-letter species code are

<i>Drosophila melanogaster</i>	<i>Dme\Orco</i>
<i>Drosophila pseudoobscura</i>	<i>Dpse\Orco</i>
<i>Anopheles gambiae</i>	<i>Agam\Orco</i>
<i>Aedes aegypti</i>	<i>Aaeg\Orco</i>
<i>Bombyx mori</i>	<i>Bmor\Orco</i>
<i>Tribolium castaneum</i>	<i>Tcas\Orco</i>

All other insect species in which *Orco* has been identified should use the same 4-letter species coding system for assigning the relevant gene name. The use of a 2-letter species code (e.g., “Ag” for *A. gambiae* and “Aa” for *A. aegypti*, etc.) is strongly discouraged because it is ambiguous.

When new insect genomes become available, the following criteria should be applied to call a gene “*Orco*”:

Orco is a member of the insect *Or* superfamily that shows a high degree of conservation across insect species. *Orco* will typically be at least 50% identical to *Orco* in other insect species.

Orco messenger RNA and protein is expressed in a majority of chemosensory neurons.

Orco has a larger predicted protein size due to an insertion in intracellular loop 2 relative to conventional *Ors* such that the full-length *Orco* protein will be approximately 475

amino acids, relative to ligand-specific Ors, which will be approximately 400 amino acids in length.

The insect olfaction field faced a similar issue in 1999 when 3 different groups, Axel (Vosshall et al. 1999), Carlson (Clyne et al. 1999), and Chess (Gao and Chess 1999), identified the first insect OR genes in *Drosophila* and gave them different names. This created great confusion and was a major barrier to the continued growth of this field as every conversation required a complicated look-up table to compare the various gene names being used for the same locus. To deal with this problem, a unified nomenclature was proposed and approved by members of the field (*Drosophila Odorant Receptor Nomenclature Committee* 2000) and has completely replaced the former names.

We recommend that the unified “*Orco*” nomenclature be used in future publications *(for further details, see Acknowledgements section).

Acknowledgements

In addition to the authors of this letter, the *Orco* nomenclature has been endorsed by the following scientists and organizations: William Gelbart, FlyBase (gelbart@morgan.harvard.edu); Matthew Cobb, University of Manchester, UK (cobb@manchester.ac.uk); Laurence J. Zwiebel, Vanderbilt University, USA (l.zwiebel@vanderbilt.edu); Jürgen Krieger, University Hohenheim, Germany (krieger@uni-hohenheim.de); John Carlson, Yale University, USA (john.carlson@yale.edu); Emmanuelle Jacquin-Joly, INRA Centre de Versailles, France (emmanuelle.jacquin@versailles.inra.fr); Rickard Ignell, SLU, Sweden (Rickard.Ignell@ltj.slu.se); Marien de Bruyne, Monash University, Australia (Marien.DeBrujne@sci.monash.edu.au); C. Giovanni Galizia, Universität Konstanz, Germany (galizia@uni-konstanz.de); Richard Newcomb, The New Zealand Institute for Plant & Food Research, New Zealand (Richard.Newcomb@plantandfood.co.nz); Kristin Scott, HHMI/University of California, Berkeley, USA (kscott@berkeley.edu); Rachel Wilson, Harvard Medical School, USA (rachel_wilson@hms.harvard.edu); Richard Benton, University of Lausanne, Switzerland (Richard.Benton@unil.ch); Richard Vogt, University of South Carolina, USA (vogt@biol.sc.edu); Walter Leal, University of California, Davis, USA (wsleal@ucdavis.edu); Bertram Gerber, University of Leipzig, Germany (bertram.gerber@uni-leipzig.de); Mattias Larsson, SLU, Sweden (mattias.larsson@ltj.slu.se); Kazushige Touhara, University of Tokyo, Japan (ktouhara@mail.ecc.u-tokyo.ac.jp); Gaiti Hasan, NCBS, India (gaiti@ncbs.res.in); Dean Smith, UTSW Medical Center, USA (Dean.Smith@UTSouthwestern.edu); Walton D. Jones, KAIST Biological Sciences, South Korea (waltonjones@kaist.edu); Craig Montell, Johns Hopkins University (cmontell@jhmi.edu); Coral Warr, Monash University, Australia (coral.warr@monash.edu); Liquan Luo, Stanford University, USA (lluo@stanford.edu); Hugh Robertson, University of Illinois Champaign-Urbana, USA

(hughrobe@life.illinois.edu); Frederic Marion-Poll, INRA Centre de Versailles, France (frederic.marion-poll@versailles.inra.fr); Marcus Stensmyr, Max Planck Institute for Chemical Ecology, Germany (mstensmyr@ice.mpg.de); Matthieu Louis, CRG Barcelona, Spain (matthieu.louis@crg.es); Anandasankar Ray, University of California Riverside, USA (anandr@ucr.edu); Scott Waddell, UMass Medical School, USA (Scott.Waddell@umassmed.edu); and Charles W. Luetje, University of Miami Miller School of Medicine, USA (cluetje@med.miami.edu).

References

Benton R, Sachse S, Michnick SW, Vosshall LB. 2006. Atypical membrane topology and heteromeric function of *Drosophila* odorant receptors *in vivo*. *PLoS Biol.* 4:e20.

Clyne PJ, Warr CG, Freeman MR, Lessing D, Kim J, Carlson JR. 1999. A novel family of divergent seven-transmembrane proteins: candidate odorant receptors in *Drosophila*. *Neuron.* 22:327–338.

Drosophila Odorant Receptor Nomenclature Committee. 2000. A unified nomenclature system for the *Drosophila* odorant receptors. *Cell.* 102: 145–146.

Gao Q, Chess A. 1999. Identification of candidate *Drosophila* olfactory receptors from genomic DNA sequence. *Genomics.* 60:31–39.

Krieger J, Klink O, Mohl C, Raming K, Breer H. 2003. A candidate olfactory receptor subtype highly conserved across different insect orders. *J Comp Physiol A.* 189:519–526.

Larsson MC, Domingos AI, Jones WD, Chiappe ME, Amrein H, Vosshall LB. 2004. *Or83b* encodes a broadly expressed odorant receptor essential for *Drosophila* olfaction. *Neuron.* 43:703–714.

Nakagawa T, Sakurai T, Nishioka T, Touhara K. 2005. Insect sex-pheromone signals mediated by specific combinations of olfactory receptors. *Science.* 307:1638–1642.

Penalva-Arana DC, Lynch M, Robertson HM. 2009. The chemoreceptor genes of the waterflea *Daphnia pulex*: many Grs but no Ors. *BMC Evol Biol.* 9:79.

Pitts RJ, Fox AN, Zwiebel LJ. 2004. A highly conserved candidate chemoreceptor expressed in both olfactory and gustatory tissues in the malaria vector *Anopheles gambiae*. *Proc Natl Acad Sci U S A.* 101: 5058–5063.

Robertson HM, Warr CG, Carlson JR. 2003. Molecular evolution of the insect chemoreceptor gene superfamily in *Drosophila melanogaster*. *Proc Natl Acad Sci U S A.* (100 Suppl)2:14537–14542.

Sato K, Pellegrino M, Nakagawa T, Nakagawa T, Vosshall LB, Touhara K. 2008. Insect olfactory receptors are heteromeric ligand-gated ion channels. *Nature.* 452:1002–1006.

Vosshall LB, Amrein H, Morozov PS, Rzhetsky A, Axel R. 1999. A spatial map of olfactory receptor expression in the *Drosophila* antenna. *Cell.* 96: 725–736.

Wicher D, Schafer R, Bauernfeind R, Stensmyr MC, Heller R, Heinemann SH, Hansson BS. 2008. *Drosophila* odorant receptors are both ligand-gated and cyclic-nucleotide-activated cation channels. *Nature.* 452:1007–1011.