

Reduction of 1,4-dichlorobut-2-yne by titanocene to a 1,2,3-butatriene. Formation of a 1-titanacyclopent-3-yne and a 2,5-dititanabicyclo[2.2.0]hex-1(4)-ene

Vladimir V. Burlakov,^a Perdita Arndt,^b Wolfgang Baumann,^b Anke Spannenberg,^b Uwe Rosenthal,^{*b} Pattiyl Parameswaran^c and Eluvathingal D. Jemmis^{*c}

^a A. N. Nesmeyanov Institute of Organoelement Compounds, Russian Academy of Sciences, Vavilov St. 28, 117813, Moscow, Russia. Fax: +7 0951355085; Tel: +7 0951359252; E-mail: vvburl@ineos.ac.ru

^b Leibniz-Institut für Organische Katalyse an der Universität Rostock e.V., Buchbinderstr. 5-6, D-18055 Rostock, Germany. Fax: +49 3814669376; Tel: (int. +49)3814669376; E-mail: uwe.rosenthal@ifok.uni-rostock.de

^c School of Chemistry, University of Hyderabad, Hyderabad 500 046, India. Fax: +91 4023010917; Tel: +91 4023010221; E-mail: edjsc@uohyd.ernet.in

Received (in Cambridge, UK) 4th May 2004, Accepted 2nd July 2004

First published as an Advance Article on the web 11th August 2004

The 2,5-dititanabicyclo[2.2.0]hex-1(4)-ene (bis-titanocene- μ -(*Z*)-1,2,3-butatriene complex) (**3**) is formed starting from $[\text{Cp}_2\text{Ti}(\eta^2\text{-Me}_3\text{SiC}_2\text{SiMe}_3)]$ by *in situ* generated titanocene and 1,4-dichlorobut-2-yne via the 1-titanacyclobut-3-yne (**2**).


The complexation of titanocene with butatriene was described by Maercker, but the suggested structures resulting from $\text{Me}_2\text{C}=\text{C}=\text{C}=\text{Me}_2$ and “ Cp_2Ti ” gave further reactions with CH-activation to yield more stable complexes.¹ Suzuki and coworkers recently reported 1-zirconacyclopent-3-yne, formed from a divalent zirconocene-equivalent “ Cp_2Zr ” (generated by the Negishi reagent, $[\text{Cp}_2\text{ZrCl}_2]$ and 2 equivalents of *n*-BuLi) with 1,4-disubstituted (*Z*)-butatrienes $\text{RHC}=\text{C}=\text{C}=\text{CHR}$ ($\text{R} = \text{Me}_3\text{Si}$, *t*-Bu).² In the reaction mixture of $[\text{Cp}_2\text{ZrCl}_2]$, 1,4-dichlorobut-2-yne and 2 equivalents of magnesium, even the coordination of an unsubstituted butatriene $\text{H}_2\text{C}=\text{C}=\text{C}=\text{CH}_2$ with the “ Cp_2Zr ” formed, to give 1-zirconacyclopent-3-yne was realized.³ Closely related 1-metallacyclopenta-2,3,4-trienes (five-membered metallacyclocumulenes)⁴ were obtained in reactions of 1,3-butadiynes $\text{RC}\equiv\text{C}\equiv\text{CR}$ by using the excellent metallocene sources $[\text{Cp}_2\text{M}(\eta^2\text{-Me}_3\text{SiC}_2\text{SiMe}_3)]$.⁵ Both types of rather exotic metallacyclopent-3-yne were discussed and compared by calculations⁶ and prompted us to report here on the first results of reactions of $[\text{Cp}_2\text{Ti}(\eta^2\text{-Me}_3\text{SiC}_2\text{SiMe}_3)]$ with $\text{ClCH}_2\text{C}\equiv\text{CCH}_2\text{Cl}$.

Compound **2** is formed by the reaction of two equivalents of $[\text{Cp}_2\text{Ti}(\eta^2\text{-Me}_3\text{SiC}_2\text{SiMe}_3)]$ (**1**) with $\text{ClCH}_2\text{C}\equiv\text{CCH}_2\text{Cl}$ together with the liberation of $\text{Me}_3\text{SiC}_2\text{SiMe}_3$ and $[\text{Cp}_2\text{TiCl}_2]$ (Scheme 1).^{†‡}

The composition of the diamagnetic complex **2** was verified by analytical and spectral data and by reaction with a second “ Cp_2Ti ”. The ¹H NMR (δ (CH_2) 3.03) and ¹³C{¹H} NMR signals (δ (CH_2): 51.2; ($\text{C}\equiv\text{C}$) 106.9 ppm and the IR data (ν ($\text{C}\equiv\text{C}$) 2029 cm^{-1}) of complex **2** correspond very well to those of the analogous zirconium complex (¹H NMR: δ (CH_2) 2.73; ¹³C{¹H} NMR: δ (CH_2): 38.64; ($\text{C}\equiv\text{C}$) 102.45 ppm and IR (ν ($\text{C}\equiv\text{C}$) 2018 cm^{-1}).³

The molecular structure of **2** represents a resonance structure between a 1-titanacyclopent-3-yne $[\text{Cp}_2\text{Ti}(\eta^2\text{-1,2,3,4-CH}_2\text{C}\equiv\text{CCH}_2)]$ and a titanocene- η^4 -(*E*)-butatriene complex $[\text{Cp}_2\text{Ti}(\eta^4\text{-}(E)\text{-H}_2\text{C}=\text{C}=\text{C}=\text{CH}_2)]$ as discussed before for the analogous zirconium complex,^{3,6} containing chelating σ -propargyl or π -allenyl structural elements.⁷

Complex **2** reacts with $[\text{Cp}_2\text{Ti}(\eta^2\text{-Me}_3\text{SiC}_2\text{SiMe}_3)]$ ⁵ to form



Scheme 1 Preparation of complexes.

complex **3** which can also be obtained by a 3 : 1 reaction of $[\text{Cp}_2\text{Ti}(\eta^2\text{-Me}_3\text{SiC}_2\text{SiMe}_3)]$ with $\text{ClCH}_2\text{C}\equiv\text{CCH}_2\text{Cl}$ (Scheme 1).^{§¶}

The X-ray crystal structure analysis of **3**[¶] (Fig. 1) revealed two bent titanocenes which are bridged symmetrically by a “zig-zag” C_4 -ligand. The four carbon and two titanium atoms are in a plane with a mean deviation of 0.0017 Å. The central bond of the C_4 -ligand is coordinated to both titanium centers, unsymmetrically for each titanium atom [$\text{C}1\text{-C}(1\text{A})$ 1.325(5); $\text{C}1\text{-Ti}1$ 2.152(3); $\text{C}(1\text{A})\text{-Ti}1$ 2.268(3) Å], whereas each of the methylene groups is coordinated to only one Ti atom [$\text{C}2\text{-Ti}1$ 2.167(3); $\text{C}1\text{-C}(2\text{A})$ 1.443(4) Å]. By this coordination a bonding mode is formed with a $\text{C}1\text{-C}(1\text{A})\text{-C}2$ angle of 135.1(3)[°].

The molecular structure of **3** is represented by the resonance structures of a butynediyl-bridged σ -propargylic complex (**A**) and a butatriene-bridged π -complex (**B**), but in contrast to complex **2** here in a bridging mode (Scheme 2).⁶ Despite that, the structure of **3** as a μ -*trans*-butatriene complex is best described as a dititanabicyclo (**C**) in analogy with μ -*trans*-butadiyne complexes (**D**), formed by analogous titanocene complexation of 1-titanacyclopenta-2,3,4-trienes (five-membered titanacyclocumulenes)⁴ (Scheme 3).⁵ **3** is not a σ -propargyl or a π -allenyl complex^{6,7} and is also different from the “bridged allylic” structures found in μ -butatriene-bis(tricarbonyliron) complexes^{8a} or substituted butatriene dianion dilithium,^{8b} μ -butyne-1,4-diyl^{8b} structures as well as μ -butadiyne complexes with μ -(η^2, η^2)-(PhC=C=C=CPh) units.^{8c}






Fig. 1 Molecular structure of complex **3**. Hydrogen atoms except the H-atoms of the C_4 -ligand are omitted for clarity. The thermal ellipsoids correspond to 30% probability.



Scheme 2 Resonance structures of complex **3**.



Scheme 3 Formation of μ -butadiyne complexes (**D**).

We have studied the structure and bonding of these molecules using Density Functional Theory calculations (B3LYP/LANL2DZ).⁹ The calculated geometrical parameters are in close agreement with the experimental structure. The bonding in **3** is best described by treating the bridging ligand as formally $[\text{H}_2\text{CCCCH}_2]^{(-4)}$ species, making Ti(+4). The C1–C(1A) π bond perpendicular to the TiC_4Ti plane does not interact substantially with the metals. The remaining eight valence electrons of the $[\text{H}_2\text{CCCCH}_2]^{(-4)}$ ligand occupy four in-plane delocalized orbitals resulting from the interaction with the Cp_2Ti fragment orbitals. The bonding here is very similar to that in the μ -*trans*-butadiyne complex $[\text{Cp}_2\text{Ti}(\text{HCCCCH})\text{TiCp}_2]$ (type **D**) except that **3** has an ethylenic π bond in place of the *trans*-butadiene of the butadiyne complex.¹⁰ A C_{2v} isomer of **3** derived directly from the complexation of the middle C1–C(1A) bond of **2** is calculated to be higher in energy by 9.00 kcal mol⁻¹. Experimental and theoretical studies on the details of this species, its conversion to **3**, and further transformations of **3** are currently in progress.

This work was supported by the SPP 1118 of the Deutsche Forschungsgemeinschaft (RO 1269/5-1) and the Russian Foundation for Basic Research (Project code 02-03-32589).

Notes and references

† General procedure for the preparation of complex **2**: complex **1** (2.040 g, 5.85 mmol) was dissolved in *n*-hexane (20 mL) under Ar. The resulting yellow-brown solution was filtered, and $\text{ClCH}_2\text{C}=\text{CCH}_2\text{Cl}$ (0.286 mL, 2.93 mmol) was added to the resulting solution under stirring. The solution rapidly became brown and a dark-red precipitate of $[\text{Cp}_2\text{TiCl}_2]$ was formed. The mixture was allowed to stand in an argon atmosphere at 20 °C. After 24 h the solution was filtered and evaporated to 10 ml under vacuum. Upon cooling to -78 °C for 1 day, brown crystals were formed, which were separated from the mother liquor by decanting, and washed with a small amount of cold *n*-hexane and dried under vacuum. Yield of **2** was 0.454 g (65%), mp 211–212 °C (dec. at slow heating (3 °C per min); at fast heating (20 °C per min) blows up at ca. 145–150 °C) under Ar.

‡ Data for **2**: elemental analysis calcd for $\text{C}_{14}\text{H}_{14}\text{Ti}$: C, 73.07; H, 6.13. Found: C, 72.43; H 6.19%. ¹H NMR (C_6D_6 , 297 K): δ 3.03 (s, 4H, CH_2); 4.68 (s, 10H, Cp). ¹³C{¹H} NMR (C_6D_6 , 297 K): δ 51.2 (CH_2); 102.4 (Cp); 106.9 (C=C). IR (Nujol mull, cm⁻¹): 2029 (weak, $\nu\text{C}=\text{C}$). MS (70 eV, *m/z*): 230 $[\text{M}]^+$, 178 $[\text{Cp}_2\text{Ti}]^+$, 113 $[\text{CpTi}]^+$.

§ General procedure for the preparation of complex **3**: Complex **1** (0.486 g, 1.38 mmol) was dissolved in toluene (7–8 mL) under Ar. The obtained

solution was filtered and added gradually to a filtrated brown solution of **2** (0.298 g, 1.29 mmol) in 7–8 mL of toluene. The resulting solution rapidly turned green and crystals of **3** appeared on the bottom and walls of the vessel. After 24 h the solution was decanted. Subsequent washing of the dark green crystals with cold toluene and drying in vacuum gave 0.485 g (92%) of **3**, mp 220–222 °C (dec.) under Ar.

¶ Data for **3**: elemental analysis calcd for $\text{C}_{24}\text{H}_{24}\text{Ti}_2$: C, 70.62; H, 5.93. Found: C, 70.14; H 5.88%. ¹H NMR (C_6D_6 , 297 K): δ 3.97 (br., 4H, CH_2); 5.18 (s, 20H, Cp). ¹³C{¹H} NMR (C_6D_6 , 297 K): δ 85.5 (CH_2); 108.5 (Cp); 153.4 (C=C). MS (70 eV, *m/z*): 408 $[\text{M}]^+$, 406 $[\text{M} - 2\text{H}]^+$, 352 $[\text{M} - \text{C}_4\text{H}_8]^+$, 178 $[\text{Cp}_2\text{Ti}]^+$.

|| X-Ray crystal structure analysis of **3**: STOE-IPDS diffractometer, graphite monochromated MoK α radiation, solution of the structure by direct methods (SHELXS-86¹¹), refinement with full-matrix least-squares techniques against F^2 (SHELXL-93¹²). Crystal data: monoclinic, space group $P2_1/n$, $a = 8.687(2)$, $b = 7.887(2)$, $c = 13.353(3)$ Å; $\beta = 90.17(3)^\circ$; $V = 914.9(4)$ Å³, $Z = 2$, $D_c = 1.482$ g cm⁻³; 2621 reflections measured, 1429 were independent of symmetry and 1221 were observed [$I > 2\sigma(I)$], $R1 = 0.036$, $wR2(\text{all data}) = 0.096$, 126 parameters. CCDC 239591. See <http://www.rsc.org/suppdata/cc/b4/b406494a/> for crystallographic data in .cif or other electronic format.

- 1 A. Maercker and A. Groos, *Angew. Chem.*, 1996, **108**, 216, *Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 210.
- 2 N. Suzuki, M. Nishiura and Y. Wakatsuki, *Science*, 2002, **295**, 660.
- 3 N. Suzuki, N. Ahihara, H. Takahara, T. Watanabe, M. Iwasaki, M. Saburi, D. Hashizume and T. Chihara, *J. Am. Chem. Soc.*, 2004, **126**, 60.
- 4 U. Rosenthal, P.-M. Pellny, F. G. Kirchbauer and V. V. Burlakov, *Acc. Chem. Res.*, 2000, **33**, 119.
- 5 (a) U. Rosenthal and V. V. Burlakov, in *Titanium and Zirconium in Organic Synthesis*, ed. I. Marek, Wiley-VCH, Weinheim, 2002, p. 355; (b) U. Rosenthal, V. V. Burlakov, P. Arndt, W. Baumann and A. Spannenberg, *Organometallics*, 2003, **22**, 884.
- 6 (a) K. C. Lam and Z. Lin, *Organometallics*, 2003, **22**, 3466; (b) E. D. Jemmis, A. K. Phukan, H. Jiao and U. Rosenthal, *Organometallics*, 2003, **22**, 4958; (c) U. Rosenthal, *Angew. Chem.*, 2004, **116**, 3972, *Angew. Chem., Int. Ed. Engl.* 2004, **43**, 3882.
- 7 (a) P. W. Blosser, J. C. Gallucci and A. Wojcicki, *J. Am. Chem. Soc.*, 1993, **115**, 2994; (b) P. W. Blosser, J. C. Gallucci and A. Wojcicki, *J. Organomet. Chem.*, 2000, **597**, 125.
- 8 (a) J. N. Gerlach, R. M. Wing and P. C. Ellgen, *Inorg. Chem.*, 1976, **15**, 2959; (b) T. Matsuo, M. Tanaka and A. Sekiguchi, *Chem. Commun.*, 2001, 503 and references cited therein; (c) Y. Wang, H. Wang, H. Wang, H.-S. Chan and Z. Xie, *J. Organomet. Chem.*, 2003, **683**, 39.
- 9 (a) A. D. Becke, *J. Chem. Phys.*, 1993, **98**, 5648; (b) A. D. Becke, *Phys. Rev. A*, 1998, **38**, 2398; (c) C. Lee, W. Yang and R. G. Parr, *Phys. Rev. B*, 1988, **37**, 785; (d) P. J. Hay and W. R. Wadt, *J. Chem. Phys.*, 1985, **82**, 299; (e) Gaussian 03, Revision B.03, M. J. Frisch, *et al.*, Gaussian, Inc., Pittsburgh, PA, 2003.
- 10 (a) E. D. Jemmis and K. T. Giju, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 606; (b) E. D. Jemmis and K. T. Giju, *J. Am. Chem. Soc.*, 1998, **120**, 6952; (c) P. N. V. Pavan Kumar and E. D. Jemmis, *J. Am. Chem. Soc.*, 1988, **110**, 125.
- 11 G. M. Sheldrick, *Acta. Crystallogr., Sect. A*, 1990, **46**, 467.
- 12 SHELXL-93, G. M. Sheldrick, University of Göttingen, Germany, 1993.