

An active ferrocenyl triarylphosphine for palladium-catalyzed Suzuki–Miyaura cross-coupling of aryl halides†

Fuk Yee Kwong,^{*a} Kin Shing Chan,^b Chi Hung Yeung^a and Albert S. C. Chan^{*a}

^a Open Laboratory of Chirotechnology of the Institute of Molecular Technology for Drug Discovery and Synthesis, Department of Applied Biology and Chemical Technology, The Hong Kong Polytechnic University, Hung Hom, Hong Kong. E-mail: bcfyk@inet.polyu.edu.hk; bcachan@polyu.edu.hk

^b Department of Chemistry, The Chinese University of Hong Kong, Shatin, New Territories, Hong Kong

Received (in Cambridge, UK) 24th May 2004, Accepted 29th July 2004


First published as an Advance Article on the web 21st September 2004

Pd-catalyzed Suzuki–Miyaura reaction of aryl chlorides was accomplished through the use of an active ferrocene-based triarylphosphine ligand. This air- and moisture-stable ligand was found to be effective for the cross-coupling of aryl halides at room temperature to 115 °C.

Recent advances in cross-coupling methodologies have made significant contribution to organic synthesis as well as pharmaceutical and agricultural chemistry.¹ Suzuki–Miyaura cross-coupling is one of the most attractive and practical protocols for carbon-carbon bond formation due to its tolerance of a wide range of functional groups and the simple separation of products.² Recently, improved catalyst systems for Suzuki–Miyaura couplings have been described, such as Pd-complexes of trialkylphosphines,³ *ortho*-biaryl or ferrocenyl dialkylphosphines,^{4,5} and other ligands,⁶ that allow couplings to be carried out under mild conditions for the preparation of substituted biaryl compounds from relatively inexpensive aryl chlorides.⁷ Despite the intensive attention this area has received, few examples are known where triarylphosphines were used in Suzuki–Miyaura couplings employing aryl chlorides.^{8,9} Since triarylphosphines are usually prepared from relatively inexpensive chlorodiphenylphosphine with Grignard or organolithium reagents, it would add a further attractive feature if these modified triarylphosphines are active and similar to some results of Pd-trialkylphosphine or -dialkylarylyphosphine catalyzed reactions. We report a simple and active ferrocenyl triarylphosphine for Suzuki–Miyaura cross-coupling of sterically hindered aryl bromides at room temperature and aryl chlorides at elevated temperatures.¹⁰

In view of the electronic and steric differences between a benzene ring and the cyclopentadienyl ring of ferrocene, we envisioned that the replacement of a benzene ring with ferrocene would generate a new class of potentially active ligand for cross-coupling reactions. To our delight, the active ligand **7** demonstrated that even Pd-triarylphosphine moiety could catalyze Suzuki–Miyaura coupling using aryl chloride as the substrates.

Triarylphosphine **7** was prepared from commercially available ferrocene in two steps (Scheme 1). Since ferrocene cannot be directly halogenated, the use of the Kagan lithiation protocol is necessary.¹¹ Ferrocene was lithiated with *t*-BuLi in THF to afford ferrocenyllithium and subsequent transmetalation with ZnCl₂ to obtain the Negishi coupling precursor (Scheme 1). Heating the mixtures in the presence of 3% Pd(PPh₃)₄ and *ortho*-bromoiodobenzene led to the formation of 2-bromophenylferrocene.



Scheme 1 Synthesis of ferrocenyl triarylphosphine ligand.

† Electronic supplementary information (ESI) available: experimental section. See <http://www.rsc.org/suppdata/cc/b4/b407661c>


Lithiation with *n*-BuLi and subsequent reaction with CIPPh₂ cleanly gave air-stable ligand **7** in good yield (Scheme 1).

To test the effectiveness of this new ligand **7**, we used it in comparison with various commercially available triarylphosphine ligands in the Suzuki–Miyaura coupling of an *ortho*-substituted aryl bromide (eqn. (1) and Table 1). While *rac*-BINAP and tri-*o*-tolylphosphine gave moderate conversions, the new monodentate ferrocenyl triarylphosphine **7** showed superior results in Suzuki coupling of sterically hindered aryl bromide at room temperature. In addition, the triarylphosphine **7** has the highest activity over the other triarylphosphines in Suzuki coupling of aryl chloride (eqn. (2), Table 1).

This interesting finding led us to the exploration of using **7** in the coupling of various substituted aryl bromides (Table 2). *ortho*-Phenyl substituted aryl bromide was coupled with phenyl boronic acid in excellent yield (Table 2, entry 2). Highly substituted aryl bromides, such as 2,6-diisopropyl or 2-*tert*-butyl-6-methoxy substituted bromobenzene required slightly higher temperature (30 °C) (Table 2, entries 4 and 5).

The combination of variety of solvents and bases were screened for coupling of aryl chlorides. Potassium phosphate monohydrate was found to be superior in toluene. Functional groups such as aldehyde, keto, nitrile and ester were found to be compatible under these reaction conditions (Table 3). Electron-deficient aryl chloride can be coupled with arylboronic acid at room temperature (Table 3, entry 4). The coupling of deactivated aryl chlorides with *ortho*-substituted arylboronic acids required slightly higher catalyst loading (Table 3, entry 6). Interestingly, about 2% of diarylmethanol was observed from GC-MS (Table 3, entry 2), possibly resulting from the addition of arylboronic acid to the carbonyl group.¹² Aryl-aryl exchange reactions between Pd-Ar and P-Ar are typical side reactions encountered when triarylphosphine ligands are employed in coupling reactions,¹³ however, no such side products were formed from under these reaction conditions when ligand **7** was employed in Suzuki–Miyaura coupling.¹⁴

Table 1 Triarylphosphine ligands screening in Pd-catalyzed Suzuki–Miyaura cross-coupling of aryl bromide and chloride



Entry	Ligand L	GC yield (%), eqn. (1)	GC yield (%), eqn. (2)
1	No ligand	8	0
2	<i>rac</i> -BINAP	32	5
3	DPPF	7	4
4	Xantphos	7	5
5	(<i>o</i> -tolyl) ₃ P	55	15
6	Ph ₃ P	18	9
7	Ligand 7	98	92

Table 2 Suzuki–Miyaura coupling of sterically hindered ArBr at RT^a

Entry	ArBr	Ar'B(OH) ₂	Product	% Yield ^b
1				98
2				96
3				95
4 ^c				93
5 ^c				87
6				90

^a Reaction conditions: ArBr (1.0 mmol), Ar'B(OH)₂ (1.5 mmol), Pd₂(dba)₃ (0.5 mol%), ligand 7 (1.2 mol%), KF (3.0 mmol) and THF (2.0 mL) were charged in a Schlenk flask under N₂ and stirred at RT. ^b Isolated yields. ^c Reactions were conducted at 30 °C for 72 h.

Heteroaryl and alkenyl chlorides were effective substrates. (Table 3, entries 8–9). Additionally, the cross-coupling of aryl chlorides with alkylboronic acids are rare.¹⁵ Gratifyingly, a preliminary study of the couplings of alkylboronic acids with both electron-poor and rich aryl chlorides was found to be successful (Table 3, entries 10–11).

In summary, we have developed a new, air-stable and active ferrocenyl triarylphosphine and have established its effectiveness in the palladium-catalyzed Suzuki–Miyaura coupling of aryl halides. The capability of this system to achieve cross-coupling of activated aryl chloride at room temperature is noteworthy. In light of the fact that 7 is a triarylphosphine, the high reactivity of Pd-7 is unexpected. In view of the ease with which the structure of 7 can be modified, we anticipate that further enhancements in reactivity as well as the design of versatile ligands for asymmetric catalysis will be a fruitful pursuit.

We thank the University Grants Committee Area of Excellence Scheme in Hong Kong (Project No. AoE/P-10/01) and the Hong Kong Polytechnic University Area of Strategic Development Fund for financial support of this study.

Notes and references

- (a) O. Baudoin, M. Cesario, D. Guénard and F. Guéritte, *J. Org. Chem.*, 2002, **67**, 1199; (b) M. Hocek, A. Holý, I. Votruba and H. Dvořáková, *J. Med. Chem.*, 2000, **43**, 1817.
- (a) N. Miyaura, *Top. Curr. Chem.*, 2002, **219**, 11; (b) S. Kotha, K. Lahiri and D. Kashinath, *Tetrahedron*, 2002, **58**, 9633.
- (a) A. F. Littke, C. Dai and G. C. Fu, *J. Am. Chem. Soc.*, 2000, **122**, 4020; (b) A. F. Littke and G. C. Fu, *Angew. Chem., Int. Ed. Engl.*, 1998, **37**, 3387.
- For dialkyl biphenyl-based phosphine ligands, see: (a) D. W. Old, J. P. Wolfe and S. L. Buchwald, *J. Am. Chem. Soc.*, 1998, **120**, 9722; (b) J. P. Wolfe and S. L. Buchwald, *Angew. Chem., Int. Ed. Engl.*, 1999, **38**, 2413; (c) H. N. Nguyen, X. Huang and S. L. Buchwald, *J. Am. Chem. Soc.*, 2003, **125**, 11818; (d) X. Bei, H. W. Turner, W. H. Weinberg, A. S. Guram and J. L. Petersen, *J. Org. Chem.*, 1999, **64**, 6797; (e) A. Zapf, R. Jackstell, F. Rataboul, T. Riermeier, A. Monsees, C. Fuhrmann, N. Shaikh, U. Dingerdissen and M. Beller, *Chem. Commun.*, 2004, 38; (f) S. D. Walker, T. E. Barder, J. R. Martinelli and S. L. Buchwald, *Angew. Chem., Int. Ed. Engl.*, 2004, **43**, 1871; (g) J. Yin, M. P. Rainka, X.-X. Zhang and S. L. Buchwald, *J. Am. Chem. Soc.*, 2002, **124**, 1162; (h) F. Y. Kwong, W. H. Lam, C. H. Yeung, K. S. Chan and A. S. C. Chan, *Chem Commun.*, 2004, 1922.
- For dialkyl ferrocenyl-based ligands, see: (a) N. Kataoka, Q. Shelby,

Table 3 Pd-catalyzed Suzuki coupling of ArCl with Fc-ligand 7^a

Entry	ArCl	Ar'B(OH) ₂	Product	% Yield ^b
1				90
2				91
3				94
4 ^d				91
5				92
6 ^c				83
7				96
8				91
9				88
10 ^e				95
11 ^e				88

^a Reaction conditions: ArCl (1.0 mmol), Ar'B(OH)₂ (1.5 mmol), Pd₂(dba)₃ (0.5 mol%), ligand 7 (1.2 mol%), K₃PO₄ • H₂O (3.0 mmol) and toluene (2.0 mL) were stirred at 110 °C under N₂ for 24 h; entry 3, 70 °C; entry 7, 90 °C. ^b Isolated yields. ^c Reactions were conducted with Pd₂(dba)₃ (2 mol%), ligand 7 (8 mol%), 115 °C, 28 h. ^d Pd₂(dba)₃ (2 mol%), ligand 7 (4 mol%), rt, 48 h. ^e K₃PO₄ (3.0 mmol) was used.

- J. P. Stambuli and J. F. Hartwig, *J. Org. Chem.*, 2002, **67**, 5553; (b) J. F. Jensen and M. Johannsen, *Org. Lett.*, 2003, **5**, 3025.
- For review of phosphinite ligands, see: (a) R. B. Bedford, *Chem. Commun.*, 2003, 1787. For the Suzuki reaction using phosphine oxide ligand, see: (b) G. Y. Li, *Angew. Chem., Int. Ed. Engl.*, 2001, **40**, 1513.
- A. F. Littke and G. C. Fu, *Angew. Chem., Int. Ed. Engl.*, 2002, **41**, 4176.
- For one example of Suzuki coupling of an activated aryl chloride by a Pd-triarylphosphine catalyst, see: P. Kočovský, S. Vyskočil, I. Cisařová, J. Sejbál, I. Tišlerová, M. Smrčina, G. C. Lloyd-Jones, S. C. Stephen, C. P. Butts, M. Murray and V. Langer, *J. Am. Chem. Soc.*, 1999, **121**, 7714. For Suzuki coupling of mainly activated ArCl by a Pd-PPh(2-CiC₆H₄)₂ catalyst, see: M. R. Pramick, S. M. Rosemeier, M. T. Beranek, S. B. Nickse, J. J. Stone, R. A. Stockland, Jr., S. M. Baldwin and M. E. Kastner, *Organometallics*, 2003, **22**, 523.
- For Ar₃P with -PPh₂ moiety attached to Cpring, see: (a) S.-Y. Liu, M. Choi and G. C. Fu, *Chem. Commun.*, 2001, 2408. For Fc₃P: (b) T. E. Pickett and C. J. Richards, *Tetrahedron Lett.*, 2001, **42**, 3767; (c) T. E. Pickett, F. X. Roca and C. J. Richards, *J. Org. Chem.*, 2003, **68**, 2592.
- During the preparation of the manuscript, a similar ferrocenyl dicyclohexylphosphine was reported, F. X. Roca and C. J. Richards, *Chem. Commun.*, 2003, 3002.
- F. Rebiere, O. Samuel and H. B. Kagan, *Tetrahedron Lett.*, 1990, **31**, 3121.
- For Rh-catalyzed addition of ArB(OH)₂ acids to aldehyde: M. Sakai, M. Euda and N. Miyaura, *Angew. Chem., Int. Ed. Engl.*, 1998, **37**, 3279.
- (a) K. C. Kong and C. H. Cheng, *J. Am. Chem. Soc.*, 1991, **113**, 6313; (b) F. E. Goodson, T. I. Wallow and B. M. Novak, *J. Am. Chem. Soc.*, 1997, **119**, 12441; (c) F. Y. Kwong, C. W. Lai and K. S. Chan, *J. Am. Chem. Soc.*, 2001, **123**, 8864.
- For aryl-aryl exchange side reactions in Suzuki–Miyaura coupling using triarylphosphines: (a) D. F. O'Keefe, M. C. Dannock and S. M. Marcuccio, *Tetrahedron Lett.*, 1992, **33**, 6679; (b) F. E. Goodson, T. I. Wallow and B. M. Novak, *Macromolecules*, 1998, **31**, 2047.
- For coupling of alkylboronic acid with ArCl using dialkylphosphine ligands, see ref. 5a and, most recent, ref. 4f.