

Millisecond long-lived charge separated state at room temperature in a flexibly linked diphenylaminopolyene-C₆₀ dyad

Yanong Han,[†] Laura Dobeck, Aijun Gong, Fanqing Meng, Charles W. Spangler and Lee H. Spangler*

Received (in Columbia, MO, USA) 7th July 2004, Accepted 4th November 2004

First published as an Advance Article on the web 11th January 2005

DOI: 10.1039/b410201k

Long-lived photoinduced charge separation involving one-step electron transfer is achieved in diphenylaminopolyene based C₆₀-donor dyads with a short, flexible linkage.

Extensive research has been directed towards the synthesis of electron donor–acceptor linked compounds in the hope of finding systems which will form long-lived charge separated (CS) states with high quantum efficiency, as is achieved in the photosynthetic reaction.^{1–5} The lifetimes of CS states are sensitive to the energy of the CS state, the type of linkage between the donor and acceptor molecules, and also the molecular topology and distance and orientation of the donor and acceptor moieties.^{6–11} Large efforts have been made to achieve a long-lived CS state by tuning these factors with the longest currently reported CS state lifetime of a dyad at room temperature solution being 260 μs for a zinc imidazoporphyrin-C₆₀ dyad.⁶ Relatively unexplored is CS lifetime dependence on the flexibility of the linkage between the donor and acceptor in the dyad.

Among a wide variety of donor molecules available for linking covalently to C₆₀, the triphenylamine group is one with good electron donating ability to excited state C₆₀ molecules.¹² We report herein photoinduced intramolecular electron transfer (ET) processes in diphenylaminopolyene (DPAP) based C₆₀-donor dyads (DPAP-C₆₀) with a short, flexible linkage. The decay mechanism of the CS state was elucidated with the use of nanosecond laser flash photolysis. Dual decay components of the CS state were observed, with first order lifetimes of about 300 ns and 1 ms respectively. To our knowledge, this millisecond slow component lifetime is the longest first order lifetime at room temperature ever reported for intramolecular charge recombination (CR) in donor–acceptor dyad systems involving one-step electron transfer. A new method was introduced to simultaneously analyze the backward electron transfer (BET) process which involves both *intra*- (first order) and *intermolecular* (second order) CR.

One and two armed DPAP-C₆₀ dyads (D1 and D2) were synthesized as shown in Scheme 1. Anhydrous tetrahydrofuran (THF) was used freshly distilled from sodium and benzophenone-dried tetrahydrofuran. Other solvents were used as received. Wittig reaction methodologies were used to extend the conjugated lengths of these molecules. The donor chromophores were synthesized with a free OH functionality for attachment to malonyl dichloride. The resulting diester was then coupled to C₆₀ using Hirsh–Bingel methodology, yielding the target dyads as dark solids in high yield.


The successful synthesis of a new kind of Wittig salt (compound 1, Scheme 1) provides an easy way to obtain DPAP-C₆₀ dyads in good yield. The dyads were characterized by ¹H NMR, ¹³C NMR and MALDI-TOF MS apparatus. The absorption bands of the C₆₀ and DPAP moieties are present in the spectra of the linked compounds D1 and D2.

The photoexcitation of both D1 and D2 in deaerated benzonitrile (BN) solution with 360 nm laser light results in fluorescence with the emission maximum at 540 nm. The emission intensities of D1 and D2 are quenched 35- and 9-fold respectively, compared to emission from the reference compound DPAP. This indicates that the excited state of DPAP is quenched efficiently by linked C₆₀. The emission intensity of the single armed D1 is four times smaller than that of D2 at the same DPAP concentration.

The fluorescence decay of the DPAP unit in BN solution was monitored from 415 to 715 nm after excitation with 400 nm light from a frequency doubled ultrafast Ti : sapphire laser (76 MHz, 150 fs pulse width). The fluorescence lifetime of the unlinked DPAP (τ₀ = 1.8 ns) was significantly reduced in D1 (τ_f = 9 ps). This is likely due to ET from the excited state of DPAP (DPAP*) to C₆₀. The rate constant of forward ET (k_{et}) from DPAP* to C₆₀ was estimated from k_{et} = 1/τ_f - 1/τ₀, where τ₀ is the fluorescence lifetime of the DPAP unit, and τ_f is that in the linked compound. The k_{et} value of D1 was evaluated to be 1.1 × 10¹¹ s⁻¹.

The time profile of the fluorescence intensity of the DPAP* in D2 shows biexponential decay, with fluorescence lifetimes of 20 and 1800 ps respectively. This may be due to the restriction that only one DPAP* could be quenched by C₆₀ if both arms are photoexcited. The k_{et} value from DPAP* to C₆₀ for D2 was estimated to be 4.9 × 10¹⁰ s⁻¹.

Transient absorption spectra of DPAP-C₆₀ were carried out following nanosecond laser excitation at 532 nm. Only absorption



Scheme 1 Synthesis of donor chromophores and dyads.

[†] Present address: Yatai Technologies Ltd., Tianzhu Road, Hefei, Anhui 230088, China.

*spangler@montana.edu




Fig. 1 Top: photoinduced absorption (PIA) spectra obtained by using 532 nm laser excitation of DPAP- C_{60} in deaerated BN, compared with the chemically induced absorption (CIA) spectrum of the cation generated by doping the donor with $SbCl_5$. Bottom: time profiles (μs range) of DPAP- C_{60} charge separated state decay. Inset are the ns range time profiles.

of the $^3C_{60}^*$ moiety was detected at 740 nm in deaerated toluene. From the decay of the time profile, the lifetime of DPAP- $^3C_{60}^*$ was evaluated to be 18 μs , which is close to that of pristine C_{60} . It indicates that the intersystem crossing process from DPAP- $^1C_{60}^*$ to DPAP- $^3C_{60}^*$ is the predominant one with no charge separation in nonpolar solvents.

The CS state was detected through laser flash photolysis (pumped by a 7 ns 532 nm laser) of DPAP- C_{60} (both D1 and D2) in deaerated benzonitrile (BN) solution as shown in Fig. 1. Following the excitation, an absorption band at 8800 cm^{-1} with a shoulder at 9600 cm^{-1} appears immediately. This is attributable to the DPAP cation and fullerene anion. The transient absorption in the near visible region ($15\ 000\text{ cm}^{-1}$ band) is also due to DPAP $^+$. The photoinduced DPAP $^+$ spectrum is confirmed by doping the donor with a strong oxidizing agent ($SbCl_5$) to chemically generate the cation.

The temporal decay profiles of both D1 and D2 show two obvious components. The time constant of the fast component is about 330 ns, while that of the slow component is about 1 ms. The ratios of populations of the slow and fast components are about 0.15 and 0.2 for D1 and D2 respectively, as shown in Fig. 1.

The CS states decay through back electron transfer (BET) to the ground state. The slow component decays for both D1 and D2 on the μs timescale are neither strictly first order nor second order, as shown in Fig. 1. The BET occurs through two processes: it is close




Fig. 2 Plot of $d[CS]/dt$ vs. $[CS]$ for 0.2 mM D2 charge separation state decay in deaerated BN. Circles: experimental data resulting from different pump powers (0.5–5 mJ); solid line: fit with second order polynomial.

to first order at low concentration ($t > 500\ \mu s$), and at high concentration it is significantly affected by *intermolecular* BET. The decay behavior follows:

$$\frac{d[CS]}{dt} = -k_{1st}[CS] - k_{2nd}[CS][CS]$$

where k_{1st} ($=1/\tau_{lifetime}$) and k_{2nd} are *intra-* and *intermolecular* decay rate constants respectively. The dependence of the charge separation (CS) state decay rate on $[CS]$ was examined by employing different laser powers over a wide range and it exhibits a good second order polynomial fit (see Fig. 2). This fitting process gives us values for both first and second order rate constants, so we can calculate the lifetimes expected for the first order process alone. The first order lifetimes of D1 and D2 were determined to be 1.1 and 0.7 ms, respectively. These values are approximately two times larger than what we get from single exponential fitting at low concentrations and long times, since the *intermolecular* BET contribution is still present at the lowest concentrations used.

To our knowledge, the millisecond lifetime is the longest CS lifetime in solution ever reported for *intramolecular* charge recombination at room temperature. Evolution of the 8800 cm^{-1} CS band shape over a 300 ns timescale suggests conformational changes resulting in some solvent separation of the linked DPAP $^+$ and C_{60}^- . This may lead to a new synthetic strategy to attain long lifetime charge separation donor–acceptor dyads by using flexible linkages and controlling the folding. Further research is under way to probe this point.

We would like to acknowledge DOE grant DE-FG02-01ER45869 and ARO grant DAAD 19-00-1-0162 for financial support and Reed Howald for helpful discussions.

Yanong Han,[†] Laura Dobeck, Aijun Gong, Fanqing Meng, Charles W. Spangler* and Lee H. Spangler*

Department of Chemistry, Montana State University, Bozeman, MT 59717, USA. E-mail: spangler@montana.edu; Fax: 01 406 994 5407; Tel: 01 406 994 4399

Notes and references

- (a) M. R. Wasielewski, *Chem. Rev.*, 1992, **92**, 435; (b) F. D. Lewis, X. Liu, J. Liu, S. E. Miller, R. T. Hayes and M. R. Wasielewski, *Nature*, 2000, **406**, 51.

-
- 2 (a) D. Gust, T. A. Moore and A. L. Moore, *Acc. Chem. Res.*, 1993, **26**, 198; (b) C. C. Page, C. C. Moser, X. Chen and P. L. Dutton, *Nature*, 1999, **402**, 47.
- 3 (a) S. Fukuzumi and D. M. Guldi, in *Electron Transfer in Chemistry*, ed. V. Balzani, Wiley-VCH, Weinheim, 2001, **vol. 2**, pp 270–337.
- 4 K. Ohkubo, H. Imahori, J. Shao, Z. Ou, K. M. Kadish, Y. Chen, G. Zheng, R. K. Pandey, M. Fujitsuka, O. Ito and S. Fukuzumi, *J. Phys. Chem. A*, 2002, **106**, 10991.
- 5 C. Luo, D. M. Guldi, H. Imahori, K. Tamaki and Y. Sakata, *J. Am. Chem. Soc.*, 2000, **122**, 6535.
- 6 H. Imahori, D. M. Guldi, K. Tamaki, Y. Yoshida, C. Luo, Y. Sakata and S. Fukuzumi, *J. Am. Chem. Soc.*, 2001, **123**, 6617.
- 7 Y. Kashiwagi, K. Ohkubo, J. A. McDonald, I. M. Blake, M. J. Crossley, Y. Araki, O. Ito, H. Imahori and S. Fukuzumi, *Org. Lett.*, 2003, **5**, 2719.
- 8 (a) D. M. Guldi, M. Maggini, G. Scorrano and M. Prato, *J. Am. Chem. Soc.*, 1997, **119**, 974; (b) P. A. Liddell, D. Kuciauska, J. P. Sumida, B. Nash, D. Nguyen, A. L. Moore, T. A. Moore and D. Gust, *J. Am. Chem. Soc.*, 1997, **119**, 1400.
- 9 N. Martin, L. Sanchez, B. Illescas and I. Perez, *Chem. Rev.*, 1998, **98**, 2527.
- 10 (a) H. Imahori, K. Tamaki, D. M. Guldi, C. Luo, M. Fujitsuka, O. Ito, Y. Sakata and S. Fukuzumi, *J. Am. Chem. Soc.*, 2001, **123**, 2607; (b) D. M. Guldi, C. Luo, T. Da Ros, M. Prato, E. Dietel and A. Hirsch, *Chem. Commun.*, 2000, 375; (c) S. Fukuzumi, H. Imahori, H. Yamada, M. E. El-Khouly, M. Fujitsuka, O. Ito and D. M. J. Guldi, *J. Am. Chem. Soc.*, 2001, **123**, 2571.
- 11 (a) F. D'Souza, G. R. Deviprasad, M. S. Rahman and J.-P. Choi, *Inorg. Chem.*, 1999, **38**, 2157; (b) T. Da Ros, M. Prato, D. M. Guldi, E. Alessio, M. Ruzzi and L. Pasimeni, *Chem. Commun.*, 1999, 635; (c) F. D'Souza, N. P. Rath, G. R. Deviprasad and M. E. Zandler, *Chem. Commun.*, 2001, 267; (d) F. D'Souza, G. R. Deviprasad, M. E. El-Khouly, M. Fujitsuka and O. Ito, *J. Am. Chem. Soc.*, 2001, **123**, 5277; (e) F. D'Souza, G. R. Deviprasad, M. E. Zandler, M. E. El-Khouly, M. Fujitsuka and O. Ito, *J. Phys. Chem. B*, 2002, **106**, 4952.
- 12 Y. Han and L. H. Spangler, *J. Phys. Chem. A*, 2002, **106**, 1701.