

Enantiospecific inclusion of chiral 1,2-dichloroethane rotamers in the crystal lattice of chiral square-pyramidal Cu(II) complexes with perfectly polar alignment of guest and host molecules

Vamsee Krishna Muppidi, Panthapally S. Zacharias^{†*} and Samudranil Pal^{*}


Received (in Cambridge, UK) 3rd February 2005, Accepted 18th March 2005

First published as an Advance Article on the web 4th April 2005

DOI: 10.1039/b501642h

The inclusion compounds, $[\text{CuL}^1_2(\text{H}_2\text{O})]\cdot(P)\text{-C}_2\text{H}_4\text{Cl}_2$ and $[\text{CuL}^2_2(\text{H}_2\text{O})]\cdot(M)\text{-C}_2\text{H}_4\text{Cl}_2$ ($\text{HL}^1 = N\text{-}(2\text{-hydroxy-5-nitrobenzyl})\text{-}(R)\text{-}\alpha\text{-methylbenzylamine}$ and $\text{HL}^2 = N\text{-}(2\text{-hydroxy-5-nitrobenzyl})\text{-}(S)\text{-}\alpha\text{-methylbenzylamine}$), crystallise in the non-centrosymmetric space group $C2$; intermolecular hydrogen bonding leads to a perfectly polar alignment of both host and guest molecules with enantioselectivity.

Polar and non-centrosymmetric organization of molecules in the crystalline state is a prerequisite for molecular materials having technologically important physical properties such as ferro-, pyro- or piezoelectricity and nonlinear optical effects.¹ Inclusion of chirality in the molecule is a common approach to ensure non-centrosymmetric organization in the crystals.² On the other hand, perfectly polar assembly in the crystal lattice is rare and requires special design strategies.³ Synthesis of inclusion compounds having polar host frameworks that can confine and align guest molecules in a polar order is one of the successful strategies.⁴ In this regard, it may well be noted that the synthesis of host frameworks with chiral cavities is a noteworthy challenge as these solids also have potential applications in the separation of enantiomers and in asymmetric catalysis.⁵ In this work, we report the enantioselective isolation of the right handed (*P*) and the left-handed (*M*) *gauche* form of 1,2-dichloroethane in the host lattices formed by the chiral square-pyramidal Cu(II) complexes having the general formula $[\text{CuL}^n_2(\text{H}_2\text{O})]$ ($\text{HL}^1 = N\text{-}(2\text{-hydroxy-5-nitrobenzyl})\text{-}(R)\text{-}\alpha\text{-methylbenzylamine}$ and $\text{HL}^2 = N\text{-}(2\text{-hydroxy-5-nitrobenzyl})\text{-}(S)\text{-}\alpha\text{-methylbenzylamine}$, H represents the dissociable phenolic proton) with perfectly polar alignment of both guest and host molecules.



HL^1 and HL^2 were prepared by condensation reactions of one mole equivalent of 5-nitrosalicylaldehyde with one mole equivalent of the corresponding enantiomerically pure α -methylbenzylamine in methanol followed by reduction with NaBH_4 . Reactions of $\text{CuCl}_2\cdot 2\text{H}_2\text{O}$, HL^n and LiOH (1:2:2 mole ratio) in methanol

produce the complexes, $[\text{CuL}^1_2(\text{H}_2\text{O})]$ (**1**) and $[\text{CuL}^2_2(\text{H}_2\text{O})]$ (**2**) in good yields.[‡] X-Ray quality single crystals were grown by slow evaporation of 1,2-dichloroethane solutions of the complexes. Both complexes crystallise as $[\text{CuL}^n_2(\text{H}_2\text{O})]\cdot\text{C}_2\text{H}_4\text{Cl}_2$ in the non-centrosymmetric space group $C2$ with essentially identical unit cell parameters.[§] The molecular structures of **1** and **2** are mirror images of each other (Fig. 1). In each complex molecule, the metal centre is in a distorted square-pyramidal N_2O_3 coordination sphere. Two bidentate chelating ligands form the N_2O_2 basal plane and the water oxygen occupies the apical site. The bidentate ligands are related by a crystallographically imposed two-fold axis passing through the Cu and the apical water O. Both atoms are at special positions on the *b*-axis. The dihedral angles between the plane defined by Cu, N1 and O1 and that defined by Cu, N1' and O1' are $18.41(4)^\circ$ and $18.97(5)^\circ$ for **1** and **2**, respectively. These values indicate a tetrahedral distortion of the N_2O_2 basal plane.




Fig. 1 Molecular structures of (a) $[\text{CuL}^1_2(\text{H}_2\text{O})]$ (**1**) and (b) $[\text{CuL}^2_2(\text{H}_2\text{O})]$ (**2**) with the atom-labelling schemes. All non-hydrogen atoms are represented by their 35 and 50% probability thermal ellipsoids for **1** and **2**, respectively.

[†] Present address: Vice Chancellor, Goa University, Goa 403 206, India.
^{*} spsc@uohyd.ernet.in (Samudranil Pal)

The bond parameters associated with the Cu(II) centres in **1** and **2** are unexceptional.⁶

Three rotamers are possible for 1,2-dichloroethane. These are the *trans* or *anti*, the *cis* or eclipsed and the *gauche* forms. The eclipsed form is energetically most unfavorable. Between the *trans* and the *gauche* forms the former is more stable (by ~ 1 kcal mol⁻¹).⁷ The *gauche* form is chiral and also polar. One of the enantiomers has the right-handed (*P*) form and the other one has the left-handed (*M*) form. In the structures of **1**·C₂H₄Cl₂ and **2**·C₂H₄Cl₂, the asymmetric unit contains one-half of the 1,2-dichloroethane molecule. The other half can be generated by rotation about a two-fold axis passing through the mid-point of the C–C bond. The molecule is in *gauche* form in each structure (Fig. 2). The torsion angles involving Cl–C–C–Cl are 63(1)° and 64.6(9)° in **1**·C₂H₄Cl₂ and **2**·C₂H₄Cl₂, respectively. The molecule is trapped in the *P*-form in **1**·C₂H₄Cl₂ (Fig. 2a). On the other hand, in the case of **2**·C₂H₄Cl₂ only the *M*-form of it exists (Fig. 2b). Thus the chiral host complexes **1** and **2** produce the appropriate frameworks in the crystal lattice for exclusive accommodation of the *P*- and *M*-forms of 1,2-dichloroethane, respectively.

Reports of inclusion crystals of optically active and polar rotamers of 1,2-dichloroethane with a chiral host compound are very rare.⁸ There are no reports on inclusion crystals of a chiral rotamer of 1,2-dichloroethane with polar alignment of the guest molecules in the host framework. The crystals of **1**·C₂H₄Cl₂ and **2**·C₂H₄Cl₂ provide examples, not only of enantioselective isolation of the *P*- and *M*-forms of the chiral rotamers, but also of the unprecedented, perfectly polar alignment of both host and guest molecules in the crystal lattice. In each case, the polar alignments of host and guest molecules are along the *b*-axis. It may be noted that for **1** and **2**, and also for (*P*)-C₂H₄Cl₂ and (*M*)-C₂H₄Cl₂, the molecular dipole axis is along the *b*-axis or parallel to the *b*-axis. The polar alignments are in opposite directions due to the enantiomeric relationship between the two host molecules and that between the two guest molecules in **1**·(*P*)-C₂H₄Cl₂ and **2**·(*M*)-C₂H₄Cl₂ (Fig. 3). In the crystal lattice of each structure, the apical water molecule of the complex is involved in hydrogen bonding interactions with the nitro groups of the neighboring molecules. Most likely these O–H···O interactions are primarily responsible for the alignment of the host molecules in a head-to-tail manner along the *b*-axis. For **1** the O4–H···O2 and O4···O2 distances and O4–H···O2 angle are 0.86(6), 2.03(7) and 2.846(3) Å and 160(9)°, respectively. For **2** the values of these parameters are 0.89(5), 1.98(6) and 2.816(3) Å and 156(7)°, respectively. In both cases, the same O-atom of the nitro group participates in an additional weak intermolecular C–H···O




Fig. 2 (a) The right-handed (*P*) and (b) the left-handed (*M*) forms of 1,2-dichloroethane in **1**·C₂H₄Cl₂ and **2**·C₂H₄Cl₂, respectively.




Fig. 3 Polar alignments of (a) **1**·(*P*)-C₂H₄Cl₂ and (b) **2**·(*M*)-C₂H₄Cl₂ viewed along the *a*-axis; H-atoms of the ligands and 1,2-dichloroethane are omitted for clarity.

interaction involving one of the methylene hydrogens.⁹ The C7···O2 distance and the C7–H···O2 angle are 3.388(4) Å and 146° and 3.336(4) Å and 147° for **1** and **2**, respectively. These O–H···O and C–H···O interactions connect the complex molecules and a polar two-dimensional layered structure is formed (Fig. 4). In the channels formed by the parallel layers of the complex molecules, the 1,2-dichloroethane molecules are trapped and oriented in a polar order. The methylene groups of the 1,2-dichloroethane are involved in intermolecular C–H···O interactions with the other O-atom of the nitro group (Fig. 5). The C16···O3 distance and the C16–H···O3 angle are 3.285(14) Å and 150° and 3.195(11) Å and 149° for **1**·(*P*)-C₂H₄Cl₂ and




Fig. 4 View of the two-dimensional layered structures of (a) **1** and (b) **2** along the *c*-axis; only the H₂O and the CH₂ group H-atoms involved in the non-covalent interactions (dashed lines) are shown.




Fig. 5 Projections of the crystal lattices of (a) **1-(P)-C₂H₄Cl₂** and (b) **2-(M)-C₂H₄Cl₂** onto the *ac*-plane.

2-(M)-C₂H₄Cl₂, respectively. These C–H···O interactions act as bridges between the parallel layers of the complex molecules and a three-dimensional network, with the polar alignment of both host and guest molecules, is formed.

Self-assembly *via* intermolecular non-covalent interactions is one of the powerful tools for designing and synthesising polar crystals as well as enantioselective chiral host frameworks. The spatial disposition of the functional groups participating in the intermolecular non-covalent interactions is expected to control the alignment of the host and guest molecules as well as spontaneous resolution of chiral guest compounds. The structures of **1-(P)-C₂H₄Cl₂** and **2-(M)-C₂H₄Cl₂** reveal this functional group directed alignment of molecules and selective trapping of guest molecules. The self-assembly of the host molecules (**1** and **2**) occurs through the O–H···O interactions and the guest molecules (1,2-dichloroethane) are held *via* the C–H···O interactions. In both intermolecular interactions, ligand nitro group O-atoms act as the acceptors. The positions and orientations of the nitro groups on the enantiomeric ligands in **1** and **2** are befitting for a perfectly polar alignment of the host molecules and for enantioselective confinement of the chiral rotamers of 1,2-dichloroethane in a polar order.

Financial support for this work was provided by the Council of Scientific and Industrial Research (CSIR), New Delhi [Grant No. 01(1880)/03/EMR-II to S. P.]. V. K. M. thanks the CSIR for a senior research fellowship. X-Ray crystallographic studies were performed at the National Single Crystal Diffractometer Facility, School of Chemistry, University of Hyderabad (funded by the Department of Science and Technology, New Delhi). We thank the University Grants Commission, New Delhi for the facilities provided under the Universities with Potential for Excellence program.

Vamsee Krishna Muppidi, Panthapally S. Zacharias^{†*} and Samudranil Pal*

School of Chemistry, University of Hyderabad, Hyderabad 500 046, India. E-mail: spsc@uohyd.ernet.in; Fax: +91 40 2301 2460; Tel: +91 40 2313 4756

Notes and references

[†] Synthesis of HL¹: a methanol solution (30 ml) of 5-nitrosalicylaldehyde (1.67 g, 10 mmol) was added to a methanol solution (30 ml) of (*R*)- α -methylbenzylamine (1.21 g, 10 mmol) and stirred at room temperature for ½ h. To the resulting yellow solution, 0.74 g (20 mmol) of NaBH₄ was added and the mixture was stirred for another ½ h until a colourless solution was obtained. The reaction mixture was evaporated to dryness on a rotary evaporator, followed by addition of 100 ml of water. The solid

suspended in water was extracted with (2 × 30 ml) CH₂Cl₂. The CH₂Cl₂ extracts were dried over anhydrous Na₂SO₄ and then evaporated to dryness. The compound (HL¹) was obtained in the form of light yellow solid (1.96 g, yield 72%, m.p. 91–93 °C). HL² was prepared in similar yield by following the same procedure as described above using (*S*)- α -methylbenzylamine instead of (*R*)- α -methylbenzylamine. The elemental analysis data and the ¹H NMR spectra (in CDCl₃) of the compounds are consistent with the expected molecular formula and structures.

Synthesis of **1**: a methanol solution (10 ml) of CuCl₂·2H₂O (89 mg, 0.5 mmol) was added to a methanol solution (20 ml) of HL¹ (272 mg, 1 mmol) and LiOH·H₂O (42 mg, 1 mmol). The mixture was stirred at room temperature for 1 h. The green solid separated was collected by filtration, washed with methanol and dried in vacuum. Yield, 230 mg (74%). Complex **2** was synthesized in similar yield and isolated as a green solid by following the same procedure as described for **1** using HL² instead of HL¹. § *Crystal data*: for **1**-C₂H₄Cl₂: CuC₃₂H₃₆N₄O₇Cl₂, *M* = 723.09, monoclinic, space group C2, *a* = 20.2834(12), *b* = 8.1769(5), *c* = 12.8799(7) Å, β = 128.6480(10)°, *V* = 1667.94(17) Å³, *T* = 293(2) K, *Z* = 2, 3842 independent reflections, 3572 reflections with *I* > 2 σ (*I*), 215 parameters, *R*₁ = 0.0411 and *wR*₂ = 0.1035 [*I* > 2 σ (*I*)], *R*₁ = 0.0443 and *wR*₂ = 0.1056 (all data), goodness-of-fit on *F*² = 1.068, Flack parameter = –0.01(1), largest peak 0.647 e Å^{–3}. For **2**-C₂H₄Cl₂: CuC₃₂H₃₆N₄O₇Cl₂, *M* = 723.09, monoclinic, space group C2, *a* = 20.1329(17), *b* = 8.0328(6), *c* = 12.7987(13) Å, β = 128.290(2)°, *V* = 1624.6(2) Å³, *T* = 293(2) K, *Z* = 2, 3815 independent reflections, 3661 reflections with *I* > 2 σ (*I*), 215 parameters, *R*₁ = 0.0433 and *wR*₂ = 0.1063 [*I* > 2 σ (*I*)], *R*₁ = 0.0451 and *wR*₂ = 0.1076 (all data), goodness-of-fit on *F*² = 1.053, Flack parameter = –0.01(1), largest peak 1.187 e Å^{–3}. Data were collected on a Bruker-Nonius SMART APEX CCD single crystal diffractometer using Mo K α radiation (λ = 0.71073 Å) and ω -scans. An absorption correction was applied to each data set with the help of the SADABS¹⁰ program. The structures were solved by direct methods and refined by least-squares methods on *F*² with all reflections (SHELX-97¹¹). CCDC 263054–263055. See <http://www.rsc.org/suppdata/cc/b5/b501642h/> for crystallographic data in CIF or other electronic format.

- 1 *Molecular Nonlinear Optics: Materials, Physics and Devices*, ed. J. Zyss, Academic Press, London, 1994; *Handbook of Laser Science and Technology*, ed. M. J. Webber, CRC Press, Boca Raton, FL, 1995.
- 2 J. Zyss, J.-F. Nicoud and M. Coquillay, *J. Chem. Phys.*, 1984, **81**, 4160; D. E. Eaton, *Science*, 1991, **253**, 281.
- 3 S. R. Marder, J. W. Perry and W. P. Scafer, *Science*, 1989, **245**, 626; S. R. Marder, J. W. Perry and C. P. Yakymyshyn, *Chem. Mater.*, 1994, **6**, 1137; S. P. Anthony and T. P. Radhakrishnan, *Chem. Commun.*, 2001, 931; P. K. Thallapally, G. R. Desiraju, M. Bagieu-Beucher, R. Masse, C. Bourgoigne and J.-F. Nicoud, *Chem. Commun.*, 2002, 1052; A. Jouaiti, M. W. Hosseini and N. Kyritsakas, *Chem. Commun.*, 2002, 1898.
- 4 W. Tam, D. F. Eaton, J. C. Calabrese, I. D. Williams, Y. Wang and A. G. Anderson, *Chem. Mater.*, 1989, **1**, 128; O. König, H.-B. Bürgi, T. Armbruster, J. Hullinger and T. Weber, *J. Am. Chem. Soc.*, 1997, **119**, 10632; A. Quintel, J. Hullinger and M. Wübbenhorst, *J. Phys. Chem. B*, 1998, **102**, 4277; K. T. Holman, A. M. Pivovar and M. D. Ward, *Science*, 2001, **294**, 1907.
- 5 X. X. Zhang, J. S. Bradshaw and R. M. Izatt, *Chem. Rev.*, 1997, **97**, 3313; E. Brunet, *Chirality*, 2002, **14**, 135; L. Pérez-García and D. B. Amabilino, *Chem. Soc. Rev.*, 2002, **31**, 342; B. Kesanli and W. Lin, *Coord. Chem. Rev.*, 2003, **246**, 305; L.-X. Dai, *Angew. Chem. Int. Ed.*, 2004, **43**, 5726; M. A. Mateos-Timoneda, M. Crego-Calama and D. N. Reinhoudt, *Chem. Soc. Rev.*, 2004, **33**, 363.
- 6 L. L. Koh, J. O. Ranford, W. T. Robinson, J. O. Svensson, A. L. C. Tan and D. Wu, *Inorg. Chem.*, 1996, **35**, 6466; C. T. Yang, H. Moubaraki, K. S. Murray and J. J. Vittal, *Dalton Trans.*, 2003, 880.
- 7 E. L. Eliel and S. H. Wilen, *Stereochemistry of Organic Compounds*, Wiley, New York, 1994, p. 606.
- 8 F. Toda, K. Tanaka and R. Kuroda, *Chem. Commun.*, 1997, 1227.
- 9 G. R. Desiraju, *Acc. Chem. Res.*, 1996, **29**, 441.
- 10 G. M. Sheldrick, *SADABS, Program for area detector adsorption correction*, Institute for Inorganic Chemistry, University of Göttingen, Germany, 1996.
- 11 G. M. Sheldrick, *SHELXL-97, Program for refinement of crystal structures*, University of Göttingen, Germany, 1997; G. M. Sheldrick, *SHELXS-97, Program for solution of crystal structures*, University of Göttingen, Germany, 1997.