

Reactivity of electrophilic μ -phosphinidene complexes with heterocumulenes: formation of the first σ - π -aminophosphamine complexes $[\text{Mn}_2(\text{CO})_8\{\mu\text{-}\eta^1, \eta^2\text{-P}(\text{N}^i\text{Pr}_2)=\text{NR}\}]$ and diazoalkane insertions into metal–phosphorus bonds†

Todd W. Graham,^{ab} Konstantin A. Udachin^a and Arthur J. Carty^{*ab}

Received (in Berkeley, CA, USA) 18th April 2005, Accepted 24th June 2005


First published as an Advance Article on the web 29th July 2005

DOI: 10.1039/b505472a

The bridging phosphinidene complexes $[\text{Mn}_2(\text{CO})_8(\mu\text{-PN}^i\text{Pr}_2)]$ and $[\text{Co}_2(\text{CO})_4(\mu\text{-dppm})(\mu\text{-PNR}_2)]$ ($\text{NR}_2 = \text{N}^i\text{Pr}_2$, TMP) react with heterocumulenes RN_3 , CH_2N_2 and $\text{Ph}_2\text{C}=\text{N}=\text{N}$ to form complexes with $\mu\text{-}\eta^1, \eta^2\text{-aminophosphamine}$, $\mu\text{-}\eta^1, \eta^2\text{-aminophosphaalkene}$ and $\mu\text{-}\eta^1, \eta^2\text{-aminophosphadiphenylmethylazaine}$ ligands, respectively.

Carbene (CX_2) and phosphinidene (PX) ligands are isolobal¹ and there are now well-established structure, bonding, and reactivity similarities^{1,2} between, for example, Schrock type carbene complexes and their nucleophilic η^1 -phosphinidene counterparts.³ More recently the isolation of thermally stable, terminal electrophilic phosphinidenes of the mid⁴ to late⁵ transition metals has stimulated interest in the chemistry of these Fischer type molecules to complement the extensive knowledge of the reaction manifolds for transient phosphinidenes.⁶

Although bridging carbenes (μ -alkylidenes)⁷ comprise an important and diverse class of organometallic molecules and intermediates, their μ -PX counterparts are relatively rare and their chemistry is virtually unexplored.^{8,9} There are also structural and electronic differences between $\mu\text{-CX}_2$ and $\mu\text{-PX}$ complexes: whereas bridging carbenes are formally saturated with tetrahedral stereochemistry at carbon as in I, in the latter, II, the phosphorus centre is planar and unsaturated,¹⁰ and reactivity may be initiated by attack at the electrophilic phosphorus atom.



In this communication we describe the synthesis of two new examples of the class of μ -phosphinidene complexes, $[\text{Mn}_2(\text{CO})_8(\mu\text{-PN}^i\text{Pr}_2)]$ (**1**) and $[\text{Co}_2(\text{CO})_4(\mu\text{-dppm})(\mu\text{-PN}^i\text{Pr}_2)]$ (**2a**) and their novel reactions with heterocumulenes RN_3 ($\text{R} = \text{Me}_3\text{Si}$, Me_3Sn , Ph, adamantyl) and diazoalkanes (CH_2N_2 ,

$\text{Ph}_2\text{C}=\text{N}=\text{N}$) which afford the first complexes of σ, π -bound aminophosphamines $[\text{Mn}_2(\text{CO})_8\{\mu\text{-}\eta^1, \eta^2\text{-P}(\text{N}^i\text{Pr}_2)=\text{NR}\}]$ (**3a–d**) and rare $\sigma\text{-}\pi$ -aminophosphaalkene ligands $[\text{Co}_2(\text{CO})_4(\mu\text{-dppm})(\mu\text{-}\eta^1, \eta^2\text{-PNR}_2=\text{CH}_2)]$ ($\text{NR}_2 = \text{N}^i\text{Pr}_2$ (**4a**), TMP (**4b**)). Diphenyldiazomethane $\text{Ph}_2\text{C}=\text{N}=\text{N}$ also inserts into one arm of the phosphinidene bridge in **1** without loss of dinitrogen to yield the novel complex $[\text{Mn}_2(\text{CO})_8\{\mu\text{-P}(\text{N}^i\text{Pr}_2)=\text{N}-\text{N}(\text{CPh}_2)\}]$ (**5**) which contains a Mn_2PN_2 five-membered ring.

The complex $[\text{Mn}_2(\text{CO})_8(\mu\text{-PN}^i\text{Pr}_2)]$ (**1**) was readily prepared by the reaction of $\text{K}[\text{Mn}(\text{CO})_5]$ with $\text{Cl}_2\text{PN}^i\text{Pr}_2$. This compound is surprisingly stable and persists for days in THF solution, whereas the related complexes $[\text{Co}_2(\text{CO})_6(\mu\text{-CO})(\mu\text{-PNR}_2)]$ ($\text{NR}_2 = \text{N}^i\text{Pr}_2$, TMP) readily lose carbonyl ligands and undergo condensation reactions to form the cluster complexes $[\text{Co}_4(\text{CO})_8(\mu\text{-CO})\{\mu\text{-PNR}_2\}_2]$ ($\text{NR}_2 = \text{TMP}$, N^iPr_2).⁸ Complex **1** has a ³¹P NMR resonance at δ 731, the downfield chemical shift of which is typical for bridging phosphinidene ligands.⁸ The X-ray structure† shows a pseudooctahedral ligand arrangement about each manganese centre and a bridging aminophosphinidene ligand across the two metal centres ($\text{Mn}(1)\text{--Mn}(2) = 2.9193(4)$ Å).

The reaction of **1** with organic azides RN_3 ($\text{R} = \text{Me}_3\text{Si}$, Me_3Sn , Ph, adamantyl) readily affords the complexes $[\text{Mn}_2(\text{CO})_8\{\mu\text{-}\eta^1, \eta^2\text{-P}(\text{N}^i\text{Pr}_2)=\text{NR}\}]$ ($\text{R} = \text{Me}_3\text{Si}$ (**3a**), Me_3Sn (**3b**), Ph (**3c**), adamantyl (**3d**)). The ³¹P NMR spectrum of **3a** has a resonance at δ 216 and the ¹H NMR spectrum shows the presence of both ⁱPr and Me_3Si groups. The X-ray structure of **3a**† (Fig. 1) shows that oxidation of the phosphinidene phosphorus has occurred with the formation of a novel $\mu\text{-}\eta^1, \eta^2\text{-aminophosphamine}$ ligand. The phosphorus atom of the phosphamine is bonded to Mn(2) ($\text{Mn}(2)\text{--P}(1) = 2.1857(3)$ Å) with an elongated P=N bond ($\text{P}(1)\text{--N}(2) = 1.5756(9)$ Å) coordinated in π -fashion to Mn(1) ($\text{Mn}(1)\text{--P}(1) = 2.4084(3)$ Å, $\text{Mn}(1)\text{--N}(2) = 2.1717(9)$ Å). To our knowledge this is the first example of a phosphamine bound in typical μ -alkenyl fashion in a binuclear complex, although free phosphamines have been previously characterised.¹²

Complexes that are isoelectronic to **3a–d** were produced in the reaction of $[\text{Co}_2(\text{CO})_6(\mu\text{-dppm})(\mu\text{-PNR}_2)]$ ($\text{NR}_2 = \text{N}^i\text{Pr}_2$ (**2a**), TMP (**2b**)) with CH_2N_2 . The ³¹P NMR spectrum of the product ($\text{NR}_2 = \text{N}^i\text{Pr}_2$ (**4a**)) shows a new resonance at δ 247 and two resonances for the $\mu\text{-dppm}$ group at δ 46 and 38. The ¹³C NMR spectrum of **4a** shows a resonance for the CH_2 fragment of the phosphalkene ligand appearing as a broad signal at δ 24. The X-ray structure of **4b**† (Fig. 1) shows that insertion of methylene into a Co–P bond of **2b** has occurred with loss of N_2 and the

^aSteele Institute for Molecular Sciences, National Research Council of Canada, 100 Sussex Drive, Ottawa, Ontario, Canada K1A 0R6

^bThe Ottawa-Carleton Chemistry Institute, University of Ottawa, Ottawa, Ontario, Canada K1N 6N5. E-mail: acarty@pco-bcp.gc.ca; Fax: 001 613 948 6667; Tel: 001 613 948 6672

† Electronic supplementary information (ESI) available: Synthetic and spectroscopic data for compounds **1**, **3a–d**, **4a–b**, **5**; crystallographic details for compounds **1**, **3a**, **3c**, **4b**, and **5**. See <http://dx.doi.org/10.1039/b505472a>





Fig. 1 ORTEP diagrams of compounds **3a**, **4b** and **5**. Selected hydrogen atoms have been eliminated for clarity and thermal ellipsoids are shown at the 50% probability level. Selected bond lengths (Å) and angles (°): **3a**: Mn(1)–Mn(2) 2.8888(3), Mn(1)–P(1) 2.4084(3), Mn(1)–N(2) 2.1717(9), Mn(2)–P(1) 2.1857(3), P(1)–N(1) 1.6512(9), P(1)–N(2) 1.5756(9), N(2)–Si(1) 1.7297(9), Mn(2)–P(1)–N(1) 128.69(3), Mn(2)–P(1)–N(2) 118.04(4), Mn(1)–P(1)–Mn(2) 77.76(1), Mn(1)–P(1)–N(1) 126.23(4), Mn(1)–P(1)–N(2) 61.97(3), N(2)–P(1)–N(1) 113.15(5); **4b**: Co(1)–Co(2) 2.7143(5), Co(1)–P(1) 2.2135(6), Co(1)–P(3) 2.1321(6), Co(2)–P(2) 2.1948(6), Co(2)–P(3) 2.1232(6), Co(1)–C(2a) 2.105(2), N(1)–P(3) 1.669(2), P(3)–C(2a) 1.737(2), Co(1)–P(3)–Co(2) 79.27(2), Co(2)–P(3)–C(2a) 108.92(8), Co(1)–P(3)–C(2a) 64.98(7); **5**: Mn(1)–Mn(2) 2.8945(4), Mn(1)–P(1) 2.1845(5), Mn(2)–N(3) 2.103(1), P(1)–N(1) 1.631(1), P(1)–N(2) 1.593(1), N(2)–N(3) 1.418(2), Mn(1)–P(1)–N(1) 127.71(5), Mn(1)–P(1)–N(2) 125.49(5), N(1)–P(1)–N(2) 106.69(7).

formation of a new $\mu\text{-}\eta^1,\eta^2$ -coordinated aminophosphaalkene complex. The principal structural features are a short Co–Co bond (Co(1)–Co(2) 2.7143(5) Å) and a coordinated phosphalkene (TMP)P=CH₂ bound to Co(2) *via* phosphorus and to Co(1) in π -fashion *via* the P=C bond (P(3)–C(2a) = 1.737(2) Å). The P=C bond length is comparable to that in [Cp₂Mo₂(CO)₄($\mu\text{-}\eta^1,\eta^2$ -P(Ph)=CHMe)], a similar phosphalkene complex synthesized *via* an entirely different route by Mays and co-workers.¹¹ The methine carbon resonance of the CHMe group in [Cp₂Mo₂(CO)₄($\mu\text{-}\eta^1,\eta^2$ -P(Ph)=CHMe)] appears at δ 42 and is similar to the =CH₂ resonance in **4a**. Although the ¹³C chemical shift of C(2a) is more indicative of an sp³ hybridized carbon atom, the short P(3)–C(2a) bond length is consistent with a metal coordinated phosphalkene. We are unaware of any other examples of this class of molecule.

The reaction of **1** with diphenyldiazomethane (Scheme 1) affords a species with a ³¹P NMR resonance at δ 285. Although the ¹H NMR spectrum of **5** shows the presence of both Ph and ⁱPr groups, it was necessary to carry out an X-ray crystallographic study to determine if the N₂ fragment of the diazoalkane was retained.[‡] The structure (Fig. 1) shows that the diazoalkane has inserted into one Mn–P bond without N₂ extrusion to form a five-membered Mn₂PN₂ ring. The dimetallacycle has a puckered conformation which results in a staggered arrangement for the three carbonyl ligands on each metal that are perpendicular to the metal–metal bond. The geometry about each metal centre is pseudooctahedral with a Mn(1)–P(1) separation of 2.1845(5) Å and a Mn(2)–N(3) bond length of 2.103(1) Å. While the stereochemistry about P(1) is trigonal planar, the P(1)–N(1) distance to the diisopropylamino group (1.630(1) Å) is distinctly longer than P(1)–N(2) (1.593(1) Å) and together with the short N(2)–N(3) bond (1.418(2) Å) indicates a significant π -character in the PN₂ portion of the metallacycle. However the P(1)–N(2) bond is slightly longer than the P=N separation of 1.545(2) Å in the free phosphimine (Me₃Si)₂NP=NSiMe₃¹² but is similar to the PN



Scheme 1

bond length in the nickel cluster [Ni₄(CN^{*i*}Bu)₄($\mu_3\text{-}\eta^2\text{-CN}^i\text{Bu}$)($\mu_3\text{-}\eta^2\text{-(Me}_3\text{Si)}_2\text{NP=NSiMe}_3$)₂].¹³

While the reaction of **2** with CH₂N₂ and **1** with Ph₂CN₂ both lead to insertion into the phosphinidene bridge, complex **5** is formed without extrusion of N₂ from the diazoalkane whereas in **4b** loss of N₂ has afforded a phosphathene ligand. Attempts to convert **5** thermally to the corresponding phosphadiphenylethylene complex (80 °C, C₆D₆, overnight) resulted in decomposition.

In conclusion, these results confirm that doubly bridging phosphinidene complexes, like their μ -alkylidene counterparts¹⁴ are highly susceptible to insertion reactions with unsaturated substrates. The electrophilic, unsaturated nature of the μ -PR bridge appears to be only partially alleviated by the presence of the electron donating NR₂ group and through back donation from the metal centres. The reactivity of the PR bridge presents a wealth of new opportunities for the synthesis of novel phosphorus ligands.

This work was supported by the National Research Council of Canada and the Natural Sciences and Engineering Research Council of Canada (to A.J.C.).

Notes and references

‡ Crystallographic data for: **1** [C₁₄H₁₄Mn₂NO₈P], *M* = 465.11, triclinic, *P* $\bar{1}$, *a* = 6.6736(7) Å, *b* = 9.5035(9) Å, *c* = 15.4285(15) Å, α = 104.344(2)°, β = 96.656(2)°, γ = 97.440(2)°, *V* = 928.81(16) Å³, *T* = 125(2) K, *Z* = 2, μ = 1.486, 5109 unique, *R*₁[*I* > 2 σ (*I*)] = 0.0232, *R*_w (all data) = 0.0633. CCDC 275485. **3a**: [C₁₇H₂₃Mn₂N₂O₈PSi], *M* = 552.31, monoclinic, *P*₂/*c*, *a* = 15.7307(13) Å, *b* = 10.5029(8) Å, *c* = 15.5917(13) Å, α = 90°, β = 113.0100(10)°, γ = 90°, *V* = 2371.1(3) Å³, *T* = 125(2) K, *Z* = 4, μ = 1.226, 5835 unique, *R*₁[*I* > 2 σ (*I*)] = 0.0198, *R*_w (all data) = 0.0543. CCDC 275484. **3c**: [C₂₀H₁₉Mn₂N₂O₈P], *M* = 556.22, monoclinic, *P*₂/*c*, *a* = 8.8228(11) Å, *b* = 9.7766(12) Å, *c* = 28.072(3) Å, α = 90°, β = 97.045(2)°, γ = 90°, *V* = 2403.1(5) Å³, *T* = 125(2) K, *Z* = 4, μ = 1.164, 6717 unique, *R*₁[*I* > 2 σ (*I*)] = 0.0236, *R*_w (all data) = 0.0634. CCDC 275483. **4b**: [C₃₉H₄₂Co₂NOP₃], *M* = 799.51, monoclinic, *C*2/*c*, *a* = 30.170(6) Å, *b* = 10.6655(19) Å, *c* = 23.493(4) Å, α = 90°, β = 102.347(3)°, γ = 90°, *V* = 7385(2) Å³, *T* = 125(2) K, *Z* = 8, μ = 1.070, 10308 unique, *R*₁[*I* > 2 σ (*I*)] = 0.0347, *R*_w (all data) = 0.0914. CCDC 269680. **5**: [C₂₉H₂₉Mn₂N₃O_{8.5}P], *M* = 696.40, monoclinic, *P*₂/*c*, *a* = 12.1027(16) Å, *b* = 20.563(3) Å, *c* = 13.6644(18) Å, α = 90°, β = 111.888(2)°, γ = 90°, *V* = 3155.5(7) Å³, *T* = 125(2) K, *Z* = 4, μ = 0.905, 7871 unique, *R*₁[*I* > 2 σ (*I*)] = 0.0322, *R*_w (all data) = 0.0887. CCDC 269679. See <http://dx.doi.org/10.1039/b505472a> for crystallographic data in CIF or other electronic format.

- 1 K. B. Dillon, F. Mathey and J. F. Nixon, *Phosphorus, The Carbon Copy*, John Wiley & Sons Ltd., Chichester, 1998.
- 2 *Multiple Bonds and Low Coordination in Phosphorus Chemistry*, ed. M. Regitz and O. J. Scherer, Georg Thieme Verlag, Stuttgart, 1991; A. T. Termaten, M. Schakel, A. W. Ehlers, M. Lutz, A. L. Spek and K. Lammertsma, *Chem. Eur. J.*, 2003, **9**, 3577.
- 3 P. B. Hitchcock, M. F. Lappert and W.-P. Leung, *J. Chem. Soc., Chem. Commun.*, 1987, 1282; T. L. Breen and D. W. Stephan, *J. Am. Chem.*

- Soc.*, 1995, **117**, 11914; A. T. Termaten, T. Nijbacker, M. Schakel, M. Lutz, A. L. Spek and K. Lammertsma, *Chem. Eur. J.*, 2003, **9**, 2200; A. H. Cowley, *Acc. Chem. Res.*, 1997, **30**, 445; C. C. Cummins, R. R. Schrock and W. M. Davis, *Angew. Chem.*, 1993, **105**, 758; J. B. Bonanno, P. T. Wolczanski and E. B. Lobkovsky, *J. Am. Chem. Soc.*, 1994, **116**, 11159; D. S. J. Arney, R. C. Schnabel, B. C. Scott and C. J. Burns, *J. Am. Chem. Soc.*, 1996, **118**, 6780.
- 4 B. T. Sterenberg, K. A. Udachin and A. J. Carty, *Organometallics*, 2001, **21**, 2657; B. T. Sterenberg, K. A. Udachin and A. J. Carty, *Organometallics*, 2003, **22**, 3927.
- 5 J. Sánchez-Nieves, B. T. Sterenberg, K. A. Udachin and A. J. Carty, *J. Am. Chem. Soc.*, 2003, **125**, 2404; R. Melenkivitz, D. J. Mindiola and G. L. Hillhouse, *J. Am. Chem. Soc.*, 2002, **124**, 3846; R. Waterman and G. L. Hillhouse, *J. Am. Chem. Soc.*, 2003, **125**, 13350; A. J. Termaten, H. Aktas, M. Schakel, A. W. Ehlers, M. Lutz, A. L. Spek and K. Lammertsma, *Organometallics*, 2003, **22**, 1827.
- 6 J. B. M. Wit, G. T. van Eijkel, M. Schakel and K. Lammertsma, *Tetrahedron*, 2000, **56**, 137; J. B. M. Wit, G. T. van Eijkel, F. J. J. de Kanter, M. Schakel, A. W. Ehlers, M. Lutz, A. L. Spek and K. Lammertsma, *Angew. Chem., Int. Ed.*, 1999, **38**, 2596; A. Marinetti and F. Mathey, *Organometallics*, 1984, **3**, 456; A. Marinetti and F. Mathey, *J. Am. Chem. Soc.*, 1982, **104**, 4484; N. H. T. Huy and F. Mathey, *J. Org. Chem.*, 2000, **65**, 652.
- 7 J. Holton, M. F. Lappert, R. Pearce and P. I. W. Yarrow, *Chem. Rev.*, 1983, **83**, 135; W. A. Herrmann, *Pure Appl. Chem.*, 1982, **54**, 65; S. A. R. Knox, *J. Organomet. Chem.*, 1990, **400**, 255; K. H. Theopold and R. G. Bergman, *J. Am. Chem. Soc.*, 1981, **103**, 2489.
- 8 J. Sánchez-Nieves, B. T. Sterenberg, K. A. Udachin and A. J. Carty, *J. Cluster Sci.*, 2004, **15**, 151; J. Sánchez-Nieves, B. T. Sterenberg, K. A. Udachin and A. J. Carty, *Can. J. Chem.*, 2004, **82**, 1507.
- 9 M. Esther García, V. Riera, M. A. Ruiz, D. Sáez, J. Vaissermann and J. C. Jeffrey, *J. Am. Chem. Soc.*, 2002, **124**, 14304; H. Lang, O. Orama and G. Huttner, *J. Organomet. Chem.*, 1985, **291**, 293; G. Huttner and K. Evertz, *Acc. Chem. Res.*, 1986, **19**, 406; A. M. Arif, A. H. Cowley and M. Pakulski, *J. Chem. Soc., Chem. Commun.*, 1985, 1707; A. H. Arif, A. H. Cowley, N. C. Norman, A. G. Orpen and M. Pakulski, *Organometallics*, 1988, **7**, 309.
- 10 K. Lammertsma, *Top. Curr. Chem.*, 2003, **229**, 119.
- 11 J. E. Davies, M. J. Mays, P. R. Raithby and A. D. Woods, *Chem. Commun.*, 1999, 2455.
- 12 S. Pohl, *Chem. Ber.*, 1979, **112**, 3159.
- 13 O. J. Scherer, R. Walter and W. S. Sheldrick, *Angew. Chem., Int. Ed. Engl.*, 1985, **24**, 115.
- 14 M. Akita, R. Hua, S. A. R. Knox, Y. Moro-oka, S. Nakanishi and M. I. Yates, *Chem. Commun.*, 1997, 51; P. M. Maitlis, H. C. Long, R. Quyoum, M. L. Turner and Z.-Q. Wang, *Chem. Commun.*, 1996, 1; C. E. Sumner, Jr., P. E. Riley, R. E. Davis and R. Pettit, *J. Am. Chem. Soc.*, 1980, **102**, 1752.