

# Polyoxygenated Steroids of Marine Origin

Maria Valeria D'Auria, Luigi Minale,\* and Raffaele Riccio

Dipartimento di Chimica delle Sostanze Naturali, Università di Napoli Federico II, Via D. Montesano 49, 80131 Napoli, Italy

Received February 4, 1993 (Revised Manuscript Received April 30, 1993)

## Contents

I. Introduction	1839
II. Polyoxygenated Steroids of Marine Algae	1839
A. Red Algae	1839
B. Brown Algae	1840
C. Microalgae	1841
III. Polyoxygenated Steroids of Porifera (Sponges)	1841
IV. Polyoxygenated Steroids of Coelenterata	1849
A. Hydrozoa	1849
B. Anthozoa	1849
1. Hexacorallia	1849
2. Octocorallia	1851
V. Polyoxygenated Steroids of Bryozoa	1858
VI. Polyoxygenated Steroids of Marine Mollusca	1858
VII. Polyoxygenated Steroids of Echinodermata	1860
A. Asteroidea (Starfish)	1861
1. Polyhydroxysteroids	1861
2. Glycosides of Polyhydroxysteroids	1870
3. Asterosaponins	1880
4. Cyclic Steroidal Glycosides	1887
B. Ophiuroidea (Brittle Stars)	1887
VIII. Polyoxygenated Steroids of Arthropoda	1889
IX. Polyoxygenated Steroids of Tunicates	1890
X. Polyoxygenated Steroids of Vertebrates (Fish)	1891

## I. Introduction

Sterols of marine organisms have been found to be comprised of most complex mixtures. Over 200 new monohydroxysterols have so far been identified from marine sources, and evidence has been presented for the occurrence of over 70 sterols from a single marine species.<sup>1</sup> Sterol patterns in marine invertebrates reflect the complexity of sterols arising through food chains. The capability of further biochemical modification of dietary sterols makes the sterol mixtures even more complex. The symbiotic relationship between organisms also complicates the sterol compositions.<sup>2</sup>

In the last 20 years many sterols with unprecedented structures have been isolated from marine sources. For many years the known carbon skeleton of sterols ranged from C<sub>27</sub> to C<sub>29</sub>, and the carbon variation occurred exclusively in the side chain at C<sub>24</sub>.<sup>3</sup> After the discovery of the C<sub>26</sub>-sterols, first detected in 1970 from the mollusk *Placopecten Magellanicus*<sup>4</sup> and later found widespread in marine invertebrates and also in a marine phytoplankton,<sup>5</sup> thus suggesting that all the marine C<sub>26</sub>-sterols originate from phytoplankton, a number of "nonconventional" sterols have been reported. These include sterols having side chains modified by the apparent loss of carbon atoms or by the addition of extra carbon atoms at biogenetically unprecedented positions of a normal C<sub>8</sub> side chain, as well sterols with

unusual nuclei. The discovery of novel sterols as major components of the extracts of marine organisms is not the rule, although there have been many exceptions, especially from sponges. Typical examples are aplysterol (9) and 24(28)-didehydroaplysterol (10), the first sterols found<sup>6</sup> with a methyl group at C-26, which have been found as the major sterols of the sponges of the genus *Aplysina* (*Verongia*); calysterol (51), the major sterol (90% of the sterol mixture) of the sponge *Calyx nicaensis*;<sup>7</sup> petrosterol (48), isolated from the sponge *Petrosia ficiformis*;<sup>8,9</sup> strongylosterol (27), the sole sterol of *Strongylophora durissima*;<sup>10</sup> and xestosterol (15) and sutinasterol (37), apparently the result of four biomethylation, isolated as the predominant sterols of *Xestospongia muta*<sup>11</sup> and a *Xestospongia* sp.,<sup>12</sup> respectively. A series of 19-nor-5 $\alpha$ -stanols and A-norstanols, both with conventional and nonconventional side chains, again from marine sponges, have been firstly isolated as the major sterol components from *Axinella polypoides*<sup>13</sup> and *A. verrucosa*,<sup>14</sup> respectively. One more interesting case of discovery of a novel sterol as major component of a mixture is dinosterol (11) from *Gonyaulax tamarensis*,<sup>15</sup> which is the most characteristic sterol of dinoflagellates and possibly the biological precursor of the cyclopropyl sterol gorgosterol (45) found in zooxanthellae-containing hosts.<sup>16</sup> Structures of the nonconventional side chains of marine monohydroxysterols are shown in Figure 1.

Several excellent reviews on the structures, biosynthesis and distribution of marine sterols are available.<sup>2,55-62</sup> A paper dedicated to the biosynthesis of marine sterols side chain is published in this issue. This review focuses on polyoxygenated sterols from marine organisms, which are a growing group of metabolites with potential interesting biological and pharmacological activities and is intended to be all inclusive with respect to its coverage. Polyhydroxysteroids have been found in algae, and virtually in every marine invertebrata phyla, *i.e.* Porifera, Coelenterata, Bryozoa, Mollusca, Echinodermata, Arthropoda, and Tunicata as well as in fish.

## II. Polyoxygenated Steroids of Marine Algae

Polyoxygenated sterols have been occasionally isolated from red (Rhodophyta) and brown (Phaeophyta) algae, but have not yet been reported from the green (Chlorophyta) ones (Figure 2).

### A. Red Algae

Dihydroxy sterols 57 and 58 along with 24,25-epoxycholesterol 59 have been isolated as minor components of *Asparagopsis armata*<sup>63</sup> and *Rissoella verruculosa*;<sup>64</sup> 57 and 58 have been also isolated from *Rhodymenia palmata*<sup>65</sup> and the diol 58 has been found



Maria Valeria D'Auria was born in Naples, Italy, in 1958. She received her degree (Laurea) in chemistry in 1981 from the Università di Napoli. In 1983 she was appointed as Ricercatore at the Dipartimento di Chimica delle Sostanze Naturali, Università di Napoli Federico II. Her research interests include the echinoderm chemistry and, recently, the study of antitumor macrolides from sponges.



Luigi Minale was born in Tripoli, Libia, in 1936 and obtained his Laurea in Chemistry (1960) and Libera Docenza (1967) from the University of Naples. From 1961 to 1969 he was at the Institute of Organic Chemistry of the University of Naples with Professors R. A. Nicolaus and M. Piattelli. In 1969, he moved to the Institute for the Chemistry of Molecules of Biological Interest, C.N.R., Arco Felice and became Director of this institute in 1973 (until 1981) as well as, almost at the same time, Professor of Organic Chemistry at the University of Catania (1975). In the 1977 he moved to the University of Napoli Federico II, Faculty of Pharmacy, and became Director of Department of the Chemistry of Natural Products in 1984 (until 1990). He has been visiting Professor at the University of Aberdeen (Scotland, U.K.; 1973) and at the Tohoku University, Sendai (Japan; 1986). Professor Minale has published more than 200 papers in the chemistry of natural products and organic chemistry. He is member of the Advisory Board of the Gazzetta Chimica Italiana.

in *Liagora distenta*<sup>66</sup> and *Scinaia furcellata*.<sup>66</sup> It has been suggested that these side chain oxygenated sterols may be artifacts caused by autoxidation during the isolation process. Francisco *et al.*<sup>63</sup> have shown that **57** and **58** can indeed be produced by the autoxidation of desmosterol (cholesta-5,24-dien-3 $\beta$ -ol); they are nevertheless present in the fresh red alga *A. armata*.

Two ecdysone-like sterols, pinnasterol (**60**) and acetylpinnasterol (**61**), both showing biological activity as molting hormones, have been isolated from *Laurencia pinnata*.<sup>67</sup> The structures were secured by X-ray crystallography of the acetylpinnasterol (**61**). In contrast with ecdysone and the many ecdysteroids isolated from arthropods and certain plant species, all having the 22*R* configuration, which is very important for high



Raffaele Riccio was born in Napoli, Italy, in 1948. He obtained his degree (Laurea) in Chemistry in 1972 from the University of Napoli and in 1973 was appointed as Ricercatore of the Consiglio Nazionale delle Ricerche at the Istituto per la Chimica di Molecole di Interesse Biologico, Arco Felice, where he began his research activity in the field of marine natural products working on the chemistry and sterol biogenesis of Mediterranean sponges. From 1976 to 1977 he spent one year in a post-doctoral position with Paul J. Scheuer at the University of Hawaii. In Napoli again he became interested in the chemistry of polyhydroxylated steroidal glycosides from echinoderms. In 1987 he moved at the Faculty of Pharmacy of the Università di Napoli Federico II as Associated Professor of Organic Chemistry. He has spent research leaves with Koji Nakanishi at Columbia University, in 1981, and with Bill Fenical at Scripps Institution of Oceanography, in 1990. He has published more than 100 papers in the field of natural products and organic chemistry.

activity, the algal pinnasterol and acetylpinnasterol possess the 22*S* configuration. The same configuration has been found in (20*R*,22*S*)-5 $\alpha$ -cholesta-9(11),24(25)-diene-3 $\beta$ ,6 $\alpha$ ,20,22-tetrol, the aglycon of the asterosaponin protoreasteroside (**497**) isolated from the starfish *Protoreaster nodosus* and *Pentaceraster alveolatus*.<sup>68</sup>

5 $\alpha$ -Cholestane-3,6-dione (**62**) has been reported from some red algae species,<sup>69</sup> and recently the novel 11 $\alpha$ -hydroxy-5 $\alpha$ -cholestane-3,6-dione (**63**), of potential utility as corticosteroid intermediate, has been isolated from *Acanthophora spicifera*.<sup>70</sup> Another diketosteroid, 5 $\beta$ -cholest-3-ene-7,11-dione (**64**) was isolated from *Hypnea musciformis*.<sup>71</sup> Recently 2 $\alpha$ -oxa-2-oxo-5 $\alpha$ -hydroxy-3,4-dinorcholestane (**65**), the first example of a ring A-dinorsteroid from a natural source, has been isolated from *Laurencia obtusa*<sup>72</sup> as a minor component. The structure was confirmed by synthesis from 2-(hydroxymethylene)cholest-4-en-3-one and the 5 $\alpha$ -hydroxy stereochemistry was determined by NMR experiments.

## B. Brown Algae

The first dihydroxysteroid isolated from a marine source was saringosterol (**66**) identified by Ikekawa *et al.*<sup>73</sup> in two brown algae. Since then **66** and 24-oxocholestane (**67**) have been isolated<sup>74-76</sup> from a number of brown algae. Recently the presence of **66** and **67** along with 24,28-epoxyfucosterol (**68**) in the brown alga *Hizikia fusiformis* has been reported.<sup>77</sup> Their origin was considered as doubtful: they could have been *bona fide* constituents of the algae or be artifacts caused by oxidation during the isolation process. Knights<sup>74</sup> showed that the presence of saringosterol (**66**) and the 24-ketosteroid **67** in the brown alga *Ascophyllum nodosum* were artifacts arising from autoxidation during the air-drying of the alga, the fresh material containing only fucosterol. An intriguing steroid diol with a novel

side chain 69 was isolated from *Desmarestia aculeata*.<sup>78</sup> An unusual sterol with a 24-(vinyl-23-ene) side chain 70 was isolated from *Sargassum thumbergii*<sup>79</sup> as a minor component, the predominant sterol of the mixture being fucosterol. Artifacts of formation of 70 appear very unlikely and the authors have suggested a biogenetically possible process involving the removal of some leaving group at C-23 of fucosterol 24,28-epoxide, with simultaneous cleavage of the C-24,28 bond.

Steroids containing a 3,5-dien-7-one and 3,5-diene 7 $\alpha$ -hydroxycholestane nucleus were isolated from *Fucus evanescens*.<sup>75</sup> Ikekawa and colleagues have suggested that the 3,5-dien-7-ones are probably formed during the isolation procedure from the 3 $\beta$ -hydroxy-5-en-7-one counterparts.<sup>75</sup> We would note that the steroid  $\Delta^5$ -3 $\beta$ ,7 $\alpha$ - and  $\Delta^5$ -3 $\beta$ ,7 $\beta$ -diols, along with the  $\Delta^5$ -3 $\beta$ -hydroxy 7-ones, are well recognized autoxidation products of  $\Delta^5$ -sterols,<sup>80</sup> thus leaving the origin of the nuclear oxygenated sterols of *F. evanescens* as doubtful.

### C. Microalgae

The Haptophyceae are microscopic unicellular algae, which are widely distributed in the ocean and often constitute a major proportion of marine phytoplankton. A novel sterol sulfate, with the dinosterol side chain, hymenosulfate, 71 has been recently isolated from the cultured marine haptophyte *Hymenomonas* sp.<sup>81</sup> Sulfation is typical in the biosynthesis of secondary metabolites in many marine invertebrates, especially echinoderms, but hymenosulfate is the first report of a sterol sulfate from marine microalgae. This steroid has a potent SR (sarcoplasmic reticulum) Ca-releasing activity.

### III. Polyoxygenated Steroids of Porifera (Sponges)

The sponges have yielded the most varied and biogenetically unprecedented array of sterols found among the invertebrate phyla. Most of the 200 new monohydroxysterols found in marine organisms have been isolated from the sponges.<sup>62</sup> The uniqueness of sterols in the cell membranes of sponges is probably related to the presence of unusual fatty acids in their phospholipids.<sup>82-85</sup>


In addition to the monohydroxylated sterols, a number of polyoxygenated sterols have been isolated from sponges; most of them have appeared in the literature in the last two years.

Halistanol sulfate (72) isolated from *Halichondria moorei*<sup>86</sup> is the first example of the growing list of sulfated polyhydroxysteroids from sponges (Figure 3), which are very attractive because of their potential activity against HIV virus.<sup>87,88</sup> Both the *tert*-butyl moiety at the end of side chain and the 2 $\beta$ ,3 $\alpha$ ,6 $\alpha$ -trihydroxy functions in halistanol are biosynthetically intriguing. Another trisulfated steroid, sokotrosterol sulfate (73), has been isolated together with the previous 72 from two Halichondriidae species.<sup>89</sup> The side chain of 73 is unprecedented and involves a "normal" alkylation at C-24 of a standard C<sub>8</sub> side chain and the addition of two extra methyl groups at C-26 and one extra methyl group at C-25. Halistanol and sokotrosterol belong to an unusual class of steroids which possess side chains with quaternary carbons. The side chain of 73 is related


to that of mutasterol (26), and the authors<sup>89</sup> have proposed a biosynthesis *via* three consecutive SAM biomethylations of codisterol (24*S*-methyl-5,25-cholestadien-3 $\beta$ -ol) or epicodisterol (24*R*). Recently in a continuing search for the biosynthesis of marine lipids, Djerassi and his group<sup>90</sup> have elucidated the biosynthesis of mutasterol in the Caribbean sponge *Xestospongia muta* by feeding selected radioactive precursors and have shown that codisterol is efficiently transformed (10 times faster than its epimer) into mutasterol (26). Interestingly, the isolation of the 26-norsokotrosterol sulfate (74) with the side chain very similar with that of mutasterol (26), the difference being in the position of the double bond, from the sponge *Trachyopsis halichondroides* (family Halichondriidae) has been reported.<sup>91</sup> Very recently Fusetani and his group<sup>92</sup> reported the isolation of four more steroids (75-78) with the halistanol sulfate nucleus from the sponge *Epipolasis* sp. The unusual side chain of halistanol sulfate B (76) is unprecedented for naturally occurring sterols.<sup>62</sup> A fifth new sulfate steroid isolated from the same sponge, halistanol sulfate E (79) is related to 72 by the introduction of an additional hydroxyl group at C-15.<sup>92</sup> Two more steroids with the halistanol sulfate nucleus (80 and 81), showing anti-HIV activity, have been isolated from the sponge *Pseudoaxinissa digitata* (order Axinellida, family Axinellidae).<sup>93</sup>

Three novel sterol sulfates (82-84), sharing the same tetracyclic nucleus with a sulfate at C-3 and the 19-methyl group oxidized to a carboxyl group, have been isolated from the sponge *Toxadocia zumi*.<sup>94</sup> Antimicrobial, antifeedant, and cytotoxic properties were reported for these steroid sulfates, and the authors suggest that they might be in part responsible for the lack of fouling organisms on *Toxadocia zumi*.<sup>94</sup> A 3 $\beta$ -sulfoxy-4 $\beta$ -hydroxypregnane (85) has been isolated from the sponge *Stylopus australis*.<sup>95</sup> A novel group of sterol sulfates (86-90) was discovered as antiviral substances in the sponge *Petrosia weinbergi*.<sup>87,88</sup> The structure of the major weinbersterol disulfate A (86) was determined by a 2D INADEQUATE NMR experiment performed on a 150-mg sample in conjunction with its HMBC, HETCOR, COLOC, and COSY data, which established the complete connectivity for the molecule.<sup>87</sup> Weinbersterol disulfate B (87) is related to A (86), but lacks the hydroxyl group at C-20 and a further hydroxyl group is located at C-18. Both compounds possess an unprecedented cyclopropane-containing side chain, showing one more examples for the diversity in the side-chain structures of sponge sterols, and exhibited *in vitro* activity against the feline leukemia virus. Weinbersterol disulfate A (86) also showed activity against the human immunodeficiency virus. Three minor sterol sulfates (ortho ester disulfates A-C, 88-90) in the same sponge were isolated, and their structures were found to have an ortho ester functionality.<sup>88</sup> They appear the first reported examples in the steroid class of this particular combination of functionalities. It is of interest to note that the trans diaxial disulfate (2 $\beta$ ,3 $\alpha$ ) and the trans AB ring juncture (5 $\alpha$ -H) have been previously encountered in the 5 $\alpha$ -cholestane 2 $\beta$ ,3 $\alpha$ ,26-triol sulfates (530) isolated from an echinoderm, the ophiuroid *Ophiaracna incrassata*.<sup>96</sup>


An array of polyoxygenated sterols have been isolated from sponges of the genus *Dysidea* (Figure 4). The




1, *Damiriana hawaiiana*<sup>17</sup>  
(sponge)




2, *Placopecten magellanicus*<sup>18</sup>  
(scallop)




3, *Placopecten magellanicus*<sup>18</sup>  
(scallop)




4, *Placopecten magellanicus*<sup>18</sup>  
(scallop)




5, ocellasterol  
*Pseudoptamilla ocellata*<sup>19</sup>  
(annelid)




6, dinoflagellates<sup>20</sup>




7, *Halichondria panicea*<sup>21</sup>  
(sponge)




8, *Pseudoaxinella lunacharia*<sup>22</sup>  
(sponge)




9, aplysterol  
*Aplysina aerophoba*<sup>6</sup>  
(sponge)




10, *Aplysina aerophoba*<sup>6</sup>  
(sponge)




11, dinosterol  
*Gonyaulax tamarensis*<sup>15</sup>  
(dinoflagellate)




12, peridinosterol  
*Peridinium foliaceum*<sup>23</sup>  
(dinoflagellate)




13, *Gonyaulax monilata*<sup>24</sup>  
(dinoflagellate)




14, ficisterol  
*Petrosia ficiformis*<sup>25</sup>  
(sponge)




15, xestosterol  
*Xestospongia muta*<sup>11</sup>  
(sponge)




16, verongulasterol  
*Verongula cauliformis*<sup>26</sup>  
(sponge)




17, *Eutreptia viridis*<sup>27</sup>  
(euglenid)




18, *Pseudoaxinissa* sp.<sup>28</sup>  
(sponge)




19, *Pseudoaxinissa* sp.<sup>28</sup>  
(sponge)




20, *Verongula cauliformis*<sup>29</sup>  
(sponge)




21, stelliferasterol  
*Jaspis stellifera*<sup>30</sup>  
(sponge)




22, isostelliferasterol  
*Jaspis stellifera*<sup>30</sup>  
(sponge)




23, reimersterol  
*Sinularia remei*<sup>31</sup>  
(soft coral)




24, siapalosterol  
*Sinularia siapalosa*<sup>32</sup>  
(soft coral)




25, *Sinularia ramulosa*<sup>33</sup>  
(soft coral)




26, mutasterol  
*Xestospongia muta*<sup>34</sup>  
(sponge)




27, strongylosterol  
*Strongylophora durissima*<sup>10</sup>  
(sponge)




28, durissimasterol  
*Strongylophora durissima*<sup>35</sup>  
(sponge)




29, pulchrasterol  
*Aciculites pulchra*<sup>36</sup> (sponge)




30, *Pseudoaxinissa* sp.<sup>37</sup>  
(sponge)



31, *Pseudoaxinissa* sp.<sup>37</sup>  
(sponge)




32, *Strongylophora durissima*<sup>35</sup>  
(sponge)



**Figure 1.** Nonconventional side chains of marine monohydroxysterols. All compounds possess the cholest-5-ene tetracyclic nucleus except 12 and 13 (4 $\alpha$ -methylcholestane nucleus), 14 (5 $\alpha$ -cholestane nucleus), 17 (cholesta-5,7-diene nucleus), and 29 (cholest-7-ene nucleus).

first example of this group of steroids is 9 $\alpha$ ,11 $\alpha$ -epoxycholest-7-ene-3 $\beta$ ,5 $\alpha$ ,6 $\beta$ ,19-tetrol 6-acetate (91) isolated from an unidentified species of *Dysidea*

collected in Guam.<sup>97</sup> The 19-hydroxyl group was previously found in one sterol from the soft coral *Lithophyton viridis*<sup>98</sup> and later in many sterols from




**Figure 2.** Polyoxygenated sterols from marine algae.

*Dysidea*. The 9,11-epoxide is the most novel aspect of this sterol. The shape of the 11-H proton signal (doublet with  $J = 4.7$  Hz) in the  $^1\text{H}$  NMR of 91 was the key argument supporting the  $9\alpha,11\alpha$  stereochemistry, as the 3.5-Hz coupling between H-6 and H-7 was considered by the authors<sup>97</sup> more compatible with a  $6\beta$ -acetoxy than with a  $6\alpha$ -acetoxy stereochemistry. The stereochemistry at C-6 has subsequently been revised by Fujimoto *et al.*<sup>99</sup> as the  $6\alpha$ -isomer through further NMR measurements involving the evaluation of the pyridine-induced shifts of the protons at C-4 of the corresponding 6-ol.

Herbasterol (92), a polyhydroxylated 9,11-secosterol, is responsible for the ichthyotoxic and antimicrobial

activities observed in the methanol extract of *Dysidea herbacea*.<sup>100</sup> The structure of 92 was elucidated by interpretation of spectral data and conversion, on either acid or basic treatment, into 19-norherbasterol (93). During the retroaldol reaction, the stereochemistry at C-6 undergoes inversion, and the authors<sup>100</sup> explained this phenomenon by assuming an A/B cis ring junction in herbasterol (92), the inversion occurring at C-10 during the retroaldol reaction. A further toxic polyoxygenated sterol 94 has been isolated from the Mediterranean *Dysidea tupa*.<sup>101</sup> A group of eight new polyhydroxysterols, all showing a common  $5\alpha$ -cholest-7-ene- $2\alpha,3\beta,5,6\beta,9\alpha,11\alpha,19$ -heptol framework 95 and various conventional side chains, have been isolated




**Figure 3.** Steroid sulfates from sponge.

from *Dysidea etheria*,<sup>102</sup> collected in Bermuda. The 6 $\beta$ -hydroxy stereochemistry has been confirmed by evaluating the pyridine-induced shifts of the protons at C-4 according to the method suggested by Fujimoto *et al.*<sup>99,103</sup> Specifically, the chemical shift of the 4 $\beta$ -proton at  $\delta$  3.12 requires the hydroxyl group at C-6 to be in the  $\beta$ -configuration. The major sterol of this group 95 with R<sub>1</sub> = R<sub>2</sub> = Ac and the cholesterol side chain was active in the cytotoxic KB assay.<sup>102</sup> Three minor polyhydroxylated sterols 96–98 in the same sponge were isolated and their structures were found to be analogous to those discovered earlier 95, but distinguished by a

5 $\beta$ -skeleton,<sup>104</sup> which is a rare feature among sterols from sponges. Herbasterol (92) from *Dysidea herbacea*<sup>100</sup> and a series of coprostanols from *Petrosia ficiformis*<sup>105</sup> are the only other examples of 5 $\beta$ -sterols from Porifera. The authors<sup>104</sup> have suggested that both the 5 $\alpha$ ,6 $\beta$ - and 5 $\beta$ ,6 $\alpha$ -hydroxylated series cooccurring in the same sponge could arise from a 5 $\alpha$ ,6 $\alpha$ -epoxide common precursor. More recently another group of polar sterols 99–102 have been discovered in a specimen of *Dysidea herbacea*,<sup>106</sup> collected near Massawa, Ethiopia. Strong support for the assignment of configurations at C-6 of the two epimers 100 and 101 was





**Figure 4.** Polyhydroxysteroids from sponges of the genus *Dysidea*.

derived from the comparison of the  $^1\text{H}$  NMR data. The small coupling constant between H-6 and H-7 (<1 Hz) in 100, when compared with  $J$  of 5.5 Hz observed in the spectrum of 101, determined the configuration 6 $\alpha$ -OAc in 100 and the 6 $\beta$ -OAc in 101. The NOE observed between 19-Me and H-6 in 100 reinforced these assignments. Extensive NOE measurements also yielded data that allow the stereochemistry of 102 to be determined.<sup>106</sup> Two other polyhydroxysteroids 103 and 104 were reported from the sponge *Dysidea fragilis*<sup>107</sup> collected in the Black Sea and their structures were found to be related to 95 earlier isolated from *D. etheria*,<sup>102</sup> but lacking the 19-hydroxyl group.

Several other polyoxygenated sterols were isolated from sponges (Figure 5). Three pregnane steroids 105–107 were isolated from *Haliclona rubens*.<sup>108</sup> They appeared to be the first example of pregnane derivatives isolated from marine sources. Two novel unusual pregnanes, agnatasterone A (108) and B (109), pregnatrienolones without much relationship on reference steroids, have recently been discovered in *Axinella agnata*<sup>109</sup> collected off Roscoff, France, and in Northern area of Ile de Batz. The pregnane structures of 105–109 immediately suggest steroid hormones, but the biogenesis and the role of these sponge-derived pregnane steroids have yet to be elucidated.



Sterols with the  $\Delta^7$ - $3\beta,5\alpha,6\beta$ -triol nucleus of **99** have previously been described from *Spongionella gracilis*,<sup>110</sup> *Hippospongia communis*,<sup>111</sup> *Spongia officinalis*,<sup>111</sup> and *Ircinia variabilis*.<sup>111</sup> Sterols with this nucleus have also been isolated from the bryozoan *Myriapora truncata*<sup>112</sup> and from the scallop *Patinopekten yessoensis*.<sup>113</sup> Very recently the 24-ethylcholestane- $3\beta,5\alpha,6\beta$ -triol and its 6-keto derivative have been isolated from the sponge *Spirastrella incostans*.<sup>114</sup> The  $6\beta$ -hydroxy stereochemistry in the  $\Delta^7$  derivatives was deduced from spectral data and supported by comparison with synthetic model compounds.<sup>113</sup> *Spongionella gracilis*<sup>115</sup> also gave  $\Delta^7$ - $3\beta,6\alpha$ -dihydroxysterols (**110**), *Spongia officinalis*<sup>116</sup> gave  $\Delta^7$ - $3\beta,5\alpha,6\beta,9\alpha$ -tetrahydroxysterols (**111**) and *Hippospongia communis*<sup>117,118</sup> has yielded 5,6-secosterols (**112**) with various conventional side chains. The  $\Delta^7$ - $3\beta,5\alpha,6\beta$ -trihydroxysterols have been isolated from sponges which contain  $\Delta^{5,7}$ -sterols as main sterol components and it has been suggested<sup>110</sup> that they could arise from  $\Delta^{5,7}$ -sterols. The 5,6-secosterols could be in turn derived from the cleavage of the C-5/C-6 bond of the  $\Delta^7$ - $3\beta,5\alpha,6\beta$ -trihydroxysterols.<sup>117</sup> We note that the polyoxygenated derivatives of ergosterol characterized by  $\Delta^7$ - $3\beta,5\alpha,6\beta$ -triol and  $\Delta^7$ - $3\beta,5\alpha,6\beta,9\alpha$ -tetrol partial structures were previously isolated from the terrestrial fungus *Polyporus versicolor*.<sup>119</sup>

More recently a group of norsterols, incisterols (**113** and conventional  $\Delta^{22}$  side chains), with an unprecedented highly degraded skeleton have been isolated from the Mediterranean sponge *Dictyonella incisa*.<sup>120</sup> The authors have proposed that they could arise from  $\Delta^{5,7}$ -sterols, through the formation of  $3\beta,5\alpha,6\beta$ -triols, both of which are present in *D. incisa*, followed by a successive cleavage of the C-5/C-6 linkage to give 5,6-secosteroids related to **112**. Subsequent cleavage of the C-9/C-10 linkage would result in the removal of the entire ring A including the 19-methyl group. The incisterols, by the authors own admission, are probably artifacts of methanol incorporation, since methanol was used in the extraction. An endoperoxide or a keto acid are suggested as the possible natural products.<sup>120</sup>




Recently a few more secosterols have been recovered from sponges. A new 9,11-secosterol **114** has been isolated from the Mediterranean sponge *Spongia officinalis*.<sup>121</sup> 9,11-Secosterols were earlier found in a gorgonian<sup>122</sup> and in a soft coral of the genus *Sinularia*<sup>123,124</sup> and later in the sponge *Dysidea herbacea*.<sup>100</sup> A 8,9-secosteroid, jereisterol A (**115**), and an 8,14-secosteroid, jereisterol B (**116**), have been isolated from the Pacific sponge *Jereicopsis graphidiophora*.<sup>125</sup> Jereisterol A and B combine the unique  $3\beta$ -methoxy group with a rare secostructure. Surprisingly this sponge lacks the usual  $3\beta$ -hydroxysterols, but contains,  $3\beta$ -methoxysteroids with normal  $\Delta^5, \Delta^0$  and the rare  $\Delta^8, \Delta^8$  and  $\Delta^{7,9(11)}$  nuclei; some others have further oxygenation in the nucleus (**117-121**).<sup>126</sup>

Anthosterones A (**122**) and B (**123**), which represent the first examples of a new type of ring A contraction in a steroid nucleus, have been isolated from the sponge *Anthoracuata graceae*.<sup>127</sup> The structure of anthosterone A (**122**) was verified by single-crystal X-ray diffraction analysis.<sup>127</sup> The authors suggested a possible biosynthesis of the anthosterone nucleus through a benzylic acid rearrangement of a 2,3-diketosteroid precursor as a ring contraction step.  $\Delta^4$ -3,6-Diketo-


steroids **124** were also isolated in the same sponge.<sup>127</sup> Steroidal  $\Delta^4$ -3,6-diketones have been then found in the sponge *Geodia cydonium*<sup>128</sup> and *Cinachyra tarentina*,<sup>129</sup> where they cooccur with the more common steroidal  $\Delta^4$ -3-ketones.  $\Delta^{4,7}$ -3,6-Diketones **125** with the conventional 24-methyl-24-ethyl and cholesterol side chains were isolated from *Raphidostila incisa*,<sup>130</sup> where they cooccur with a major amount of 5,8-epidioxy- $\Delta^6$ -sterols. Three unusual steroidal  $\Delta^{5,8(9)}$ - $3\beta$ -hydroxy 7-ketones (**126**) have been isolated from the sponge *Clathrina clathrus*.<sup>131</sup> Contignasterol (**127**), a highly oxygenated steroid with the "unnatural"  $14\beta$ -proton configuration and cyclic hemiacetal functionality in its side chain, has been isolated from the sponge *Petrosia contignata*,<sup>132</sup> collected in Papua New Guinea. The authors acknowledge that the  $14\beta$ -epimers of a number of semisynthetic 15-ketosteroids were found to be more stable than the corresponding  $14\alpha$ -epimers, and they attempted to epimerize contignasterol (**127**) with base in order to determine the relative stability of its  $14\alpha$ - and  $14\beta$ -epimers. Unfortunately all attempts failed. The extraction and chromatography conditions are unlikely to have caused epimerization, and the authors have assumed that **127** exists exclusively as the  $14\beta$ -epimer in the sponge. Thus, contignasterol (**127**) represents the first naturally occurring steroid with the  $14\beta$ -configuration, although steroids with a  $14\beta$ -hydroxyl group (e.g. digitoxin) are well known from nature. A few months later Shoji *et al.*<sup>133</sup> reported the isolation of xestobergsterol A (**128**) and B (**129**), which are the second examples of steroids with the  $14\beta$ -H proton configuration, from the sponge *Xestospongia bergquistia* collected in Okinawa. The structures of xestobergsterol A (**128**) and B (**129**) are unique also in that they are pentacyclic steroids with the fifth ring linking C-16 to C-23. Extensive NOE measurements indicated a boat conformation of ring C. Both **128** and **129** strongly inhibited histamine release from rat mast cells induced by anti-IgE.

New  $\Delta^8(14)$ - $3\beta,7\alpha$ -dihydroxysterols (**130**) have been isolated from the Mediterranean *Pellina semitubulosa*.<sup>134</sup> Rare D-ring unsaturated steroid  $3\beta,16\alpha$ -diols (**131**) have been isolated from the Mediterranean sponge *Topsentia aurantiaca*.<sup>135</sup> Sterols with a D-ring unsaturation are rare, having been until now discovered only in the sponge *Homaxinella trachys*<sup>136</sup> and from cultured marine dinoflagellates.<sup>137</sup> A series of  $\Delta^5$ - $3\beta,7\beta$ - and  $\Delta^5$ - $3\beta,7\alpha$ -diols, along with the  $3\beta$ -hydroxy  $\Delta^5$ -7-ones, well-recognized autoxidation products of  $\Delta^5$ -sterols, have been isolated from *Haliclona oculata*<sup>138</sup> from the bay of Funds, Canada, and *Stelodoryx chlorophylla*<sup>139</sup> from New Caledonia. They were accompanied by side chain oxygenated sterols, such as 24-oxo-,  $\Delta^{22}$ -24-oxo-,  $\Delta^{24(28)}$ -25-hydroxysteroids,<sup>138,139</sup> and short side chain ketones, such as the pregnane derived 20-ketones and the 26,27-bisnor  $\Delta^{22}$ -24-ketones,<sup>139</sup> also found in a sponge of genus *Hirtio*<sup>140</sup> and in *Damiriana hawaiiiana*,<sup>17</sup> *Haliclona rubens*,<sup>108</sup> and *Psammaphysilla purpurea*.<sup>141</sup> Also these side chain oxygenated sterols could be of abiotic origin, deriving through oxidation.


In the echinoderms, sea cucumber and starfish are known to produce saponins. Besides these echinoderms, only a limited number of marine organisms have been shown to contain steroidal or triterpenoid glycosides. Recently, glycosides with cholestane- and

105, R,R<sup>1</sup> =  $\text{C}=\text{O}$ 106, R=H R<sup>1</sup>=OH107, R=OH R<sup>1</sup>=H108,  $\Delta^2$ 109,  $\Delta^3$ 


110




111




112




113




114




115




116




117




118




119



120






121



122, R =

123, R =





**Figure 5.** Polyoxygenated steroids from other sponge species.

lanostane-type aglycons have also been isolated from some sponges (Figure 6). Several norlanostane triterpenoid glycosides, sarasinosides A<sub>1</sub>, B<sub>1</sub>, C<sub>1</sub> (132) have been isolated from *Asteropus sarasinosum*.<sup>142,143</sup> They are penta- (A<sub>1</sub>, B<sub>1</sub>) and tetraglycosides (C<sub>1</sub>) containing two amino sugars (NAcGlc and NAcGal) each and have been reported to be ichthyotoxic and to inhibit cell division of fertilized starfish eggs.<sup>142</sup> A 4-methyl steroidal diglycoside 133 has been isolated from *Erylus lendenfeldi*,<sup>144</sup> collected in the Red Sea. More recently two novel glycosides eryloside C (134) and D (135) have been isolated from a sponge of the genus *Erylus*<sup>145</sup> collected at depth of 500 m in south of New Caledonia. The novel lanostane derived aglycon of 134 and 135 features a rare 14-carboxyl group and a 24-methylene-25-methyl side chain. It appears closely related to penasterol, a lanostane-derived metabolite with the 14-carboxy group and the cholestane side chain, isolated from the Okinawan sponge *Penares* sp.<sup>146</sup> Both penasterol and erylosides C and D may have biosynthetic implications concerning the synthesis of sterols in the sponges, in which the loss of the 14-methyl group could involve the intermediacy of a carboxylic acid rather than an aldehyde. A novel steroidal saponin, pachastrelloside A (136), has been isolated from the sponge *Pachastrella* sp.<sup>147</sup> This glycoside inhibits cell division of fertilized starfish eggs.

#### IV. Polyoxygenated Steroids of Coelenterata

The phylum Coelenterata is subdivided into the classes Hydrozoa, Cubozoa, Scyphozoa, and Anthozoa. The latter class is subdivided into Hexacorallia and Octocorallia.

Polyhydroxysteroids have been found in Hydrozoa and Anthozoa.

##### A. Hydrozoa


Marine hydroids are the simplest Cnidarians, generally characterized by an alternation of generation; they occur in most habitats from the shore to the deep sea. The sessile animals (polyps) grow as bushlike colonies on stones, shells, or seaweeds. Only a few reports on their chemical constituents, including sterols, have appeared, due largely to the difficulties of collection and identifying the biological material. Cholesterol is usually the predominant sterol in hydroids<sup>2,148</sup> and polyhydroxysteroids have been discovered in species of the genus *Eudendrium* sp. (Figure 7). Cimino *et al.*<sup>149</sup> first found 3-oxocholest-4-ene-4,16 $\beta$ ,18,22(*R*)-tetrol 16,18-diacetate (137) in Mediterranean hydroids of the genus *Eudendrium* (sp. *rameum*, *racemosus*, and *ramosum*) as well in their predators, several species of nudibranchs (*e.g.* *Hervia peregrina*, *Flabellina affinis*, *Coryphella lineata*). Other C-18 oxygenated sterols 138–141, containing the rare 2 $\alpha$ ,3 $\alpha$ -diol functionality, have been isolated from *Eudendrium glomeratum*,<sup>150–152</sup> collected in the Bay of Napoli. The 2 $\alpha$ -deoxy analog 142 has been isolated from the same organism.<sup>152</sup>

##### B. Anthozoa

###### 1. Hexacorallia

The subclass Hexacorallia comprises organisms such as sea anemones (order Actinaria) and zoanthids (order Zoanthidea), which are characterized by the production

sugar chains:



**Figure 6.** Polyoxygenated steroid glycosides of sponges.

of very powerful toxins.<sup>153</sup> Sea anemone toxins are polypeptides or proteins whose amino acid sequences exhibit a high degree of homology and have become very useful tools for studying the voltage-dependent  $\text{Na}^+$  channel in nerve, cardiac, and muscle cells. Toxins isolated from several *Palythoa* spp. were the most toxic marine natural products before the discovery of maitotoxin,<sup>154</sup> one of the causative agent of ciguatera, a poisoning caused by ingestion of coral reef fishes. Palytoxin is uniquely distinct from other molecules with which organic chemists have previously dealt in terms of molecular size and of structural complexity; it has molecular weight of about 3000 Da but totally lacks repeating units, such as amino acid, sugar, and fatty acid residues, commonly found in molecules of that size.<sup>155</sup>

Publications on sterols from Hexacorallia are quite limited and report on the occurrence of conventional  $\Delta^5$ -sterols.<sup>2</sup> Recent papers from Pietra and colleagues<sup>156-158</sup> account for the discovery of the zoanthid

*Gerardia savaglia* as an unexpected marine source of moulting hormones (Figure 8). The authors first reported the isolation of ecysterone (143),<sup>156</sup> and then of ecdysone (144) and ajugasterone C (145),<sup>157</sup> the latter an ecdysteroid previously thought to be an exclusive and rare plant product. All three compounds were obtained in large amounts. The fourth ecdysteroid isolated as minor component from *Gerardia savaglia* is the new gerardiasterone 146.<sup>158</sup> Ecdysteroids have been identified, in trace amounts, in arthropods,<sup>159</sup> nematodes and annelids,<sup>160</sup> mollusks,<sup>161</sup> and, in relative large amounts, in terrestrial plants.<sup>159</sup> The occurrence of ecdysteroids in such large amounts in the zoanthid *G. savaglia* poses a number of intriguing questions about their origin and role. The authors have noted that *G. savaglia*, after it had been kept for 15 months in an aquarium, still gave ecdysteroids in roughly the same large amounts as immediately after its collection,<sup>157,158</sup> and they suggested that ecdysteroids can be, at least in part, synthesized within the zoanthid from, possibly,




Figure 7. Polyhydroxysteroids from hydroids of genus *Eudendrium*.

dietary cholesterol. The role of ecdysteroids as hormones in the zoanthid is ruled out because of the high concentration of the steroid, and a possible defensive role has been suggested. Finally, the authors conclude that it is likely that ecdysteroids will prove to be much more widely distributed in the marine environment than was thought.<sup>156,157</sup> The finding that pinnasterol (60), an ecdysteroid-like sterol, is a constituent of the red alga *Laurencia pinnata*<sup>67</sup> might support the hypothesis.

## 2. Octocorallia

The subclass of Octocorallia comprises different orders among which Gorgonaceae and Alcyonacea are distinguished by their high content of polyhydroxylated sterols, which often are the major sterols. The structures of the polyoxygenated sterols from Gorgonaceae are shown in Figure 9.

5 $\alpha$ -Cholestane-3 $\beta$ ,5,6 $\beta$ ,9-tetrol (147) one of first polyhydroxysteroids isolated from marine species was obtained from *Pseudopterogorgia elisabethae*.<sup>162</sup> After the isolation of 147 it was suggested<sup>56</sup> that a more detailed examination of the polar constituents of gorgonians will result in the isolation of others oxygenated sterols.

The first unique 9,11-secocholestane system was 3 $\beta$ ,11-dihydroxy-9,11-secogorgost-5-en-9-one (148), which




















Figure 8. Ecdysteroids from the zoanthid *Gerardia savaglia* (Anthozoa, Hexacorallia).

was isolated from the gorgonian *Pseudopterogorgia americana* together with the known gorgosterol 45.<sup>122</sup> The structure and the absolute configuration was established by X-ray analysis. Hippurin-1 (149), with a unique oxygenation pattern and a spiroketal ring structure on the side chain of a 24-methylcholestane skeleton, was isolated from *Isis hippuris*,<sup>163</sup> a common gorgonian found on the Great Barrier Reef. A single-crystal X-ray analysis of the monoacetate secured the structure and relative stereochemistry of hippurin-1. ORD studies of the 11-keto and 3,11-diketo derivatives established the absolute configuration shown in 149, which is the normal configuration found in sterols. Two related sterols of further complex structural features in the same gorgonian were isolated, and their structures





**Figure 9.** Polyoxygenated steroids of Gorgonaceae (Anthozoa, Octocorallia).

were determined as  $3\alpha$ -acetoxy- $11\beta$ -hydroxy-24-methyl-22,25-epoxy- $5\alpha$ -furostan-18,20 $\beta$ -lactone (150) and  $3\alpha$ -acetoxy- $11\beta$ ,18:18,20:22,25-triepoxy- $5\alpha$ -furostan (151).<sup>164</sup> The eptacyclic steroid 150 was converted to the octacyclic one 151 on reduction with  $\text{LiAlH}_4$  at reflux followed by acetylation. A minor compound, presumably the 18-epimer was also formed. The 18 $R$ -anomer, which requires less strained *cis* junction of the two five-membered rings composed of the bicyclic acetal, is the most stable epimer, as confirmed by treatment of 151 with *p*-toluenesulfonic acid in THF at room temperature for 30 h, which showed no signs of epimerization.<sup>164</sup> The authors remark that naturally occurring 18-substituted steroids are uncommon. They were only found in the aglycons of holothurins,<sup>165</sup> glycosides of sea cucumbers with an 18,20 lactone functionality on the lanostane skeleton, and in the highly active hormone aldosterone and in its metabolic products.<sup>166</sup> More recently C-18 oxygenated sterols have been isolated from hydroids (Figure 7), sponges of the genus *Dysidea* (Figure 4), gorgonians (Figure 9) and soft corals of the genus *Sinularia* (Figure 10). Several related desacetoxy hippurins (*e.g.* 152) have also been described;<sup>167</sup> five more hippurins closely related to 149 were found in specimens of *Isis hippuris* collected in the Andaman Islands, India.<sup>168</sup> The same species also yielded a tetrahydroxylated steroid, which was identified as gorgost-5-ene- $3\beta$ ,7 $\alpha$ ,11 $\alpha$ -12 $\beta$ -tetrol 12-monoacetate (153).<sup>169</sup>

A pregnane derivative 154 has been isolated from the Mediterranean gorgonian *Eunicella cavolini*.<sup>170</sup> Four new metabolites, muricin-1 through muricin-4 (155–158) which are unique aminogalactose saponins containing pregnane-derived aglycons have been isolated from the Pacific gorgonian *Muricea fruticosa*.<sup>171</sup> They were supposed to be responsible in reducing fouling on *M. fruticosa*, possibly through effective growth inhibition of the pennate diatom *Phaeodactylum tricorutum*. The ecologies and life histories of *M. fruticosa* in comparison with those of *M. californica*, the two major *Muricea* in California, give support to the potential role of these compounds in contributing to the reduced fouling of *M. fruticosa*. *M. californica*, which is found growing together with the morphologically very similar *M. fruticosa*, was consistently overgrown with typical encrusting plants and animals such as red algae, hydroids, ectoprocts, zoanthids, ascidians, mollusks, and bryozoans. In contrast *M. fruticosa* was found virtually free of surface fouling organisms. While each contained mixtures of triglycerides, sterols, and fatty acids, only the extracts of *M. fruticosa* were found to contain muricin-1 to muricin-4.<sup>171</sup> Another pregnane

glycoside,  $3\beta$ -pregna-5,20-dienyl  $\beta$ -D-galactopyranoside (159) has been isolated from the gorgonian *Pseudoplexaura wagnaari*.<sup>172</sup> The pregna-1,4,20-trien-3-one was previously isolated from an alcyonacean coral *Gersemia rubiformis*.<sup>173</sup>


A reactive sterol, 3,16-dioxocholesta-4,14-diene-15,20-diol (160) has been isolated from the Mediterranean gorgonian *Leptogorgia sarmentosa*.<sup>174</sup> On treatment with acetic anhydride and pyridine 160 suffers the loss of the entire side chain through a retro-aldol cleavage. Three polyoxygenated steroids, one being guggulsterol III (161) so far thought to be a plant product, the remaining 162 and 163 being closely related to 161, have been discovered in the same species, *L. sarmentosa*.<sup>175</sup> The unusual upfield resonance of C-17 in guggulsterol III (161) ( $\delta_C$  60.1 ppm) when compared with its acetate derivative weakly downshifted at  $\delta_C$  60.4 ppm and of its 16-epimer 163 at  $\delta_C$  68.1 ppm is explained as due to the hydrogen linkage between the hydroxyl groups at C-20 and C-16 in 161. A series of five C-18 hydroxylated homologues (164) of guggulsterol III (161) have been isolated from the same organism, *L. sarmentosa*.<sup>176</sup>

A novel secosterol, astrogorgiadiol (165) has been isolated from a gorgonian *Astrogorgia* sp.,<sup>177</sup> as an inhibitor of cell division in fertilized starfish eggs. The authors suggested that 165 is likely biosynthesized from cholesterol via dienol-phenol rearrangement and cleavage of the 9,10-bond. Two more secosterols, closely related to astrogorgiadiol (165) have been reported from the gorgonian *Caligorgia* sp.<sup>178</sup>

Two novel glycosides named dimorphoside A (166) and B (167) have been isolated from the gorgonian *Anthoplexaura dimorpha*,<sup>179</sup> widely distributed along the southern coast of Japan, as the major active constituents (inhibition of the development of fertilized sea urchin eggs) of that organism. The oxidation of the C-19 methyl group to carboxylate is a rare feature, although related steroid sulfates have been isolated from a sponge.<sup>94</sup>

*Sinularia* sp. (Alcyonacea) has given three new 9,11-secosterols with the same 9,11-seconucleus as 148 and conventional 24-methylene, 24-methyl, and cholesterol side chains (168–170).<sup>123,124</sup> The same species also yielded a very minor component, which proved to be the C-8 epimer of  $3\beta$ ,11-dihydroxy-24-methylene-9,11-secocholest-5-en-9-one (171).<sup>124</sup> According to the authors it is unlikely that the less stable  $8\alpha$ -H epimer is an artifact of the isolation procedure. The fact that a group of 9,11-secosterols with a variety of side chains has been discovered suggest that *Sinularia* species probably cleaves ring C of exogenous sterols similar





**Figure 10.** Polyoxygenated steroids from Alcyonacea (Anthozoa, Octocorallia) of genus *Sinularia*.

to the situation observed in some sponges, e.g. 5,6-secosterols in *Hippospongia communis*,<sup>117</sup> incisterols from *Dictyonella incisa*<sup>120</sup> and the ring A contraction of *Achantella acuta* sterols.<sup>180</sup>

In addition to the 9,11-secosterols, the soft corals of the genus *Sinularia* have been shown to produce a series of new polyhydroxylated sterols (Figure 10). The common 3 $\beta$ ,5 $\alpha$ ,6 $\beta$ -trihydroxy sterols were isolated from

*Sinularia dissecta*.<sup>181</sup> A novel group of eight polyhydroxylated sterols, all of them possessing an  $11\alpha$ -hydroxy substituent of potential utility as corticosteroid intermediates have been then isolated from the same specie.<sup>182</sup> Seven of these compounds possessed identical  $1\alpha,3\beta,11\alpha$ -trihydroxycholest-5-ene nuclei (172), but differed in the side chains, which, in addition to the conventional  $C_8$  and  $C_9$  side chains, include those of gorgosterol (45) and 23-demethylgorgosterol (44).<sup>183</sup> In continuation with the examination of the extracts of *S. dissecta* the same authors isolated a major group of five sterols with the same  $1\alpha,3\beta,11\alpha$ -hydroxylation pattern as well as an acetylated group located at C-18, 173.<sup>183</sup> In addition the same organism also yielded the analogous 13-formyl, 174, and 13-carboxyl, 175, derivatives.<sup>183</sup> The corresponding  $5\alpha,6$ -dihydro derivatives of the major 24-methylene side chain sterols have also been found 176–179. The isolation of three groups of C-18 functionalized sterols from the same organism is so far unprecedented. The authors remark that the fact that similar nuclei possessing diverse side chains are encountered suggests the existence of enzyme systems—either in the coral or in some symbiont—that introduce the  $1\alpha$ - and  $11\alpha$ -hydroxyl groups into a dietary precursor and that may also functionalize the C-18 angular methyl group.<sup>183</sup> Eventual isolation of the enzyme systems may provide a possible practical route to otherwise rare sterols—notably those with oxygenated C-11 and C-18 positions.



Two new cholestadienone derivatives, 180 and 181, which possess epimeric cyclic ketals of the C-22 side chain position involving the oxidation of the C-18 methyl group, have been isolated from an apparently undescribed species of *Sinularia* genus.<sup>184</sup> Accurate  $^1\text{H}$  NMR studies indicated the conformations of the six-membered ketal rings to be a twist boat in 180 and a chair in 181. The variation in the ring conformation was then related to the stereochemistry of the ketal carbon C-22. Similar sterols were described from the gorgonian *Isis hippuris*.<sup>163,164</sup>

Two novel polyhydroxysterols, numersterols A and B, have been isolated from the South China Sea *Sinularia numerosa*, and their structures were determined as 24-methylenecholestane- $1\alpha,3\beta,5\alpha,6\beta$ -tetrol (182) and 25-methylene-22-homocholestane- $1\beta,3\beta,5\alpha$ -triol (183).<sup>185</sup> The structure of numersterol A (182) with nuclear hydroxylation pattern differing from that encountered in many polyoxygenated sterols from soft coral (e.g. *Sarcophyton glaucum*,<sup>186–188</sup> *Lobophytum pauciflorum*,<sup>189</sup> and *Sclerophytum* sp.<sup>190</sup>) only in the stereochemistry at C-1, was confirmed by X-ray analysis.<sup>185</sup> The side chain of numersterol B (183) is apparently the result of two biomethylations at C-27 of normal cholesterol side chain. The corresponding 6-keto derivative of numersterol A (182) has been recently isolated from *Sinularia microclavata*,<sup>191</sup> one of the most typical species found in Indo-Pacific coral reefs, collected at Ishigaki Islands, Okinawa. *Sinularia crispa* from the East coast of Sri Lanka has been shown to contain a novel steroid glycoside (184), showing spermatostatic activity.<sup>192</sup>



Figure 11 shows the structures of polyoxygenated sterols found in Alcyonaceans other than those found in the genus *Sinularia*.

A common feature of many polyhydroxysterols from soft corals is the  $3\beta,5\alpha,6\beta$ -trihydroxy moiety. The first of the soft coral sterols found to have this pattern was 185 isolated from *Sarcophyton elegans*,<sup>193</sup> then found as 25-hydroxy analog 186 also in *Sclerophytum* sp.<sup>190</sup> *Sclerophytum* sp. have been shown to contain also the rare (24*S*)-24-methylcholestane- $3\beta,5\beta,6\alpha,25$ -tetrol (187),<sup>190</sup> isomeric with the common 186 and differing from that in the stereochemistry at C-5 and C-6. On pyridinium chlorochromate (PCC) oxidation followed by dehydration with thionyl chloride in pyridine, both 186 and 187 afforded the same  $\Delta^4$ -3,6-diketo derivative.<sup>190</sup> The polyhydroxysteroid 188 was isolated from the same soft coral *Sarcophyton elegans*.<sup>194</sup> After that a series of  $1\beta,3\beta,5\alpha,6\beta$ -tetrahydroxysteroids (189–194) have been isolated from *Sarcophyton glaucum*,<sup>186–188</sup> one of the most common species found in Indo-Pacific coral reefs, collected at Ishigaki Islands, Okinawa, and *Lobophytum pauciflorum*,<sup>189</sup> from the same Ishigaki Islands, and from a species of the genus *Sclerophytum*,<sup>190</sup> collected in the Andaman and Nicobar Seas of the Indian Ocean. Except for some of the  $5\beta$ -sapogenols and for the  $\Delta^5$ -ruscogenin,<sup>195</sup>  $1\beta$ -hydroxysteroids are rare in nature. It is also to be noted that 194 is the first report on the isolation of a polyoxygenated androstane derivative from marine invertebrates.<sup>187</sup> Among the complex mixture of polyhydroxysteroids of *S. glaucum* Kobayashi and Mitsuhashi<sup>196</sup> also isolated a minor polyhydroxysteroid and determined the structure of (24*S*)-24-methyl-5 $\alpha$ -cholestane- $3\beta,5,6\beta,25\epsilon,26$ -pentol (195). Hydroxylation at C-26 is common among polyhydroxysteroids from starfish<sup>197</sup> and the 24-methyl 25-, 26-diol side chain of 195 have been found in a steroidal nonaol from the starfish *Archaster typicus*,<sup>198</sup> but never found before in Anthozoa. More recently new additions to the group of  $1\beta,3\beta,5\alpha,6\beta$ -tetrahydroxysterols have been found in *Sarcophyton subviride*,<sup>199</sup> soft coral collected from the Katchal Islands of Andaman and Nicobar Coasts. These include the  $\Delta^{25(26)}$ -analog of 192, the C-18 hydroxy analog of 189 (196), the 25-*O*-acetyl derivative of 195, and 197 with the gorgostane skeleton. A new  $3\beta,5\alpha,6\beta$ -trihydroxy steroid with further hydroxylation at C-22 and C-24 (198) have been isolated from the soft coral *Asterospicularia randalli*<sup>200</sup> collected at Guam Island. Hydroxylation at C-22 was also found in lobophytosterol (199), depresosterol (200), and three other closely related sterols,  $5\beta,6\beta$ -epoxysterols 201–203 from the Red Sea soft coral *Lobophytum depressum*.<sup>201</sup> The occurrence of C-28 oxygenated sterols, which may be intermediate in the demethylation pathway, might be of biogenetic interest.


The soft coral *Minabea* sp. contains sterol lactones of the withanolide class, three are new  $C_{28}$  compounds, minabeolides 1–3 (204–206), and five are  $C_{27}$  derivatives, minabeolides 4–8 (207–211).<sup>202</sup> Instead of the  $\Delta^2$ -1-keto ring A, very common in plant withanolides, the marine-derived withanolides have a  $\Delta^4$ -3-keto ring A. Further sterols with the common  $3\beta,5\alpha,6\beta$ -trihydroxy moiety feature and a fourth hydroxyl group found in the  $7\beta$ -position have been isolated in the soft coral *Anthelia glauca*<sup>203</sup> (212) collected at Laing Island, Papua, New Guinea, and also as  $7\beta$ -acetate analogs 213–215 in a species of the genus *Xenia* collected at Zamami-jima, Okinawa.<sup>204</sup> Gorgosta- $3\beta,5\alpha,6\beta$ -triol has been also isolated from the same species.<sup>204</sup> The mixture con-

185, R=H R<sup>1</sup>=OAc186, R=OH R<sup>1</sup>=OH188, R=OH R<sup>1</sup>=OAc


187

189, R=Me R<sup>1</sup>=OAc190, R=Me R<sup>1</sup>=OH191, R=Me R<sup>1</sup>=H192, R=CH<sub>2</sub> R<sup>1</sup>=H


194




195




196




197




198




199




200




201



202




203




204, R=H

205, R=OAc

206, R=H 1,2-dihydro

207, R=H R<sup>1</sup>=H208, R=OAc R<sup>1</sup>=H209, R=H R<sup>1</sup>=H 1,2-dihydro210, R=OAc R<sup>1</sup>=H 1,2-dihydro211, R=H R<sup>1</sup>=OAc 1,2-dihydro



**Figure 11.** Polyhydroxysteroids from Alcyonacea (Anthozoa, Octocorallia) other than genus *Sinularia*.

taining the above four sterols exhibited a growth-inhibitory activity against B-16 melanoma cells.<sup>204</sup>

*Litophyton viridis* is the source of the rare 19-oxygenated sterols 216-218.<sup>98,205</sup> Compound 216 with

the 7 $\beta$ -acetoxy functionality appeared to be the first report of a naturally occurring 19-hydroxy steroid.<sup>98</sup> The 5 $\beta$ ,6 $\beta$ -epoxide 218 showed antileukemic activity (IC<sub>50</sub> 0.5  $\mu$ g/mL) against P<sub>388</sub> leukemia cells *in vitro*.<sup>205</sup>

The 19-hydroxylated sterols are very rare in nature only found in a sponge<sup>97</sup> and in the soft coral *L. viridis*<sup>98,205</sup> and are of interest as a possible intermediate in the biosynthesis of 19-norsterols. Examination of less polar fractions of the extract of *L. viridis* led Bortolotto *et al.*<sup>206</sup> to isolate 4 $\alpha$ -methyl-3 $\beta$ ,8 $\beta$ -dihydroxy-5 $\alpha$ -ergost-24(28)-en-23-one (219). The structure was secured by X-ray of the *p*-bromobenzoate derivative. At that time the 8 $\beta$ -hydroxy function appeared unique in marine sterols; several polyhydroxysteroids have been later isolated from starfish and the majority of them possess the 8 $\beta$ -hydroxy substituent.<sup>197</sup>

The 3,4,5,6-oxygenated pattern discovered in lobosterol (220), isolated first from *Lobophytum pauciflorum*<sup>207</sup> and later in species of the genus *Sclerophytum*,<sup>208,209</sup> is unprecedented in natural sterols. This was also the first marine sterol reported to have an A/B *cis* ring fusion. The available amount of lobosterol was insufficient for a complete chemical elucidation of the structure, and this determination was obtained by X-ray diffraction analysis of the 4-*O*-*p*-bromobenzoate.<sup>207</sup> In addition to lobosterol (220) another *Sclerophytum* sp. of soft coral collected off the Andaman and Nicobar coasts has been shown to contain two new polyhydroxysterols, andamansterol (221) and nicobarsterol (222).<sup>209</sup> Both compounds 221 and 222 bear an oxygenated C-21, which is rare in marine sterols except for those found in echinoderm ophiuroids and one starfish. The structure of andamansterol (221) was secured by X-ray analysis, thus confirming the conventional C-20 configuration. Nicobarsterol (222) is a new type of 9,11-secoesterol with the C-11 oxidized at a carboxyaldehyde level and forms a seven-membered hemiacetal ring with the C-21 hydroxy group. A sterol with the 9,11-glycolated structure like 221 could likely be a precursor of 222. Lead tetraacetate treatment of andamansterol (221) gave a 9,11-seco derivative having the same seven-membered hemiacetal ring as 222, including the 11*R* configuration.<sup>209</sup> Examination of another *Sclerophytum* sp. resulted in the isolation of a new steroid 223, which is the first example to have C-21 oxidized to carboxylate.<sup>210</sup>

In a continuing investigation of the soft corals of the Andaman and Nicobar coasts of the Indian Ocean by Kobayashi and colleagues, three new polyhydroxysterol glycosides were isolated from one sample identified as *Alcyonium* sp.,<sup>211</sup> and their structures were determined as 24-methylenecholest-5-ene-3 $\beta$ ,16 $\beta$ -diol 3-*O*- $\alpha$ -L-fucoside (224) and its 7 $\beta$ - (225) and 7 $\alpha$ -hydroxy (226) derivatives. From the chemotaxonomic viewpoint it may be of interest that this is the second example of *Alcyonium* sp. containing steroidal glycosides, which are rare in soft corals. Pregnene-type steroidal glycosides were previously isolated from an Okinawan coral-reef *Alcyonium* sp. by Kobayashi *et al.*<sup>212</sup> Two of these glycosides, pregnediosides A and B, were characterized as 4-*O*- $\beta$ -D-arabinopyranosyl (227) and 4-*O*- $\beta$ -D-xylopyranosyl (228) of 3 $\beta$ ,4 $\alpha$ -dihydroxy-5 $\alpha$ -pregn-20-ene, respectively, the remaining three being their monoacetates.<sup>212</sup> The role of the *Alcyonium* sp. steroidal glycosides is still obscure.

Kingston and Fallis<sup>213</sup> have reported the isolation of two unusual C<sub>26</sub> sterols, 12 $\beta$ -hydroxy-24-norcholesta-1,4,22-trien-3-one (229) and its acetate 230 from the sea raspberry *Gersemia rubiformis*, an alcyonacean

coral from the cold waters off Newfoundland and Labrador. A  $\Delta^{1,4}$ -3-ketopregnane 231 was also found in the same organism.<sup>214</sup>


Very few polyhydroxysteroids have been found in species of the class Anthozoa other than those found in the order Gorgonaceae and Alcyonacea and are shown in Figure 12. Four new cytotoxic steroids, named stoloniferones were isolated from the stoloniferan Okinawan soft coral *Clavularia viridis*<sup>215</sup> (order Stolonifera). They possess the same 11 $\alpha$ -hydroxy-5 $\beta$ ,6 $\beta$ -epoxy 2-en-1-one steroidal skeleton, but differ in their side-chain structures. A X-ray crystallographic analysis of stoloniferone D (235) confirmed the structures and the relative stereochemistries derived from spectroscopic studies, and CD measurements of the 2,3-dihydrostoloniferone C (234) secured the absolute stereostructures. Stoloniferones exhibited growth inhibition against P<sub>388</sub> leukemia cells. Unusual 20-epicholanolic acid derivatives 236–239 were isolated from the sea pen *Ptilosarcus guerneyi*<sup>216</sup> (order Pennatulaceae). Characteristic properties of these compounds are (a) the shorter retention time than those of their natural C-20 isomer (20 $\alpha$ -H) on GLC analysis and (b) the C-21 methyl protons resonance at 0.1 ppm higher field in the NMR spectra than the 20 $\alpha$ -H compounds. The isolated steroids, as methyl esters, were identical with the synthetic 20-epicholanolic methyl esters.<sup>217</sup> The authors analyzed also the sea pen's free sterols and found them to possess the "normal" C-20 stereochemistry.<sup>217</sup> Thus the free sterols are not the immediate biosynthetic precursor of the 20-episteroids, and their origin is still an open question. Cholestane-3 $\beta$ ,5 $\alpha$ ,6 $\beta$ -triol has been isolated from the sea pen *Pteroides esperi*<sup>218</sup> (order Pennatulaceae) and found to have steatotic and cytotoxic activity. A series of highly oxygenated sterols 240–246 have been isolated from *Anthipathes subpinnata* (order Antipatharia) commonly named black coral.<sup>219,220</sup> The  $\Delta^5$ -3 $\beta$ ,7 $\beta$ ,19-hydroxylation pattern of 246 has been previously found in 216 from the soft coral *Litophyton viridis*.<sup>98</sup> Interesting *A. subpinnata* produces both 18- and 19-hydroxylated sterols; the hydroxylation at C-20 is rare in marine sterols except for the aglycons of the asterosaponins found in starfishes.<sup>197</sup>

## V. Polyoxygenated Steroids of Bryozoa

Bryozoa are minute filter feeders animals which contain biological active alkaloids and macrolides. Despite the considerable interest in anticancer activities of bryostatins from *Bugula neritina*,<sup>221</sup> the chemical studies of bryozoans are proceeding slowly, probably because of the difficulties experienced in collecting sufficient material for analysis. Only two papers on sterols of bryozoans have appeared in the literature, one of which describes the isolation of five polyhydroxysterols, all possessing a common  $\Delta^7$ -3 $\beta$ ,5 $\alpha$ ,6 $\beta$ -trihydroxy steroidal skeleton and saturated and  $\Delta^2$ -unsaturated conventional C<sub>8</sub>, C<sub>9</sub>, and C<sub>10</sub> side chains from *Myriapora truncata*.<sup>112</sup> These sterols have also been found in a sponge<sup>110</sup> and in a mollusk.<sup>113</sup>

## VI. Polyoxygenated Steroids of Marine Mollusca


More species of Mollusca have been analyzed for their sterol content and composition than have those of any other phylum, in part because of their nutritional



**Figure 12.** Polyhydroxysteroids from Octocorallia (Anthozoa) other than Gorgonacea and Alcyonacea.

value.<sup>2,60,62</sup> Even so, the occurrence of polyoxygenated sterols in mollusks (Figure 13) is relatively rare. The highly oxygenated steroid 137, isolated from three species of the hydroids of the genus *Eudendrium*, was also found in their predators, the nudibranchs (class Gastropoda) *Hervia peregrina*, *Flabellina affinis*, and *Coryphella lineata*.<sup>149</sup> Recent studies have shown that most dorid nudibranchs utilize organic metabolites obtained from their diets as chemical defences against predation.<sup>222</sup> The steroid 137 seems to be one of the nudibranch metabolites to which no defensive role is ascribed. Two cholanic acid derivatives 247 and 248, with significant antifeedant activity, were isolated from the dorid nudibranch *Aldisa sanguinea cooperi*,<sup>223</sup> but were not found in the sponge *Anthoarcuata gracieae* on which it feeds. This nudibranch is apparently obtaining inactive metabolites from its diet and chemically modifying them to produce an active antifeedant.<sup>223</sup> Two new sterols, dialusterols A (249) and B (250) were isolated from the skin extracts of the dorid nudibranch *Diaulula sandiegensis* and were identified

by interpretation of spectral data.<sup>224</sup> Dialusterols are related to pinnasterol (60) and acetylpinnasterol (61), which were isolated from the red alga *Laurencia pinnata*.<sup>67</sup> All four sterols share structural features with ecdysones. The  $2\alpha,3\alpha$ -diol array of dialusterols are not commonly encountered in naturally occurring sterols. They have only been isolated from the hydroid *Eudendrium glomeratum*.<sup>150-152</sup> A new cytotoxic epoxysterol 251, active against the L<sub>1210</sub> cell line, was isolated from the marine gastropod *Planaxis sulcatus*.<sup>225</sup> This is very similar to the  $9,11\alpha$ -epoxysterol 91 isolated from a sponge of *Dysidea* sp. and therefore 251 is suspected to be the precursor of 91.<sup>225</sup>  $\Delta^7$ - $3\beta,5\alpha,6\beta$ -Trihydroxysteroids have also been found in the scallop *Patinopecten yessoensis* (class Bivalvia), including one, 252, with the rare  $9\alpha$ -hydroxylation.<sup>113</sup> The occurrence of  $\Delta^5$ - $3\beta$ -hydroxy-7-ketosterols 253 in the prosobranch mollusk *Patingera magellanica* has been reported.<sup>228</sup> The possible abiotic origin of  $\Delta^5$ -7-ketones, well-recognized autooxidation products of  $\Delta^5$ -sterols, can be suspected. A series of highly degraded sterols,



**Figure 13.** Polyhydroxysteroids of marine Mollusca.

named aplykurodins, has been isolated from two ophistobranchs of the genus *Aplysia* (class Gastropoda). The 4 $\beta$ -hydroxy lactones **254** and **255** have been isolated from *A. kurodai*;<sup>227</sup> the 4 $\beta$ -acetoxy derivative **256** and the 4-keto analog **257** have been then isolated from the Mediterranean *A. fasciata*.<sup>228</sup> The relative stereostructure of aplykurodine A (**255**) was deduced by X-ray crystallography, whereas the absolute configuration was determined by application of CD data.<sup>227</sup> The interconversion of the 1-9  $\delta$ -lactones, aplykurodins, and their isomeric 1-4  $\gamma$ -lactones has been investigated<sup>227,228</sup> and the authors concluded that the  $\delta$ -lactones are the kinetic products, whereas the  $\gamma$ -lactones are the thermodynamic ones.<sup>228</sup> Aplykurodin B (**256**) and aplykurodinone (**257**) which were isolated from the external parts of the body of the animals and showed ichthyotoxicity and antifeedant activity (feeding deterrence) were considered by the authors to be the defense allomones of *A. fasciata* against predators.<sup>228</sup>

The endocrine system of steroid hormones in mollusks has been established, and many reports on the biosynthesis and metabolism of progesterone, estrogens, and androgens have been published.<sup>229</sup> *Mytilus edulis* (class Bivalvia), the mussel, had been reported to contain 0.2  $\mu\text{g g}^{-1}$  by weight of a substance, which showed ecdysone activity.<sup>230</sup> The probable role of ecdysteroids in relation to calcification of the shell in mollusk has been suggested.<sup>161,231</sup>

### VII. Polyoxygenated Steroids of Echinodermata

The phylum Echinodermata, which comprises about 6000 living species, is divided in five classes: Crinoidea

(sea lilies and feather stars), Holothuroidea (sea cucumbers or holothurians), Echinoidea (sea urchins), Asteroidea (sea stars or starfish), and Ophiuroidea (brittle stars).

Among the echinoderms, starfish and sea cucumbers invariably contain saponins, which are responsible of their general toxicity. Chemically, saponins derived from sea cucumbers are triterpenoid glycosides whereas those from starfish are steroidal glycosides. The presence of oligoglycosides in both Holothuroidea and Asteroidea classes gives support to the opinion that sea cucumbers and starfish are phylogenetically closely related. In starfish and sea cucumbers,  $\Delta^7$ -sterols, which are probably a consequence of the presence of haemolytic saponins, are also predominant, whereas the other three classes contain the usual  $\Delta^5$ -sterols.<sup>2</sup> Haemolysis is a consequence of the abstraction of membrane cholesterol by the saponins and saponins are known to show a much lower affinity for  $\Delta^7$ -sterols, thus the presence of  $\Delta^7$ -sterols might explain the apparent immunity of starfish and sea cucumbers to their own saponins.<sup>232</sup> Ophiuroidea, which have received moderate attention by chemists as compared to the above classes, have been reported to contain a series of sulfated polyhydroxysteroids and two steroidal glycosides. In the past few years a large number of metabolites have been isolated from echinoderms, mainly steroidal glycosides and polyhydroxylated steroids from starfish and triterpene glycosides from sea cucumbers, with cytotoxic, antifungal, and antineoplastic activity. The interest in these compounds has resulted in a number of monographs entirely or in part devoted to this subject.



We should mention the works by Hashimoto (1979),<sup>233</sup> Burnell and ApSimon (1983),<sup>232</sup> Krebs (1986),<sup>153</sup> Minale, Riccio, Pizza, and Zollo (1986),<sup>234</sup> Quinn (1988),<sup>235</sup> Stonik and Elyakov (1988),<sup>165</sup> Habermehl and Krebs (1990),<sup>236</sup> and Minale, Riccio, and Zollo (1993).<sup>197,237</sup> The present paper focuses on free and sulfated polyhydroxysteroids and on their oligoglycosides occurring in starfish and ophiuroids.


### A. Asteroidea (Starfish)

Most of the work on chemical constituents of starfish has been initially prompted by the discovery of their toxic saponins and interest in their biological properties. In the recent years the structural studies of these molecules has grown up rapidly, largely exceeding the biological studies on individual compounds and almost 250 steroidal constituents, which include free and sulfated polyhydroxysteroids and steroidal glycosides, have been isolated from *ca.* 50 different starfish species, belonging to 14 families and representative of the three major orders (Paxilloidea, Valvatida, and Forcipulata) of the class Asteroidea. According to their chemical structures, the steroidal glycosides from starfish have been subdivided into three main groups: the astero-saponins, which are sulfated steroidal penta- and hexaglycosides; the cyclic glycosides, so far only found in two species of the genus *Echinaster*; and the glycosides of polyhydroxysteroids, which, although unnoticed for a long time, are as widespread as astero-saponins among starfish.<sup>238</sup> These molecules, which usually occur in minute amounts, consist of a polyhydroxylated steroidal aglycon linked to one or two sugar units and can be found in both sulfated and nonsulfated forms. Analysis of the polar extractives of the starfish *Tremaster novaecaledoniae* has recently led to the discovery of a new class of steroidal glycosides, in which the polyhydroxylated steroidal aglycon present also a phosphate conjugation to which the sugars are glycosidically attached.<sup>239</sup>

In the present paper we will discuss polyhydroxysteroids, glycosides of polyhydroxysteroids, astero-saponins, and cyclic glycosides, in that order.

#### 1. Polyhydroxysteroids

Starfish appear as the richest source of polyhydroxysteroids. In contrast with other marine phyla, in which polyhydroxysteroids have been isolated from only a limited number of species sometime belonging to the same genus, polyhydroxysteroids are widespread in starfish, where they have been found, usually as complex mixtures, in almost all species examined. More than 80 polyhydroxysteroids from starfish have been reported so far 258–343. The large majority of them possess a  $3\beta,6\alpha$ (or  $\beta$ ), $8,15\alpha$ (or  $\beta$ ), $16\beta$ -pentahydroxycholestane nucleus, sometimes with additional hydroxyl groups at one or more of positions  $4\beta,5\alpha,7\alpha$ (or  $\beta$ ) and occasionally  $14\alpha$ . A  $26$ -hydroxyl function with  $25S$ -configuration is usually present in the side chain; less commonly the side chain is hydroxylated at  $C-24$  with  $24S$ -configuration or functionalized in various ways. All hydroxyl groups are disposed on one side of the tetracyclic nucleus inducing an amphiphilic character in the molecules. Up to nine hydroxyl groups have been found in the nonols 310–312, 315, and 316, that to the best of our knowledge represent the most highly




**Figure 14.** 3,21-Dihydroxysteroids 3,21-disulfates from the starfish *Euretaster insignis*.

hydroxylated sterols isolated from a natural source. Polyhydroxysteroids occur sometimes in sulfated forms, with the sulfate group located at position  $3\beta$ ,  $6\alpha$ ,  $15\alpha$ , or  $24$ , and rare examples of phosphated derivatives have been recently isolated from the deep-water starfish *Tremaster novaecaledoniae*.<sup>240</sup>



*Euretaster insignis* is a unique example among the many starfish species examined, it is apparently devoid of both common astero-saponins and glycosides of polyhydroxysteroids and also lacks the type of polyhydroxysteroids occurring in all other species. The analysis of its polar extract yielded instead the  $3\beta,21$ -dihydroxysteroid 3,21-disulfates 258–263 (Figure 14) as a mixture resistant to further attempts of separation. After solvolysis to remove the sulfate groups, the mixture was fractionated by HPLC to afford pure desulfated 259 and two more fractions, still containing mixed dihydroxysteroids, that were resolved in the individual components by acetylation followed by argentic silica gel column chromatography.<sup>241</sup>

Although the large majority of polyhydroxysteroids from starfish (Table 1) possess a  $8\beta$ -hydroxyl group, the compounds 264–272 lack this distinctive feature. These simpler polyhydroxylated nuclei are easily identifiable by their characteristic  $^1\text{H}$  NMR spectral data (Tables 2 and 3). The  $3\beta,5,6\beta$ -trihydroxy moiety in 265 is a common element in marine polyhydroxysteroids and is encountered in a large number of polyhydroxysteroids from starfish. Besides the relatively low field shifts of the  $19$ -methyl protons, the occurrence of a  $5\alpha$ -hydroxyl group in  $3\beta$ -hydroxy- $5\alpha$ -steroidal nuclei is well evidenced by an *ca.* 0.5 ppm downfield shift experienced by the  $3\alpha$ -H resonance. Hydroxylation of  $C-26$  is a further common feature in the large majority of polyhydroxysteroids from starfish.  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectral data of epimeric  $26$ -hydroxysteroids show such small differences<sup>248</sup> that direct comparison of both stereoisomer is required for unequivocal assignment of configuration at  $C-25$ . For instance in the 500-MHz  $^1\text{H}$  NMR spectra of 265 and 269 the signals of the  $27$ -methyl protons are observed at  $\delta$  0.934 ( $25S$ -isomer) and 0.925 ( $25R$ -isomer) ppm, respectively and equally small differences are observed for carbon signals  $C-24$ ,  $C-26$ , and  $C-27$  in the  $^{13}\text{C}$  NMR spectra [ $\delta_{\text{C}}$  34.1, 68.4, and 17.3 for the  $25S$ -isomer 265 and 34.7, 68.6, and 17.1 for the  $25R$ -isomer 269].<sup>240</sup> Since it was observed that

Table 1. Polyhydroxysterols from Starfish

Basic Nuclear Hydroxylation Pattern: 3 $\beta$ ,6 $\beta$ ,15 $\alpha$ ,16 $\beta$ 




no.	side chain	R	R <sup>1</sup>	R <sup>2</sup>	sources and references
264		H	H	H	<i>Hacelia attenuata</i> <sup>242</sup>
265		OH	H	H	<i>Luidia maculata</i> , <sup>243</sup> <i>Solaster borealis</i> , <sup>244</sup> <i>Myxoderma platyacanthum</i> <sup>245</sup>
266		OH	H	SO <sub>3</sub> -Na <sup>+</sup>	<i>Myxoderma platyacanthum</i> , <sup>245</sup> <i>Rosaster</i> sp. <sup>246</sup>
267		H	OH	H	<i>Luidia maculata</i> <sup>243</sup>
268		OH	OH	H	<i>Luidia maculata</i> <sup>243</sup>
269		OH	H	H	<i>Tremaster novaecaledoniae</i> <sup>240</sup>
270		OH	H	SO <sub>3</sub> -Na <sup>+</sup>	<i>Tremaster novaecaledoniae</i> <sup>240</sup>
271		H	H	H	<i>Sphaerodiscus placenta</i> <sup>247</sup>


272, from *Tremaster novaecaledoniae*<sup>240</sup>Basic Nuclear Hydroxylation Pattern: 3 $\beta$ ,6 $\alpha$ ,8,15 $\alpha$ ,16 $\beta$ 

no.	side chain	R	R <sup>1</sup>	sources and references
273		H	H	<i>Protoreaster nodosus</i> , <sup>250</sup> <i>Poraster superbus</i> , <sup>251</sup> <i>Pentaceraster alveolatus</i> , <sup>252</sup> <i>Asterina pectinifera</i> , <sup>253,254</sup> <i>Patiria pectinifera</i> , <sup>255</sup> <i>Patiria miniata</i> , <sup>256</sup> <i>Rosaster</i> sp. <sup>246</sup>
274		OH	H	<i>Protoreaster nodosus</i> , <sup>257</sup> <i>Patiria moniata</i> , <sup>258</sup> <i>Pentaceraster alveolatus</i> , <sup>252</sup> <i>Solaster borealis</i> <sup>244</sup>
275		H	OH	as 273, <i>Pycnopodia helianthoides</i> , <sup>258</sup> <i>Solaster borealis</i> <sup>244</sup>
276		OH	OH	as 273, <i>Pycnopodia helianthoides</i> , <sup>258</sup> <i>Solaster borealis</i> <sup>244</sup>
277		OH	OH	<i>Oreaster reticulatus</i> <sup>259</sup>

6-O-sulfate

Table 1 (Continued)

no.	side chain	R	R <sup>1</sup>	sources and references
278		H	H	<i>Protoreaster nodosus</i> <sup>257</sup>
279		OH	H	<i>Protoreaster nodosus</i> <sup>257</sup>
280		H	H	<i>Poraster superbus</i> <sup>251</sup>

Basic Nuclear Hydroxylation Pattern: 3 $\beta$ ,6 $\alpha$ ,8,15 $\beta$ ,16 $\beta$ 

















no.	side chain	R	R <sup>1</sup>	R <sup>2</sup>	sources and references
281		H	H	H	<i>Halityle regularis</i> , <sup>264</sup> <i>Nardoa gomophia</i> , <sup>265</sup> <i>Pycnopodia helianthoides</i> , <sup>268</sup> <i>Dermasterias imbricata</i> , <sup>266</sup> <i>Astropecten scoparius</i> <sup>287</sup>
282		OH	H	H	<i>Halityle regularis</i> <sup>264</sup>
283	 3-O-sulfate	OH	H	H	<i>Coscinasterias tenuispina</i> <sup>268</sup>
284		H	OH	H	<i>Pycnopodia helianthoides</i> , <sup>258</sup> <i>Asterina pectinifera</i> , <sup>254</sup> <i>Astropecten scoparius</i> <sup>287</sup>
285		H	H	OH	<i>Dermasterias imbricata</i> <sup>269</sup>
286		OH	OH	H	<i>Asterina pectinifera</i> , <sup>253,254</sup> <i>Patiria miniata</i> , <sup>258</sup> <i>Solaster borealis</i> , <sup>244</sup> <i>Astropecten scoparius</i> <sup>287</sup>
287	 6-O-sulfate	OH	OH	H	<i>Asterina pectinifera</i> <sup>253</sup>
288		H	H	H	<i>Dermasterias imbricata</i> <sup>266</sup>
289		H	H	H	<i>Dermasterias imbricata</i> <sup>266</sup>
290		OH	H	H	<i>Culcita novaeguineae</i> <sup>269</sup>
291		H	H	OH	<i>Dermasterias imbricata</i> <sup>266</sup>
292		OH	OH	H	<i>Patiria miniata</i> , <sup>256</sup> <i>Astropecten scoparius</i> , <sup>267</sup> <i>Solaster borealis</i> <sup>244</sup>
293	 6-O-sulfate	OH	OH	H	<i>Patiria miniata</i> <sup>256</sup>

Table 1 (Continued)

294 from *Dermasterias imbricata*<sup>266</sup>Basic Nuclear Hydroxylation Pattern: 3 $\beta$ ,6 $\beta$ ,8,15 $\alpha$ ,16 $\beta$ 

no.	side chain	R	R <sup>1</sup>	R <sup>2</sup>	sources and references
295		H	H	H	<i>Sphaerodiscus placenta</i> , <sup>247</sup> <i>Crossaster papposus</i> , <sup>270</sup> <i>Culcita novaeguineae</i> <sup>288</sup>
296		H	H	OH	<i>Rosaster</i> sp., <sup>246</sup> <i>Culcita novaeguineae</i> <sup>288</sup>
297		OH	H	OH	<i>Rosaster</i> sp. <sup>246</sup>
298		H	H	H	<i>Sphaerodiscus placenta</i> <sup>247</sup>
299		H	H	H	<i>Hacelia attenuata</i> <sup>271</sup>
300		OH	H	H	<i>Hacelia attenuata</i> <sup>271</sup>
301		OH	H	H	<i>Hacelia attenuata</i> <sup>271</sup>

Basic Nuclear Hydroxylation Pattern: 3 $\beta$ ,6 $\beta$ ,8,15 $\beta$ ,16 $\beta$ 

no.	R	side chain	no.	R	side chain	sources and references
302	H		304	OH		<i>Solaster borealis</i> <sup>244</sup>
303	OH		305	OH		<i>Solaster borealis</i> <sup>244</sup>

26-O derivatization of a signal natural stereoisomer with chlorides of (*R*)-(+)- and (*S*)-(-)- $\alpha$ -methoxy- $\alpha$ -(trifluoromethyl)phenylacetic acid (MTPA, the Mosher's reagent) results in significant differences in the <sup>1</sup>H NMR spectra of the two diastereomeric derivatives,<sup>249</sup> the assignment of configuration at C-25 has been systematically made by comparison of the diastereotopic 26-








H<sub>2</sub> signals in the spectra of the (*R*)-(+)- and (*S*)-(-)-MTPA esters.<sup>197</sup> As exemplified by 265 and 269, in the spectra of the MTPA esters of the 25*S*-isomer 265 the 26-methylene proton signals appear much closer in the spectrum of the (*R*)-(+)-MTPA ester ( $\delta$  4.21 ppm, bd) than in that of the (*S*)-(-)-MTPA derivative ( $\delta$  4.13-4.29 ppm, dd), while the reverse occurs for MTPA esters

Table 2. <sup>1</sup>H NMR Data of Representative Polyhydroxysterols from Starfish

no.	hydroxylation pattern	3-H	4-H	6-H	7β-H	15-H	16-H	18-H <sub>3</sub>	19-H <sub>3</sub>	others
264	3β,6β,15α,16β	3.57 m		3.76 br s		3.80 dd (10.5, 3.0)	4.00 dd (8.5, 3.0)	0.95 s	1.07 s	
265	3β,5,6β,15α,16β	4.04 m		3.50 br s		3.76 dd (10.5, 3.0)	4.00 dd (8.5, 3.0)	0.94 s	1.21 s	
266	3β,5,6β,15α,16β 15-O-sulfate	4.08 m		3.50 br s		4.38 dd	4.33 dd	1.00 s	1.20 s	
267	3β,6β,7α,15α,16β	3.60 m		3.62 t (3.5)	3.91 t (3.5)	3.89 dd (10.5, 3.0)	4.04 dd (7.5, 3.0)	0.96 s	1.04 s	
268	3β,5,6β,7α,15α,16β	4.05 m		3.55 d (3.0)	4.02 t (3.0)	3.88 dd (10.5, 3.0)	4.04 dd (7.5, 3.0)	0.96 s	1.17 s	
273	3β,6α,8,15α,16β	3.59 m		3.66 dd (10.0, 3.0)	2.43 dd (12.0, 3.0)	4.06 dd (11.5, 2.5)	4.00 dd (8.5, 2.5)	1.15 s	1.03 s	
274	3β,4β,6α,8,15α,16β	3.46 m	4.28 br s	4.24 m	2.50 dd (12.0, 3.0)	4.09 dd (10.5, 3.0)	4.01 dd (7.5, 3.0)	1.15 s	1.21 s	
275	3β,6α,7α,8,15α,16β	3.58 m		3.81 m	3.81 m	4.16 dd (11.5, 2.5)	4.03 dd (8.5, 2.5)	1.15 s	1.03 s	
276	3β,4β,6α,7α,8,15α,16β	3.46 m	4.22 m	4.25 dd (11.5, 2.5)	3.88 d (2.5)	4.18 dd (10.5, 2.5)	4.03 dd (8.5, 2.5)	1.15 s	1.21 s	
277	3β,4β,6α,7α,8,15α,16β 6-O-sulfate	3.52 m	4.26 br s	5.03 dd (11.5, 2.5)	4.18 d (2.5)	4.18 dd (10.0, 2.5)	4.02 dd (7.0, 2.5)	1.15 s	1.30 s	
281	3β,6α,8,15β,16β	3.62 m		3.74 td (10.5, 4.0)	2.42 dd (12.5, 4.0)	4.40 dd (6.8, 5.6)	4.25 t (6.8)	1.27 s	1.02 s	
282	3β,4β,6α,8,15β,16β	3.50 m	4.29 br s	4.22 td (10.5, 4.0)	2.49 dd (12.5, 4.0)	4.40 dd (6.8, 5.6)	4.25 t (6.8)	1.27 s	1.19 s	
284	3β,6α,7α,8,15β,16β	3.53 m		3.87 td (12.5, 2.5)	3.90 d (2.5)	4.50 dd (6.5, 5.5)	4.25 t (6.5)	1.28 s	1.02 s	14-H: 1.43 d (5.5)
285	3β,6α,8,14α,15β,16β	3.50 m		3.77 td (10.5, 4.0)	2.14 dd (12.5, 4.0)	4.00 d (6.5)	4.37 t (6.5)	1.35 s	1.04 s	
286	3β,4β,6α,7α,8,15β,16β	3.47 m	4.22 br s	4.32 td (11.0, 2.5)	3.98 d (2.5)	4.51 dd (6.5, 5.5)	4.26 t (6.5)	1.27 s	1.19 s	5-H: 1.52 dd (11.2, 3.0) 14-H: 1.42 d (5.5)
295	3β,6β,8,15α,16β	3.43 m		3.89 br s	2.46 dd (15.0, 2.5)	4.17 dd (11.2, 7.5)	4.02 dd (7.5, 2.5)	1.15 s	1.20 s	
296	3β,6β,7α,8,15α,16β			3.72 dd (3.1, 2.9)	3.87 d (3.1)	4.21 dd (10.0, 2.5)	4.00 dd (7.5, 2.5)	1.15 s	1.15 s	
297	3β,4β,6β,7α,8,15α,16β	3.50 m	4.10 br s	4.04 dd (3.1, 2.9)	3.85 d (3.0)	4.21 dd (10.0, 2.5)	4.00 dd (7.5, 2.5)	1.15 s	1.43 s	
302	3β,6β,8,15β,16β	3.60 m		3.60 dd (3.0, 2.9)	4.03 d (3.0)	4.50 dd (5.0, 6.2)	4.25 t (6.2)	1.30 s	1.18 s	
303	3β,4β,6β,8,15β,16β	3.55 m	4.08 br s	4.00 dd	4.02 d (3.0)	4.50 dd (5.0, 6.2)	4.23 t (6.2)	1.30 s	1.45 s	

<sup>a</sup> Data (δ, ppm) were mostly obtained at 250 MHz from solution in CD<sub>3</sub>OD and are referred to the central line of the CHD<sub>2</sub>OD signal (δ 3.34 ppm).

Table 3. Selected <sup>1</sup>H NMR Data of Side Chains of Polyhydroxysterols from Starfish

	no.	21-H	22, 23-H's	26-H <sub>2</sub>	27-H <sub>3</sub>	28-H's	other
	273	0.96 d (7.0)		3.46 dd (10.5, 6.0) ca. 3.32 <sup>b</sup>	0.94 d (7.0)		
	288	1.06 d (6.5) 5.51 dt (16.0, 6.5)	5.60 dd (6.5, 16.0)	3.46 dd (10.5, 6.0) ca. 3.32 <sup>b</sup>	0.92 d (7.0)		
	289	1.00 d (7.5)		3.41 dd (11.0, 7.5) 3.63 dd (11.0, 6.0)	1.10 d (7.5)	4.78 br s 4.85 br s	
	278	0.95 d (7.0)		3.40 dd (10.5, 6.0) ca. 3.32 <sup>b</sup>	0.82 d (6.5)	0.84 d (6.5)	
	279	1.04 (7.0)	5.45 m	3.41 dd (11.0, 6.0) 3.56 dd (11.0, 6.0)	0.91 d (7.0)	0.99 d (7.0)	20-H: 2.55 m 24-H: 2.10 m
	293	1.07 (7.0)	5.54 m	3.42 d (11.0) 3.50 d (11.0)	1.10 s	1.01 d (7.0)	
	280	0.96 d (7.0)		0.89 d (7.0)	0.92 d (7.0)		29-H <sub>2</sub> : 4.06 m (desulfated: 3.60 m)


<sup>a</sup> Data (δ, ppm) were mostly obtained at 250 MHz from solution in CD<sub>3</sub>OD and are referred to the central line of the CHD<sub>2</sub>OD signal (δ 3.34 ppm). <sup>b</sup> Partially overlapping with solvent signal.

of the 25*R*-isomer 269, the resonance being closer in the (*S*)-(-)-MTPA ester (δ 4.19–4.23 ppm, dd) and more separated in the (*R*)-(+)-MTPA (δ 4.14–4.28 dd). It has also been observed that derivatization on ring D, as sulfate group at C-15 in 266 and 270, can affect the extent of the separation of the 26-methylene proton signals, thus it is always advisable to assign the configuration by direct comparison of both (*R*)-(+)- and (*S*)-(-)-MTPA derivatives.<sup>240,245</sup>

Compounds 269, 270, and 272, all isolated from the deep-water starfish *Tremaster novaecaledoniae*, represent the only examples of 26-hydroxysteroids from starfish with 25*R* configuration. 272 is also the only one to exhibit a 3α-hydroxy-5β-steroidal nucleus, as indicated by the low field signal of the angular methyl

carbon-19 at 26.1 ppm and by the multiplet at δ 3.54 ppm, having the typical shape for a 3β-hydroxymethine signal in a 5β-steroid.

Compounds 273, 275, and 276, with the basic 3β,6α,8,15α,16β-hydroxylation pattern, were the first polyhydroxysteroids to be isolated from a starfish, the Pacific Ocean species *Protoreaster nodosus*;<sup>250</sup> they were later shown to be the most common polyhydroxysteroids from starfish. They exhibited moderate cytotoxic and anticancer activities.<sup>253,260</sup> The structures of these first steroids were determined on the results of NMR and mass spectral analysis, chemical transformations and related spectroscopic data. Acetylation of 273 followed by oxidation with Jones reagent afforded 3β,6α,15α,26-tetrakis(acetyloxy)-8-hydroxy-5α-cholestan-16-



**Figure 15.** Unusual polyhydroxysteroids with a side-chain methyl group oxidized to carboxyl group from the starfish *Myxoderma platyacanthum*.

one. The elimination of the side chain with migration of one hydrogen ( $m/z$  464, McLafferty rearrangement) followed by 18-methyl fission ( $m/z$  449) in the mass spectrum was diagnostic of a 16-ketosteroid. Acetylation of 275 and 276 afforded likewise the corresponding tetraacetates [3 $\beta$ ,6 $\alpha$ ,15 $\alpha$ ,26-tetrakis(acetyloxy)], which, on oxidation with Jones reagent, gave the 3 $\beta$ ,6 $\alpha$ ,15 $\alpha$ ,26-tetrakis(acetyloxy)-7 $\alpha$ ,8-dihydroxy-5 $\alpha$ -cholestan-16-one and 3 $\beta$ ,6 $\alpha$ ,15 $\alpha$ ,26-tetrakis(acetyloxy)-7 $\alpha$ ,8-dihydroxy-5 $\alpha$ -cholestan-4,16-dione, respectively. Treatment of 273, 275, and 276 with *p*-bromobenzoyl chloride in pyridine led to the formation of the corresponding 3 $\beta$ ,6 $\alpha$ ,15 $\alpha$ ,26-tetrakis(*p*-bromobenzoates) with CD curves displaying strong positive first and negative second Cotton effects ( $\Delta\epsilon_{252} +37.2/\Delta\epsilon_{235} -20.0$ ) in agreement with the expected positive chirality of the three major dibenzoates interactions (3 $\beta$ /6 $\alpha$ , 3 $\beta$ /15 $\alpha$ , 6 $\alpha$ /15 $\alpha$ ) in a cholestane skeleton with 5 $\alpha$ -H absolute configuration. The 25S configuration was assigned, as described before, by  $^1\text{H}$  NMR pattern of the 26- $\text{H}_2$  signals of the (*R*)-(+)- and (*S*)-(-)-MTPA derivatives.<sup>249</sup>

Steroids 274, 278, and 279 were later isolated from the same starfish species, *Protoreaster nodosus*,<sup>257</sup> has since been isolated also from other species. The stereochemistry of the side chains of 278 and 279 is suggested after the stereoselective synthesis of the 24-methyl-26-hydroxysteroid models.<sup>261</sup> Compound 280 is a quite rare example of non-glycosylated C-29 polyhydroxysteroid, these compounds have been more commonly found as aglycons in several glycosides of polyhydroxysteroids (see section VII.A.2). The 24R configuration is suggested on the basis of the reported chemical shifts of the isopropyl methyls in both  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra and comparison with model compounds.<sup>262,263</sup>


Steroids 281–293 share the common 3 $\beta$ ,6 $\alpha$ ,8,15 $\beta$ ,16 $\beta$ -hydroxylation pattern, while 294 is a rare example of a 26-hydroxysteroid missing the 16 $\beta$ -hydroxyl function. The inversion of stereochemistry at C-15, with respect to the previous ones, is connected with a slight downfield shift of the 18-methyl proton signal and with major changes of the 7-, 15-, and 16-proton signals, whose shapes and chemical shifts become distinctive features of this structural subgroup. The *cis*-15,16-dihydroxy stereochemistry was supported by the formation of a 15,16-monoacetonide on treatment of the hexol 281 with  $\text{Me}_2\text{CO}$  and  $\text{TsOH}$  and by the formation of a 3,4:15,16-bisacetonide on similar treatment of the heptol 282.<sup>264</sup> The chemical shift differences of the 26-methylene protons in the 26-*O*-(*R*)-(+)- and 26-*O*-(*S*)-(-)-MTPA esters ( $\delta$  4.26–4.31 dd and 4.20–4.41 dd,

respectively) were used to establish the 25S configuration of 292.<sup>256</sup> The 14 $\alpha$ -hydroxyl group in 285 and 291 induces substantial modifications in both  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra *vs* their 14-deoxy analogs 281 and 289, most significant in the  $^{13}\text{C}$  NMR spectrum are the large  $\gamma$ -upfield shifts observed for the C-7 (–4.6 ppm), C-9 (–8.9 ppm), C-12 (–5.4 ppm), and C-17 (–10.0 ppm) signals. The absolute configuration of the side chain in 291 has been suggested by comparison of  $^1\text{H}$  and  $^{13}\text{C}$  NMR data with those of the steroid 315 from *Archaster tipicus*.

The steroids 295–301 originate the structural group having the 3 $\beta$ ,6 $\beta$ ,8,15 $\alpha$ ,16 $\beta$ -hydroxylation pattern. Most significant modifications of  $^1\text{H}$  NMR signals *vs* their 6 $\alpha$ -counterparts refer to the shape and chemical shift of the 6-H signal and to the 19-methyl proton signal, downfield shifted by *ca.* 0.17 ppm. Compounds 299–301 are very minor components of the Mediterranean starfish *Hacelia attenuata* and their structures with truncated side chains were mainly deduced by interpretation of NMR spectra. The occurrence of these truncated steroids may be of some interest as indicators of the capability of the starfish to oxidize dietary sterols.

Steroids 302–305, all isolated recently from the starfish *Solaster borealis*,<sup>244</sup> represent a further and minor variation in the stereochemical disposition of the nuclear hydroxyl groups. Compounds 302 and 303 are epimeric with the previous 284, 286, 296 and 297 from which they differ alternatively for the stereochemistry at C-6 or at C-15. The relative *threo* configuration at C-24 and C-25 of 305 was assigned by comparison of the NMR spectral data of the side chain with those of synthetic model compounds, while the absolute (24*R*,25*S*) configuration followed from derivatization with (*R*)-(+)-MTPA chloride and comparison of the  $^1\text{H}$  NMR spectrum with those of the 26-*O*-(*R*)-(+)-MTPA esters of the two *threo* synthetic models (*i.e.* the 24*R*,25*S*- and 24*S*,25*R*-isomeric pair).

Polyhydroxysteroids 306–309 (Figure 15), isolated from the starfish *Myxoderma platyacanthum*<sup>245</sup> together with 265 and 266, have the same 3 $\beta$ ,5,6 $\beta$ ,15 $\alpha$ -tetrahydroxycholestane nucleus with different side chains: 26-acid;  $\Delta^{22}$ -24-methyl 26-acid;  $\Delta^{22}$ -27-nor-24-methyl 26-acid; and 24-carboxymethyl. In 306 and 307 the carboxylic acid function is found as the amide derivative of taurine. They are examples of the structural variety of steroids cooccurring in the same organism and also constitute the first reported isolation from starfish of steroids with a methyl group oxidized to carboxyl. The absolute configuration of the side chain stereogenic centers in 307–309 was determined




**Figure 16.** Polyhydroxysteroids from the starfish *Archaster typicus*.

by spectral comparison with stereoisomeric  $\Delta^{22(E)}$ -24-methyl 26-acid,<sup>261</sup> 24-methyl-27-nor 26-acid,<sup>245</sup> and 24-carboxymethyl<sup>263</sup> steroidal models obtained by stereoselective synthesis. Only very small differences were observed in the <sup>1</sup>H NMR spectra of the two epimeric 24-methyl-27-nor 26-acid synthetic models, mainly dealing with the shifts of the olefinic protons: double doublets centered at  $\delta$  5.32 with the internal lines coincident in the 24*S*-isomer and separated by 1 Hz in the 24*R*-isomer. The olefinic pattern in the spectrum of the natural 308 was coincident with that of the 24*S*-


model, thus suggesting the reported absolute configuration.

Figure 16 illustrates the variety of highly hydroxylated steroids encountered in *Archaster typicus*,<sup>198,272</sup> these steroids have been isolated in relatively large amounts as compared to the very limited fraction of steroidal glycosides in this starfish. Compounds 313 and 318 were shown to be moderately cytotoxic and to cause inhibition of growth of human lymphoma cells at a dose of 0.05  $\mu$ g/mL.<sup>260</sup> The structures of the nonols 310–312, 315, and 316, the most highly hydroxylated





#	side chain	R	R <sup>1</sup>	R <sup>2</sup>	#	side chain
325		H	OH	H	330	
326		OH	H	H	331	
327		OH	H	OH	332	
328		OH	H	H		
329		OH	H	H		





#	R	R <sup>1</sup>
333	OH	
334	OH	
335	H	

Figure 17. Polyoxygenated steroids from the starfish *Styrcaster caroli*.


steroids isolated from natural sources, were determined essentially upon accurate spectral analysis. Major

support in the elucidation of structures 310–312 came from 2D <sup>1</sup>H-<sup>13</sup>C NMR heterocorrelation experiments

Chart I




#	R	sources and references	side chain	#	R	sources and references
336	H	<i>Asterina pectinifera</i> <sup>253</sup>		338	H	<i>Gomophia watsoni</i> <sup>276</sup> , <i>Nardoa gomophia</i> <sup>265</sup> , <i>Dermasterias imbricata</i> <sup>266</sup> , <i>Culcita novaeguineae</i> <sup>269</sup>
337	H	<i>Aphelasterias japonica</i> <sup>274</sup>		339	OH	<i>G. watsoni</i> <sup>276</sup> , <i>C. novaeguineae</i> <sup>269</sup>
				340	H	<i>Astropecten scoparius</i> <sup>267</sup>
				341	H	<i>Dermasterias imbricata</i> <sup>266</sup>



342

source

*Coscinasterias tenuispina*<sup>268</sup>

343

(HETCOR and COLOC) that, through the long-range correlations with quaternary carbons C-5 and C-14, allowed firm location of tertiary hydroxyl groups and consequent build up of the tetracyclic framework. Stereochemical arrangement of hydroxyl groups was deduced by NOEDS experiments showing through-space interactions between 6-H and 19-Me, 9-H and 7-H, 15-H and 18-Me. Furthermore, on treatment with acetone and TsOH, compound 310 afforded a 3 $\beta$ ,4 $\beta$ :14 $\alpha$ ,15 $\alpha$ -bisacetone, thus supporting the 14 $\alpha$ -hydroxy stereochemistry. The application of the Horeau's method of kinetic resolution to this bisacetone allowed the assignment of the 24*R* configuration to C-24. The assignment of the stereochemistry at C-24 and C-25 in 315 and 316 required NMR spectral comparison with synthetic model compounds.<sup>198</sup> The structure 310, which has the remarkable feature of eight sequential hydroxyl groups protruding from the same side of the molecule, has been confirmed by a single-crystal X-ray diffraction study.<sup>273</sup> The crystal packing of molecules was shown to be a consequence of the amphiphilic character; the hydroxyl groups develop an intricate and extensive network of hydrogen bonds connecting the

molecules through their hydrophilic moieties and giving rise to double layers, which interact through the hydrophobic surfaces.

A group of 3 $\beta$ ,6 $\alpha$ -disulfated steroids 319–322 has been more recently isolated from the "living fossil" species *Tremaster novaecaledoniae* collected at a depth of 530 m during exploration of the bathyal zone off New Caledonia.<sup>240</sup> Furthermore, analysis of polar extracts led to the isolation of the steroid 323 and of a new group of glycosides of polyhydroxysteroids, the tremasterols A–C (459–461), all possessing both phosphate and sulfate conjugation.<sup>239,240</sup> To the best of our knowledge they constitute the first reported isolation of phosphated steroids from a natural source. The presence of the phosphate group at C-6, at first indicated by the shape of the 6-proton signal, was confirmed by the proton noise-decoupled <sup>13</sup>C NMR spectrum in which the C-5 and C-6 signals appeared as doublets due to the <sup>31</sup>P–<sup>13</sup>C couplings through two and three linkages and finally by <sup>31</sup>P NMR. *Tremaster novaecaledoniae* contains also the uncommon polyhydroxysteroids with 25*R* stereochemistry 269, 270, and 272.

A further  $3\beta,6\alpha$ -disulfated steroid **324**, closely related to the aglycons of the asterosaponins also by the presence of an oxygenated function at C-23, has been isolated from the starfish *Aphelasterias japonica*.<sup>274</sup>

The very recent investigation of *Styracaster caroli*, a deep-water starfish collected between Thio and Lifou (New Caledonia) at a depth of 2000 m, revealed the occurrence of a very complex mixture of unprecedented polyhydroxysteroids (Figure 17) with unusual side chains.<sup>275</sup> Compounds **327** and **331** show a unique 24-ethyl-25-hydroxy-26-sulfoxycholestane skeleton, while compound **328** presents a  $\Delta^{22-23,24}$ -dimethyl substitution pattern typical of dinosterol (**11**) and **329** is its 29-hydroxy derivative. An unusual oxygenation pattern is found in the C<sub>29</sub> side chain of **332**, with a sulfoxy group at C-28 and a hydroxyl group at C-29. Compounds **333–335** are further examples of polyhydroxysteroids possessing an amido function in the side chain, with a D-(-)-cysteinolic acid (2*S*-amino-3-hydroxypropanesulfonic acid) linked to a C-24 carboxyl group.

In a more limited number of cases polyhydroxysteroids with hydroxyl or sulfate groups at C-24 have been isolated, **336–343** (Chart I). The hydroxylation at C-24 is a structural feature commonly encountered in the aglycons of the many steroidal glycosides isolated from starfish and indeed **336** is the aglycon of the glycosides **351–359**, while **338** and **339** have been found as the native aglycons in the glycosides **369–383** and **384–388**, respectively. The unsaturated **341** is the aglycon of **391–393** and the 15-*O*-sulfated of **343** has been found in **434**. The isomeric relationship at C-15 between **336** and **338** was confirmed by formation of an 8,15-phenylboronate in **338**. The 24*S* configuration was suggested in **338** by <sup>13</sup>C NMR data in comparison with those reported for 24(*S*)- and 24(*R*)-hydroxycholestanol. In fact differences observed between the spectra of the two epimers are very small<sup>277,278</sup> and in the desulfated **340** the 24*S* configuration was later confirmed, upon derivatization with (*R*)-(+)- and (*S*)-(-)-MTPA, by the Mosher's method for determination of absolute configuration of chiral secondary carbinols.<sup>279,280</sup> The steroid **341** was only isolated in admixture, with its 22-, 23-dihydro congeners **338** resistant to further attempts of separation.

## 2. Glycosides of Polyhydroxysteroids


This group of steroidal glycosides from starfish shows a large degree of structural variability. Most of these compounds usually occur in minute amounts and are also widespread among starfish, having been found, usually as complex mixtures, in almost all the species investigated. They are composed of a polyhydroxylated steroidal aglycon and a carbohydrate portion mostly made up from one or two monosaccharide units (Figure 18) often linked to each other and glycosidically attached at C-3 or C-24 of the aglycon. Only very recently cytotoxic triglycosides (**418**, **419**, and **421**) were isolated from the New Caledonian species *Fromia monilis*.<sup>281</sup> They constitute the only examples of triglycosides among more than 100 different mono and diglycosides of polyhydroxysteroids isolated so far. The most common monosaccharides are D-xylopyranose, often methylated at position 2 and/or 4 and occasionally at position 3, and L-arabinose, found in its furanose form. Rare examples of xylofuranosides (**371**, **372**, and

**403**), galactofuranosides (**400**, **441**, and **444**), fucofuranosides (**389** and **390**), and arabinopyranosides (**417** and **420**) have also been found.

The first representative of such compounds, the cytotoxic nodoside **431**, was first isolated from Pacific *Protoreaster nodosus*<sup>282</sup> and later from other Valvatida species. This was followed by the structures of more than 100 different glycosides of polyhydroxysteroids. Structural variations originate from the hydroxylation pattern of the steroidal tetracyclic nucleus, the functionalization of the side chain, the presence of sulfate, and the nature and location of the saccharide moiety. Beside the invariable  $3\beta$ -hydroxylation, hydroxyl groups are commonly found at the positions  $6\alpha$ (or  $\beta$ ), 8,  $15\alpha$ (or  $\beta$ ), and 24 of the aglycon with additional hydroxyl group(s) at one or more of the positions  $4\beta$ ,  $5\alpha$ ,  $7\alpha$ , and  $16\beta$ . The glycosides of polyhydroxysteroids often occur as complex mixtures with free and sulfated polyhydroxysteroids and asterosaponins; for example the polar steroids from *Coscinasterias tenuispina* have been resolved into 19 constituents.<sup>268</sup> Illustrative examples of the structural variety of steroidal glycosides cooccurring in the same organism are the 11 glycosides of polyhydroxysteroids isolated from the starfish *Henricia laeviuscola*,<sup>283</sup> the eight steroidal diglycosides, the halitylosides, isolated from the starfish *Halityle regularis*<sup>284</sup> along with two polyhydroxysteroids and four asterosaponins, and the 11 glycosides and five polyhydroxysteroids isolated during a recent reinvestigation of the polar extractives from the starfish *Culcita novaeguineae*.<sup>269</sup>

The structures of the currently known glycosides of polyhydroxysteroids are listed in Table 4 with their sources and references. Compounds **344–350** are uncommon examples of steroidal glycosides lacking the hydroxyl group at C-8 of the aglycon, a structural feature present in all other glycosides. Glycosides **351–363** are characterized by the steroidal aglycons with the basic  $3\beta,6\alpha,8,15\alpha$ -hydroxylation pattern, while in **364–368** the tetracyclic nucleus also bears a  $16\beta$ -OH. The largest group of compounds presents aglycons with the basic  $3\beta,6\alpha,8,15\beta$ -hydroxylation pattern, **369–402**, with a single case of  $\Delta^4$ -unsaturation in **402**. The (2*S*)- $5\alpha$ -cholestane- $3\beta,6\alpha,8,15\beta,24$ -pentol aglycon, first encountered in attenuoside A II (**369**) from *Hacelia attenuata*, is the most common steroidal aglycon found in the glycosides from starfish. In the aglycons with the  $3\beta,6\alpha,8,15\beta,16\beta$ -hydroxylation pattern **403–422** the oxygenated C-28 and C-29 side chains prevail on the common 24-hydroxyl side chain only found in indicoside C (**403**). Compounds **423–436** constitute the group of glycosides whose aglycons possess the basic  $3\beta,6\beta,8,15\alpha$ -hydroxylation pattern; this group of compounds comprises several examples of those glycosides in which the two monosaccharides are not linked to each other but at positions 3 and 24 of the aglycon. Also in the group with  $3\beta,6\beta,8,15\alpha,16\beta$ -hydroxylation pattern **437–452** the C-28 and C-29 aglycons are predominant; moreover in the  $\Delta^4$ -unsaturated compounds **447–452** the xylosyl unit is constantly found linked at C-3 of the aglycon. The smallest group comprising compounds **453–455** has the basic  $3\beta,6\beta,8,15\beta$ -hydroxylation pattern.

Compounds **456–458** cannot be classified in any of the above structural groups; they contain aglycons with



**Figure 18.** Saccharide chains found in glycosides of polyhydroxysterols from starfish.

unsaturated tetracyclic nuclei, a quite unusual feature among glycosides of polyhydroxysteroids from starfish,


only found in the above mentioned compounds with Δ<sup>4</sup>-steroidal skeleton.

Table 4. Glycosides of Polyhydroxysteroids<sup>a</sup>

Basic Nuclear Hydroxylation Pattern: 3 $\beta$ ,6 $\alpha$ ,15 $\alpha$							
name	no.	side chain	R	R <sup>1</sup>	R <sup>2</sup>	sources and references	
amurensoside A	344		H	SO <sub>3</sub> -Na <sup>+</sup>	X <sub>1</sub>	<i>Asterias amurensis</i> <sup>284</sup>	
amurensoside B	345		OH	SO <sub>3</sub> -Na <sup>+</sup>	X <sub>1</sub>	<i>Asterias amurensis</i> <sup>284</sup>	
amurensoside C	346		H	H	X <sub>4</sub>	<i>Asterias amurensis</i> <sup>284</sup>	
Basic Nuclear Hydroxylation Pattern: 3 $\beta$ ,6 $\beta$ ,15 $\alpha$							
name	no.	side chain	R	R <sup>1</sup>	R <sup>2</sup>	sources and references	
granuloside B	347		X <sub>2</sub>	H	A <sub>1</sub>	<i>Choriaster granulatus</i> <sup>285</sup>	
laeviuscoside F	348		X <sub>4</sub>	OH	A <sub>1</sub>	<i>Henricia laeviuscola</i> <sup>283</sup>	
3 $\beta$ ,6 $\alpha$ ,15 $\beta$							
amurensoside D	349	<i>Asterias amurensis</i> <sup>284</sup>					
3 $\beta$ ,6 $\beta$ ,15 $\beta$							
moniloside C	350	<i>Fromia monilis</i> <sup>281</sup>					
Basic Nuclear Hydroxylation Pattern: 3 $\beta$ ,6 $\alpha$ ,8,15 $\alpha$							
name	no.	side chains	R	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	sources and references
asterosaponin P-1	351		H	H	H	A <sub>2</sub>	<i>Patiria pectinifera</i> , <sup>266</sup> <i>Oreaster reticulatus</i> , <sup>269</sup> <i>Patiria miniata</i> <sup>268</sup>

Table 4 (Continued)

name	no.	side chains	R	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	sources and references
	352		H	H	H	A <sub>3</sub>	<i>Oreaster reticulatus</i> , <sup>259</sup> <i>Patiria miniata</i> <sup>256</sup>
miniatoside B	353		H	H	H	XA <sub>1</sub>	<i>Patiria miniata</i> <sup>256</sup>
crossasteroside B	354		H	H	H	XX <sub>3</sub>	<i>Crossaster papposus</i> <sup>287</sup>
crossasteroside C	355		H	H	H	XA <sub>4</sub>	<i>Crossaster papposus</i> <sup>287</sup>
borealoside A	356		H	H	H	XA <sub>3</sub>	<i>Solaster borealis</i> <sup>244</sup>
borealoside B	357		H	H	H	XA <sub>2</sub>	<i>Solaster borealis</i> <sup>244</sup>
borealoside C	358		H	H	H	X <sub>3</sub>	<i>Solaster borealis</i> <sup>244</sup>
crossasteroside A	359		H	H	OH	XX <sub>3</sub>	<i>Crossaster papposus</i> <sup>288</sup>
crossasteroside D	360		H	H	OH	XX <sub>2</sub>	<i>Crossaster papposus</i> <sup>288</sup>
attenuatoside C	361		OH	H	H	A <sub>1</sub>	<i>Hacelia attenuata</i> <sup>289</sup>
borealoside D	362		OH	H	H	X <sub>3</sub>	<i>Solaster borealis</i> <sup>244</sup>
6- <i>epi</i> -nodososide	363		H	OH	H	XA <sub>5</sub>	<i>Pentaceraster alveolatus</i> <sup>252</sup>

Basic Nuclear Hydroxylation Pattern: 3 $\beta$ ,6 $\alpha$ ,8,15 $\alpha$ ,16 $\beta$ 

name	no.	side chains	R	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	sources and references
	364		X <sub>2</sub>	H	H		<i>Poraster superbus</i> <sup>251</sup>
	365		X <sub>3</sub>	H	OH		<i>Poraster superbus</i> <sup>251</sup>
	366		H	H	H	A <sub>1</sub>	<i>Patiria pectinifera</i> <sup>290</sup>
miniatoside A	367		H	H	H	A <sub>4</sub>	<i>Patiria miniata</i> <sup>256</sup>

Table 4 (Continued)



















name	no.	side chains	R	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	sources and references
asterosaponin P-2	368		H	OH	H	A <sub>2</sub>	<i>Patiria pectinifera</i> <sup>291</sup>
Basic Nuclear Hydroxylation Pattern: 3β,6α,8,15β							
							
name	no.	side chain	R	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	sources and references
attenuatoside A-II	369		H	H	H	A <sub>1</sub>	<i>Hacelia attenuata</i> <sup>271</sup>
scoparioside A	370		H	H	H	A <sub>5</sub>	<i>Astropecten scoparius</i> <sup>287</sup>
scoparioside B	371		H	H	H	Xf <sub>1</sub>	<i>Astropecten scoparius</i> <sup>287</sup>
indicoside B	372		H	H	H	Xf <sub>2</sub>	<i>Astropecten indicus</i> <sup>292</sup>
pycnopodioside A	373		H	H	H	X <sub>1</sub>	<i>Pycnopodia helianthoides</i> <sup>258</sup>
pycnopodioside B	374		SO <sub>3</sub> <sup>-</sup> Na <sup>+</sup>	H	H	X <sub>1</sub>	<i>Pycnopodia helianthoides</i> <sup>258</sup>
pycnopodioside C	375		H	H	H	Glc <sub>2</sub>	<i>Pycnopodia helianthoides</i> <sup>258</sup>
luridoside A	376		H	H	H	X <sub>5</sub>	<i>Comasterias lurida</i> <sup>293</sup>
scoparioside C	377		H	H	H	X <sub>6</sub>	<i>Astropecten scoparius</i> <sup>287</sup>
attenuatoside A-I	378		H	H	H	XA <sub>5</sub>	<i>Hacelia attenuata</i> <sup>294</sup>
culcitoside C <sub>4</sub>	379		H	H	H	XA <sub>4</sub>	<i>Culcita novaeguineae</i> <sup>289</sup>
halityloside E	380		H	H	H	XA <sub>6</sub>	<i>Halityle regularis</i> , <sup>264</sup> <i>Nardoa gomophia</i> , <sup>285</sup> <i>Sphaerodiscus placenta</i> , <sup>247</sup> <i>Culcita novaeguineae</i> <sup>288</sup>
glacialoside A	381		X <sub>1</sub>	H	H	SO <sub>3</sub> <sup>-</sup> Na <sup>+</sup>	<i>Marthasterias glacialis</i> <sup>295</sup>
distolasteroside D <sub>1</sub>	382		X <sub>1</sub>	H	H	X <sub>1</sub>	<i>Distolasterias nippon</i> <sup>296</sup>



Table 4 (Continued)

name	no.	side chain	R	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	sources and references
distolasteroside D <sub>3</sub>	383		S <sub>1</sub>	H	H	Glc <sub>1</sub>	<i>Distolasterias nippon</i> <sup>286</sup>
attenuatoside B-2	384		H	OH	H	A <sub>1</sub>	<i>Hacelia attenuata</i> <sup>289</sup>
attenuatoside B-1	385		H	OH	H	XA <sub>5</sub>	<i>Hacelia attenuata</i> <sup>289</sup>
culcitoside C <sub>5</sub>	386		H	OH	H	XA <sub>4</sub>	<i>Culcita novaeguineae</i> <sup>289</sup>
halytiloside D (culcitoside C <sub>1</sub> )	387		H	OH	H	XA <sub>6</sub>	<i>Halityle regularis</i> , <sup>284</sup> <i>C. novaeguineae</i> , <sup>289,297</sup> <i>N. gomphia</i> , <i>N. novaecaledoniae</i> , <sup>285</sup> <i>Linkia guildingi</i> <sup>297</sup>
glacialoside B	388		SO <sub>3</sub> <sup>-</sup> Na <sup>+</sup>	OH	H	X <sub>1</sub>	<i>Marthasterias glacialis</i> <sup>295</sup>
imbricatoside B	389		H	H	OH	QuiFucf	<i>Dermasterias imbricata</i> <sup>289</sup>
imbricatoside A	390		H	OH	OH	QuiFucf	<i>Dermasterias imbricata</i> <sup>289</sup>
luridoside B	391		H	H	H	X <sub>5</sub>	<i>Comasterias lurida</i> <sup>293</sup>
22-dehydrohalytiloside E (poranoside A)	392		H	H	H	XA <sub>6</sub>	<i>Sphaerodiscus placenta</i> , <sup>247</sup> <i>Porania pulvillus</i> <sup>298</sup>
distolasteroside D <sub>2</sub>	393		X <sub>1</sub>	H	H	X <sub>1</sub>	<i>Distolasterias nippon</i> <sup>296</sup>
22-dehydrohalytiloside D	394		H	OH	H	XA <sub>6</sub>	<i>Sphaerodiscus placenta</i> <sup>247</sup>
coscinasteroside F	395		H	OH	H	X <sub>1</sub>	<i>Coscinasterias tenuispina</i> <sup>288</sup>
halytiloside I (6-O-sulphate)	396		H	OH	H	XX <sub>4</sub>	<i>Halityle regularis</i> , <sup>284</sup> <i>Nardoa gomphia</i> <sup>285</sup>
placentoside A	397		H	OH	H	XX <sub>4</sub>	<i>Sphaerodiscus placenta</i> <sup>247</sup>
culcitoside C <sub>6</sub>	398		H	OH	H	XA <sub>6</sub>	<i>Coscinasterias tenuispina</i> <sup>288</sup>
culcitoside C <sub>7</sub> 6-sulfate	399		H	OH	H	XA <sub>6</sub>	<i>Coscinasterias tenuispina</i> <sup>288</sup>
indicoside A	400		H	H	OH	Galf	<i>Astropecten indicus</i> <sup>299</sup>
gomphioside B	401		H	OH	H	XA <sub>6</sub>	<i>Gomphia watsoni</i> <sup>276</sup>

Table 4 (Continued)


pisasteroside E 402 *Pisaster giganteus*<sup>300</sup>Basic Nuclear Hydroxylation Pattern; 3 $\beta$ ,6 $\alpha$ ,8,15 $\beta$ ,16 $\beta$ 

name	no.	side chain	R	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	sources and references
indicoside C	403		H	H	H	Xf <sub>2</sub>	<i>Astropecten indicus</i> <sup>292</sup>
coscinasteroside E	404		H	OH	H	X <sub>1</sub>	<i>Coscinasterias tenuispina</i> <sup>268</sup>
coscinasteroside D	405		H	H	H	Glc <sub>2</sub>	<i>Coscinasterias tenuispina</i> <sup>268</sup>
coscinasteroside A	406		X <sub>1</sub>	H	OH	SO <sub>3</sub> -Na <sup>+</sup>	<i>Coscinasterias tenuispina</i> <sup>268</sup>
culcitoside C <sub>3</sub>	407		H	H	H	XX <sub>4</sub>	<i>Culcita novaeguineae</i> <sup>268</sup>
culcitoside C <sub>3</sub>	408		H	H	H	XA <sub>6</sub>	<i>Culcita novaeguineae</i> <sup>301</sup>
culcitoside C <sub>2</sub>	409		H	OH	H	XA <sub>6</sub>	<i>Culcita novaeguineae</i> <sup>301</sup>
coscinasteroside C	410		H	H	H	Glc <sub>3</sub>	<i>Coscinasterias tenuispina</i> <sup>268</sup>
pisasteroside A	411		H	H	H	Glc <sub>2</sub>	<i>Pisaster ochraceus</i> and <i>Pisaster brevispinus</i> <sup>302</sup>
pisasteroside F	412		H	H	H	Glc <sub>2</sub>	<i>Pisaster giganteus</i> <sup>300</sup>
halityloside B	413		H	H	H	XX <sub>4</sub>	<i>Halityle regularis</i> , <sup>264</sup> <i>Nardoia novaecaledoniae</i> and <i>N. gomophia</i> , <sup>265</sup> <i>Sphaerodiscus placenta</i> , <sup>247</sup> <i>Culcita novaeguineae</i> <sup>269</sup>

Table 4 (Continued)

name	no.	side chain	R	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	sources and references
halityloside A	414		H	OH	H	XX <sub>4</sub>	<i>Halityle regularis</i> , <sup>264</sup> <i>Nardoa novaecaledoniae</i> and <i>N. gomophia</i> , <sup>265</sup> <i>Sphaerodiscus placenta</i> , <sup>247</sup> <i>Culcita novaeguineae</i> <sup>268</sup>
halityloside H	415		H	OH	H	XA <sub>6</sub>	<i>Halityle regularis</i> , <sup>264</sup> <i>N. gomophia</i> <sup>265</sup>
halityloside H 6-O-sulfate	416		H	OH	H	XA <sub>6</sub>	<i>Halityle regularis</i> , <sup>264</sup> <i>N. gomophia</i> <sup>265</sup>
moniloside E	417		H	OH	H	AX	<i>Fromia monilis</i> <sup>284</sup>
moniloside I	418		H	OH	H	XXX <sub>1</sub>	<i>Fromia monilis</i> <sup>284</sup>
moniloside G	419		H	OH	H	XXX <sub>2</sub>	<i>Fromia monilis</i> <sup>284</sup>
moniloside F	420		H	OH	H	AX	<i>Fromia monilis</i> <sup>284</sup>
moniloside H	421		H	OH	H	XXX <sub>2</sub>	<i>Fromia monilis</i> <sup>284</sup>
pisasteroside C	422		H	H	H	X <sub>5</sub>	<i>Pisaster brevispinus</i> <sup>302</sup>


Basic Nuclear Hydroxylation Pattern: 3β,6β,8,15α



name	no.	side chain	R	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	sources and references
coscinasteroside B	423		H	H	H	SO <sub>3</sub> -Na <sup>+</sup>	X <sub>1</sub>	<i>Coscinasterias tenuispina</i> , <sup>269</sup> <i>Pycnopodia helianthoides</i> <sup>256</sup>
pisasteroside B	424		X <sub>1</sub>	H	H	H	SO <sub>3</sub> -Na <sup>+</sup>	<i>Pisaster ochraceus</i> <sup>302</sup>
aphelasteroside A	425		SO <sub>3</sub> -Na <sup>+</sup>	H	H	H	X <sub>1</sub>	<i>Aphelasterias japonica</i> <sup>274</sup>
5-deoxyisonodososide	426		X <sub>2</sub>	H	H	H	A <sub>1</sub>	<i>Acanthaster planci</i> , <sup>303</sup> <i>Choriaster granulatus</i> <sup>285</sup>





Table 4 (Continued)


name	no.	side chain	R	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	sources and references
laeviuscoloside I forbeside I	427		X <sub>4</sub>	OH	H	H	H	<i>Henricia laeviuscola</i> , <sup>283</sup> <i>Asterias forbesi</i> <sup>304</sup>
granuloside A	428		X <sub>2</sub>	OH	H	H	A <sub>1</sub>	<i>Choriaster granulatus</i> , <sup>285</sup> <i>Thromidia catalai</i> <sup>305</sup>
laeviuscoloside G	429		X <sub>4</sub>	OH	H	H	A <sub>1</sub>	<i>Henricia laeviuscola</i> , <sup>283</sup> <i>Asterias forbesi</i> <sup>304</sup>
isonodososide	430		X <sub>2</sub>	H	OH	H	A <sub>1</sub>	<i>Acanthaster planci</i> <sup>303</sup>
nodososide	431		H	H	OH	H	XA <sub>5</sub>	<i>Protoreaster nodosus</i> , <sup>282</sup> <i>Pentaceraster alveolatus</i> , <sup>68</sup> <i>A. planci</i> , <i>Linkia Laevigata</i> <sup>306</sup>
echinasteroside B <sub>2</sub>	432		H	OH	H	H	XA <sub>5</sub>	<i>Echinaster sepositus</i> <sup>307</sup>
echinasteroside B <sub>1</sub>	433		H	OH	H	COCH <sub>3</sub>	XA <sub>5</sub>	<i>Eschinaster sepositus</i> <sup>307</sup>
scoparioside D	434		H	H	H	SO <sub>3</sub> -NA <sup>+</sup>	X <sub>1</sub>	<i>Astropecten scoparius</i> <sup>287</sup>
laeviuscoloside H	435		X <sub>4</sub>	OH	H	H	H	<i>Henricia laeviuscola</i> <sup>288</sup>
forbeside K	436		X <sub>2</sub>	OH	H	H	A <sub>1</sub>	<i>Asterias forbesi</i> <sup>304</sup>







Basic Nuclear Hydroxylation Pattern: 3 $\beta$ ,6 $\beta$ ,8,15 $\alpha$ ,16 $\beta$ 


name	no.	side chain	R	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	sources and references
laeviuscoloside B	437		X <sub>4</sub>	H	SO <sub>3</sub> -Na <sup>+</sup>		<i>Henricia laeviuscola</i> <sup>288</sup>
	438		X <sub>2</sub>	H	H		<i>Poraster superbus</i> <sup>251</sup>
thromidioside	439		X <sub>2</sub>	OH	H		<i>Thromidia catalai</i> <sup>305</sup>
laeviuscoloside C	440		X <sub>2</sub>	H	SO <sub>3</sub> -NA <sup>+</sup>		<i>Henricia laeviuscola</i> <sup>288</sup>
crossasteroside P <sub>1</sub>	441		H	H	H	XGalF	<i>Crossaster papposus</i> <sup>306</sup>
attenuatoside S-I	442		H	H	SO <sub>3</sub> -NA <sup>+</sup>	X <sub>1</sub>	<i>Hacelia attenuata</i> <sup>309</sup>

Table 4 (Continued)

name	no.	side chain	R	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	sources and references
laeviuscoloside E	443		X <sub>2</sub>	H	SO <sub>3</sub> -Na <sup>+</sup>	H	<i>Henricia laeviuscola</i> <sup>288</sup>
crossasteroside P <sub>2</sub>	444		H	OH	H	XGalF	<i>Crossaster papposus</i> <sup>308</sup>
attenuatoside S-II	445		H	H	SO <sub>3</sub> -Na <sup>+</sup>	X <sub>1</sub>	<i>Hacelia attenuata</i> <sup>309</sup>
attenuatoside S-III	446		H	H	SO <sub>3</sub> -Na <sup>+</sup>	X <sub>1</sub>	<i>Hacelia attenuata</i> <sup>309</sup>

Basic Nuclear Hydroxylation Patterns:  $\Delta^4$ -3 $\beta$ ,6 $\beta$ ,8,15 $\alpha$  and  $\Delta^4$ -3 $\beta$ ,6 $\beta$ ,8,15 $\alpha$ ,16 $\beta$ 

name	no.	side chain	R	R <sup>1</sup>	R <sup>2</sup>	sources and references
pisasteroside D	447		X <sub>1</sub>	H	H	<i>Pisaster giganteus</i> <sup>300</sup>
aphelasteroside B	448		X <sub>1</sub>	SO <sub>3</sub> -Na <sup>+</sup>	OH	<i>Aphelasterias japonica</i> <sup>274</sup>
echinasteroside A	449		X <sub>2</sub>	SO <sub>3</sub> -Na <sup>+</sup>	OH	<i>Echinaster sepositus</i> , <sup>310</sup> <i>Henricia laeviuscola</i> <sup>288</sup>
laeviuscoloside D	450		X <sub>2</sub>	SO <sub>3</sub> -Na <sup>+</sup>	OH	<i>Henricia laeviuscola</i> <sup>288</sup>
echinasteroside B	451		X <sub>2</sub>	SO <sub>3</sub> -Na <sup>+</sup>	OH	<i>Echinaster sepositus</i> , <sup>310</sup> <i>Henricia laeviuscola</i> <sup>288</sup>
forbeside L	452		X <sub>2</sub>	H	OH	<i>Asterias forbesi</i> <sup>304</sup>

Basic Nuclear Hydroxylation Pattern: 3 $\beta$ ,6 $\beta$ ,8,15 $\beta$ 


name	no.	side chain	R	R <sup>1</sup>	R <sup>2</sup>	sources and references
halityloside F	453		H	H	XA <sub>6</sub>	<i>Halityle regularis</i> , <sup>284</sup> <i>Gomophia watsoni</i> , <sup>276</sup> <i>Nardoia gomophia</i> , <sup>285</sup> <i>Calcita novaeguineae</i> <sup>289</sup>

Table 4 (Continued)

name	no.	side chain	R	R <sup>1</sup>	R <sup>2</sup>	sources and references
gomophioside A	454		H	OH	XA <sub>6</sub>	<i>Gomphia watsoni</i> , <sup>276</sup> <i>C. novaeguineae</i> <sup>268</sup>
moniloside D	455		X <sub>2</sub>	OH	H	<i>Fromia monilis</i> <sup>281</sup>
laeviuscoloside A	456					<i>Henricia laeviuscola</i> <sup>283</sup>
moniloside A	457		R = H			<i>Fromia monilis</i> <sup>281</sup>
moniloside B	458		R = OH			<i>Fromia monilis</i> <sup>281</sup>
tremasterol A	459		H	H		<i>Tremaster novaecaledoniae</i> <sup>239</sup>
tremasterol B	460		COCH <sub>3</sub>	COCH <sub>3</sub>		
tremasterol C	461		H	COCH <sub>3</sub>		

<sup>a</sup> Abbreviations for R groups are defined in Figure 18.

Further recent additions are the phosphated steroids tremasterols A-C (459-461), isolated from the deep-water species *Tremaster novaecaledoniae*.

### 3. Asterosaponins

The term asterosaponin was originally coined to generally designate toxic steroidal saponins occurring in starfish, by analogy with the term holothurin used for triterpenoid saponins of holothuroids. Following the elucidation of first structures and the subsequent discovery of different structural type of steroidal glycosides from starfish, the term has been used to indicate those highest molecular weight compounds unified by a number of structural analogies involving both the aglycons and the oligosaccharide chains. The aglycons are  $\Delta^{9(11)}$ - $3\beta,6\alpha$ -dihydroxysteroids bearing a sulfate group at C-3 and often a 23-oxo function. The oligosaccharide chains, commonly made up by five or six sugar units, are always glycosidically linked at C-6. A close resemblance is also evident in this saccharide portions: sugars are in their pyranose form with  $\beta$ -anomeric configurations ( $\alpha$  for L-arabinose) and linked with a constant pattern of interglycosidic linkages. A

branching point is always located on the second monosaccharide (usually xylose or quinovose) starting from the aglycon and a terminal quinovose is always found 2-linked to the branched sugar. The more common sugars are D-fucose, D-quinovose, D-xylose, D-galactose, and D-glucose. Other less common monosaccharides are D-6-deoxy-xylo-hex-4-ulose and L-arabinose. The single exception to the general pattern of interglycosidic linkages is represented by the recently isolated santiagoside (488),<sup>311</sup> with a 4-substituted glucose unit linked to the aglycon, instead of the 3-substituted unit present in all asterosaponins. We would note that the <sup>13</sup>C NMR data of santiagoside (488) are very close to those assigned to related asterosaponins [e.g. marthasteroide C (489)], including the unusually downfield shifted signal at 91.0 ppm, which is a distinctive feature of <sup>13</sup>C NMR spectra of all asterosaponins and has been assigned to the glycosylated carbon-3 of the monosaccharide unit ( $\beta$ -glucopyranosyl or  $\beta$ -quinovopyranosyl) directly attached to the aglycon.

Also asterosaponins are widespread among starfish, having been found in the majority of species examined. The toxic properties of starfish have been known for many years, but it was only in 1960 that Hashimoto





Figure 19. Steroidal aglycons found in asterosaponins from starfish.

and Yasumoto recognized that the toxicity is associated with compounds similar to plant saponins.<sup>312</sup> Astero-saponins also possess ichthyotoxicity,<sup>312,313</sup> and toxicity has also been noted toward annelids, mollusks, arthropods and vertebrates.<sup>314</sup> Because of their general toxicity, it is probable that saponins act primarily as chemical defense agents, rejecting infectious aquatic fungi, protists, parasites, and predators. Makie *et al.* found that asterosaponins are the substances eliciting the escape response in some species of molluscs when in presence of starfishes.<sup>315</sup> In addition, these oligoglycosides, in some species of starfish, participate in reproduction processes. Ikegami *et al.* identified saponins as the spawning inhibitors in the ovaries of *Asterias amurensis*.<sup>316</sup> Fujimoto *et al.* observed that three steroidal saponins, designated Co-ARIS I, II, and III, isolated from the egg jelly of the starfish *Asterias amurensis*, are essential for inducing the acrosome reaction.<sup>317</sup> Starfish extracts and purified saponin fractions have shown a variety of pharmacological activities: hemolytic activity,<sup>314,318</sup> *in vitro* cytotoxicity toward tumor cells,<sup>319-321</sup> antiviral activity,<sup>322</sup> blockage of neuromuscular transmission in mammals,<sup>323</sup> and antiinflammatory, analgesic, and hypotensive activities.<sup>324</sup> Starfish saponins are also known to inhibit development of fertilized sea urchin eggs.<sup>318,325</sup> A more recent study of the biological activity of representative saponins and related steroids from starfish confirmed a high incidence of cytotoxicity and inhibition of Gram-positive bacteria but only weak antiviral activity and no inhibition of Gram-negative bacteria.<sup>260</sup>

Astero-saponins, as do all these polyhydroxysteroids and glycosides from starfish, also occur as complex mixtures that are fairly difficult to separate in the individual components. This has probably been the main reason for which most of the initial papers were

concerned with the analysis of aglycons obtained by acid hydrolysis of partially purified saponin mixtures. This procedure resulted in the production of several artifacts, such as asterone (462),<sup>326,327</sup> an artifact obtained by retro-aldol cleavage of the genuine thornasterol A sulfate (463),<sup>328</sup> which has been the most widely reported steroid obtained by acid hydrolysis of astero-saponins.  $\Delta^{20(22)}$ - and  $\Delta^{17(20)}$ -steroids, along with a rearranged  $\Delta^{13(14)}$ -17-methyl-18-norsteroid, certainly artifacts generated during acid hydrolysis, have also been reported.<sup>329,330</sup> An excellent review of previous work on the aglycons can be found in the work of Burnell and ApSimon.<sup>232</sup>



The first complete structure of an astero-saponin, thornasteroside A (464), was reported in 1978 by

Table 5. Asterosaponins from Starfish\*

Name	#	oligosaccharide chain	sources and references
Asterosaponins containing thomasterol A, aglycone A			
thomasteroside A	464	$\begin{array}{ccccccc} \text{Fuc} & \xrightarrow{1-2} & \text{Gal} & \xrightarrow{1-4} & \text{Xyl} & \xrightarrow{1-3} & \text{Qui} \xrightarrow{1-6} \text{A} \\ & & & & \uparrow & & \\ & & & & \text{Qui} & & \end{array}$	<i>Acanthaster planci</i> <sup>331</sup> , <i>Asterias amurensis</i> <sup>332</sup> , <i>Luidia maculata</i> <sup>333</sup> , <i>Ophidiaster ophidianus</i> <sup>334</sup> , <i>Linkia laevigata</i> <sup>335</sup> , <i>Protoreaster nodosus</i> and <i>Pentacaster alveolatus</i> <sup>68</sup> , <i>Halityle regularis</i> <sup>336</sup> , <i>Nordoa gomphia</i> <sup>265</sup> , <i>Coscinasterias tenuispina</i> <sup>268</sup> , <i>Thromidia catalai</i> <sup>305</sup> , <i>Pisaster ochraceus</i> and <i>Pisaster brevispinus</i> <sup>303</sup> , <i>Pisaster giganteus</i> <sup>300</sup> , <i>Pycnopodia helianthoides</i> <sup>258</sup>
glycoside B <sub>2</sub> = forbeside B	465	$\begin{array}{ccccccc} \text{Qui} & \xrightarrow{1-2} & \text{Gal} & \xrightarrow{1-4} & \text{Xyl} & \xrightarrow{1-3} & \text{Qui} \xrightarrow{1-6} \text{A} \\ & & & & \uparrow & & \\ & & & & \text{Qui} & & \end{array}$	<i>Asterias amurensis</i> <sup>284, 337</sup> , <i>Asterias forbesi</i> <sup>338</sup>
acanthaglycoside C	466	$\begin{array}{ccccccc} \text{Fuc} & \xrightarrow{1-2} & \text{Glc} & \xrightarrow{1-4} & \text{Xyl} & \xrightarrow{1-3} & \text{Qui} \xrightarrow{1-6} \text{A} \\ & & & & \uparrow & & \\ & & & & \text{Qui} & & \end{array}$	<i>Astropecten latespinosus</i> <sup>339</sup> , <i>Acanthaster planci</i> <sup>340</sup> , <i>Asterina pectinifera</i> <sup>341</sup> , <i>Patiria miniata</i> <sup>342</sup>
maculoside = luidiaglycoside B	467	$\begin{array}{ccccccc} \text{Fuc} & \xrightarrow{1-2} & \text{Qui} & \xrightarrow{1-4} & \text{Xyl} & \xrightarrow{1-3} & \text{Qui} \xrightarrow{1-6} \text{A} \\ & & & & \uparrow & & \\ & & & & \text{Qui} & & \end{array}$	<i>Luidia maculata</i> <sup>333</sup> , <i>Linkia laevigata</i> <sup>335</sup>
ophidianoside F	468	$\begin{array}{ccccccc} \text{Fuc} & \xrightarrow{1-2} & \text{Xyl} & \xrightarrow{1-4} & \text{Xyl} & \xrightarrow{1-3} & \text{Qui} \xrightarrow{1-6} \text{A} \\ & & & & \uparrow & & \\ & & & & \text{Qui} & & \end{array}$	<i>Ophidiaster ophidianus</i> <sup>334</sup> , <i>Linkia laevigata</i> <sup>335</sup> , <i>Thromidia catalai</i> <sup>305</sup>
regularoside B	469	$\begin{array}{ccccccc} \text{Fuc} & \xrightarrow{1-2} & \text{Fuc} & \xrightarrow{1-4} & \text{Xyl} & \xrightarrow{1-3} & \text{Qui} \xrightarrow{1-6} \text{A} \\ & & & & \uparrow & & \\ & & & & \text{Qui} & & \end{array}$	<i>Halityle regularis</i> <sup>336</sup> , <i>Coscinasterias tenuispina</i> <sup>268</sup> , <i>Thromidia catalai</i> <sup>305</sup>
laevigatoside	470	$\begin{array}{ccccccc} \text{Fuc} & \xrightarrow{1-2} & \text{Ara} & \xrightarrow{1-4} & \text{Qui} & \xrightarrow{1-3} & \text{Qui} \xrightarrow{1-6} \text{A} \\ & & & & \uparrow & & \\ & & & & \text{Qui} & & \end{array}$	<i>Linkia laevigata</i> <sup>335</sup>
pectinoside A	471	$\begin{array}{ccccccc} \text{Fuc} & \xrightarrow{1-2} & \text{Glc} & \xrightarrow{1-4} & \text{Qui} & \xrightarrow{1-3} & \text{Qui} \xrightarrow{1-6} \text{A} \\ & & & & \uparrow & & \\ & & & & \text{Qui} & & \end{array}$	<i>Asterina pectinifera</i> <sup>343</sup>
ovarian asterosaponin I = Co-Aris I = forbeside C	472	$\begin{array}{ccccccc} & & & & \begin{array}{c} \text{HO} \quad \text{CH}_3 \\   \quad   \\ \text{O} \quad \text{O} \quad \text{O} \quad \text{O} \\   \quad   \quad   \quad   \\ \text{HO} \quad \text{OH} \quad \text{OH} \quad \text{O} \end{array} & \xrightarrow{1-6} & \text{A} \\ \text{Fuc} & \xrightarrow{1-2} & \text{Fuc} & \xrightarrow{1-4} & \text{Qui} & & \\ & & & & \uparrow & & \\ & & & & \text{Qui} & & \end{array}$	<i>Asterias amurensis</i> <sup>317, 284, 344, 345</sup> , <i>Asterias forbesi</i> <sup>346</sup> , <i>Asterias vulgaris</i> <sup>346</sup>
marthasteroside A <sub>2</sub> = luidiaglycoside A	473	$\begin{array}{ccccccc} \text{Fuc} & \xrightarrow{1-3} & \text{Fuc} & \xrightarrow{1-2} & \text{Qui} & \xrightarrow{1-4} & \text{Xyl} & \xrightarrow{1-3} & \text{Qui} & \xrightarrow{1-6} & \text{A} \\ & & & & & & \uparrow & & & & \\ & & & & & & \text{Qui} & & & & \end{array}$	<i>Luidia maculata</i> <sup>333, 347</sup> , <i>Marthasterias glacialis</i> <sup>348, 349</sup>



Table 5 (Continued)




Name	#	oligosaccharide chain	sources and references
marthasteroside A <sub>1</sub>	474	$\begin{array}{c} \text{Fuc} \xrightarrow{1-3} \text{Fuc} \xrightarrow{1-2} \text{Gal} \xrightarrow{1-4} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \mathbf{A} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Marthasterias glacialis</i> <sup>348, 349</sup> , <i>Astropecten latespinosus</i> <sup>339</sup> , <i>Linkia laevigata</i> <sup>335</sup> , <i>Nardoa gomophia</i> <sup>265</sup> , <i>Pisaster brevispinus</i> <sup>303</sup> , <i>Achantaster planci</i> <sup>340</sup> , <i>Astropecten scoparius</i> <sup>267</sup>
versicoside A = forbeside A	475	$\begin{array}{c} \text{Gal} \xrightarrow{1-3} \text{Fuc} \xrightarrow{1-2} \text{Gal} \xrightarrow{1-4} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \mathbf{A} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Asterias amurensis versicolor</i> <sup>332</sup> , <i>Pisaster ochraceus</i> and <i>Pisaster brevispinus</i> <sup>303</sup> , <i>Pisaster giganteus</i> <sup>300</sup> , <i>Asterias forbesi</i> <sup>338</sup>
pectinoside F	476	$\begin{array}{c} \text{Gal} \xrightarrow{1-4} \text{Glc} \xrightarrow{1-4} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \mathbf{A} \\ \uparrow 1-2 \quad \uparrow 1-2 \\ \text{Fuc} \quad \text{Qui} \end{array}$	<i>Asterias pectinifera</i> <sup>321</sup>
pectinoside E	477	$\begin{array}{c} \text{Fuc} \xrightarrow{1-4} \text{Glc} \xrightarrow{1-4} \text{Qui} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \mathbf{A} \\ \uparrow 1-2 \quad \uparrow 1-2 \\ \text{Xyl} \quad \text{Qui} \end{array}$	>>
pectinoside G	478	$\begin{array}{c} \text{Ara} \xrightarrow{1-4} \text{Glc} \xrightarrow{1-4} \text{Qui} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \mathbf{A} \\ \uparrow 1-2 \quad \uparrow 1-2 \\ \text{Fuc} \quad \text{Qui} \end{array}$	<i>Asterina pectinifera</i> <sup>254</sup> , <i>Patiria miniata</i> <sup>342</sup>
forbeside H	479	$\text{Qui} \xrightarrow{1-2} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \mathbf{A}$	<i>Asterias forbesi</i> <sup>350</sup>
forbeside G	480		>>
forbeside F	481		>>
myxodermoside A	482	$\begin{array}{c} \text{Gal} \xrightarrow{1-4} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \mathbf{A} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Myxoderma platyacanthum</i> <sup>245</sup>
Asterosaponins containing 24,25-dehydro-thomasterol A, aglycone B			
achantaglycoside A	483	$\begin{array}{c} \text{Fuc} \xrightarrow{1-2} \text{Qui} \xrightarrow{1-4} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \mathbf{B} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Achantaster planci</i> <sup>351</sup>
achantaglycoside B	484	$\begin{array}{c} \text{Fuc} \xrightarrow{1-2} \text{Glc} \xrightarrow{1-4} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \mathbf{B} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Achantaster planci</i> <sup>340</sup>
achantaglycoside D	485	$\begin{array}{c} \text{Fuc} \xrightarrow{1-2} \text{Gal} \xrightarrow{1-4} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \mathbf{B} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Achantaster planci</i> <sup>340</sup>


Table 5 (Continued)

Name	#	oligosaccharide chain	sources and references
Asterosaponins containing marthasterone, aglycone C			
marthasteroside B	486	$\begin{array}{c} \text{Fuc} \xrightarrow{1-2} \text{Fuc} \xrightarrow{1-4} \text{Qui} \xrightarrow{1-3} \text{Glc} \xrightarrow{1-6} \text{C} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Marthasterias glacialis</i> <sup>348, 349</sup> , <i>Luidia maculata</i> <sup>333</sup> , <i>Coscinasterias tenuispina</i> <sup>268</sup>
luidiaglycoside C	487	$\begin{array}{c} \text{Fuc} \xrightarrow{1-2} \text{Qui} \xrightarrow{1-4} \text{Qui} \xrightarrow{1-3} \text{Glc} \xrightarrow{1-6} \text{C} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Luidia maculata</i> <sup>352</sup>
Asterosaponins containing 24,25-dihydromarthasterone, aglycone D			
santiagoside	488	$\begin{array}{c} \text{Fuc} \xrightarrow{1-4} \text{Qui} \xrightarrow{1-4} \text{Glc} \xrightarrow{1-6} \text{D} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Neosmilaster georgianus</i> <sup>311</sup>
marthasteroside C	489	$\begin{array}{c} \text{Fuc} \xrightarrow{1-2} \text{Fuc} \xrightarrow{1-4} \text{Qui} \xrightarrow{1-3} \text{Glc} \xrightarrow{1-6} \text{D} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Marthasterias glacialis</i> <sup>348, 349</sup> , <i>Luidia maculata</i> <sup>333</sup> , <i>Coscinasterias tenuispina</i> <sup>268</sup>
luidiaglycoside D	490	$\begin{array}{c} \text{Fuc} \xrightarrow{1-2} \text{Qui} \xrightarrow{1-4} \text{Qui} \xrightarrow{1-3} \text{Glc} \xrightarrow{1-6} \text{D} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Luidia maculata</i> <sup>352</sup>
Asterosaponins containing 24-nor-thomasterol A, aglycone E			
ophidianoside B	491	$\begin{array}{c} \text{Fuc} \xrightarrow{1-2} \text{Gal} \xrightarrow{1-4} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \text{E} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Ophidiaster ophidianus</i> and <i>Hacelia attenuata</i> <sup>334</sup>
ophidianoside C	492	$\begin{array}{c} \text{Fuc} \xrightarrow{1-2} \text{Xyl} \xrightarrow{1-4} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \text{E} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	>>
asterosaponins containing 22,23-epoxy-aglycone F			
tenuispinoside A	493	$\begin{array}{c} \text{Fuc} \xrightarrow{1-2} \text{Gal} \xrightarrow{1-4} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \text{F} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Coscinasterias tenuispina</i> <sup>268</sup>
tenuispinoside B	494	$\begin{array}{c} \text{Fuc} \xrightarrow{1-2} \text{Fuc} \xrightarrow{1-4} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \text{F} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	>>
asteroside A	495	$\begin{array}{c} \text{Qui} \xrightarrow{1-2} \text{Gal} \xrightarrow{1-4} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \text{F} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Asterias amurensis</i> <sup>284</sup>

Table 5 (Continued)

Name	#	oligosaccharide chain	sources and references
asterosaponin containing 22,23-epoxy-24-nor-aglycone <b>G</b>			
asteroside B	496	$\begin{array}{c} \text{Qui} \xrightarrow{1-2} \text{Gal} \xrightarrow{1-4} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \mathbf{G} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Asterias amurensis</i> <sup>284</sup>
asterosaponins containing the less common aglycones <b>H, I, J, K</b> and <b>L</b>			
protoreasteroside	497	$\begin{array}{c} \text{Fuc} \xrightarrow{1-2} \text{Qui} \xrightarrow{1-4} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \mathbf{H} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Protoreaster nodosus</i> and <i>Pentacaster alveolatus</i> <sup>68</sup>
ovarian asterosaponin-4 = Co-Aris III	498	$\begin{array}{c} \text{Qui} \xrightarrow{1-2} \text{Glc} \xrightarrow{1-4} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \mathbf{I} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Asterias amurensis</i> <sup>317, 284, 353</sup>
solasteroside A	499	$\begin{array}{c} \text{Fuc} \xrightarrow{1-2} \text{Fuc} \xrightarrow{1-4} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \mathbf{I} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Solaster borealis</i> <sup>244</sup>
Co-Aris II	500	$\begin{array}{c} \text{Fuc} \xrightarrow{1-2} \text{Fuc} \xrightarrow{1-4} \text{Qui} \xrightarrow{1-6} \mathbf{J} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$ 	<i>Asterias amurensis</i> <sup>317</sup>
tenuispinoside C	501	$\begin{array}{c} \text{Fuc} \xrightarrow{1-2} \text{Gal} \xrightarrow{1-4} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \mathbf{K} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Coscinasterias tenuispina</i> <sup>268</sup>
asteroside D	502	$\begin{array}{c} \text{Qui} \xrightarrow{1-2} \text{Glc} \xrightarrow{1-4} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \mathbf{L} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Asterias amurensis</i> <sup>284</sup>
asterosaponins containing (24R)-24-methylthomasterol A, aglycone <b>M</b>			
versicoside C = thornasteroside B	503	$\begin{array}{c} \text{Fuc} \xrightarrow{1-2} \text{Gal} \xrightarrow{1-4} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \mathbf{M} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Asterias amurensis</i> [cf.] <i>versicolor</i> <sup>345</sup> , <i>Coscinasterias tenuispina</i> <sup>268</sup>
asteroside C	504	$\begin{array}{c} \text{Qui} \xrightarrow{1-2} \text{Gal} \xrightarrow{1-4} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \mathbf{M} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Asterias amurensis</i> <sup>284</sup>
versicoside B	505	$\begin{array}{c} \text{Gal} \xrightarrow{1-3} \text{Fuc} \xrightarrow{1-2} \text{Gal} \xrightarrow{1-4} \text{Xyl} \xrightarrow{1-3} \text{Qui} \xrightarrow{1-6} \mathbf{M} \\ \uparrow 1-2 \\ \text{Qui} \end{array}$	<i>Asterias amurensis</i> [cf.] <i>versicolor</i> <sup>345</sup>





#### 4. Cyclic Steroidal Glycosides

Toxic saponins of a completely different structural type have been discovered in two species of the genus *Echinaster* (Figure 20). They have a number of unusual features when compared to the more common asterosaponins: there is no sulfate group and charge is due to a glucuronic acid unit in the saccharide moiety, the  $\Delta^7$ - $3\beta,6\beta$ -dihydroxysteroidal nucleus is unprecedented, and, the most remarkable feature, the trisaccharide chain is cyclized between C-3 and C-6 of the aglycon, giving rise to a macrocyclic ring reminiscent of a crown ether. Sepositoside A (513), the major saponin from the Mediterranean starfish *Echinaster sepositus*,<sup>357</sup> is accompanied by smaller amounts of three related saponins 514–516, which differ only in the structure of the side chain of the aglycons, all having a 22,23-epoxy functionality.<sup>358</sup> The key step during structural study of sepositoside A (513) was the mild acid hydrolysis, which, by cleavage of the allylic ether linkage, gave rise to the opening of the macrocyclic ring

and formation of the UV-active glycoside, 517. A further representative of this class of glycosides was isolated from a Pacific starfish belonging to the same genus, *E. luzonicus*, and accordingly named luzonicoside A (518).<sup>359</sup> Sepositoside A (513) is moderately toxic ( $LD_{50} = 43 \text{ mg kg}^{-1}$  by ip injection in mouse) and showed cytotoxic activity toward bovine turbinate cells up to a level of  $1 \mu\text{g mL}^{-1}$ .<sup>260</sup> Both sepositoside A (513) and luzonicoside A (518) were slightly effective in the inhibition of cell division of fertilized sea urchin eggs (ca. 30% inhibition at  $10^{-5} \text{ M}$ ) and showed antifungal activity.

#### B. Ophiuroidea (Brittle Stars)

Until a few years ago brittle stars (ophiuroids) had only received moderate attention as compared to the echinoderm classes of asteroids and holothuroids and only sporadic papers dealing with their sterols content had appeared in the literature. Quite recently we had the occasion to investigate a number of ophiuroid species from which we could isolate several sulfated polyhydroxysteroids and two steroidal glycosides (Figure 21).

The steroid glycosides, longicaudoside A (519) and B (520) have been isolated from the Mediterranean *Ophioderma longicaudum*.<sup>360</sup> Both have a  $5\alpha$ -cholestane- $3\alpha,6\beta,12\beta,21$ -tetrol aglycon bearing a sulfate group at C-3 and a monosaccharide residue at C-12 ( $\beta$ -D-xylopranosyl in 519 and  $\beta$ -D-glucopyranosyl in 520). In contrast with the hydroxylation at C-26, commonly encountered among polyhydroxysteroids from starfish, the polar steroids from ophiuroids are characterized by the hydroxylation at C-21, only found among





Figure 20. Cyclic steroidal glycosides from starfish of the genus *Echinaster*.



**Figure 21.** Polyhydroxysteroids from ophiuroids.

starfish metabolites, in steroids from *Euretaster insignis*.<sup>241</sup> <sup>13</sup>C NMR spectroscopy proved to be the better tool for differentiation between the two hydroxylated side chains. Noticeable differences are observed in the methyl and hydroxymethyl resonances: the C-21 hydroxylated side chain exhibits the typical signals for C-26 and C-27 at 23.0 and 23.1 ppm and the CH<sub>2</sub>OH signal at 63.4 ppm, while the C-26 hydroxylated side

chain exhibits methyl carbon signals at 17.3 and 19.2 ppm and the CH<sub>2</sub>OH signal at 68.4 ppm. The narrow signal at a rather low field ( $\delta$  4.72 ppm,  $W_{1/2}$  = 7 Hz) in the <sup>1</sup>H NMR spectra of both 519 and 520, shifted to  $\delta$  4.10 in the spectra of their desulfated derivatives, was indicative of a 3 $\alpha$ -sulfato-5 $\alpha$ -stanol structure. The alternative 3 $\beta$ -sulfato-5 $\beta$ -stanol structure could be eliminated mainly because of the chemical shift of C-19

at 15.3 ppm. *Ophioderma longicaudum* also furnished unusual sulfated polyhydroxysterols together with a mixture of common  $\Delta^5$ - $3\beta$ -hydroxysterols.<sup>361</sup> The more polar sulfated sterol was characterized as  $5\beta$ -cholestane- $3\alpha,4\alpha,11\beta,12\beta,21$ -pentol 3,21-disulfate (521), featuring the *cis*-A/B ring fusion, later also found in the polar sterols from the further species analyzed.<sup>96,362</sup> A second group of components consisted in a mixture of disulfated  $3\alpha,21$ -dihydroxysteroids 522–525, whose composition was established after solvolytic removal of the sulfate groups.<sup>361</sup> Additional sulfated polyhydroxysteroids have been then isolated recently from three further species of ophiuroids, *Ophiocoma dentata*, *Ophiarthrum elegans*, and *Ophiorachna incrassata*, all collected off Noumeà (New Caledonia).<sup>96</sup> The major compound in all three species has been shown to be  $5\beta$ -cholestane- $3\alpha,4\alpha,11\beta,21$ -tetrol 3,21-disulfate (526). Two minor components of *Ophiocoma dentata* possess the same nucleus as 526 but differ in the side chain, 527 and 528. Along with minor amounts of 528 and major amounts of the disulfated tetraol 526, *Ophiarthrum elegans* also contains the 11-keto derivative 529. *Ophiorachna incrassata* also contains major amounts of the steroid disulfate 526 along with minor amounts of the two more polar steroids 530 and 531, bearing three sulfate groups, which represent the first occurrence of 26-hydroxylation in ophiuroids. Major support for the presence of both 25*S*- and 25*R*-epimers is found in the <sup>13</sup>C NMR spectra, where every side-chain carbon signal appears split into two peaks separated by 0.04–0.2 ppm. The pentol disulfate 521, the tetrol disulfate 526, and the  $\Delta^5$ -steroidal sulfates 523 and 525 have moderate cytotoxic activity.<sup>260</sup>

The investigation of the Pacific ophiuroid *Ophiolepis superba*, collected at Okinawa, Japan, has led to the isolation of seven sulfated polyhydroxysteroids (532–538), all with  $3\alpha,21$ -disulfoxy- $4\alpha$ -hydroxy substituents and the A/B *cis* ring junction.<sup>362</sup> The stereochemistry at C-24 and C-25 in 533 and 535 have been determined after the synthesis of model compounds.<sup>281,363</sup> The

$3\alpha,4\alpha,5\beta$ -trihydroxycholestane structure of 538 was derived from analysis of <sup>1</sup>H and <sup>13</sup>C NMR data and confirmed by the synthesis of  $5\beta$ -cholestane- $3\alpha,4\alpha,5$ -triol and the alternative  $5\alpha$ -cholestane- $3\beta,4\beta,5$ -triol models. A comparison of the <sup>1</sup>H and <sup>13</sup>C NMR spectra of 538 with those of its desulfated derivative allowed the placement of the sulfoxy group at C-3. The introduction of an additional hydroxyl group at C-2 $\beta$  in the  $3\alpha$ -sulfoxy- $4\alpha$ -hydroxy- $5\beta$ -steroidal nucleus, as in structures 536 and 537, leads to the appearance in the <sup>1</sup>H NMR spectrum of a signal at  $\delta$  4.03 overlapping with the 3-H signal. When the spectrum of the desulfated 536 was measured, the three hydroxymethine protons appeared as isolated signals at  $\delta$  3.81 (m), 3.23 (dd,  $J = 9.5, 3.5$  Hz) and 3.94 (t,  $J = 3.5$  Hz) ppm. Decoupling experiments proved that they were located in a sequential arrangement and allowed the inference of a  $2\beta,3\alpha,4\alpha$ -trihydroxy- $5\beta$ -steroidal structure. The 24*S* configuration was assigned to 537 after the stereoselective synthesis of (24*R*,22*E*)- and (24*S*,22*E*)-24-(hydroxymethyl)cholesta-5,22-dien-3 $\beta$ -ol.<sup>364</sup>

Stonik and Elyakov reported to have isolated cholest-5-ene- $3\alpha,4\beta,21$ -triol 3-sulfate (539) from *Ophiura sarsi*.<sup>365</sup>

More recent additions to the sulfated polyhydroxysteroids from ophiuroids are a new  $5\beta$ -steroid,  $3\beta,21$ -disulfate 540 with a unique  $4\alpha,9\alpha$ -ether bridge, and a minor related steroid 541 from *Ophiomastix annulosa*.<sup>366</sup> *O. annulosa* also contained the more common tetrol 526 and its keto derivative 529.

### VIII. Polyoxygenated Steroids of Arthropoda

In arthropods, the molting process is under the control of a group of polyhydroxysteroid hormones referred as ecdysones. Although much of the interest of ecdysone biochemistry has centered on Insecta, the role of these sterols in crustaceans has also received considerable attention. The structures of ecdysones reported in crustaceans are shown in Figure 22. In 1960, the




Figure 22. Structures of ecdysones found in Crustacea.













Figure 23. Polyhydroxysteroids from Tunicata.



presence of a substance which showed ecdysone (542) activity was detected in crabs by Karlson and Skinner.<sup>367</sup> In 1966, crustecdysone (543) was first isolated from the crayfish, *Jasus lalandei*, by Horn *et al.*<sup>368</sup> and subsequently 2-deoxycrustecdysone (544)<sup>369</sup> was isolated. The 20-hydroxyecdysteroids, callinecdysone A (inokosterone) (545) and callinecdysone B (makisterone A) (546) were isolated from *Calinectes sapidus*.<sup>370</sup> Ecdysteroids metabolism in crustaceans has been reviewed by Goad<sup>2,229</sup> and Ikekawa.<sup>60</sup>

### IX. Polyoxygenated Steroids of Tunicates

Several species of ascidians have been analyzed for their sterol content and were shown to contain typical mixtures of sterols with cholesterol as the major component.<sup>2</sup> Coprostanols along with  $\Delta^4$ -3-ketosteroids and 4 $\alpha$ -methylsterols have been described from *Ascidia nigra*.<sup>371</sup> 5 $\beta$ -Stanols have later been found in *Halicynthia papillosa*, *Microcosmus sulcatus*, *Microcosmus savignyi*, and *Styela plicata*.<sup>372</sup> The tunicates *Phallusia*

Figure 24. Polyoxygenated steroids from fish.



*mamillata* and *Ciona intestinalis* were reported to contain 5,8-endoperoxides of several  $\Delta^{5,7,9(11)}$ -sterols (547)<sup>373</sup> and the 24-hydroperoxy-24-vinylcholesterol (548) along with its corresponding 24-hydroxy derivative,<sup>374</sup> this latter previously found in an alga (Figure 23).

### X. Polyoxygenated Steroids from Vertebrates (Fish)

The bile of vertebrates characteristically contains bile salts, and in most animals these are largely steroidal C<sub>24</sub> carboxylic acid derivatives. In the case of some fish the bile salts include C<sub>27</sub> alcohols present as their sodium sulfate esters.

Scymnol (549, Figure 24) is the bile alcohol from shark bile and the first of such alcohols to be isolated.<sup>375</sup> The history of the structure determination of this alcohol has been reviewed by Scheuer.<sup>376</sup> The final structure 549 was deduced by two groups, Cross<sup>377</sup> and Briggs and Haslewood.<sup>378</sup> The two bile alcohols myxinol (550) and deoxymyxinol (551) were both isolated from hagfish. For myxinol, the first of the two to be isolated,<sup>379</sup> the 5 $\beta$ -structure was initially proposed, but later work established the 5 $\alpha$ -structure 550.<sup>380</sup> Deoxymyxinol was then assigned the structure 551 by the same research group.<sup>381</sup>

Certain fish have a self-defense mechanism consisting of the secretion of toxic substances that repel their predators. Among them are the Red Sea Moses sole *Pardachirus marmoratus* and its congener, the peacock sole *Pardachirus pavoninus*, which repel sharks by emission of their toxic secretion at the moment when they are about to be bitten. The chemical nature of the shark repellents has been shown by Tachibana, Nakanishi, and co-workers to consist of mixture of the peptidic paradaxins and two group of steroid glycosides, pavonins 1-6 (552-557) from *P. pavoninus*,<sup>382,383</sup> and mosesins 1-5 (558-562) from *P. marmoratus*.<sup>384</sup> The chemical structure of both pavonins and mosesins has been established by spectroscopic studies and chemical conversions. Mosesin 4 (560) has been successfully obtained by synthesis.<sup>385</sup> The biological activity of these toxins is believed to be related to their surfactant properties, that in the case of glycosides arises from a hydrophobic "top" and a hydrophilic "bottom" region. The respective roles of the steroidal glycosides and of the peptidic paradaxins in defense mechanism still remain obscure. It has been suggested that pavonins and mosesins are repellents that act on the shark's olfactory sense, whereas the proteinaceous toxin is possibly an antifeedant that acts on its gustatory sense. A review on chemical defense in fishes, which includes the fascinating history of the shark repellents from *Pardachirus* sp. has been published by Tachibana.<sup>386</sup>

### References

- Itoh, T.; Sica, D.; Djerassi, C. *J. Chem. Soc. Perkin Trans. 1* 1983, 147.
- Goad, L. J. The sterols of marine invertebrates: Composition, Biosynthesis, and Metabolites. In *Marine Natural Products: Chemical and Biological Perspectives*; Scheuer, P. J., Ed.; Academic Press: New York, 1978; Vol II, pp 75-172.
- Lederer, E. *Quart. Rev.* 1969, 23, 453.
- Idler, D. R.; Wiseman, P. M.; Safe, L. M. *Steroids* 1970, 16, 451.
- Ferezon, J. P.; Devys, M.; Allois, J. P.; Barbier, M. *Phytochemistry* 1974, 13, 593.
- De Luca, P.; De Rosa, M.; Minale, L.; Sodano, G. *J. Chem. Soc., Perkin Trans. 1* 1972, 2132.
- Fattorusso, E.; Magno, S.; Mayol, L.; Santacroce, C.; Sica, D. *Tetrahedron* 1975, 31, 1715.
- Sica, D.; Zollo, F. *Tetrahedron Lett.* 1978, 837.
- Ravi, B. N.; Kokke, W. C. M. C.; Delseth, C.; Djerassi, C. *Tetrahedron Lett.* 1978, 4379.
- Bortolotto, M.; Braekman, J. C.; Daloze, D.; Tursch, B. *Bull. Soc. Chem. Belg.* 1978, 87, 539.
- Kokke, W. C. M. C.; Tarchini, C.; Stierle, D. B.; Djerassi, C. *J. Org. Chem.* 1979, 44, 3385.
- Kerr, R. G.; Kerr, S. L.; Pettit, G. R.; Harold, D. L.; Groy, T. L.; Djerassi, C. *J. Org. Chem.* 1991, 56, 58.
- Minale, L.; Sodano, G. *J. Chem. Soc. Perkin Trans. 1* 1974, 1888.
- Minale, L.; Sodano, G. *J. Chem. Soc. Perkin Trans. 1* 1974, 2380.
- Shimizu, Y.; Alam, M.; Kobayashi, A. *J. Am. Chem. Soc.* 1976, 98, 1059.
- Kokke, W. C. M. C.; Fenical, W.; Bohlin, L.; Djerassi, C. *Comp. Biochem. Physiol.* 1981, 68B, 281.
- Delseth, C.; Carlson, R. M. K.; Djerassi, C.; Erdman, T. R.; Scheuer, P. J. *Helv. Chim. Acta* 1978, 61, 1470.
- Idler, D. R.; Khalil, M. W.; Gilbert, J. D.; Brooks, C. J. W. *Steroids* 1976, 27, 155.
- Kobayashi, M.; Mitsuhashi, H. *Steroids* 1974, 24, 399.
- Kokke, W. C. M. C.; Withers, N. W.; Massey, I. J.; Fenical, W.; Djerassi, C. *Tetrahedron Lett.* 1979, 38, 3601.
- Zielinski, J.; Li, H.; Milkova, T. S.; Popov, S.; Marenkov, N. L.; Djerassi, C. *Tetrahedron Lett.* 1981, 22, 2345.
- Sjörstrand, U.; Kornprost, J. M.; Djerassi, C. *Steroids* 1981, 38, 355.
- Swenson, W.; Targle, B.; Clardy, J.; Withers, N. W.; Kokke, W. C. M. C.; Fenical, W.; Djerassi, C. *Tetrahedron Lett.* 1980, 21, 4663.
- Wengrovitz, R.; Sanduja, R.; Alam, M. *Comp. Biochem. Physiol., B* 1981, 69, 535.
- Khalil, M. W.; Durham, L. J.; Djerassi, C.; Sica, D. *J. Am. Chem. Soc.* 1980, 102, 2133.
- Kokke, W. C. M. C.; Fenical, W.; Pak, C. S.; Djerassi, C. *Tetrahedron Lett.* 1978, 4373.
- Zielinski, J.; Kokke, W. C. M. C.; Fenical, W.; Djerassi, C. *Steroids* 1982, 40, 403.
- Hofheinz, W.; Oesterhelt, G. *Helv. Chim. Acta* 1979, 62, 1307.
- Kokke, W. C. M. C.; Pak, C. S.; Fenical, W.; Djerassi, C. *Helv. Chim. Acta* 1979, 62, 1310.
- Theobald, N.; Wells, R. J.; Djerassi, C. *J. Am. Chem. Soc.* 1978, 100, 1677.
- Long, K.; Lou, Y.; Fang, Z. *Zhong. Daux. Xuebao, Ziran Kexue* 1982, 4, 71.
- Long, K.; Zheng, L.; Zheng, H. *Zhong. Daux. Xuebao, Ziran Kexue* 1981, 3, 105.
- Wu, Z.; Long, K. *Zhong. Daux. Xuebao, Ziran Kexue* 1981, 3, 83.
- Li, L. N.; Sjörstrand, U.; Djerassi, C. *J. Am. Chem. Soc.* 1981, 103, 115.
- Li, L. N.; Djerassi, C. *Tetrahedron Lett.* 1981, 22, 4639.
- Crist, B. V.; Li, X.; Bergquist, P. R.; Djerassi, C. *J. Org. Chem.* 1983, 48, 4472.
- Li, X.; Djerassi, C. *Tetrahedron Lett.* 1983, 24, 665.
- Li, L. N.; Djerassi, C. *J. Am. Chem. Soc.* 1981, 103, 3606.
- Li, L. N.; Sjörstrand, U.; Djerassi, C. *J. Org. Chem.* 1981, 46, 3867.
- Shubina, L. K.; Makarieva, T. N.; Kalinovsky, A. I.; Stonik, V. A. *Khim. Prir. Soedin.* 1985, 232.
- Catalan, C. A. N.; Lakshmi, V.; Schmitz, F. J.; Djerassi, C. *Steroids* 1982, 40, 455.
- Kobayashi, M.; Mitsubashi, H. *Steroids* 1982, 40, 665.
- Blanc, P. A.; Djerassi, C. *J. Am. Chem. Soc.* 1980, 102, 7113.
- Francisco, C.; Combaut, G.; Teste, J.; Maume, B. F. *Biophys. Acta* 1977, 487, 115.
- Schmitz, F. J.; Pattabhiraman, T. *J. Am. Chem. Soc.* 1970, 92, 6073.
- Bergmann, W.; Mc lean, M. J.; Lester, D. *J. Org. Chem.* 1943, 8, 271.
- Ling, N. C.; Hale, R. L.; Djerassi, C. *J. Am. Chem. Soc.* 1970, 92, 5281.
- Li, L. N.; Li, H.; Lang, R. W.; Itoh, T.; Sica, D.; Djerassi, C. *J. Am. Chem. Soc.* 1982, 104, 6726.
- Proudfoot, J. R.; Li, X.; Djerassi, C. *J. Org. Chem.* 1985, 50, 2026.
- Kokke, W. C. M. C.; Shoolery, J. N.; Fenical, W.; Djerassi, C. *J. Org. Chem.* 1984, 49, 3742.
- Cho, J. H.; Djerassi, C. *J. Chem. Soc. Perkin Trans. 1* 1987, 1307.
- Doss, G. A.; Djerassi, C. *J. Am. Chem. Soc.* 1983, 110, 8124.
- Steiner, E.; Djerassi, C.; Fattorusso, E.; Magno, S.; Mayol, L.; Santacroce, C.; Sica, D. *Helv. Chim. Acta* 1977, 60, 475.
- Theobald, N.; Shoolery, J. N.; Djerassi, C.; Erdman, T. R.; Scheuer, P. J. *J. Am. Chem. Soc.* 1978, 100, 5574.
- Minale, L.; Sodano, G. Non-Conventional Sterols of Marine Origin. In *Marine Natural Products Chemistry*; Faulkner, D. J., Fenical, W. H., Eds.; Plenum Publishing Corporation: New York; pp 87-109.
- Schmitz, F. J. Uncommon Marine Steroids. In *Marine Natural Products—Chemical and Biological Perspectives*; Scheuer, P. J., Ed.; Academic Press: New York, 1978; Vol. I, pp 241-297.
- Djerassi, C.; Theobald, N.; Kokke, W. C. M. C.; Pak, C. S.; Carlson, R. M. K. *Pure Appl. Chem.* 1979, 51, 211.

- (58) Djerassi, C. *Pure Appl. Chem.* 1981, 53, 873.
- (59) Withers, N. In *Marine Natural Products-Chemical and Biological Perspective*; Scheuer, P. J., Ed.; Academic Press: New York, 1983; Vol. V, pp 88-127.
- (60) Ikekawa, N. Structures, biosynthesis and function of sterols in invertebrates. In *Steroids and Bile Acids*; Danielsson, H., Sjövall, J., Eds.; Amsterdam, 1985; pp 199-230.
- (61) Garson, M. *J. Nat. Prod. Rep.* 1989, 6, 143.
- (62) Kerr, R. G.; Baker, B. *J. Nat. Prod. Rep.* 1991, 8, 465.
- (63) Francisco, C.; Combaut, G.; Teste, J.; Tarchini, C.; Djerassi, C. *Steroids* 1979, 34, 163.
- (64) Kabore, S. A.; Combaut, G.; Vidal, J. P.; Codomier, L.; Passet, J.; Girard, J.-P.; Rossi, J. C. *Phytochemistry* 1983, 22, 1239.
- (65) Morisaki, M.; Kidoova, S.; Ikekawa, N. *Chem. Pharm. Bull.* 1976, 24, 3214.
- (66) Fattorusso, E.; Magno, S.; Santacroce, C.; Sica, D.; Impellizzeri, G.; Mangiafico, S.; Oriente, G.; Piattelli, G.; Sciuto, S. *Phytochemistry* 1978, 14, 1579.
- (67) Fukuzawa, A.; Kumagai, Y.; Masumune, T.; Furusaki, A.; Katayama, C.; Matsumoto, T. *Tetrahedron Lett.* 1981, 22, 4085.
- (68) Riccio, R.; Zollo, F.; Finamore, E.; Minale, L.; Laurent, D.; Bargibant, G.; Pusset, J. *J. Nat. Prod.* 1985, 48, 266.
- (69) Wahidulla, S.; D'Souza, L.; Patel, J. *Phytochemistry* 1987, 26, 2864.
- (70) Prakash, O.; Roy, R.; Bhakuni, D. S.; Wahidulla, S.; Kamat, S. Y. *J. Nat. Prod.* 1989, 52, 686.
- (71) Babu, M. J.; Trivedi, G. K.; Mathur, H. H. *Phytochemistry* 1989, 28, 3237.
- (72) Kobayashi, M.; Murata, O. *Tetrahedron Lett.* 1992, 33, 519.
- (73) Ikekawa, N.; Tsuda, K.; Morisaki, N. *Chem. Ind. (London)* 1966, 14, 1179.
- (74) Knights, B. A. *Phytochemistry* 1970, 9, 903.
- (75) Ikekawa, N.; Morisaki, M.; Hirayama, K. *Phytochemistry* 1972, 11, 2317.
- (76) Newburger, J.; Vebel, J. J.; Ikawa, M.; Andersen, K. K.; Gagosian, R. B. *Phytochemistry* 1979, 18, 2042.
- (77) Okano, M.; Fukamiya, N.; Fujita, T.; Matsumoto, H.; Aratani, T. *Bull. Jpn. Soc. Sci. Fish.* 1985, 51, 1305.
- (78) Findlay, J. A.; Patil, A. D. *Phytochemistry* 1985, 24, 366.
- (79) Kobayashi, M.; Hasegawa, A.; Mitsuhashi, H. *Chem. Pharm. Bull.* 1985, 33, 4012.
- (80) Smith, L. L. *Cholesterol Autoxidation*; Pergamon Press: Oxford; 1981; pp 13-41.
- (81) Kobayashi, J.; Ishibashi, M.; Nakamura, H.; Ohizumi, Y.; Hirata, Y. *J. Chem. Soc., Perkin Trans. 1* 1989, 101.
- (82) Carballera, N.; Thomson, J. E.; Ayanoglu, Z.; Djerassi, C. *J. Org. Chem.* 1986, 51, 2751 and references cited there.
- (83) Hahn, S.; Stoilov, I. L.; Tam Ha, T. B.; Raedersdorff, D.; Doss, G. A.; Li, H.-T.; Djerassi, C. *J. Am. Chem. Soc.* 1988, 110, 8117.
- (84) Lam, W.-K.; Hahn, S.; Ayanoglu, E.; Djerassi, C. *J. Org. Chem.* 1989, 54, 3428.
- (85) Hahn, S.; Lam, W.-K.; Wu, L.; Silva, J. C.; Djerassi, C. *J. Biol. Chem.* 1989, 264, 21043.
- (86) Fusetani, N.; Matsunaga, S.; Konosu, S. *Tetrahedron Lett.* 1981, 22, 1985.
- (87) Sun, H. H.; Cross, S. S.; Gunasekera, M.; Koehn, F. E. *Tetrahedron* 1991, 47, 1185.
- (88) Kohen, F. E.; Gunasekera, M.; Cross, S. S. *J. Org. Chem.* 1981, 56, 1322.
- (89) Makarieva, T. N.; Shubina, L. K.; Kalinovskiy, A. I.; Stonik, V. A.; Elyakov, G. B. *Steroids* 1983, 42, 267.
- (90) Kerr, R. G.; Kerr, S. L.; Djerassi, C. *J. Org. Chem.* 1991, 56, 63.
- (91) Makar'eva, T. M.; Dmitrenak, P. S.; Shubina, L. K.; Stonik, V. A. *Chem. Nat. Prod.* 1988, 24, 312.
- (92) Kanazawa, S.; Fusetani, N.; Matsunaga, S. *Tetrahedron* 1992, 48, 5467.
- (93) Bruno, I.; Cardellina, J. H., II; Minale, L.; Riccio, R. Unpublished results.
- (94) Nakatsu, T.; Walker, R. P.; Thompson, J. E.; Faulkner, D. J. *Experientia* 1983, 39, 759.
- (95) Prinsep, M. R.; Blunt, J. W.; Munro, M. H. G. *J. Nat. Prod.* 1989, 52, 657.
- (96) D'Auria, M. V.; Riccio, R.; Minale, L.; La Barre, S.; Pusset, J. *J. Org. Chem.* 1987, 52, 3947.
- (97) Gunasekera, S. P.; Schmitz, F. J. *J. Org. Chem.* 1983, 48, 885.
- (98) Bartolotto, M.; Braekman, J. C.; Dalozze, D.; Losman, D.; Tursch, B. *Steroids* 1976, 28, 461.
- (99) Fujimoto, Y.; Yamada, T.; Ikegawa, N. *Chem. Pharm. Bull.* 1985, 33, 3129.
- (100) Capon, R. J.; Faulkner, D. J. *J. Org. Chem.* 1985, 50, 4771.
- (101) Braekman, J. C.; Dalozze, D.; Moussiaux, B.; Vandevyver, G.; Riccio, R. *Bull. Soc. Chim. Belg.* 1988, 97, 293.
- (102) West, R. R.; Cardellina, J. H., II. *J. Org. Chem.* 1988, 53, 2782.
- (103) Fujimoto, Y.; Miyasaka, S.; Ikeda, T.; Ikegawa, N.; Ohnishi, E.; Mizuno, T.; Watanabe, K. *J. Chem. Soc., Chem. Commun.* 1985, 10.
- (104) West, R. R.; Cardellina, J. H., II. *J. Org. Chem.* 1989, 54, 3234.
- (105) Seidel, S. B.; Proudfoot, J. R.; Djerassi, C.; Sica, D.; Sodano, G. *Steroids* 1986, 47, 49.
- (106) Isaacs, S.; Berman, R.; Kashman, Y.; Gebreyesus, T.; Yosief, T. *J. Nat. Prod.* 1991, 54, 83.
- (107) Milkova, Ta. S.; Mikhova, B. P.; Nikolov, N. M.; Popov, S. S.; Andreev, S. N. *J. Nat. Prod.* 1992, 55, 974.
- (108) Ballantine, J. A.; Williams, K.; Burke, B. A. *Tetrahedron Lett.* 1977, 1547.
- (109) Guella, G.; Pietra, F. *Helv. Chim. Acta* 1988, 71, 62.
- (110) Piccialli, V.; Sica, D. *J. Nat. Prod.* 1987, 50, 915.
- (111) Madaio, A.; Piccialli, V.; Sica, D. *J. Nat. Prod.* 1989, 52, 952.
- (112) Cafieri, F.; Fattorusso, E.; Gavagnin, M.; Santacroce, C. *J. Nat. Prod.* 1985, 48, 944.
- (113) Iorizzi, M.; Minale, L.; Riccio, R.; Lee, J.-S.; Yasumoto, T. *J. Nat. Prod.* 1988, 51, 1098.
- (114) Das, B.; Srinivas, K. V. N. S. *J. Nat. Prod.* 1992, 55, 974.
- (115) Piccialli, V.; Sica, D. *J. Nat. Prod.* 1987, 50, 915.
- (116) Migliuolo, A.; Notaro, G.; Piccialli, V.; Sica, D. *J. Nat. Prod.* 1990, 53, 1414.
- (117) Madaio, A.; Piccialli, V.; Sica, D. *Tetrahedron Lett.* 1988, 29, 5999.
- (118) Madaio, A.; Notaro, G.; Piccialli, V.; Sica, D. *J. Nat. Prod.* 1990, 53, 565.
- (119) Valisolo, Ji.; Luu, B.; Ourisson, G. *Tetrahedron* 1983, 39, 2779.
- (120) Ciminiello, P.; Fattorusso, E.; Magno, S.; Mangoni, A.; Pansini, M. *J. Am. Chem. Soc.* 1990, 112, 3505.
- (121) Migliuolo, A.; Piccialli, V.; Sica, D. *Tetrahedron* 1991, 47, 7937.
- (122) Enwall, E. L.; Van Der Helm, D.; Nan Hsu, I.; Pattabhiraman, T.; Schmitz, F. J.; Spraggins, R. L.; Weinheimer, A. J. *J. Chem. Soc., Chem. Commun.* 1972, 215.
- (123) Kazlauskas, R.; Murphy, P. T.; Ravi, B. N.; Sanders, R. L.; Wells, R. J. *Aust. J. Chem.* 1982, 35, 69.
- (124) Bonini, C.; Cooper, C. B.; Kazlauskas, R.; Wells, R. J.; Djerassi, C. *J. Org. Chem.* 1983, 48, 2108.
- (125) D'Auria, M. V.; Gomez Paloma, L.; Minale, L.; Riccio, R.; Debitus, C. *Tetrahedron Lett.* 1991, 32, 2149.
- (126) D'Auria, M. V.; Gomez Paloma, L.; Minale, L.; Riccio, R.; Debitus, C.; Lévi, C. *J. Nat. Prod.* 1992, 55, 311.
- (127) Tishler, M.; Ayer, S. W.; Andersen, R. J.; Mitchell, J. F.; Clardy, J. *Can. J. Chem.* 1988, 66, 1173.
- (128) Migliuolo, A.; Piccialli, V.; Sica, D. *J. Nat. Prod.* 1990, 53, 1262.
- (129) Aiello, A.; Fattorusso, E.; Magno, S.; Menna, M. M.; Pansini, M. *J. Nat. Prod.* 1991, 54, 281.
- (130) Malorni, A.; Minale, L.; Riccio, R. *Nouv. J. Chem.* 1978, 2, 351.
- (131) Aiello, A.; Ciminiello, P.; Fattorusso, E.; Magno, S. *Steroids* 1988, 52, 533.
- (132) Burgoyne, D. L.; Andersen, R. J.; Allen, T. M. *J. Org. Chem.* 1992, 57, 525.
- (133) Shoji, N.; Umeiyama, A.; Shin, K.; Takeda, K.; Arihara, S.; Kobayashi, J.; Takei, M. *J. Org. Chem.* 1992, 57, 2996.
- (134) Notaro, G.; Piccialli, V.; Sica, V.; Pronzato, R. *J. Nat. Prod.* 1992, 55, 773.
- (135) Ciminiello, P.; Fattorusso, E.; Magno, S.; Mangoni, A.; Pansini, M. *Steroids* 1992, 57, 62.
- (136) Eggersdorfer, M. L.; Kokke, W. C. M. C.; Crandell, C. W.; Hochlowsky, J. E.; Djerassi, C. *J. Org. Chem.* 1982, 45, 5304.
- (137) Kokke, W. C. M. C.; Fenical, W.; Djerassi, C. *Phytochemistry* 1981, 20, 127.
- (138) Findlay, J. A.; Patil, A. D. *Can. J. Chem.* 1985, 63, 2406.
- (139) De Riccardis, F.; Minale, L.; Iorizzi, M.; Debitus, C.; Lévi, C. *J. Nat. Prod.* 1993, 56, 282.
- (140) Koch, P.; Djerassi, C.; Lakshimi, V.; Schmitz, F. *Helv. Chim. Acta* 1983, 66, 2431.
- (141) Ayanoglu, E.; Djerassi, C.; Erdman, T. R.; Scheuer, P. J. *Steroids* 1978, 31, 815.
- (142) Kitagawa, J.; Kobayashi, M.; Okamoto, Y.; Yoshikawa, M.; Hamamoto, Y. *Chem. Pharm. Bull.* 1987, 35, 5036.
- (143) Schmitz, F. J.; Ksebat, M. B.; Gunasekera, S. P.; Agarwal, S. J. *Org. Chem.* 1988, 53, 5941.
- (144) Carmely, S.; Roll, M.; Loya, Y.; Kashman, Y. *J. Nat. Prod.* 1989, 52, 167.
- (145) D'Auria, M. V.; Gomez Paloma, L.; Minale, L.; Riccio, R. *Tetrahedron* 1992, 48, 491.
- (146) Cheng, J.; Kobayashi, J.; Nakamura, H.; Ohizumi, Y.; Hirata, Y.; Sasaki, T. *J. Chem. Soc., Perkin Trans. 1* 1988, 2403.
- (147) Hirota, H.; Takayama, S.; Miyashiro, S.; Ozaki, Y.; Ikegami, S. *Tetrahedron Lett.* 1990, 31, 3321.
- (148) Fattorusso, E.; Lanzotti, V.; Magno, S.; Novellino, E. *Biochem. System. Ecol.* 1985, 13, 167.
- (149) Cimino, G.; De Rosa, S.; De Stefano, S.; Sodano, G. *Tetrahedron Lett.* 1980, 21, 3303.
- (150) Fattorusso, E.; Lanzotti, V.; Magno, S.; Novellino, E. *J. Org. Chem.* 1985, 50, 2868.
- (151) Fattorusso, E.; Lanzotti, V.; Magno, S.; Novellino, E. *J. Nat. Prod.* 1985, 48, 784.
- (152) Aiello, A.; Fattorusso, E.; Magno, S. *J. Nat. Prod.* 1987, 50, 191.
- (153) Krebs, H. Chr. *Progress in the Chemistry of Organic Natural Products*; Herz, W., Grisebach, H., Kirby, G. W., Tamm, Ch., Eds.; Springer-Verlag: Wien, New York, 1986; Vol. 49, pp 151-363.
- (154) Murata, M.; Naoki, H.; Iwashita, T.; Matsunaga, S.; Sasaki, M.; Yokoyama, A.; Yasumoto, T. *J. Am. Chem. Soc.* 1993, 115, 2060.
- (155) Moore, R. E. *Progress in the Chemistry of Organic Natural Products*; Herz, W., Grisebach, H., Kirby, G. W., Tamm, Ch., Eds.; Springer-Verlag: New York, 1985; Vol. 48, pp 81-202.

- (156) Sturaro, A.; Guerriero, A.; De Clauser, R.; Pietra, F. *Experientia* 1982, 38, 1184.
- (157) Guerriero, A.; Pietra, F. *Comp. Biochem. Physiol.* 1985, 80B, 277.
- (158) Guerriero, A.; Traldi, P.; Pietra, F. *J. Chem. Soc., Chem. Commun.* 1986, 40.
- (159) Hoffman, J. A. *Progress in Ecdysozoan Research*; Elsevier: Amsterdam, 1980.
- (160) Gilbert, L. J.; Bollenbacher, W. E.; Gronger, N. A. *Annu. Rev. Physiol.* 1980, 42, 497.
- (161) Whitehead, D. L.; Seellheyner, K. *Experientia* 1982, 38, 1249.
- (162) Schmitz, F. J.; Campell, D. C.; Kubo, I. *Steroids* 1976, 28, 211.
- (163) Kazlauskas, R.; Murphy, P. T.; Quinn, R. J.; Wells, R. J.; Schonholzer, P. *Tetrahedron Lett.* 1977, 4439.
- (164) Higa, T.; Tanaka, J.; Tachibana, K. *Tetrahedron Lett.* 1981, 22, 2777.
- (165) For review: Stonik, V. A.; Elyakov, G. B. In *Biorganic Marine Chemistry*; Scheuer, P. J., Ed.; Springer-Verlag: Berlin, 1988; Vol. 2, pp 43-86.
- (166) Kelly, W. G.; Baudi, L.; Schoolery, J. N.; Lieberman, S. *Biochemistry* 1962, 1, 173.
- (167) Higa, T.; Tanaka, J.; Tsukitani, Y.; Kikuchi, H. *Chem. Lett.* 1981, 1647.
- (168) Rao, Ch. B.; Ramana, K. V.; Rao, D. V.; Fahy, E.; Faulkner, D. J. *J. Nat. Prod.* 1988, 81, 954.
- (169) Tanaka, J.; Higa, T.; Tachibana, K.; Iwashita, T. *Chem. Lett.* 1982, 1295.
- (170) Cimino, G.; Desiderio, B.; De Stefano, S.; Sodano, G. *Experientia* 1979, 35, 299.
- (171) Bandurraga, M. M.; Fenical, W. *Tetrahedron* 1985, 41, 1057.
- (172) Wasyluk, J. M.; Martin, G. E.; Weinheimer, A. J.; Alam, M. *J. Nat. Prod.* 1989, 52, 391.
- (173) Kingston, J. F.; Gregory, B.; Fallis, A. G. *Tetrahedron Lett.* 1977, 4261.
- (174) Cimino, G.; De Rosa, S.; De Stefano, S.; Scognamiglio, G.; Sodano, G. *Tetrahedron Lett.* 1981, 22, 3013.
- (175) Benvegnù, R.; Cimino, G.; De Rosa, S.; De Stefano, S. *Experientia* 1982, 38, 1443.
- (176) Cimino, G.; De Rosa, S.; De Stefano, S.; Sodano, G. *Experientia* 1984, 40, 246.
- (177) Fusetani, N.; Nagata, H.; Hirota, H.; Tsuyuki, T. *Tetrahedron Lett.* 1989, 30, 7079.
- (178) Ochi, M.; Yamada, K.; Kotsuki, M.; Shibata, K. *Chem. Lett.* 1991, 427.
- (179) Fusetani, N.; Yasukawa, K.; Matsunaga, S.; Hashimoto, K. *Tetrahedron Lett.* 1987, 28, 1187.
- (180) Bohlin, L.; Sjostrand, U.; Sodano, G.; Djerassi, C. *J. Org. Chem.* 1982, 47, 5309.
- (181) Bortolotto, M.; Braekman, J. C.; Dalozze, D.; Tursch, B. *Bull. Soc. Chim. Belg.* 1976, 85, 27.
- (182) Jagodzinska, B. M.; Trimmer, J. S.; Fenical, W.; Djerassi, C. *J. Org. Chem.* 1985, 50, 1435.
- (183) Jagodzinska, B. M.; Trimmer, J. S.; Fenical, W.; Djerassi, C. *J. Org. Chem.* 1985, 50, 2988.
- (184) Ciminiello, P.; Fenical, W.; Paul, V. J. *Experientia* 1990, 46, 980.
- (185) Su, J.; Yu, X.; Zeng, L.; Mak, T. C. W. *J. Nat. Prod.* 1989, 52, 934.
- (186) Kobayashi, M.; Hayashi, T.; Nakajima, F.; Mitsuhashi, H. *Steroids* 1979, 34, 285.
- (187) Kobayashi, M.; Mitsuhashi, H. *Steroids* 1982, 40, 673.
- (188) Kobayashi, M.; Hayashi, T.; Hayashi, K.; Tanabe, M.; Nakagawa, T.; Mitsuhashi, H. *Chem. Pharm. Bull.* 1983, 31, 1848.
- (189) Yamada, Y.; Suzuki, S.; Iguchi, K.; Kikuchi, H.; Tsukitani, Y.; Horiai, H.; Nakanishi, H. *Chem. Pharm. Bull.* 1980, 28, 473.
- (190) Kobayashi, M.; Kanda, F.; Rao, C. V. L.; Kumar, S. M. D.; Trimurlulu, G.; Rao, C. B. *Chem. Pharm. Bull.* 1990, 38, 1724.
- (191) Li, R.; Huang, Z.; Long, K. *Steroids* 1992, 57, 3.
- (192) Tillekeratne, L. M. V.; Liyanage, G. K.; Ratnassoriya, W. D.; Ksebati, M. B.; Schmitz, F. J. *J. Nat. Prod.* 1989, 52, 1143.
- (193) Maldowan, J. M.; Tursch, B. M.; Djerassi, C. *Steroids* 1974, 24, 387.
- (194) Maldowan, J. M.; Tan, L. W.; Djerassi, C. *Steroids* 1975, 26, 107.
- (195) Kato, H.; Sakuma, S.; Tada, A.; Kawanishi, S.; Shoji, J. *Jakugakuzasshi* 1968, 88, 710.
- (196) Kobayashi, M.; Mitsuhashi, H. *Chem. Pharm. Bull.* 1983, 31, 4127.
- (197) For review: Minale, L.; Riccio, R.; Zollo, F. *Progress in the Chemistry of Organic Natural Products*; Herz, W., Kirby, G. W., Steglich, W., Tamm, Ch., Eds.; Springer-Verlag: Wien, in press.
- (198) Riccio, R.; Santaniello, M.; Squillace Greco, O.; Minale, L. *J. Chem. Soc. Perkin Trans. 1* 1989, 823.
- (199) Lakshamana, B. R.; Subbaraju, G. V.; Reddy, M. C.; Rao, D. V.; Rao, Ch. B.; Raju, S. V. *J. Nat. Prod.* 1992, 55, 904.
- (200) Ksebati, M. B.; Schmitz, F. J. *Steroids* 1984, 43, 639.
- (201) Carmely, S.; Kashman, Y. *Tetrahedron* 1981, 37, 2397.
- (202) Ksebati, M. B.; Schmitz, F. J. *J. Org. Chem.* 1988, 53, 3926.
- (203) Sjostrand, U.; Bohlin, L.; Fisher, L.; Colin, M.; Djerassi, C. *Steroids* 1981, 38, 347.
- (204) Kitagawa, I.; Kobayashi, M.; Cui, Z.; Kiyota, Y.; Ohnishi, M. *Chem. Pharm. Bull.* 1986, 34, 4590.
- (205) Igushi, K.; Saitoh, S.; Yamada, Y. *Chem. Pharm. Bull.* 1989, 37, 2553.
- (206) Bortolotto, M.; Braekman, J. C.; Dalozze, D.; Tursch, B.; Karlsson, R. *Steroids* 1977, 30, 159.
- (207) Tursch, B.; Hootelè, C.; Kaisin, M.; Losman, D.; Karlsson, R. *Steroids* 1976, 27, 137.
- (208) Kobayashi, M.; Kanda, F.; Rao, C. V. L.; Kumar, S. M. D.; Rao, D. V.; Rao, Ch. B. *Chem. Pharm. Bull.* 1991, 39, 297.
- (209) Kobayashi, M.; Kobayashi, K.; Ramana, K. V.; Rao, Ch. V. L.; Rao, V. D.; Rao, Ch. B. *J. Chem. Soc., Perkin Trans. 1* 1991, 493.
- (210) Kobayashi, M.; Haribabu, B.; Anjaneyulu, V. *Chem. Pharm. Bull.* 1992, 40, 233.
- (211) Kobayashi, M.; Kanda, F.; Damarla, S. R.; Rao, D. V.; Rao, Ch. B. *Chem. Pharm. Bull.* 1990, 38, 2400.
- (212) Kobayashi, M.; Kiyota, Y.; Orito, S.; Kyogoku, Y.; Kitagawa, I. *Tetrahedron Lett.* 1984, 25, 373.
- (213) Kingston, J. F.; Fallis, A. G. *Can. J. Chem.* 1982, 60, 820.
- (214) Kingston, J. F.; Gregory, B.; Fallis, A. G. *Tetrahedron Lett.* 1977, 4261.
- (215) Kobayashi, M.; Lee, N. K.; Son, B. W.; Yamagi, K.; Kyogoku, Y.; Kitagawa, I. *Tetrahedron Lett.* 1984, 25, 5925.
- (216) Vanderah, D. J.; Djerassi, C. *Tetrahedron Lett.* 1977, 683.
- (217) Vanderah, D. J.; Djerassi, C. *J. Org. Chem.* 1978, 43, 1442.
- (218) Datta, P. K.; Ray, A. K.; Barua, A. K.; Chowdhuri, S. K.; Patra, A. *J. Nat. Prod.* 1990, 53, 1347.
- (219) Aiello, A.; Fattorusso, E.; Menna, M. L. *Steroids* 1991, 96, 513.
- (220) Aiello, A.; Fattorusso, E.; Menna, M. L. *J. Nat. Prod.* 1992, 55, 321.
- (221) Pettit, G. R. *Progress in the Chemistry of Organic Natural Products*; Herz, W., Kirby, G. W., Steglich, W., Tamm, Ch., Eds.; Springer-Verlag: Wien, 1991; Vol. 57, pp 153-195.
- (222) Cimino, G.; Sodano, G. *Chem. Sci.* 1989, 29, 389.
- (223) Ayer, S. W.; Andersen, R. J. *Tetrahedron Lett.* 1982, 23, 1039.
- (224) Williams, D. E.; Ayer, S. W.; Andersen, R. J. *Can. J. Chem.* 1986, 64, 1527.
- (225) Alam, M.; Sanduja, R.; Weinheimer, A. J. *Steroids* 1988, 52, 45.
- (226) Romero, M. S.; Seides, A. M. *Comp. Biochem. Physiol.* 1986, 84B, 125.
- (227) Miyamoto, T.; Higuchi, R.; Komori, T.; Fujioka, T.; Mhashi, K. *Tetrahedron Lett.* 1986, 27, 1153.
- (228) Spinella, A.; Gavagnin, M.; Crispino, A.; Cimino, G.; Martinez, E.; Ortea, J.; Sodano, G. *J. Nat. Prod.* 1992, 55, 989.
- (229) Goad, L. J. *Biochemical and Biophysical Perspectives. In Marine Biology*; Mahns, D. C., Sargent, J. R., Eds.; Academic Press: London, 1976; Vol. 3, pp 213-318.
- (230) Takemoto, T.; Ogawa, S.; Nishimoto, Hoffmeister, H. Z. *Naturforsch.* 1967, 22b, 68.
- (231) Horn, D. H. S. In *Naturally occurring insecticides*; Tocobsen, M., Crosby, D. G., Eds.; Marcel Dekker: New York, 1971; p 337.
- (232) Burnell, D. J.; ApSimon, J. W. *Echinoderm Saponins. In Marine Natural Products, Chemical and Biological Perspectives*; Scheuer, P. J., Ed.; Academic Press: New York, 1983; Vol. V, pp 287-389.
- (233) Hashimoto, Y. In *Marine Toxins and Other Bioactive Marine Metabolites*; Japan Scientific Societies Press: Tokyo, 1979; pp 268-288.
- (234) Minale, L.; Riccio, R.; Pizza, C.; Zollo, F. Steroidal Olygglycosides of Marine Origin. In *Natural Products and Biological Activities. A NATO Foundation Symposium*; Imura, H., Goto, T., Murachi, T., Nakajima, T., Eds.; University of Tokyo Press: Tokyo, 1986; pp 59-74.
- (235) Quinn, R. J. Chemistry of Aqueous Marine Extracts: Isolation Techniques. In *Biorganic Marine Chemistry*; Scheuer, P. J., Ed.; Springer Verlag: Berlin, 1988; Vol. 2, pp 1-39.
- (236) Habermehl, G. G.; Krebs, H. Chr. Toxins of Echinoderms. In *Studies in Natural Products Chemistry*; Atta-Ur-Rahman, Ed.; Elsevier Science Publishers B. V.: Amsterdam, 1990; Vol. 7, pp 265-316.
- (237) Minale, L.; Riccio, R.; Zollo, F. Structural Studies on Chemical Constituents of Echinoderms. In *Bioactive Natural Products—Studies in Natural Products Chemistry*; Atta-ur-Rahman, Basha, F. Z., Eds.; Elsevier Science Publishers B. V.: Amsterdam, in press.
- (238) Minale, L.; Pizza, C.; Riccio, R.; Zollo, F. *Pure Appl. Chem.* 1982, 54, 1935.
- (239) De Riccardis, F.; Iorizzi, M.; Minale, L.; Riccio, R.; Debitus, C. *Tetrahedron Lett.* 1992, 33, 1097.
- (240) De Riccardis, F.; Minale, L.; Riccio, R.; Giovannitti, B.; Iorizzi, M.; Debitus, C. *Gazz. Chim. Ital.* 1993, 122, 79.
- (241) D'Auria, M. V.; Finamore, E.; Minale, L.; Pizza, C.; Riccio, R.; Zollo, F.; Pusset, M.; Tirard, P. *J. Chem. Soc., Perkin Trans. 1* 1984, 2277.
- (242) Minale, L.; Pizza, C.; Zollo, F.; Riccio, R. *Tetrahedron Lett.* 1982, 23, 1841.
- (243) Minale, L.; Pizza, C.; Riccio, R.; Squillace Greco, O.; Zollo, F.; Pusset, J.; Menou, J. L. *J. Nat. Prod.* 1984, 47, 784.
- (244) Iorizzi, M.; Minale, L.; Riccio, R.; Yasumoto, T. *J. Nat. Prod.* 1992, 55, 866.
- (245) Finamore, F.; Minale, L.; Riccio, R.; Rinaldo, G.; Zollo, F. *J. Org. Chem.* 1991, 56, 1146.
- (246) Bruno, I.; Minale, L.; Riccio, R.; La Barre, S.; Laurent, D. *Gazz. Chim. Ital.* 1990, 120, 449.
- (247) Zollo, F.; Finamore, E.; Minale, L. *J. Nat. Prod.* 1987, 50, 794.

- (248) Seo, S.; Yoshimura, Y.; Satoh, T.; Uomori, A.; Takeda, K. *J. Chem. Soc., Perkin Trans. 1* 1986, 411.
- (249) D'Auria, M. V.; Minale, L.; Pizza, C.; Riccio, R.; Zollo, F. *Gazz. Chim. Ital.* 1984, 114, 469.
- (250) Riccio, R.; Minale, L.; Pagonis, S.; Pizza, C.; Zollo, F.; Pusset, J. *Tetrahedron* 1982, 38, 3615.
- (251) Riccio, R.; Iorizzi, M.; Squillace Greco, O.; Minale, L.; Laurent, D.; Barbin, Y. *Gazz. Chim. Ital.* 1985, 115, 505.
- (252) Zollo, F.; Finamore, E.; Minale, L.; Laurent, D.; Bargibant, G. *J. Nat. Prod.* 1986, 49, 919.
- (253) Higuchi, R.; Noguchi, Y.; Komori, T.; Sasaki, T. *Liebigs Ann. Chem.* 1988, 1185.
- (254) Iorizzi, M.; Minale, L.; Riccio, R. *Gazz. Chim. Ital.* 1990, 120, 147.
- (255) Rashkes, Ya. V.; Kicha, A. A.; Levina, E. V.; Stonik, V. A. *Chem. Nat. Comp.* 1985, 337.
- (256) D'Auria, M. V.; Iorizzi, M.; Minale, L.; Riccio, R.; Uriarte, E. *J. Nat. Prod.* 1990, 53, 94.
- (257) Minale, L.; Pizza, C.; Riccio, R.; Sorrentino, C.; Zollo, F.; Pusset, J.; Bargibant, G. *J. Nat. Prod.* 1984, 47, 790.
- (258) Bruno, I.; Minale, L.; Riccio, R. *J. Nat. Prod.* 1989, 52, 1022.
- (259) Segura De Correa, R.; Riccio, R.; Minale, L.; Duque, C. *J. Nat. Prod.* 1985, 48, 751.
- (260) Andersson, L.; Bohlin, L.; Iorizzi, M.; Riccio, R.; Minale, L.; Moreno-Lopez, W. *Toxicol.* 1989, 27, 179.
- (261) D'Auria, M. V.; De Riccardis, F.; Minale, L.; Riccio, R. *J. Chem. Soc., Perkin Trans. 1* 1990, 2889.
- (262) Preus, M. W.; McMorris, T. C. *J. Am. Chem. Soc.* 1979, 101, 3066.
- (263) Anastasia, M.; Allevi, P.; Ciuffreda, P.; Riccio, R. *Tetrahedron* 1986, 42, 4843.
- (264) Iorizzi, M.; Minale, L.; Riccio, R.; Debray, M.; Menou, J. L. *J. Nat. Prod.* 1986, 49, 67.
- (265) Riccio, R.; Squillace Greco, O.; Minale, L.; Duhet, D.; Laurent, D.; Pusset, J.; Chauviere, G.; Pusset, M. *J. Nat. Prod.* 1986, 49, 1141.
- (266) Bruno, I.; Minale, L.; Riccio, R. *J. Nat. Prod.* 1990, 53, 366.
- (267) Iorizzi, M.; Minale, L.; Riccio, R.; Kamiya, H. *J. Nat. Prod.* 1990, 53, 1225.
- (268) Riccio, R.; Iorizzi, M.; Minale, L. *Bull. Soc. Chim. Belg.* 1986, 95, 869.
- (269) Iorizzi, M.; Minale, L.; Riccio, R.; Higa, T.; Tanaka, J. *J. Nat. Prod.* 1991, 54, 1254.
- (270) Kicha, A. A.; Kalinovskii, A. I.; Stonik, V. A. *Chem. Nat. Comp.* 1989, 25, 376.
- (271) Minale, L.; Pizza, C.; Zollo, F.; Riccio, R. *J. Nat. Prod.* 1983, 46, 736.
- (272) Riccio, R.; Squillace Greco, O.; Minale, L.; Laurent, D.; Duhet, D. *J. Chem. Soc., Perkin Trans. 1* 1986, 665.
- (273) Mattia, C. A.; Mazzarella, L.; Puliti, R.; Riccio, R.; Minale, L. *Acta Crystallogr.* 1988, C44, 2170.
- (274) Finamore, E.; Zollo, F.; Minale, L.; Yasumoto, T. *J. Nat. Prod.* 1992, 55, 767.
- (275) De Riccardis, F.; Iorizzi, M.; Minale, L.; Riccio, R.; Debitus, C.; Duhet, D.; Monnot, C. Presented at the Seventh International Symposium on Marine Natural Products, Capri, Italy, July 1992; abstract P50.
- (276) Riccio, R.; D'Auria, M. V.; Iorizzi, M.; Minale, L.; Laurent, D.; Duhet, D. *Gazz. Chim. Ital.* 1985, 115, 405.
- (277) Koizumi, N.; Morisaki; Ikekawa, N. *Tetrahedron Lett.* 1975, 26, 2203.
- (278) Kuizumi, N.; Fujimoto, Y.; Takeshita, T.; Ikekawa, N. *Chem. Pharm. Bull.* 1979, 27, 38.
- (279) Dale, J. A.; Mosher, H. S. *J. Am. Chem. Soc.* 1973, 95, 512.
- (280) Yamaguchi, S. In *Asymmetric Synthesis*; Morrison, J. D., Ed.; Academic Press: New York, 1983; Vol 1, pp 125-152.
- (281) Casapullo, A.; Finamore, E.; Minale, L.; Zollo, F.; Carrè, J. B.; Debitus, C.; Laurent, D.; Folgore, A.; Galdiero, F. *J. Nat. Prod.* 1993, 56, 105.
- (282) Riccio, R.; Minale, L.; Pizza, C.; Zollo, F.; Pusset, J. *Tetrahedron Lett.* 1982, 23, 2899.
- (283) D'Auria, M. V.; Fontana, A.; Minale, L.; Riccio, R. *Gazz. Chim. Ital.* 1990, 120, 155.
- (284) Riccio, R.; Iorizzi, M.; Minale, L.; Oshima, Y.; Yasumoto, T. *J. Chem. Soc., Perkin Trans 1* 1988, 1337.
- (285) Pizza, C.; Minale, L.; Laurent, D.; Menou, J. L. *Gazz. Chim. Ital.* 1985, 115, 585.
- (286) Kicha, A. A.; Kalinovskiy, A. J.; Levina, E. V.; Stonik, V. A.; Elyakov, G. B. *Tetrahedron Lett.* 1983, 24, 3893.
- (287) Andersson, L.; Bohlin, L.; Riccio, R.; Minale, L. *J. Chem. Res.* 1987, (S) 246, (M) 2085.
- (288) Andersson, L.; Bano, S.; Bohlin, L.; Riccio, R.; Minale, L. *J. Chem. Res.* 1985, (S) 366, (M) 3873.
- (289) Minale, L.; Pizza, C.; Riccio, R.; Zollo, F. *Experientia* 1983, 39, 569.
- (290) Kicha, A. A.; Kalinovskii, A. I.; Levina, E. V.; Rashkes, Ya. B.; Stonik, V. A.; Elyakov, G. B. *Chem. Nat. Comp. (English)* 1985, 332.
- (291) Kicha, A. A.; Kalinovskiy, A. I.; Levina, E. V.; Rashkes, Ya. V.; Stonik, V. A. *Khim. Prir. Soedin.* 1987, 21, 356; quoted in Ref. 165.
- (292) Riccio, R.; Minale, L.; Bano, S.; Bano, N.; Uddin Ahmad, V. *Gazz. Chim. Ital.* 1987, 117, 755.
- (293) Maier, M. S.; Roccatagliata, A. J.; Seldes, A. Presented at the Seventh International Symposium on Marine Natural Products, Capri, Italy, July 1992; abstract P51.
- (294) Minale, L.; Pizza, C.; Riccio, R.; Zollo, F. *Experientia* 1983, 39, 567.
- (295) Riccio, R.; Squillace Greco, O.; Minale, L. *J. Nat. Prod.* 1988, 51, 989.
- (296) Kapustina, I. I.; Kalinovskiy, A. I.; Polonik, S. G.; Stonik, V. A. *Khim. Prir. Soedin.* 1987, 23, 250; *Chem. Abstr.* 1987, 106, 404500.
- (297) Kicha, A. A.; Kalinovskiy, A. I.; Levina, E. V.; Andriyashchenko, P. V. *Khim. Prir. Soedin.* 1985, 21, 801; *Chem. Abstr.* 1986, 104, 204151.
- (298) Andersson, L.; Nasir, A.; Bohlin, L.; Kenne, L. *J. Nat. Prod.* 1987, 50, 994.
- (299) Riccio, R.; Minale, L.; Bano, S.; Uddin Ahmad, V. *Tetrahedron Lett.* 1987, 28, 2291.
- (300) Zollo, F.; Finamore, E.; Martuccio, C.; Minale, L. *J. Nat. Prod.* 1990, 53, 1000.
- (301) Kicha, A. A.; Kalinovskiy, A. I.; Andriyashchenko, P. V.; Levina, E. V. *Khim. Prir. Soedin.* 1986, 22, 592; *Chem. Abstr.* 1987, 106, 116759.
- (302) Zollo, F.; Finamore, E.; Riccio, R.; Minale, L. *J. Nat. Prod.* 1989, 52, 693.
- (303) Pizza, C.; Pezzullo, P.; Minale, L.; Breitmaier, E.; Pusset, J.; Tirard, P. *J. Chem. Res.* 1985, (S) 76, (M) 969.
- (304) Findlay, J. A.; Zheng-Quan He. *J. Nat. Prod.* 1991, 54, 428.
- (305) Riccio, R.; Squillace Greco, O.; Minale, L.; La Barre, S.; Laurent, D. *J. Nat. Prod.* 1988, 51, 1003.
- (306) Minale, L.; Pizza, C.; Riccio, R.; Zollo, F.; Pusset, J.; Laboute, P. *J. Nat. Prod.* 1984, 47, 558.
- (307) Levina, E. V.; Kalinovskii, A. I.; Andriyashchenko, P. V.; Kicha, A. A. *Khim. Prir. Soedin.* 1987, 23, 246; *Chem. Abstr.* 1987, 106, 404499.
- (308) Kicha, A. A.; Kalinovskii, A. I.; Stonik, V. A. *Chem. Nat. Comp.* 1989, 25, 569.
- (309) Minale, L.; Pizza, C.; Plomitello, A.; Riccio, R.; Zollo, F.; Mellon, F. A. *Gazz. Chim. Ital.* 1984, 114, 143.
- (310) Zollo, F.; Finamore, E.; Minale, L. *Gazz. Chim. Ital.* 1985, 115, 303.
- (311) Vázquez, M. J.; Quiñoa, E.; Riguera, R.; San Martín, A.; Dárias, J. *Tetrahedron* 1992, 48, 6739.
- (312) Hashimoto, Y.; Yasumoto, T. *Bull. Jpn. Soc. Scient. Fish.* 1960, 26, 1132.
- (313) Makie, A. M.; Singh, H. T.; Owen, Y. M. *Comp. Biochem. Physiol.* 1977, 56B, 9.
- (314) Yasumoto, T.; Wanatabe, T.; Hashimoto, Y. *Bull. Jpn. Soc. Sci. Fish.* 1964, 30, 357.
- (315) Mackie, A. M.; Lasker, R.; Grant, P. T. *Comp. Biochem. Physiol.* 1968, 36B, 415.
- (316) Ikegami, S.; Kamiya, Y.; Tamura, S. *Agr. Biol. Chem.* 1972, 36, 2005.
- (317) Fujimoto, Y.; Yamada, T.; Ikekawa, N.; Nishiyama, I.; Matsui, T.; Hoshi, M. *Chem. Pharm. Bull.* 1987, 35, 1829.
- (318) Fusetani, N.; Kato, Y.; Hashimoto, K.; Komori, T.; Itakura, Y.; Kawasaki, T. *J. Nat. Prod.* 1984, 47, 997.
- (319) Ruggeri, G. D.; Nigrelli, R. F. Physiologically Active Substances from Echinoderms. In *Bioactive Compounds from the Sea*; Humm, H. J., Lane, C. E., Eds.; Marcel Dekker: New York, 1974; Marine Science Series, Vol. 1, pp 183-195.
- (320) Pettit, G. R.; Day, J. F.; Hartwell, J. L.; Wood, H. B. *Nature* 1970, 227, 962.
- (321) Dubois, M. A.; Higuchi, R.; Komori, T.; Sasaki, T. *Liebigs Ann. Chem.* 1988, 845.
- (322) Shimizu, Y. *Experientia* 1971, 27, 1188.
- (323) Friess, S. L. *Fed. Proc.* 1972, 31, 1146.
- (324) Goldsmith, L. A.; Carlson, G. P. *Food-Drugs Sea, Proc. Conf. Drugs Sea 1974*; Webber, H. H., Ruggeri, G. D. Eds.; Marine Technol. Soc.: Washington, 1976; p 354.
- (325) Ruggeri, G. D.; Nigrelli, R. F. *Am. Soc. Biol.* 1966, 6, 592.
- (326) Sheikh, Y. M.; Tursch, B. M.; Djerassi, C. *J. Am. Chem. Soc.* 1972, 94, 3278.
- (327) ApSimon, J. W.; Buccini, J. A.; Badripersaud, S. *Can. J. Chem.* 1973, 51, 850.
- (328) Kitagawa, I.; Kobayashi, K.; Sugawara, T. *Chem. Pharm. Bull.* 1978, 26, 1852.
- (329) De Simone, F.; Dini, A.; Minale, L.; Pizza, C.; Riccio, R. *Tetrahedron Lett.* 1979, 959.
- (330) De Simone, F.; Dini, A.; Finamore, E.; Minale, L.; Pizza, C.; Riccio, R. *Comp. Biochem. Physiol.* 1979, 64B, 25.
- (331) Kitagawa, I.; Kobayashi, M. *Chem. Pharm. Bull.* 1978, 23, 1864.
- (332) Itakura, Y.; Komori, T.; Kawasaki, T. *Liebig Ann. Chem.* 1983, 2079.
- (333) Minale, L.; Riccio, R.; Squillace Greco, O.; Pusset, J.; Menou, J. L. *Comp. Biochem. Physiol.* 1985, 80B, 113.
- (334) Riccio, R.; Pizza, C.; Squillace Greco, O.; Minale, L. *J. Chem. Soc., Perkin Trans. 1* 1985, 655.
- (335) Riccio, R.; Squillace Greco, O.; Minale, L.; Pusset, J.; Menou, J. L. *J. Nat. Prod.* 1985, 48, 97.
- (336) Riccio, R.; Iorizzi, M.; Squillace Greco, O.; Minale, L.; Debray, M.; Menou, J. L. *J. Nat. Prod.* 1985, 48, 756.
- (337) Ikegami, S.; Okano, K.; Muragaki, H. *Tetrahedron Lett.* 1979, 1769.
- (338) Findlay, J. A.; Jaseja, M.; Burnell, D. J.; Brisson, J. R. *Can. J. Chem.* 1987, 65, 1386.

- (339) Itakura, Y.; Komori, T.; Kawasaki, T. *Liebigs Ann. Chem.* 1983, 56.  
(340) Itakura, Y.; Komori, T. *Liebigs Ann. Chem.* 1986, 499.  
(341) Dubois, M. A.; Noguchi, Y.; Higuchi, R.; Komori, T. *Liebigs Ann. Chem.* 1988, 495.  
(342) D'Auria, M. V.; Iorizzi, M.; Minale, L.; Riccio, R. *J. Chem. Soc. Perkin Trans. 1* 1990, 1019.  
(343) Noguchi, Y.; Higuchi, R.; Marubayashi, N.; Komori, T. *Liebigs Ann. Chem.* 1987, 341.  
(344) Okano, K.; Nakamura, T.; Kamiya, Y.; Ikegami, S. *Agric. Biol. Chem.* 1981, 45, 805.  
(345) Itakura, Y.; Komori, T. *Liebigs Ann. Chem.* 1986, 359.  
(346) Findlay, J. A.; Jaseja, M.; Brisson, J. R. *Can. J. Chem.* 1987, 65, 2605.  
(347) Komori, T.; Krebs, H. Chr.; Itakura, Y.; Higuchi, R.; Sakamoto, K.; Taguchi, S.; Kawasaki, T. *Liebigs Ann. Chem.* 1983, 2092.  
(348) Dini, A.; Mellon, F. A.; Minale, L.; Pizza, C.; Riccio, R.; Self, R.; Zollo, F. *Comp. Biochem. Physiol.* 1983, 76B, 839.  
(349) Bruno, I.; Minale, L.; Pizza, C.; Zollo, F.; Riccio, R.; Mellon, F. A. *J. Chem. Soc., Perkin Trans. 1* 1984, 1875.  
(350) Findlay, J. A.; Zheng-Quan He; Blackwell, B. *Can. J. Chem.* 1990, 68, 1215.  
(351) Komori, T.; Nanri, H.; Itakura, Y.; Sakamoto, K.; Taguchi, S.; Higuchi, R.; Kawasaki, T.; Higuchi, T. *Liebigs Ann. Chem.* 1983, 37.  
(352) Krebs, H. Chr.; Komori, T.; Kawasaki, T. *Liebigs Ann. Chem.* 1984, 296.  
(353) Okano, K.; Ohkawa, N.; Ikegami, S. *Agric. Biol. Chem.* 1985, 49, 2823.  
(354) Findlay, J. A.; Zheng-Quan He; Jaseja, M. *Can. J. Chem.* 1989, 67, 2078.  
(355) ApSimon, J. W.; Badripersaud, S.; Buccini, J. A.; Eenkhoorn, J.; Gilgan, M. W. *Can. J. Chem.* 1980, 58, 2703.  
(356) Kicha, A. A.; Kalinovskii, A. I.; Stonik, V. A. *Chem. Nat. Comp.* 1992, 27, 452.  
(357) De Simone, F.; Dini, A.; Finamore, E.; Minale, L.; Pizza, C.; Riccio, R.; Zollo, F. *J. Chem. Soc., Perkin Trans. 1* 1981, 1855.  
(358) Riccio, R.; De Simone, F.; Dini, A.; Minale, L.; Pizza, C.; Senatore, F.; Zollo, F. *Tetrahedron Lett.* 1981, 22, 1557.  
(359) Riccio, R.; Dini, A.; Minale, L.; Pizza, C.; Zollo, F.; Sevenet, T. *Experientia* 1982, 38, 68.  
(360) Riccio, R.; D'Auria, M. V.; Minale, L. *J. Org. Chem.* 1986, 51, 533.  
(361) Riccio, R.; D'Auria, M. V.; Minale, L. *Tetrahedron* 1985, 41, 6041.  
(362) D'Auria, M. V.; Riccio, R.; Uriarte, E.; Minale, L.; Tanaka, J.; Higa, T. *J. Org. Chem.* 1989, 54, 234.  
(363) D'Auria, M. V.; Gomez Paloma, L.; Minale, L.; Riccio, R. *J. Chem. Soc., Perkin Trans. 1* 1990, 2895.  
(364) Riccio, R.; Finamore, E.; Santaniello, M.; Zollo, F. *J. Org. Chem.* 1990, 55, 2548.  
(365) Levina, E. V.; Fedorov, S. N.; Stonik, V. A.; Adriyashchenko, P. V.; Kalinovskii, A. I.; Isakov, V. V. *Chem. Nat. Comp.* 1990, 26, 408.  
(366) D'Auria, M. V.; Gomez Paloma, L.; Minale, L.; Riccio, R.; Zampella, A.; Tanaka, J.; Higa, T. *Tetrahedron Lett.* 1992, 33, 4641.  
(367) Karlson, P.; Skinner, D. M. *Nature* (London) 1960, 185, 543.  
(368) Horn, D. H. S.; Middleton, E. J.; Wunderlich, J. A. *J. Chem. Soc., Chem. Commun.* 1966, 339.  
(369) Gallbraith, M. N.; Horn, D. H. S.; Middleton, E. J.; Haekney, R. *J. J. Chem. Soc., Chem. Commun.* 1968, 83.  
(370) Faux, A.; Horn, D. H. S.; Middleton, E. J.; Fales, H. M.; Lowe, M. E. *J. Chem. Soc., Chem. Commun.* 1969, 175.  
(371) Tam Ha, T. B.; Kokke, W. C. M. C.; Djerassi, C. *Steroids* 1982, 40, 433.  
(372) Zollo, F.; Finamore, E.; Gargiulo, D.; Riccio, R.; Minale, L. *Comp. Biochem. Physiol.* 1986, 85B, 559.  
(373) Guyot, M.; Durgeat, M. *Tetrahedron Lett.* 1981, 22, 1391.  
(374) Guyot, M.; Davoust, D.; Belaud, C. *Tetrahedron Lett.* 1982, 23, 1905.  
(375) Hammarsten, O. *Hoppe-Seyler's Z. Physiol. Chem.* 1898, 24, 322.  
(376) Scheuer, P. J. *Chemistry of Marine Natural Products*; Academic Press: New York 1973; pp 67-68.  
(377) Cross, A. D. *J. Chem. Soc.* 1961, 2817.  
(378) Briggs, T.; Haslewood, G. A. D. *Biochem. J.* 1961, 78, 58.  
(379) Haslewood, G. A. D. *Biochem. J.* 1961, 78, 308.  
(380) Anderson, I. G.; Haslewood, G. A. D.; Cross, A. D.; Tokes, L. *Biochem. J.* 1967, 104, 1061.  
(381) Anderson, I. G.; Haslewood, G. A. D. *Biochem. J.* 1969, 112, 763.  
(382) Tachibana, K.; Sakaitani, M.; Nakanishi, K. *Science* 1984, 226, 703.  
(383) Tachibana, K.; Sakaitani, M.; Nakanishi, K. *Tetrahedron* 1985, 41, 1027.  
(384) Tachibana, K.; Gruber, S. H. *Toxicon* 1988, 26, 839.  
(385) Gargiulo, D.; Blizzard, T. A.; Nakanishi, K. *Tetrahedron* 1989, 45, 5423.  
(386) Tachibana, K. In *Biorganic Marine Chemistry*; Scheuer, P. J., Ed.; Springer-Verlag: Berlin, 1988; Vol. 2, pp 117-138.