

# Recent Advances in the Chemistry of Dinitrogen Complexes

Masanobu Hidai\* and Yasushi Mizobe

Department of Chemistry and Biotechnology, Graduate School of Engineering, The University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113, Japan

Received January 19, 1995 (Revised Manuscript Received April 10, 1995)

## Contents

I. Introduction	1115
II. Advances in Syntheses and Structures of Dinitrogen Complexes	1116
A. Metals	1116
B. Ancillary Ligands	1117
C. Coordination Modes	1118
D. Formation of Dinitrogen Ligands	1119
III. Protonation Reactions of Coordinated Dinitrogen	1120
A. Protonation of N <sub>2</sub> Coordinated to a Single Transition Metal Center	1120
B. Protonation of N <sub>2</sub> Coordinated to More than Two Transition Metals	1121
IV. Reactions of Hydrazines Relevant to Nitrogen Fixation	1122
V. Conversion of Coordinated Dinitrogen into Organonitrogen Ligands and Compounds	1123
A. C–N Bond Formation at Coordinated N <sub>2</sub> by Reaction with Organic Halides	1123
B. C–N Bond Formation at Coordinated N <sub>2</sub> via Hydrazido(2–) or Diazenido Complexes	1124
C. Synthesis of Nitrogen-Containing Organic Compounds from Coordinated N <sub>2</sub>	1126
D. Organic Synthesis Using Nitrido Complexes Derived from Molecular N <sub>2</sub> as Key Intermediate	1127
E. Silylation and Germylation of Coordinated N <sub>2</sub> and Catalytic Conversion of Molecular N <sub>2</sub> into Silylamines	1128
VI. Reactions of Coordinated N <sub>2</sub> with Lewis Acids or Organometallic Reagents	1129
VII. Conclusions	1131
VIII. Abbreviations	1131
IX. References	1131

## I. Introduction

Development of the chemical N<sub>2</sub>-fixing system converting a quite inert N<sub>2</sub> molecule into nitrogenous compounds under mild conditions is a challenging topic in chemistry. Industrial NH<sub>3</sub> production from N<sub>2</sub> and H<sub>2</sub> gases (Haber–Bosch process) has successfully been carried out for more than 80 years by the use of Fe-based heterogeneous catalysts, but it requires extremely drastic conditions. Although a Ru-on-graphite catalyst developed recently in the Kellogg advanced ammonia process allows some milder operating conditions (70–105 bar, 350–470 °C),<sup>1</sup> this apparently awaits further improvement.

In contrast, biological N<sub>2</sub>-fixing system can reduce N<sub>2</sub> gas into NH<sub>3</sub> at ambient temperature and pres-



Masanobu Hidai, born in 1940, received his doctorate for work on oligomerization catalyzed by transition metal complexes from the Department of Industrial Chemistry at the University of Tokyo in 1968. After that, he joined the staff of the department and became a Professor in the Department of Synthetic Chemistry at the same University in 1986. He spent about one year from 1976 to 1977 at the University of Sussex, U.K., to work with Professor J. Chatt. His scientific activities are centered around the chemistry of nitrogen fixation, organic syntheses catalyzed by transition metal complexes, and syntheses and reactivities of polynuclear metal complexes or metal cluster complexes. He is now the leader of the research project Activation of Small Inert Molecules (1992–1995), which is one of the Scientific Researches on Priority Areas supported by the Ministry of Education, Science and Culture, Japan.




Yasushi Mizobe was born in Yamaguchi, Japan, in 1953. He received the Dr. Eng. degree from the University of Tokyo in 1981 under the supervision of Professors Masanobu Hidai and Yasuzo Uchida. After two postdoctoral years with Professor R. H. Holm at Harvard University, he joined the University of Tokyo, where he is currently Associate Professor of Department of Chemistry and Biotechnology. His research interest is centered on the syntheses, structures, and reactivities of organometallic complexes which can be used for the activation of small molecules.

sure (eq 1), and it has long been recognized that Mo



is involved in the catalytic component as an element essential to display an activity. More recent studies




**Figure 1.** Structural model of the FeMo cofactor site of resting state in Mo-containing nitrogenases determined by X-ray crystallography. The third bridging ligand Y, which was unknown according to the initial reports,<sup>2</sup> is now considered to be a third sulfide.<sup>4</sup>

have clarified the presence of two associated proteins in the enzyme nitrogenase, *viz.* FeMo and Fe proteins, and it is believed that the substrate binding and reduction occur at the multimetallic site involved in the former, which consists of Mo and Fe atoms bridged by sulfide ligands. Quite attractive is the FeMo cofactor structure model deduced recently from single-crystal X-ray analysis for FeMo proteins of *Azotobacter vinelandii*<sup>2</sup> and *Clostridium pasteurianum*,<sup>3</sup> which contains the cuboidal Fe<sub>4</sub>S<sub>3</sub> and Fe<sub>3</sub>MoS<sub>3</sub> units bridged by three sulfides (Figure 1). EXAFS analysis also confirmed the presence of this core in both the isolated *A. vinelandii* FeMo protein and cofactor, with a slight relaxation of the cluster structure in the latter upon removal from the protein matrix.<sup>4</sup> How the N<sub>2</sub> molecule binds to this active site is still uncertain. At a glance, a trigonal prismatic cavity surrounded by six coordinatively unsaturated Fe atoms seems to be susceptible to N<sub>2</sub> insertion, giving the  $\mu_6$ -N<sub>2</sub> ligand, although the cavity size demonstrated at least for this dithionite reduced form is considered to be too small to accommodate N<sub>2</sub>.<sup>2a</sup> Alternatively, extended Hückel type calculations using a model [HFe( $\mu_3$ -S)<sub>3</sub>Fe<sub>3</sub>( $\mu_2$ -S)<sub>3</sub>-Fe<sub>3</sub>( $\mu_3$ -S)<sub>3</sub>MoH<sub>3</sub>]<sup>-</sup> suggested the coordination of N<sub>2</sub> rather to the edge or the face of the Fe<sub>6</sub> trigonal prism as a bridge between two cuboidal units.<sup>5</sup> On the other hand, coordination of CN<sup>-</sup> to the isolated FeMo cofactor has been reported to take place at the Mo atom from the EXAFS criteria.<sup>4</sup>

Albeit this ambiguity of the binding and reduction mechanism of N<sub>2</sub> in the biological system, it is apparent that the transition metals play an important role in promoting this transformation under mild conditions. Studies of the syntheses and reactions of N<sub>2</sub> complexes are therefore of particular interest. The recent discovery of the nitrogenases containing either V in place of Mo or only Fe as a transition metal component<sup>6</sup> may imply that a wide range of transition metals possibly facilitate the intriguing reactions of N<sub>2</sub> in their coordination sphere.

Two pioneering findings associated with the chemical N<sub>2</sub>-fixing system using transition metal compounds are the isolation of the first dinitrogen complex [Ru(NH<sub>3</sub>)<sub>5</sub>(N<sub>2</sub>)]<sup>2+</sup><sup>7</sup> and the formation of NH<sub>3</sub> by treatment of N<sub>2</sub> gas with a mixture of transition metal salt and organomagnesium, -lithium, or -aluminum followed by hydrolysis,<sup>8</sup> demonstrating that even N<sub>2</sub> can coordinate to the metal to give a stable N<sub>2</sub> complex and N<sub>2</sub> can be converted readily into nitrogenous compounds in the presence of transition metal species. Stimulated by these observations, syntheses of N<sub>2</sub> complexes have been attempted for a wide range of transition metals, and

now unequivocally characterized N<sub>2</sub> complexes are known for almost all transition metals. However, despite the occurrence of numerous N<sub>2</sub> complexes, those which react in a well-defined manner to give nitrogen-containing ligands and/or compounds are relatively limited. It may be emphasized that Mo and W dinitrogen complexes containing tertiary phosphines as ancillary ligands are most intensively studied since the initial preparation of *trans*-[Mo(N<sub>2</sub>)<sub>2</sub>(dppe)<sub>2</sub>] (**1a**) in this laboratory<sup>9</sup> because of, at least in part, their possible relevance to the active site of FeMo nitrogenases. Interestingly, these Mo and W complexes, especially *trans*-[M(N<sub>2</sub>)<sub>2</sub>(dppe)<sub>2</sub>] (**1**) and *cis*-[M(N<sub>2</sub>)<sub>2</sub>(PMe<sub>2</sub>Ph)<sub>4</sub>] (**2**), have actually proved to undergo numerous transformations of their N<sub>2</sub> ligands, demonstrating the rich chemistry of N<sub>2</sub>.



This review deals with the recent progress in the syntheses and reactions of N<sub>2</sub> complexes. Significant emphasis is also placed upon the organic chemistry of the hydrazido(2-) (MNNH<sub>2</sub>) complexes readily derived from N<sub>2</sub> complexes of Mo and W and that of the nitrido (MN) complexes prepared from gaseous N<sub>2</sub>. In relation to these reactions, some intriguing reactivities displayed at the M-N multiple bonds in certain imido complexes are also cited regardless of whether the origin of the nitrogen atom in the organonitrogen ligand comes from molecular nitrogen or not. As for the older works, readers can refer to several comprehensive reviews which appeared previously.<sup>10-13</sup> More reviews are also available, which cover some limited areas in this chemistry.<sup>14-21</sup>


## II. Advances in Syntheses and Structures of Dinitrogen Complexes

Since syntheses and structures of N<sub>2</sub> complexes were extensively surveyed previously,<sup>10</sup> this chapter deals mainly with N<sub>2</sub> complexes reported more recently. Furthermore, herein are cited only the complexes which are considered to be of significant importance due to involvement of new metals or coligands, the novel coordination mode displayed by N<sub>2</sub> ligand, or the new synthetic route.


### A. Metals

Dinitrogen complexes of V have been attracting considerable attention because of the presence of V-containing nitrogenases. However, it was in 1989 that the structurally characterized V dinitrogen complex was reported for the first time, although the N<sub>2</sub> fixing ability of V(II) species had been well demonstrated much earlier. The first dinitrogen complex [ {(o-Me<sub>2</sub>NCH<sub>2</sub>C<sub>6</sub>H<sub>4</sub>)<sub>2</sub>V(py)}<sub>2</sub>( $\mu$ -N<sub>2</sub>) ] (**3**)<sup>22</sup> was prepared from [V(TMEDA)<sub>2</sub>Cl<sub>2</sub>] with o-Me<sub>2</sub>NCH<sub>2</sub>C<sub>6</sub>H<sub>4</sub>-Li, which contains a  $\mu$ - $\eta^1$ : $\eta^1$ -N<sub>2</sub> ligand with essentially linear V-N-N-V linkage. Several V complexes

have subsequently been isolated, which include mononuclear complexes with terminal end-on  $N_2$  ligands, e.g.  $[Na(THF)][V(N_2)_2(dppe)_2]$  (**4**)<sup>23</sup> and dinuclear complexes with a  $\mu-\eta^1:\eta^1-N_2$  ligand such as  $[V(CH_2Bu^t)_3]_2(\mu-N_2)$  (**5**),<sup>24</sup>  $[{(Pr^i)_2N}_3V]_2(\mu-N_2)$ ,<sup>25</sup>  $[{(Me_3Si)NC(Ph)N(SiMe_3)_2V}]_2(\mu-N_2)$ ,<sup>26</sup> and  $[Na(diglyme)][Na\{(2,4,6-Me_3C_6H_2)_3V\}_2(\mu-N_2)]$  (**6**).<sup>27</sup> In the latter complex, a  $\mu-N_2$  ligand further interacts with a  $Na^+$  cation in a side-on manner.



By the finding of these V complexes, only Pd and Pt are left as the metals involved in groups 4 to 10 for which no stable  $N_2$  complexes have been isolated. With respect to f-block elements, two Sm complexes with a  $\mu-\eta^2:\eta^2-N_2$  ligand,  $[(Cp^*Sm)_2(\mu-N_2)]$  (**7**)<sup>28</sup> and  $[{(THF)_2Li(OEPG)Sm}_2(N_2)Li_4]$  (**8**),<sup>29</sup> are known. Another  $N_2$  complex to be noted is a  $[{(THF)_3Li}_2(\mu-N_2)]^{2+}$  cation (**9**).<sup>30</sup> Although interaction of  $Li^+$  cation with  $N_2$  ligand has been demonstrated already in certain transition metal complexes, the  $N_2$  ligand bound only to non-transition metals had been unprecedented.




## B. Ancillary Ligands

It is well known that the formation and structures of  $N_2$  complexes are very sensitive to the nature of ligands such as phosphines. For example, the cis or trans geometry of Mo and W dinitrogen complexes of the type  $[M(N_2)_2(P)_4]$ , prepared from  $WCl_6$ ,  $MoCl_5$ , or  $MCl_4(P)_2$  by the reduction with Mg in the presence of an appropriate phosphine, is controlled by subtle factors. Thus the  $N_2$ -diphosphine complexes with  $(P)_4 = (dppe)_2$ ,  $(dppp)_2$ ,  $(dppe)(dppm)$ ,  $(dppe)(dppp)$ ,  $(dppe)(depe)$ , etc.,<sup>9,31,32</sup> as well as the  $PPh_2Me$  complexes<sup>33</sup> are trans, while the  $N_2$  complexes containing  $dppm$ <sup>34</sup> and  $PMe_3$ <sup>35</sup> are cis. Interestingly,  $[Mo(N_2)_2(dppm)_2]$  obtained from  $Mo(acac)_3/AlEt_3$  under  $N_2$  proved to be trans.<sup>9b</sup> It is to be noted that for  $Na[V(N_2)_2(dmpe)_2]$ , which is isoelectronic with these Mo(0) and W(0) complexes, both the cis and trans isomers are formed from the reaction of  $[VCl_3(THF)_3]$  with Na in the presence of dmpe under  $N_2$ . The cis and trans product ratio depends on the temperature, and the ratio of 12:1 at 195 K changes to 4:1 at 250 K and then to 3:1 at 290 K. The interconversion between two isomers with varying temperature is reversible.<sup>23c</sup>


The recent study of the syntheses of W(0) complexes containing  $PR_2Ph$  ( $R = Et, Pr^i$ ) ligands disclosed the formation of three different types of  $N_2$  complexes:  $[W(N_2)_2(PR_2Ph)_3]_2(\mu-N_2)$ , *trans*- $[W(N_2)_2(PR_2Ph)_4]$ , and  $[W(\eta^6-C_6H_5PR_2)(N_2)(PR_2Ph)_2]$ .<sup>36</sup>

This presents a sharp contrast to the fact that, for  $R = Me$ , only *cis*- $[W(N_2)_2(PMe_2Ph)_4]$  (**2b**) and its less common *trans* isomer<sup>37</sup> are known to be formed by similar reactions. Interestingly, the Mo complex *trans*- $[Mo(N_2)_2(PPr^i_2Ph)_4]$  has proved to be converted into a novel mononuclear tris(dinitrogen) complex *mer*- $[Mo(N_2)_3(PPr^i_2Ph)_3]$  on being dissolved in THF.<sup>38</sup>

Typical ancillary ligands widely utilized for the syntheses of  $N_2$  complexes may be organophosphines, amines, and cyclopentadienyl groups. However, recent studies have revealed that more diversified types of ligands can be used to isolate stable  $N_2$  complexes. One important class of ligands is those containing sulfur atoms. The importance of sulfur coligands apparently arises from the structure of  $N_2$  binding site proposed for nitrogenases, which consists of the metal centers surrounded by sulfur atoms. However, well-characterized  $N_2$  complexes having sulfur donor ligands are still limited, and coordination of  $N_2$  to the multimetallic core with sulfide ligand(s) has not yet been demonstrated. The complexes reported to date include the relatively low-valent complexes *trans*- $[Mo(N_2)_2(Me_8[16]aneS_4)]$  (**10**)<sup>39</sup> and  $[OsCl(SC_6F_5)(N_2)(PMe_2Ph)_3]$ <sup>40</sup> as well as the higher valent complexes  $[Re(N_2)(SC_6H_2Pr^i-2,4,6)_3(PPh_3)]$ ,<sup>41</sup>  $[{Nb(S_2CNET_2)_3}_2(\mu-N_2)]$  (**11a**),<sup>42</sup>  $[Cp^*WMe_2(SR)]_2(\mu-N_2)$ ,<sup>43</sup> and  $[Ta(SR)_3(THF)]_2(\mu-N_2)$ .<sup>44</sup> Comparison of electrochemical, spectral, and structural data for **10** with those of **1a** suggests that the  $\{Mo(Me_8[16]aneS_4)\}$  unit can exhibit the stronger  $\pi$ -basic character than the related phosphine moiety  $\{Mo(dppe)_2\}$  and high reactivities observed for the  $N_2$  ligand in **10** described later may, at least in part, be ascribed to the strong back-donation from the metal to the  $N_2$  ligand. As for the reactivities of the other complexes shown above, few are reported to date except for the protonation of **11a** (vide infra).




Other  $N_2$  complexes containing biologically important ancillary ligands are those with porphyrins and pyrazolylborate. Mono- and dinuclear Ru complexes  $[Ru(TMP)(N_2)(L)]$  ( $L = N_2, THF, DMF, NEt_3$ )<sup>45</sup> and  $[Ru_2(DPB)(Im^*)_2(\mu-N_2)]$  (**12**)<sup>46</sup> have been isolated, and




the structure of the former ( $L = THF$ ) has been determined in detail by the X-ray analysis, while two

dinuclear complexes  $[\{[\text{HB}(3,5\text{-Me}_2\text{C}_3\text{N}_2\text{H})_3]\text{Tc}(\text{CO})_2\}_2(\mu\text{-N}_2)]$  (**13**)<sup>47</sup> and  $[\{[\text{HB}(3\text{-Pr}^i\text{-5-MeC}_3\text{N}_2\text{H})_3]\text{Co}\}_2(\mu\text{-N}_2)]$ <sup>48</sup> have appeared recently as the first examples of the  $\text{N}_2$  complex with pyrazolylborate ligand. It is also to be noted that several  $\text{N}_2$  complexes of early transition metals such as V (vide supra) and Ti  $[\{[(\text{Me}_3\text{Si})_2\text{N}]_2\text{Ti}\}_2(\mu\text{-N}_2)]^-$  (**14**)<sup>49</sup> are available by employment of sterically congested amido ligands, which demonstrate another new class of complexes containing N-donor ligands solely or predominantly. Novel mono- and dinuclear  $\text{N}_2$  complexes of Mo with triamidoamine ligand  $[\text{C}_6\text{F}_5\text{-N}_3\text{N}]$  have also been prepared recently, although the X-ray structure is available only for the silyldiazenido derivative  $[[\text{C}_6\text{F}_5\text{-N}_3\text{N}]\text{Mo}(\text{NNSiPr}^i_3)]$  obtained from  $[[\text{C}_6\text{F}_5\text{-N}_3\text{N}]\text{Mo}(\text{N}_2)\{\text{Na}(\text{ether})_x\}]$  and  $\text{Pr}^i_3\text{SiCl}$ .<sup>50</sup> Quite recently the dinuclear  $\text{N}_2$  complex containing a triamidoamine ligand with  $\text{Bu}^t\text{Me}_2\text{Si}$  substituent  $[\{[\text{Bu}^t\text{Me}_2\text{Si-N}_3\text{N}]\text{Mo}\}_2(\mu\text{-N}_2)]$  (**15**) was successfully isolated and characterized by X-ray crystallography.<sup>51</sup>




In addition to the thioether, porphyrin, and pyrazolylborate complexes shown above, a significant number of  $\text{N}_2$  complexes having tridentate or quadridentate ligands have appeared recently. Most of these contain polyphosphines or modified polyphosphines in which one P atom is replaced by other heteroatoms such as N, O, or S. Among the complexes of this type, those exhibiting interesting properties or reactivities will be referred later.

Another  $\text{N}_2$  complex to be noted with respect to the nature of coligand is the resin-supported  $\text{N}_2$  complex  $\text{trans-}[\text{Mo}(\text{N}_2)_2(\text{PS-diphos})(\text{PPh}_2\text{Me})_2]$  (**16**).<sup>52</sup> The  $\text{N}_2$  ligand attached to Mo anchored to polystyrene-divinylbenzene (2%) resin through a  $\text{CH}_2\text{P}(\text{Ph})\text{CH}_2\text{-CH}_2\text{PPh}_2$  moiety is susceptible to protonation, affording hydrazido(2-) ligand or hydrazine. Polymer-bound  $\text{N}_2$  complexes of Mo,<sup>53</sup> Co,<sup>54</sup> and Mn<sup>55</sup> were also reported earlier, but the reactions of coordinated  $\text{N}_2$  had not been uncovered in a well-defined manner.




### C. Coordination Modes

Among the various binding modes of  $\text{N}_2$  to transition metals summarized in Figure 2, a terminal end-on type (i) is most ubiquitous. There exist many



**Figure 2.** Binding modes of  $\text{N}_2$  in mononuclear and multinuclear complexes.


structurally characterized complexes of this type, and their N-N bond distances are essentially similar at around 1.12 Å irrespective of the nature of metals and coligands.<sup>10,17</sup> The other well-known coordination mode is an end-on bridging type with the almost linear M-N-N-M linkage (ii). For group 4 and 5 metal complexes, type ii is commonly observed and type i is rather rare. Most of these  $\mu\text{-}\eta^1\text{:}\eta^1\text{-N}_2$  complexes are homometallic, having a symmetrical structure,<sup>18</sup> but some heterometallic complexes are also known. In the latter are included the Re/Mo,<sup>56</sup> W/Mo,<sup>57</sup> W/Ta,<sup>58</sup> and W/Zr<sup>59</sup> complexes analyzed by X-ray crystallography. The  $\mu\text{-}\eta^1\text{:}\eta^1\text{-N}_2$  complexes may formally be classified into three types, viz. those containing the dinitrogen ( $\text{N}_2$ )<sup>0</sup>, diazenido(2-) ( $\text{N}_2$ )<sup>2-</sup>, or hydrazido(4-) ( $\text{N}_2$ )<sup>4-</sup> ligand, which may be represented by the form ii-a, ii-b, or ii-c, respectively. Although the choice of the formalism often becomes arbitrary, complexes formulated as ii-a show relatively short N-N distances (1.12–1.20 Å), while the N-N bonds in those represented by ii-c are much longer and fall in the range of 1.25–1.34 Å.<sup>18</sup> For  $[\text{W}(\text{PMe}_2\text{Ph})_3(\text{py})(\mu\text{-N}_2)\text{ZrCp}_2\text{Cl}]$  (**17**), the electron count around two metals as well as the bonding parameters associated with the M-N-N-M moiety might suggest that the  $\text{N}_2$  binding mode in **17** can be described by ii-b as a predominant form.<sup>59</sup>



Complexes with a side-on bridging  $\text{N}_2$  ligand are also known. Examples of this type initially reported were Ni complexes consisting of a folded  $\text{M}_2\text{N}_2$  ring (iii).<sup>60</sup> More recently dititanium complex **14** has been isolated, whose X-ray study has revealed the presence of two  $\mu\text{-}\eta^2\text{:}\eta^2\text{-N}_2$  ligands of this type iii.<sup>49</sup> The N-N bond in this complex is significantly long at 1.38(2) Å. Several side-on bridging  $\text{N}_2$  complexes with a

planar  $M_2N_2$  moiety (**iv**) have appeared recently. In the Sm complex **7**,<sup>28</sup> the N–N distance (1.09(1) Å) is comparable to those of free  $N_2$  (1.10 Å) and the  $N_2$  bound only to two  $Li^+$  cations in **9** (1.06(1) Å).<sup>30</sup> In contrast, the Zr complex  $[\{(Pr^i_2PCH_2SiMe_2)_2N\}ZrCl]_2(\mu-N_2)$  (**18**) has a remarkably long N–N bond at 1.55(1) Å.<sup>61</sup> This and the Sm complex **8**<sup>29</sup> with the N–N bond at 1.525(4) Å, in which the planar  $Sm_2N_2$  core is present but the  $\mu-\eta^2:\eta^2-N_2$  ligand is further bound to four  $Li^+$  cations, are those containing the most elongated N–N units reported to date. The unusually long feature of these N–N bonds is apparent if compared with the N–N distance of 1.47 Å in  $N_2H_4$ .

For the  $\mu-\eta^2:\eta^2-N_2$  complex **18** and its Cp derivative containing the  $\mu-\eta^1:\eta^1-N_2$  ligand  $[\{(Pr^i_2PCH_2SiMe_2)_2N\}ZrCp]_2(\mu-N_2)$  (**19**), it has been shown that the




resonance Raman spectroscopy can be used as a good measure to distinguish the two modes of  $N_2$  coordination, *viz.* side-on bridging for the former and end-on bridging for the latter. Full assignment of the peaks associated with the  $N_2$  moiety has been done by comparing the spectra of **18** and **19** with those of  $^{15}N$  analogues. In contrast to the spectrum of **18** exhibiting a single intense peak at  $1211\text{ cm}^{-1}$ , which can be ascribed to the symmetrical  $\nu(NN)$ , that of **19** is more complicated due to the presence of many isotope-sensitive peaks, among which the strong  $731\text{ cm}^{-1}$  peak can be assigned to a totally symmetric mode composed primarily of  $\nu(NN)$  character.<sup>62</sup> Apparent differences in the spectral features between these two types of  $N_2$  ligands may indicate that the resonance Raman spectroscopy provides a convenient method for determining the  $N_2$  bridging mode unambiguously.

*Ab initio* quantum calculations performed on the  $M_2N_2$  unit have demonstrated that the unusually long N–N distance observed in **18** can be interpreted in terms of the metal interaction with both the in-plane and the out-of-plane  $\pi^*$ -orbitals of  $N_2$ . It has also been concluded that such type of state is stable only for the metals on the very left of the periodic table and not the first-row metals, *i.e.* for Zr and Y.<sup>63</sup>

Another fully characterized coordination mode of  $N_2$  was found in the Ti complex  $[\{(C_{10}H_8)Cp_2Ti_2\}\{(C_5H_4)Cp_3Ti_2\}(\mu_3-N_2)]$  (**20**) reported much earlier, in which the  $N_2$  unit with a N–N distance at 1.30(1) Å exists as a  $\mu_3-\eta^1:\eta^1:\eta^2$  ligand (**v**).<sup>64</sup> Side-on coordination of  $N_2$  to a single metal center (**vi**) has not yet been demonstrated, although such a species was proposed as the intermediate of the end-to-end rotation of coordinated  $\eta^1-N_2$  ligand in  $[Cp^*Re(CO)_2(N_2)]$ .<sup>65</sup> The Zr complex  $[Cp_2Zr(CH_2SiMe_3)(N_2)]$  was

suggested to contain this  $\eta^2-N_2$  ligand on the basis of EPR and IR data,<sup>66</sup> but details are still uncertain.



## D. Formation of Dinitrogen Ligands


Not only  $N_2$  gas but numerous nitrogenous moieties can be used to synthesize  $N_2$  complexes. In fact, the  $N_2$  ligand in the first  $N_2$  complex  $[Ru(NH_3)_5(N_2)]^{2+}$  had its origin in hydrazine. Although  $N_2$  complexes derivable from molecular  $N_2$  seem more attractive in view of the development of effective chemical  $N_2$ -fixing systems, those prepared by indirect methods are also of importance in their own right since they also can provide much fundamental information associated with the  $N_2$  chemistry.

Some terminal end-on  $N_2$  ligands have been derived from certain organonitrogen ligands. Thus  $[RhCl(N_2)(PPr^i_3)_2]$  has been prepared by thermal decomposition of the diazoalkane complex  $[RhCl(N_2-CPh_2)(PPr^i_3)_2]$ ,<sup>67</sup> while  $[Cp^*Re(CO)(L)(N_2)]$  has been obtained by treatment of the aryldiazenido complex  $[Cp^*Re(CO)(L)(NNC_6H_4OMe-p)] [BF_4]$  with  $Bu^tLi$  ( $L$  = phosphines, phosphites).<sup>68</sup> Conversion of an organodiazenido ligand into coordinated  $N_2$  had been manifested earlier for the syntheses of a series of Re complexes of the type  $[ReCl(N_2)(L)_4]$  from  $[ReCl_2(NNCOPh)(PPh_3)_2]$ .<sup>69</sup>

For  $\mu-\eta^1:\eta^1-N_2$  complexes, several new synthetic routes have been developed. These include firstly the coupling of two  $N_1$  moieties; Os complexes  $[(pic)_2Cl_3-Os(\mu-N_2)OsCl(pic)_4]^+$ <sup>70</sup> and  $[\{Os(NH_3)_4(MeCN)\}_2(\mu-N_2)]^{5+}$ <sup>71</sup> have been obtained by dissolving  $[OsNCl_4]^-$  in neat pic for the former and upon photoirradiation of a mixture of  $[Os(NH_3)_4N]^{3+}$  and hexamethylbenzene in MeCN for the latter, respectively, while the oxidative coupling of two ammine ligands in  $[(NH_3)_2-Ru_2(DPB)(Im^*)_2]$  has afforded a  $N_2$  complex of a cofacial metallodiporphyrin (**12**).<sup>46b</sup>

On the other hand, hydrazine and azine are also used as the sources of the  $\mu-\eta^1:\eta^1-N_2$  ligand, although formation of a terminal  $N_2$  ligand from  $N_2H_4$  has been well demonstrated, for example, for the Ru complex  $[Ru(NH_3)_5(N_2)]^{2+}$  shown above and  $[CpM(CO)_2(N_2)]$  ( $M = Mn$ ,<sup>72</sup>  $Re$ <sup>73</sup>) obtained from  $[CpM(CO)_2(N_2H_4)]/H_2O_2/Cu^{2+}$ . Thus the reaction of  $N_2H_4$  with alkoxide complex  $[W(PhCCPh)(OBu^t)_4]$  gives a  $\mu-N_2$  complex  $[\{W(PhCCPh)(OBu^t)_2\}_2(\mu-N_2)]$  accompanied by the liberation of  $Bu^tOH$ .<sup>74</sup> Essentially the same method has been used for preparing  $[(Cp^*MoMe_3)_2(\mu-N_2)]$  (**21a**) from  $[Cp^*MoMe_3(OC_6H_3Pr^{i-2,6})][PF_6]$  and  $N_2H_4$ ,<sup>57</sup> which has further been extended to the reaction of this aryloxy complex with a hydrazido-(2-) complex  $[Cp^*WMe_3(NNH_2)]$  to give a heterometallic complex  $[(Cp^*MoMe_3)(\mu-N_2)(Cp^*WMe_3)]$ .<sup>57</sup> As closely related reactions, formation of  $[(Cp^*WMe_3)_2-$

( $\mu$ -N<sub>2</sub>) (**21b**) or [(Cp\*WMe<sub>3</sub>)( $\mu$ -N<sub>2</sub>)(Cp\*TaMe<sub>2</sub>)] from a diazenido complex [{Cp\*WMe<sub>3</sub>(NNH)}<sub>2</sub>{ $\mu$ -Mg(T-MEDA)}] with [Cp\*WMe<sub>4</sub>][PF<sub>6</sub>] or [Cp\*TaMe<sub>3</sub>Cl], respectively, has been reported,<sup>58</sup> in which the diazenido proton is eliminated as CH<sub>4</sub>. Preparation of  $\mu$ -N<sub>2</sub> complexes by the use of the azine as a N<sub>2</sub> source was reported earlier, as shown, for example, in eq 2.<sup>75</sup>



Employment of the electroreductive conditions for the synthesis of low-valent complexes is less common, compared with the use of chemical reduction. Most of the low-valent N<sub>2</sub> complexes are prepared from the higher valent precursors in the presence of reductants such as Na, Mg, and Zn under N<sub>2</sub>. For complexes **1**, it has been shown that the controlled potential electrolysis of the solutions containing a range of precursors including oxide,<sup>76</sup> imide,<sup>77</sup> hydrazido(2-),<sup>78</sup> halide, and thiolate<sup>79</sup> complexes under N<sub>2</sub> affords the N<sub>2</sub> complexes in satisfactory yields.


### III. Protonation Reactions of Coordinated Dinitrogen

As expected, molecular nitrogen bound to the metal center(s) is activated more or less and in certain complexes it becomes susceptible to protonation under mild conditions. Such reactions have been extensively studied in relation to the biological N<sub>2</sub> fixation. A significant number of mono- and multinuclear N<sub>2</sub> complexes are now available which can facilitate the conversion of N<sub>2</sub> into NH<sub>3</sub> and in some cases into N<sub>2</sub>H<sub>4</sub> upon protonation by acid or, less commonly, by water. Although biological N<sub>2</sub> reduction gives NH<sub>3</sub> exclusively as the nitrogen-containing product, it should be noted that the enzyme during turnover has proved to liberate N<sub>2</sub>H<sub>4</sub> upon quench by acid or base.<sup>80</sup>

#### A. Protonation of N<sub>2</sub> Coordinated to a Single Transition Metal Center


Almost quantitative conversion of one N<sub>2</sub> ligand in **2b** into NH<sub>3</sub> by treatment with sulfuric acid in MeOH at room temperature was observed by Chatt and his co-workers in 1975,<sup>81</sup> and since then intensive studies have been made for clarifying the mechanism of N<sub>2</sub> reduction operating in this and related systems using complexes of the type [M(N<sub>2</sub>)<sub>2</sub>(P)<sub>4</sub>] (M = W, Mo; P = tertiary phosphine). The M–N<sub>2</sub> interactions in end-on N<sub>2</sub> complexes are ascribed to the synergistic  $\sigma$ -bonding and  $\pi$ -back bonding, as is well recognized for CO ligand. Molecular orbital and charge distribution analyses using the model compounds [M(N<sub>2</sub>)<sub>2</sub>-(PH<sub>3</sub>)<sub>4</sub>] have suggested stronger M–N<sub>2</sub> interactions for group 6 metals than the later transition metals with the order of W > Mo > Cr, along with the presence of significant negative charges on the N<sub>2</sub> ligand, especially on the terminal N atom.<sup>82</sup> The latter feature is in good agreement with the fact that the protonation of these W and Mo complexes pro-

Scheme 1



ceeds not at the metal center but at the terminal N atom in the N<sub>2</sub> ligand.

Although formation of NH<sub>3</sub> and/or N<sub>2</sub>H<sub>4</sub> upon protonation by acid or water has been demonstrated for a considerable number of N<sub>2</sub> complexes, these Mo and W systems are unique in that several intermediate stages have been isolated and characterized unambiguously. Accumulated data for these systems indicate that the protonation of the N<sub>2</sub> ligand proceeds in a stepwise manner, accompanied by the transfer of required electrons from the central metal to the nitrogenous ligand (Scheme 1). Considerably high yields of nitrogen hydrides presumably arise from the strong reducing power of the W and Mo atoms. The hydrazido(2-) complexes isolated widely<sup>31,83</sup> demonstrate the most important intermediate stage involved in the reduction pathway, and structural details of the hydrazido(2-) ligand have already been clarified by X-ray analyses for more than ten complexes containing dppe,<sup>83a-d</sup> PMe<sub>2</sub>Ph,<sup>83e,f</sup> or depe<sup>83g,h</sup> as coligands. All of these complexes contain essentially linear M–N–N linkages in which the N–N bond lengths are of the bond order between 1 and 2 and the M–N distances are suggestive of the multiple bonding nature. Diazenido complexes probably formed prior to the hydrazido(2-) complexes have not yet been isolated directly from the corresponding N<sub>2</sub> complexes but were derived from hydrazido(2-) complexes and weak base (eq 3).<sup>84</sup> Because of the lack of X-ray structure determination, a detailed structure of the diazenido ligand in these complexes is still unknown.



After the formation of the hydrazido(2-) complexes, reduction is believed to follow the different reaction courses, depending upon the nature of the metal and coligand as well as the reaction conditions. Thus, as proposed for the reaction of **2b** with sulfuric


acid in MeOH, further protonation at the terminal nitrogen atom gives a hydrazidium complex and then the following N–N bond scission affords a nitrido complex and NH<sub>3</sub>. Additional NH<sub>3</sub> is produced by subsequent acidolysis of the nitrido complex with concurrent formation of M(VI) species.<sup>81b</sup> Recently the first hydrazidium complex *trans*-[WCl(NNH<sub>3</sub>)(PMe<sub>3</sub>)<sub>4</sub>]Cl<sub>2</sub> was isolated from the reaction of either *cis*-[W(N<sub>2</sub>)<sub>2</sub>(PMe<sub>3</sub>)<sub>4</sub>] or [WCl(NNH<sub>2</sub>)(PMe<sub>3</sub>)<sub>4</sub>]Cl with HCl gas and fully characterized, although the reaction of this hydrazidium complex with HCl in MeOH gives neither NH<sub>3</sub> nor N<sub>2</sub>H<sub>4</sub>.<sup>85</sup>

In the second reaction course, protonation of the hydrazido(2–) complex occurs at the central metal, generating a hydrido–hydrazido(2–) complex. Subsequent protonation finally results in the predominant formation of N<sub>2</sub>H<sub>4</sub> and M(IV) species. Such a mechanism may operate, for example, in the reaction of **2** (M = W, Mo) with HCl gas,<sup>86</sup> and several hydrido–hydrazido(2–) complexes have been isolated and fully characterized.<sup>86,87</sup>

The third reaction course involves the disproportionation of two hydrazido(2–) ligands, forming NH<sub>3</sub> and N<sub>2</sub> together with M(III) product. In contrast to the first and second mechanisms above, this mechanism is essentially intermolecular, where one hydrazido(2–) complex acts as a reducing agent for the other. Reactions of Mo complexes, *e.g.* *trans*-[Mo(N<sub>2</sub>)<sub>2</sub>(dpepp)(PPh<sub>3</sub>)<sub>2</sub>]<sup>88</sup> and *trans*-[Mo(N<sub>2</sub>)<sub>2</sub>(dppe)(PPh<sub>2</sub>Me)<sub>2</sub>] (**22**), with HBr in THF and protonolysis of the hydrazido(2–) complexes derived from these<sup>44</sup> are considered to follow this pathway. This can be supported by the facts that the N<sub>2</sub> complex **16** as well as the hydrazido(2–) complex [MoF(NNH<sub>2</sub>)(PS-diphos)(PPh<sub>2</sub>Me)<sub>2</sub>][BF<sub>4</sub>], in which the direct interaction between the N<sub>2</sub> ligands or the hydrazido(2–) ligands hardly takes place, give no NH<sub>3</sub> by the reaction with HBr in THF, whereas the addition of soluble Mo or W hydrazido(2–) complexes as the reductant results in the production of NH<sub>3</sub> from the polymer-bound hydrazido(2–) moiety.<sup>52</sup>

These findings have subsequently led to the observation that, in the reactions of both the polymer-bound and the soluble Mo dinitrogen complexes **16**, **22**, and *trans*-[Mo(N<sub>2</sub>)<sub>2</sub>(dppe)(PPh<sub>3</sub>)] with HBr, addition of two electron reducing agents such as Sn(II) and Ge(II) halides considerably increases the yields of NH<sub>3</sub>, since all of the hydrazido(2–) ligand generated in this system can be used as the NH<sub>3</sub> source due to the presence of the added reductant.<sup>83c,89</sup> As a related reaction, the two-electron reduction of [W(OTs)(NNH<sub>2</sub>)(dppe)<sub>2</sub>]<sup>+</sup> under the conditions of controlled potential electrolysis have been shown to give NH<sub>3</sub>. An interesting feature of the present system is the recovery of the parent N<sub>2</sub> complex *trans*-[W(N<sub>2</sub>)<sub>2</sub>(dppe)<sub>2</sub>] (**1b**) in high yield, which in principle allows for repetition of the cycle involving (a) the protonation of the N<sub>2</sub> complex to the hydrazido(2–) complex and (b) electroreduction of the hydrazido(2–) complex forming NH<sub>3</sub> accompanied by regeneration of the N<sub>2</sub> complex (eq 4). However, the total amount of NH<sub>3</sub> produced after three cycles in one vessel has been reported to be 0.72 mol/W atom.<sup>78</sup> Further, it should be noted that the evolution of a

large amount of H<sub>2</sub> takes place during the electrochemistry when acid is present, *e.g.* in the cycle.



The V(–1) complexes **4** and its dmpe analogue, which are isoelectronic with Mo(0) and W(0) complexes shown above, have been reported to give 1 mol of NH<sub>3</sub> per mole of V by treatment with HBr gas.<sup>23b</sup> Concomitant oxidation of the V(–1) center to the V(II) state has been suggested.

Recent elucidation of the nitrogenase active site structure cited above has evoked renewed interest in the chemistry of Fe dinitrogen complexes. Mono- and dinuclear complexes of Fe containing a terminal end-on or bridging N<sub>2</sub> ligand have been known since the early 1970s, but reactivities of coordinated N<sub>2</sub> in these complexes have not been demonstrated except for the replacement by certain  $\pi$ -acceptor ligands. Only the poorly defined complex [{FeEt(N<sub>2</sub>)(PPh<sub>3</sub>)<sub>2</sub>]<sub>2</sub>Mg(THF)<sub>4</sub>] was previously shown to give N<sub>2</sub>H<sub>4</sub> (0.11 mol/Fe atom) and NH<sub>3</sub> (0.11 mol/Fe atom).<sup>90</sup> Recent investigation has clarified that the Fe(0) complex [Fe(N<sub>2</sub>)(dmpe)<sub>2</sub>] generated in situ from [FeH(N<sub>2</sub>)(dmpe)<sub>2</sub>] and KOBu<sup>t</sup> produces NH<sub>3</sub> (0.06–0.12 mol/Fe atom) upon protonation by H<sub>2</sub>SO<sub>4</sub> or HCl.<sup>91</sup> The resultant major Fe species was assigned as [FeCl<sub>2</sub>(dmpe)<sub>2</sub>] for the latter system. Formation of comparable amounts of NH<sub>3</sub> was also observed for [Fe(N<sub>2</sub>)(depe)<sub>2</sub>] analogously obtained in situ from [FeH(N<sub>2</sub>)(depe)<sub>2</sub>]. Very recently, isolation and full characterization of [Fe(N<sub>2</sub>)(depe)<sub>2</sub>] have been carried out by a different group, and somehow the isolated complex gave no NH<sub>3</sub> but only N<sub>2</sub> and H<sub>2</sub> gases by acidolysis.<sup>92</sup>

## B. Protonation of N<sub>2</sub> Coordinated to More Than Two Transition Metals


Since the complete reduction of coordinated N<sub>2</sub> into NH<sub>3</sub> or N<sub>2</sub>H<sub>4</sub> by protonolysis requires the concomitant transfer of six or four electrons, formation of these nitrogen hydrides in high yields from mononuclear N<sub>2</sub> complexes may be expected only for N<sub>2</sub> bound to a strongly reducing metal, which can supply the multielectrons to the coordinated N<sub>2</sub>. This presumably accounts, at least in part, for the highly effective NH<sub>3</sub> and N<sub>2</sub>H<sub>4</sub> production observed for zero-valent W and Mo complexes described above. On the other hand, multimetallic activation of N<sub>2</sub> may require less electrons per one metal for the formation of nitrogen hydrides, and therefore more diversified metals are possibly used for this transformation.

Actually a considerable number of multinuclear N<sub>2</sub> complexes have been proven to afford NH<sub>3</sub> or N<sub>2</sub>H<sub>4</sub> by protonolysis under mild conditions. Thus the previous reports described the formation of NH<sub>3</sub> from a binuclear Ni complex dimer [{(LiPh)<sub>3</sub>(Et<sub>2</sub>O)<sub>1–1.5</sub>Ni}]<sub>2</sub>( $\mu$ - $\eta^2$ : $\eta^2$ -N<sub>2</sub>)<sub>2</sub><sup>93</sup> and a tetranuclear  $\mu_3$ -N<sub>2</sub> Ti complex **20**<sup>64</sup> as well as the formation of N<sub>2</sub>H<sub>4</sub> from a binuclear Zr complex [{Cp\*<sub>2</sub>Zr(N<sub>2</sub>)<sub>2</sub>( $\mu$ -N<sub>2</sub>)]<sup>94</sup> and a series of binuclear Ta complexes, *e.g.* [{TaCl<sub>3</sub>(THF)<sub>2</sub>]<sub>2</sub>( $\mu$ -N<sub>2</sub>)].<sup>75</sup> In addition, recent studies have demonstrated the formation of NH<sub>3</sub> from **21** (M = Mo,


**Table 1. Protonation of Bridging Dinitrogen Ligands Forming Ammonia and/or Hydrazine**

complex	acid/solvent	product (%) <sup>a</sup>
$[\{(C_{10}H_8)Cp_2Ti_2\}(\{C_5H_4\}Cp_3Ti_2)(\mu_3-N_2)]$ ( <b>20</b> )	H <sub>2</sub> O	N <sub>2</sub> H <sub>4</sub> (major) + NH <sub>3</sub> (total 90)
$[\{Cp^*Zr(N_2)_2(\mu-N_2)\}]$	HCl/toluene	N <sub>2</sub> H <sub>4</sub> (33) <sup>b</sup>
$[\{(o-Me_2NCH_2C_6H_4)_2V(py)\}_2(\mu-N_2)]$ ( <b>3</b> )	HCl	NH <sub>3</sub> (33)
$[Na\{(2,4,6-Me_3C_6H_2)_3V\}_2(\mu-N_2)]^-$ ( <b>6</b> )	HCl/H <sub>2</sub> O	N <sub>2</sub> H <sub>4</sub> (36) + NH <sub>3</sub> (11)
$[\{Nb(S_2CNEt_2)_3\}_2(\mu-N_2)]$ ( <b>11b</b> )	HCl/solvent <sup>c</sup>	N <sub>2</sub> H <sub>4</sub> (>90)
$[\{TaCl_3(THF)_2\}_2(\mu-N_2)]$	HCl/ether	N <sub>2</sub> H <sub>4</sub> (80)
$[\{Ta(CH_2Bu^t)_3(THF)_2\}_2(\mu-N_2)]$	HCl/ether	N <sub>2</sub> H <sub>4</sub> (86)
$[(Cp^*MoMe_3)_2(\mu-N_2)]$ ( <b>21a</b> )	HCl/THF	NH <sub>3</sub> (16)
$[(Cp^*WMe_3)_2(\mu-N_2)]$ ( <b>21b</b> )	HCl/THF	NH <sub>3</sub> (17)
$[\{(LiPh)_3(Et_2O)_{1-1.5}Ni\}_2(\mu-N_2)]_2$	H <sub>2</sub> O	NH <sub>3</sub> (33)
$[WCl(PMe_2Ph)_4(\mu-N_2)TiCp_2Cl]$	H <sub>2</sub> SO <sub>4</sub> /MeOH	N <sub>2</sub> H <sub>4</sub> (8) + NH <sub>3</sub> (43)
$[Wl(PMe_2Ph)_4(\mu-N_2)ZrCp_2Cl]$	H <sub>2</sub> SO <sub>4</sub> /MeOH	N <sub>2</sub> H <sub>4</sub> (16) + NH <sub>3</sub> (17)

<sup>a</sup> Based on total nitrogen in the N<sub>2</sub> complex. <sup>b</sup> A half of N<sub>2</sub>H<sub>4</sub> results from the initial  $\mu$ -N<sub>2</sub> ligand. <sup>c</sup> THF, CH<sub>2</sub>Cl<sub>2</sub>, or MeCN.

**Scheme 2**

11a: M = Nb,  
11b: M = Ta



W)<sup>43,57</sup> and V complex **3**,<sup>95</sup> the formation of N<sub>2</sub>H<sub>4</sub> from  $[\{M(S_2CNEt_2)_3\}_2(\mu-N_2)]$  (M = Nb, **11a**; M = Ta, **11b**)<sup>96</sup> and Zr complexes **18** and **19**,<sup>61b</sup> and the formation of both N<sub>2</sub>H<sub>4</sub> and NH<sub>3</sub> from V complex **6**<sup>27</sup> and heterobimetallic complexes  $[WX(PMe_2Ph)_4(\mu-N_2)MCp_2Cl]$  (M = Ti, X = Cl; M = Zr, X = I).<sup>59</sup> These results are summarized in Table 1.

Interestingly, by reaction with HCl, the V alkyl complex **5** liberates the whole coordinated N<sub>2</sub> as a N<sub>2</sub> gas with concurrent formation of neopentane,<sup>24</sup> indicating the attack by proton occurs on the  $\alpha$ -carbon atom, while related Ta complex  $[\{Ta(CH_2Bu^t)_3(THF)_2\}_2(\mu-N_2)]$  is treated similarly to afford N<sub>2</sub>H<sub>4</sub> in moderate yield.<sup>75</sup> It should also be noted that the multinuclear Ni and Ti dinitrogen complexes give NH<sub>3</sub> by treatment with such a weak acid as water (Table 1), and in the case of **20** the reaction with HCl instead of water surprisingly results in the evolution of N<sub>2</sub> gas exclusively.<sup>64</sup>

Despite the appearance of many multinuclear N<sub>2</sub> complexes giving nitrogen hydrides, none of these have given any isolable intermediate stages, and the mechanism of the protonation operating on the multimetallic center is still uncertain. It may be quite important to clarify until which stage the multinuclearity of the complex is retained. However, even such a fundamental feature has been specified for only a few of the systems reported to date. For only the Nb and Ta complexes **11** has it been suggested by spectroscopic data that the protonation of N<sub>2</sub> proceeds in a stepwise manner with retention of a dinuclear core until the  $\mu$ -NHNH<sub>2</sub> stage (Scheme 2).<sup>18,96</sup> Another important finding associated with this system is that the proton transfer to the bridging N<sub>2</sub> in **11** is much slower than that of the terminal N<sub>2</sub> in mononuclear  $[M(N_2)_2(P)_4]$ .<sup>97</sup> This significant dif-


ference in the reactivity has been accounted for by attack by a proton at the electrons delocalizing over the M-N-N-M moiety in the former, which is in contrast to the proton attack at the lone-pair electrons localized on the terminal N atom in the latter.

**IV. Reactions of Hydrazines Relevant to Nitrogen Fixation**

Nitrogenase is a catalyst capable of reducing numerous substrates, and it has been shown that free N<sub>2</sub>H<sub>4</sub> is also reduced to NH<sub>3</sub>, *i.e.* N<sub>2</sub>H<sub>4</sub> + 2H<sup>+</sup> + 2e<sup>-</sup> → 2NH<sub>3</sub>.<sup>98</sup> Reactivities of N<sub>2</sub>H<sub>4</sub> toward transition metal complexes are therefore of much interest.


The high-valent mononuclear hydrazine complex  $[Cp^*WMe_3(\eta^2-NH_2NH_2)][OTf]$  (**23**) prepared from  $[Cp^*WMe_3(OTf)]$  and N<sub>2</sub>H<sub>4</sub> undergoes reduction by Na/Hg to give NH<sub>3</sub> and  $[Cp^*WMe_3(NH)]$ . Almost stoichiometric formation of NH<sub>3</sub> has been observed when **23** is reduced by Zn/Hg or Cp<sub>2</sub>Co in the presence of lutidine hydrochloride as a proton source. This system has successfully been extended to the catalytic reaction converting N<sub>2</sub>H<sub>4</sub> into NH<sub>3</sub>. Thus, when **23** is reduced in the presence of 10 equiv of N<sub>2</sub>H<sub>4</sub>, the yield of NH<sub>3</sub> reaches ca. 16 mol/W atom (Scheme 3).<sup>99</sup> Essentially the same result has been obtained also for the Mo analogue.<sup>100</sup>

In certain systems without external reduction, coordinated N<sub>2</sub>H<sub>4</sub> can be converted into NH<sub>3</sub> by the disproportionation reaction, *i.e.* 3N<sub>2</sub>H<sub>4</sub> → 4NH<sub>3</sub> + N<sub>2</sub>. Recently we have shown that the coordinatively unsaturated diruthenium complex  $[Cp^*Ru(\mu-SPR^d)_2-RuCp^*]$  catalyzes this reaction effectively.<sup>101</sup> Spec-

**Scheme 3**




Scheme 4



trosopic data of the reaction mixture suggest the involvement of the  $\mu$ -diazene complex  $[\text{Cp}^*\text{Ru}(\mu\text{-SP}^{\text{r}})_2(\mu\text{-NHNH})\text{RuCp}^*]$  as an intermediate. The catalytic cycles proposed for this Ru system are shown in Scheme 4.

The reaction of silylated 2-pyridinethiol 2-HSC<sub>5</sub>H<sub>3</sub>N-3-SiMe<sub>3</sub> with MoCl<sub>5</sub> or [MoCl<sub>4</sub>(MeCN)<sub>2</sub>] has been shown to give mononuclear 1*H*-pyridine-2-thione complex [MoCl<sub>4</sub>(2-SC<sub>5</sub>H<sub>3</sub>NH-3-SiMe<sub>3</sub>)<sub>2</sub>] or a dinuclear complex with bridging disulfide and the thione ligands, [Mo<sub>2</sub>Cl<sub>4</sub>( $\mu$ -S<sub>2</sub>)( $\mu$ -2-SC<sub>5</sub>H<sub>3</sub>NH-3-SiMe<sub>3</sub>)(2-SC<sub>5</sub>H<sub>3</sub>N-3-SiMe<sub>3</sub>)<sub>2</sub>], respectively. These two complexes catalyze not only the disproportionation of hydrazine in the absence of an external reductant and a proton source but also the reduction of hydrazine in the presence of Zn/Hg and lutidine hydrochloride. Experimental evidence suggests that for the catalytic NH<sub>3</sub> production both systems using the mono- and dinuclear complexes as the catalyst precursor involve the same intermediate [Mo<sub>2</sub>Cl<sub>3</sub>(2-SC<sub>5</sub>H<sub>3</sub>N-3-SiMe<sub>3</sub>)<sub>3</sub>].<sup>102</sup>

A more sophisticated model system which relates to the reduction of N<sub>2</sub>H<sub>4</sub> in nitrogenase is found in reactions using MoFe<sub>3</sub>S<sub>4</sub> single cubane clusters [(Cl<sub>4</sub>-cat)MoFe<sub>3</sub>S<sub>4</sub>Cl<sub>3</sub>(MeCN)]<sup>2-</sup> (**24**) and [(citr)MoFe<sub>3</sub>S<sub>4</sub>Cl<sub>3</sub>]<sup>3-</sup> (**25**).<sup>103</sup> In the absence of a reducing agent and a



proton source, the disproportionation of N<sub>2</sub>H<sub>4</sub> is slowly catalyzed by the former but not by the latter. On the other hand, by addition of Cp<sub>2</sub>Co and 2,6-lutidine hydrochloride, both clusters catalyze the reduction of N<sub>2</sub>H<sub>4</sub> into NH<sub>3</sub> with good turnovers. The EPR study confirms the retention of the cluster core during the whole turnovers. From the reaction of **24** containing an easily displaceable MeCN ligand with PhNHNH<sub>2</sub>, [(Cl<sub>4</sub>-cat)MoFe<sub>3</sub>S<sub>4</sub>Cl<sub>3</sub>(NH<sub>2</sub>NHPh)]<sup>2-</sup> has been isolated<sup>103,104</sup> and the stoichiometric and catalytic reduction of the terminally bound PhNHNH<sub>2</sub> in this adduct has afforded PhNH<sub>2</sub> and NH<sub>3</sub>, while the N<sub>2</sub>H<sub>4</sub>-bridged double cubane [(Cl<sub>4</sub>-cat)MoFe<sub>3</sub>S<sub>4</sub>Cl<sub>3</sub>]<sub>2</sub>( $\mu$ -NH<sub>2</sub>NH<sub>2</sub>)<sup>4-</sup><sup>105</sup> available from **24** and N<sub>2</sub>H<sub>4</sub> exhibits only a low catalytic activity. In addition, [Fe<sub>4</sub>S<sub>4</sub>Cl<sub>4</sub>]<sup>2-</sup>

shows no catalytic activity. These results suggest that the Mo atom in the MoFe<sub>3</sub>S<sub>4</sub> single cubanes is directly involved as the reaction site in this catalysis. Furthermore, the finding that the activity observed for **25** containing the Mo atom coordinated by tridentate citrate ligand is similar to that of **24** implies that in the structurally resembling FeMo cofactor (Figure 1) the coordinatively saturated Mo atom by bidentate homocitrate as well as one N and three S atoms may also serve as the site of substrate binding and reduction. The hapticity change of the citrate ligand upon protonation, for example, might provide the coordination site for substrate reduction. It has been clarified subsequently that the double cubane cluster [MoFe<sub>3</sub>S<sub>4</sub>Cl<sub>3</sub>{SCH(CH<sub>3</sub>)COO}]<sub>2</sub><sup>4-</sup>, which contains the Mo atom coordinated by carboxylate, also exhibits a high catalytic activity. The active species in this system has also been characterized to be a single cubane cluster.<sup>106</sup>


### V. Conversion of Coordinated Dinitrogen into Organonitrogen Ligands and Compounds

One-pot synthesis of nitrogen-containing organic compounds from molecular nitrogen may open a new field in organic syntheses. Initial study associated with this area appeared in 1966, which showed the formation of PhNH<sub>2</sub> along with NH<sub>3</sub> by treatment of Cp<sub>2</sub>TiCl<sub>2</sub> or Cp<sub>2</sub>TiPh<sub>2</sub> with excess PhLi in ether at room temperature and under an atmospheric pressure of N<sub>2</sub> followed by hydrolysis. An increase in the N<sub>2</sub> pressure up to 80–100 bar considerably raised the yield of PhNH<sub>2</sub> (0.10–0.15 mol/Ti atom).<sup>107</sup> However, the details of this reaction are unknown due to the lack of any information about the intermediate species. Several C–N bond-forming reactions that proceed in well-defined N<sub>2</sub> complexes have subsequently been clarified, and most of these have been observed for the Mo and W complexes.

### A. C–N Bond Formation at Coordinated N<sub>2</sub> by Reaction with Organic Halides

Formation of a C–N bond at the N<sub>2</sub> ligand was initially observed in the reactions of Mo and W complexes **1** with RCOCl.<sup>108</sup> The products were characterized to be acyl- and aroyldiazenido complexes *trans*-[MCl(NNCOR)(dppe)<sub>2</sub>], and the X-ray structure is available for *trans*-[MoCl(NNCOPh)(dppe)<sub>2</sub>].<sup>109</sup> A similar reaction also proceeds for *trans*-[ReCl(N<sub>2</sub>)(PMe<sub>2</sub>Ph)<sub>4</sub>].<sup>108a</sup> Another C–N bond-forming reaction defined in detail is the formation of alkyl-diazenido complexes *trans*-[MX(NNR)(dppe)<sub>2</sub>] from **1** and alkyl halides under irradiation of a W lamp.<sup>108a,110</sup> The mechanism of this alkylation involves attack of the alkyl radical on the N<sub>2</sub> ligand, which is generated by homolysis of RX occurring within the coordination sphere.<sup>111</sup> In contrast, the former acylation is now believed to proceed via nucleophilic attack of the terminal N atom on the acyl carbon.<sup>17</sup> The terminal N atoms in these diazenido ligands are significantly nucleophilic, and a series of acyl-, aroyl-, and alkyl-hydrazido(2-) complexes *trans*-[MCl(NNHCOR)(dppe)<sub>2</sub>]<sup>+</sup> and *trans*-[MX(NNHR)(dppe)<sub>2</sub>]<sup>+</sup> are readily derived upon protonation by acid. Analogously, dialkylhydrazido(2-) complexes *trans*-[MBr(NNMeEt)-

Scheme 5



$(R_2PCH_2CH_2PR_2)_2^+$  ( $M = Mo, W$ ) have been prepared for  $R = Et$  and various aryl groups from the reaction of  $trans$ - $[MBr(NNEt)(R_2PCH_2CH_2PR_2)_2]$  and  $MeI$  that proceeds via an  $S_N2$  mechanism.<sup>112</sup> One-pot dialkylation observed upon treatment of **1b** with  $MeBr$  to give  $trans$ - $[WBr(NNMe_2)(dppe)_2]Br$  is also believed to follow these two steps. A series of dialkylhydrazido(2<sup>-</sup>) complexes of the type  $trans$ - $[MBr$ - $\overline{[NN(CH_2)_nCH_2]}(dppe)_2]Br$  ( $M = Mo, W; n = 2-4$ ) are similarly available by the use of  $\alpha,\omega$ -dibromides,<sup>113</sup> while extension of this reaction to that with *gem*-dibromide has led to the isolation of diazoalkane complexes  $trans$ - $[WBr(NNCR'R')(dppe)_2]Br$  ( $R, R' = H, Me$ ).<sup>114</sup> It should be noted that the C-N bond formation at coordinated  $N_2$  by reaction with alkyl halides is only observed for diphosphine complexes such as **1** but not for monophosphine complexes such as **2**.


The much higher reactivity of the Mo tetrathioether  $N_2$  complex **10** toward organic halides compared with the phosphine complex **1a** is noteworthy. Thus **10** reacts not only with  $MeI$  but also with  $PhCH_2Br$  to give dialkylhydrazido(2<sup>-</sup>) complexes  $trans$ - $[MoX(NNR_2)(Me_8[16]aneS_4)]X$ . Furthermore, even aryl halides ( $ArX$ ) such as  $PhBr, PhI,$  and  $p$ - $MeOCOC_6H_4I$  react readily with **10** to afford novel aryldiazenido complexes  $trans$ - $[MoX(NNAr)(Me_8[16]aneS_4)]$ .<sup>115</sup> Neither benzylation nor arylation of the coordinated  $N_2$  in Mo and W phosphine complexes has been successful except for the arylation using a unique bimetallic system (vide infra).

The C-N bond-forming reactions described in this section are summarized in Scheme 5.

## B. C-N Bond Formation at Coordinated $N_2$ via Hydrazido(2<sup>-</sup>) or Diazenido Complexes

The Mo and W hydrazido(2<sup>-</sup>) complexes such as  $trans$ - $[MF(NNH_2)(dppe)_2][BF_4]$  (**26**) and  $cis,mer$ - $[MX_2(NNH_2)(PMe_2Ph)_3]$  (**27**;  $X = Cl, Br, I$ ), easily derived from the reactions of **1** with aqueous  $HBF_4$  in THF or those of **2** with aqueous  $HX$  in MeOH, are not only the important intermediate stages for the reduction of coordinated  $N_2$  but serve as the versatile precursors to prepare organodinitrogen ligands. Owing to the significant nucleophilicity of the terminal N atom, the hydrazido(2<sup>-</sup>) ligands in **26** and **27** react with various substrates containing a  $>C=O$  group,

Scheme 6



giving a series of organodinitrogen ligands (Scheme 6).

The reaction of **26b** ( $M = W$ ) with succinyl chloride results in the formation of cyclic diacylhydrazido(2<sup>-</sup>)


complex  $trans$ - $[WF(NNCOCH_2CH_2CO)(dppe)_2][BF_4]$  (nucleophilic substitution),<sup>116</sup> while the reaction of **27** ( $X = Cl$ ) with diphenylketene gives the monoacylhydrazido(2<sup>-</sup>) complexes  $[MCl_2(NNHCOCHPh_2)(PMe_2Ph)_3]$  (nucleophilic addition).<sup>117</sup> Similar reaction of  $[MBr(NNH_2)(dppe)_2]Br$  ( $M = Mo, W$ ) with phenyl isocyanate seems to occur, but the products containing the  $NNHCONHPh$  ligand were only poorly characterized.<sup>116</sup>

More intriguing reactions of the hydrazido(2<sup>-</sup>) ligand are the condensation with ketones and aldehydes to give a numerous body of diazoalkane complexes such as  $trans$ - $[MF(NN=CRR')(dppe)_2][BF_4]$  (**28**) and  $cis,mer$ - $[MX_2(NN=CRR')(PMe_2Ph)_3]$  (**29**).<sup>118</sup> The reactions are markedly enhanced by the addition of catalytic amounts of acid,  $HBF_4$  or  $HX$ . Related diazoalkane complexes  $trans$ - $[WBr(NN=CRR')(dppe)_2]Br$  are accessible from the reactions of **1b** with *gem*-dibromides as described already.<sup>114</sup> However, this condensation method may provide a more convenient route to derive diazoalkane ligands since the carbonyl compounds with various substituents are more easily available than *gem*-dibromides. Furthermore, the present method is applicable to both *dppe* and *PMe*<sub>2</sub>-*Ph* complexes **1** and **2**, whereas the reactions of **2** with organic halides giving diazoalkane complexes are unknown.

Formation of an alkenyldiazenido ligand from a  $N_2$  ligand has been demonstrated for the reaction of **2b** with acetylacetone in MeOH. The product has been spectroscopically characterized to be *mer*- $[W(acac)(NNCMe=CHCOMe)(PMe_2Ph)_3]$  (**30**), which has been supported by the X-ray structure determination of its  $HBr$  adduct *mer*- $[W(acac)(NNHCMe=CHCOMe)(PMe_2Ph)_3]Br$ .<sup>119</sup> Formation of the alkenyldiazenido ligand in **30** may proceed via initial protonation presumably by acetylacetone forming hydrazido(2<sup>-</sup>) ligand and its successive condensation with acetylacetone followed by deprotonation.


Free diazoalkanes are highly reactive species, but their instability sets the limitation in utilizing these potential reagents for organic synthesis. Reactivities of coordinated diazoalkanes in **28** and **29** derived from a variety of aldehydes and ketones are therefore of particular interest,<sup>120</sup> the study of which may lead

## Scheme 7




to the development of new synthetic pathways toward organonitrogen compounds from molecular  $N_2$ . The reactions of **28** and **29** demonstrated recently in this context are as follows.

Reactions of diazoalkane complexes **28** formulated as *trans*-[MF(NN=CR<sup>1</sup>CHR<sup>2</sup>R<sup>3</sup>)(dppe)<sub>2</sub>][BF<sub>4</sub>]<sup>-</sup> with LDA or NaN(SiMe<sub>3</sub>)<sub>2</sub> afford alkenyldiazenido complexes *trans*-[MF(N=NCR<sup>1</sup>=CR<sup>2</sup>R<sup>3</sup>)(dppe)<sub>2</sub>] (**31**) quantitatively.<sup>121</sup> This presents a potential route to a series of alkenyldiazenido complexes. Reactions giving alkenyldiazenido complexes reported earlier had been the formation of **30** from **1b** (vide supra) and the synthesis of *trans*-[MF(N=NCR<sup>1</sup>=CHCOMe)(dppe)<sub>2</sub>] from **28** (R = Me, R' = CH<sub>2</sub>COMe) and NaOMe.<sup>119</sup> The complexes containing the alkenyldiazenido ligand highly substituted with electron-withdrawing groups were also obtained from the *N*-vinylation reaction of *trans*-[WX(NNH<sub>2</sub>)(dppe)<sub>2</sub>]<sup>+</sup> (X = F, Br) with equimolar amounts of the corresponding alkenes in the presence of NEt<sub>3</sub> (eq 5).<sup>122</sup> The probable nucleophile




in this system is not the hydrazido(2<sup>-</sup>) complex but the diazenido complex [WX(NNH)(dppe)<sub>2</sub>] produced in situ from the hydrazido(2<sup>-</sup>) complex and the base. More recently, another route toward alkenyldiazenido complexes has been clarified, which involves the  $\beta$ -alkylation of  $\alpha,\beta$ -unsaturated diazoalkane complexes by LiCuR<sub>2</sub> (eq 6).<sup>123</sup>



Complexes **31** are susceptible to electrophilic attack at the terminal alkenyl carbon by RX, PhNCO,


RCHO, etc., yielding new diazoalkane complexes containing ligands such as NN=CR<sup>1</sup>CR<sup>2</sup>R<sup>3</sup>R, NN=CMeCH<sub>2</sub>CONHPh, NN=CMeCH(CONHPh)<sub>2</sub>, and NN=CHCH=CHR. On the other hand, oxidation of **31** (R<sup>1</sup> = R<sup>2</sup> = H, R<sup>3</sup> = Me; R<sup>1</sup> = Me, R<sup>2</sup> = R<sup>3</sup> = H) by I<sub>2</sub> or CuCl<sub>2</sub> gave dinuclear bis(diazo)alkane complexes [ { *trans*-MF(dppe)<sub>2</sub> }<sub>2</sub>( $\mu$ -NN=CR<sup>1</sup>CR<sup>2</sup>R<sup>3</sup>CR<sup>2</sup>R<sup>3</sup>-CR<sup>1</sup>=NN)].<sup>121a</sup> Transformations starting from **28** described herein are summarized in Scheme 7 for the case of *trans*-[MF(NN=CMe<sub>2</sub>)(dppe)<sub>2</sub>]<sup>+</sup>.

In **29**, the PMe<sub>2</sub>Ph ligand *trans* to the halide can be replaced by a range of  $\pi$ -acceptor molecules L at elevated temperatures, and the resultant complexes containing these molecules *cis,trans*-[MX<sub>2</sub>(NN=CRR')(PMe<sub>2</sub>Ph)<sub>2</sub>(L)] (**32**) have been isolated (L = CO, alkene, aldehyde, Bu<sup>t</sup>NC) (eq 7).<sup>124</sup> From complexes **32** (L = Bu<sup>t</sup>NC) were derived the carbene-diazoalkane complexes *cis,trans*-[MX<sub>2</sub>(NN=CMePh)(PMe<sub>2</sub>Ph)<sub>2</sub>(=CMeNHBu<sup>t</sup>)] upon treatment with AlMe<sub>3</sub> followed by hydrolysis.<sup>125</sup> In contrast to the formation of **32** from the thermal reaction with CO, **29** (M = W, X = Cl) gives cationic diazoalkane complex *mer*-[WCl(NN=CMe<sub>2</sub>)(PMe<sub>2</sub>Ph)<sub>3</sub>(CO)]<sup>+</sup> (**33**) when reacted with CO in the presence of ZnCl<sub>2</sub>(dioxane).<sup>126</sup>




The reactivities of the coordinated L in **32** and CO in **33** toward the adjacent diazoalkane ligand is of particular interest, since this might open the alternative novel routes for preparing a variety of organonitrogenous ligands and compounds. Recently, the highly reactive nature of the metal-imido bonds has been clarified in coordinatively unsaturated Zr and

Ir complexes such as  $[\text{Cp}_2\text{Zr}=\text{NR}]$  generated from either  $[\text{Cp}_2\text{Zr}(=\text{NR})(\text{THF})]$  or  $[\text{Cp}_2\text{Zr}(\text{NHR})_2]$  and  $[\text{Cp}^*\text{Ir}=\text{NBU}^t]$ . Thus reactions of the former Zr species with alkynes afforded azametallacyclobutene complexes (eq 8),<sup>127</sup> while treatment of the latter Ir



complex with *e.g.*  $\text{Bu}^t\text{NC}$ ,  $\text{CO}$ ,  $\text{CO}_2$ , and alkynes also resulted in the addition of these molecules to the Ir–N multiple bond as depicted in Scheme 8.<sup>128</sup>

Scheme 8




However, in spite of intensive investigation of the reactivities of the  $\text{M}=\text{N}$  bond in **32**, its carbene-diazoalkane derivatives, and **33**, any reactions have not yet been found in which the diazoalkane ligand interacts with the substrate species in *cis* position. It might be noteworthy that reduction of the cationic complex **33** by Mg under CO resulted in the isolation of the dinuclear bis(diazenido) complex  $[\{\text{trans},\text{trans}\text{-WCl}(\text{CO})_2(\text{PMe}_2\text{Ph})_2\}_2(\mu\text{-N}=\text{NCMe}_2\text{CMe}_2\text{N}=\text{N})]$ .<sup>126</sup>

Complexes **29** ( $\text{M} = \text{W}$ ) also react with excess  $\text{CH}_2\text{X}_2$  in hot toluene, resulting in the formation of paramagnetic diazoalkane complexes *mer,trans*- $[\text{WX}_3(\text{NN}=\text{CRR}')(\text{PMe}_2\text{Ph})_2]$  ( $\text{X} = \text{Br}, \text{I}$ ).<sup>129</sup> The bonding parameters associated with the diazoalkane ligand in this W(V) complex ( $\text{X} = \text{Br}, \text{R} = \text{Me}, \text{R}' = \text{Ph}$ ) are quite similar to those in the W(IV) complexes.

Recent extension of the condensation reaction of hydrazido(2-) complexes with monocarbonyl compounds to that with dialdehydes has led to the formation of nitrogen heterocyclic ligands. Thus the reaction of **26** with 2,5-dimethoxytetrahydrofuran, a succinaldehyde equivalent, yields 1-pyrrolylimido

complexes *trans*- $[\text{MF}(\text{NNCH}=\text{CHCH}=\text{CH})(\text{dppe})_2]\text{[BF}_4\text{]}^-$  (**34**).<sup>130</sup> Interestingly, the pyrrole ring in **34b** ( $\text{M} = \text{W}$ ) undergoes  $\beta$ -regioselective electrophilic substitution reactions; treatment of **34b** with NBS gives only the  $\beta$ -bromo derivative. Exclusive  $\beta$ -cyanation and  $\beta$ -benzoylation also occur, when reacted with chlorosulfonyl isocyanate and  $\text{PhCOCl}/\text{AlCl}_3$ , respectively (eq 9). This presents a sharp contrast to the electrophilic substitutions of free pyrrole, which are known to proceed preferentially at the  $\alpha$ -position. This remarkable difference in the reactivity between the pyrrole ring in **34b** and free

pyrrole is presumably ascribed to the steric effect of the dppe ligands, protecting the  $\alpha$ -position of the pyrrole ring.



The reaction of **27b** ( $\text{X} = \text{Cl}$ ) with phthalaldehyde results in the formation of a phthalimidine ring. For the dppe complex **26b**, only the condensation of the hydrazido(2-) ligand with one aldehyde group proceeds to give the diazoalkane complex *trans*- $[\text{WF}(\text{NN}=\text{CHC}_6\text{H}_4\text{CHO}-o)(\text{dppe})_2]^+$  in high yield. However, this diazoalkane ligand can also be converted into the phthalimidine ring by subsequent reaction with  $\text{AlCl}_3$  in THF at reflux (Scheme 6).


### C. Synthesis of Nitrogen-Containing Organic Compounds from Coordinated $\text{N}_2$

As shown in the previous sections, considerably diversified routes are now available to obtain the organodinitrogen ligands from coordinated  $\text{N}_2$ , although most of these are still limited to the Mo and W complexes with tertiary phosphine ligands. The products characterized in a well-defined manner include organodiazenido, organohydrazido(2-), or diazoalkane ligands. Much effort has been given to liberating these ligands as the nitrogen-containing organic compounds, since such reactions may provide important information in designing the chemical  $\text{N}_2$ -fixing system capable of producing organonitrogen compounds catalytically from gaseous  $\text{N}_2$ . It is now apparent that some organodinitrogen complexes referred above can produce nitrogenous compounds under certain conditions. In this section, the reactions which appeared in fairly old references have also been cited to cover the whole reactivities of organonitrogen ligands.

Initial observation concerning the liberation of organonitrogen compounds was the reductive destruction of the Mo and W dialkylhydrazido(2-)

complexes *trans*- $[\text{MBr}\{\text{NN}(\text{CH}_2)_3\text{CH}_2\}(\text{dppe})_2]\text{Br}$  by  $\text{LiAlH}_4$  at 80 °C for 65 h. Pyrrolidine was obtained in moderate yields after workup of the reaction mixtures with MeOH and then HBr. Analogous treatment of *trans*- $[\text{WBr}(\text{NNMe}_2)(\text{dppe})_2]\text{Br}$  afforded  $\text{Me}_2\text{NH}$  in high yield. Direct base distillation of these complexes also resulted in the formation of the amines, although the yields were lower, while treatment of the latter complex with  $\text{H}_2\text{SO}_4$  afforded  $\text{MeNH}_2$  and  $\text{Me}_2\text{NH}$  in low yields.<sup>131</sup> In the reaction of **34** with  $\text{LiAlH}_4$  followed by treatment with MeOH, pyrrole was obtained in high yield along with a small amount of *N*-aminopyrrole.<sup>130</sup> The reaction of **34** proceeds readily at room temperature and completes

Scheme 9




in only 2 h. More importantly,  $[\text{MH}_4(\text{dppe})_2]$  can be isolated in moderate yields from the resultant reaction mixtures. Since the conversion of these hydrido complexes to the parent  $\text{N}_2$  complexes **1** has been reported previously, the present pyrrole synthesis can be represented as the cyclic system shown in Scheme 9.

On the other hand, the reaction with acid also proceeds cleanly, when the hydrazido(2-) complex reduced by two electrons is used. Thus  $[\text{Mo}\{\text{NN}(\text{CH}_2)_4\text{CH}_2\}(\text{dppe})_2]$ , prepared from *trans*- $[\text{MoBr}\{\text{NN}(\text{CH}_2)_4\text{CH}_2\}(\text{dppe})_2]\text{Br}$  (**35**) by either electroreduction or reaction with  $\text{LiBu}^t$ , reacted with excess  $\text{HBr}$  in THF to give piperidine and *trans*- $[\text{MoBr}(\text{NH})(\text{dppe})_2]\text{Br}$  in 74 and 67% yields, respectively.<sup>132</sup>

As for organodiazenido complexes, the reaction of *trans*- $[\text{MoBr}(\text{NNBu}^n)(\text{dppe})_2]$  with  $\text{NaBH}_4$  or  $\text{NaOMe}$  in benzene/MeOH at elevated temperature was carried out, which gave  $\text{NH}_3$  together with butylamines ( $\text{Bu}^n\text{NH}_2$  and  $\text{Bu}^n\text{NMeH}$ ).<sup>110</sup>

Treatment of the diazoalkane complexes **29** ( $\text{X} = \text{Br}$ ,  $\text{R} = \text{R}' = \text{Me}$ ) with  $\text{HX}$  results in the formation of a mixture of  $\text{N}_2\text{H}_4$  and acetone azine, while that with  $\text{LiAlH}_4$  gives  $\text{Pr}^t\text{NH}_2$  together with  $\text{NH}_3$ .<sup>118b</sup> Both reactions proceed at room temperature. In relevance to the former system, formation of acetone azine has been observed by treatment of **2b** or *trans*- $[\text{W}(\text{N}_2)_2(\text{PPh}_2\text{Me})_4]$  with a MeOH/acetone mixture at 50 °C.<sup>133</sup> Details of this reaction course are uncertain, but the mechanism involving the formation of a hydrazido(2-) complex and the subsequent condensation of this species with acetone forming a diazoalkane complex has been suggested. The Mo complex **2a** gives no azine, but the Mo tetrathioether complex **10** does afford the azine under similar conditions.<sup>115</sup> It is to be noted that conversion of coordinated  $\text{N}_2$  into acetone azine was also observed for the reactions of dinuclear Ta complexes such as  $[\{\text{TaCl}_3(\text{PR}_3)_2\}_2(\mu\text{-N}_2)]$  and  $[\{\text{TaL}_3(\text{THF})\}_2(\mu\text{-N}_2)]$  ( $\text{L} = \text{CH}_2\text{Bu}^t$ ,  $\text{OBu}^t$ ) with acetone. In these reactions, formation of acetone azine presumably results from the double bond metathesis between the  $\text{Ta}=\text{N}$  and  $>\text{C}=\text{O}$  units,

Scheme 10




which may support the diimide structure proposed for these ditantalum complexes.<sup>75</sup>

Reactions of organodinitrogen complexes under electroreductive conditions were also investigated previously. An important feature of these systems is that the elimination of the nitrogenous compound may be accompanied by the recovery of the parent  $\text{N}_2$  complex in the case of the organodinitrogen complexes containing appropriate coligands. Thus controlled potential electrolysis of **35** ( $\text{M} = \text{Mo}$ ) in THF under  $\text{N}_2$  gave *N*-aminopyrrolidine and the  $\text{N}_2$  complex **1a** in 60–70 and 45% yields, respectively (Scheme 10).<sup>134</sup> As a related reaction, formation of 5-amino-4-cyanopyrazole in ca. 40% yield from dicyanovinylhydrazido(2-) complex *trans*- $[\text{WF}\{\text{NNHCH}=\text{C}(\text{CN})_2\}(\text{dppe})_2][\text{BF}_4]$  under electroreductive conditions has been reported.<sup>135</sup> Dicyanovinylhydrazine may be liberated by electrolysis prior to the cyclization to give the pyrazole. The W species present in the reaction mixture under  $\text{N}_2$  was characterized to be the hydrazido(2-) complex **26b**. In these reactions using organohydrazido(2-) complexes, the  $\text{M}-\text{N}$  bond cleavage occurs favorably rather than the  $\text{N}-\text{N}$  bond fission, which presents a sharp contrast to the almost exclusive  $\text{NH}_3$  formation from *trans*- $[\text{W}(\text{OTs})(\text{NNH}_2)(\text{dppe})_2]^+$  by electrolysis (vide supra).<sup>78</sup>


## D. Organic Synthesis Using Nitrido Complexes Derived from Molecular $\text{N}_2$ as Key Intermediate

In certain systems, molecular nitrogen is incorporated into organometallic compounds with concurrent  $\text{N}-\text{N}$  bond scission, although in most cases the resultant nitrido species are poorly defined. The complex obtained from the reaction of  $\text{N}_2$  with a  $\text{TiCl}_3/\text{Mg}/\text{THF}$  system is believed to be a nitrido species having a composition of  $[\text{TiNMg}_2\text{Cl}_2(\text{THF})]$  (**36**),<sup>136</sup> which further reacts with  $\text{PhCOCl}$  or  $\text{CO}_2$  to give the Ti compounds formulated as  $[\text{TiNCl}(\text{PhCO})_{1.5}]$ <sup>136</sup> or  $[\text{Ti}(\text{NCO})\text{Mg}_2\text{Cl}_2\text{O}(\text{THF})_3]$  (**37**),<sup>137</sup> respectively. Recently synthesis of a variety of organonitrogen compounds using in situ generated **36** and **37** has been reported.

## Scheme 11



## Scheme 12



Treatment of **36**, prepared from  $\text{TiCl}_3$  or  $\text{TiCl}_4$ , with excess aryl chloride in THF at room temperature or at reflux followed by hydrolysis gives a mixture of  $\text{RCONH}_2$  and  $(\text{RCO})_2\text{NH}$ . From the reaction of **37** with phthaloyl dichloride or phthalic anhydride in py, phthalimide is obtained. Alternatively, when this reaction using **37** is combined with the carbonylation reaction of aryl halide using  $[\text{Pd}(\text{PPh}_3)_4]$  catalyst, the imides are produced directly from the corresponding aryl halides, e.g.  $(\text{PhCO})_2\text{NH}$  from  $\text{PhBr}$  and phthalimide from  $o\text{-BrC}_6\text{H}_4\text{COOH}$ .<sup>138</sup> This combination of carbonylation and nitrogenation has further been extended to the one-step synthesis of isoindolinones from  $o$ -halophenyl alkyl ketones and that of quinazolinones from  $o$ -haloanilides (Scheme 11).<sup>139</sup> The related reactions forming indole and quinoline derivatives have also been reported.<sup>140</sup>


More recently, catalytic synthesis of  $\text{PhCONH}_2$  from  $\text{PhCOCl}$  has been demonstrated by the use of a mixture of  $\text{TiCl}_4/\text{Li}/\text{Me}_3\text{SiCl}$  under  $\text{N}_2$  (2.43 mol/Ti atom). Phthalic anhydride was also converted into phthalimide catalytically in the presence of  $\text{CsF}$  along with  $\text{TiCl}_4/\text{Li}/\text{Me}_3\text{SiCl}$  under  $\text{N}_2$  (2.04 mol/Ti atom).<sup>141</sup> The mechanism proposed for these reactions involves the catalytic formation of  $\text{N}(\text{SiMe}_3)_3$  on the reduced Ti species, which subsequently reacts with  $\text{HCl}(\text{aq})$  or  $\text{CsF}$ , respectively, and is finally trapped as the amide or imide (Scheme 12). Catalytic formation of  $\text{N}(\text{SiMe}_3)_3$  was observed earlier by treatment of  $\text{Me}_3\text{SiCl}$  and  $\text{Li}$  in the presence of several transition metal salts such as  $\text{CrCl}_3$  under  $\text{N}_2$ .<sup>142</sup> This catalytic nitrogenation reaction using  $\text{TiCl}_4/\text{Li}/\text{Me}_3\text{SiCl}/\text{CsF}$  has also applied to the synthesis of a variety of heterocyclic compounds including indole and pyrrole derivatives using numerous diketo compounds.<sup>143</sup>

Stepwise transformation of nitrido ligand in  $\text{trans}[\text{MoCl}(\text{N})(\text{dppe})_2]$  toward glycine and alanine methyl esters has been demonstrated in a well-defined manner.<sup>144</sup> The nitrido complex may be prepared from **1a** by dialkylation of a coordinated  $\text{N}_2$  and subsequent  $\text{N}-\text{N}$  bond cleavage upon acidolysis.<sup>132b</sup>

The pathway converting the nitrido ligand into amino acid methyl esters involves initial  $\text{N}-\text{C}$  bond formation by treatment with  $\text{ICH}_2\text{COOMe}$ , giving imido complex  $\text{trans}[\text{MoCl}(\text{NCH}_2\text{COOMe})(\text{dppe})_2]\text{I}$ . Removal of one proton attached to the  $\alpha$ -carbon atom gives  $\text{trans}[\text{MoCl}(\text{N}=\text{CHCOOMe})(\text{dppe})_2]$ , which is further treated with  $\text{MeI}$  to give another imido complex,  $\text{trans}[\text{MoCl}(\text{NCH}(\text{Me})\text{COOMe})(\text{dppe})_2]\text{I}$ . The  $\text{Mo}-\text{N}$  bond cleavage in these imido complexes takes place under electroreductive conditions, resulting in the liberation of glycine and alanine methyl esters in 70 and 80% yields, respectively.


E. Silylation and Gerymylation of Coordinated  $\text{N}_2$  and Catalytic Conversion of Molecular  $\text{N}_2$  into Silylamines

The Mo and W complexes **1** and **2** react with  $\text{Me}_3\text{SiI}$  at  $50^\circ\text{C}$  in benzene under rigorously dry conditions to give the silyldiazenido complexes  $[\text{M}(\text{NNSiMe}_3)(\text{dppe})_2]$  and  $\text{trans}[\text{M}(\text{NNSiMe}_3)(\text{PMe}_2\text{Ph})_4]$  (**38**) (eq 10). In the presence of adventitious moisture, the




silyldiazenido(2-) complexes  $[\text{M}(\text{NNSiMe}_3)(\text{dppe})_2]\text{I}$  and  $\text{cis,mer}[\text{M}_2(\text{NNSiMe}_3)(\text{PMe}_2\text{Ph})_3]$  are also formed.<sup>145</sup> Silylation of the terminal nitrogen in **1b** and **2b** also proceeds by the use of  $\text{Me}_3\text{SiOTf}$ , yielding the corresponding OTf complexes.<sup>146</sup> Although  $\text{Me}_3\text{SiCl}$  does not react with **1** and **2** under similar conditions, treatment of **2b** with  $\text{Me}_3\text{SiCl}$  in the presence of excess  $\text{NaI}$  results in the formation of **38b** ( $\text{M} = \text{W}$ ) in moderate yield. Following this procedure, a variety of silyldiazenido complexes are readily derived from **2b** by the use of numerous  $\text{R}_3\text{SiCl}$  (eq 10).<sup>147</sup> It is to be noted that the further silylation of the silyldiazenido ligands did not proceed in any reactions using excess amounts of monosilyl halides. However, treatment of **2** with a mixture of  $\text{ClSiMe}_2\text{CH}_2\text{CH}_2\text{SiMe}_2\text{Cl}/\text{NaI}$  afforded disilyldiazenido(2-)

complexes  $\text{cis,mer}[\text{M}_2(\text{NNSiMe}_2\text{CH}_2\text{CH}_2\text{SiMe}_2)(\text{PMe}_2\text{Ph})_3]$ , and a subsequent replacement of one  $\text{PMe}_2\text{Ph}$  ligand by a series of  $\pi$ -acceptor molecules  $\text{L}$  gave numerous  $\text{cis,trans}[\text{M}_2(\text{NNSiMe}_2\text{CH}_2\text{CH}_2\text{SiMe}_2)(\text{PMe}_2\text{Ph})_2(\text{L})]$  (eq 11).<sup>148</sup> Moreover, the gerymydi-




azenido complexes  $\text{trans}[\text{W}(\text{NNGeR}_3)(\text{PMe}_2\text{Ph})_4]$  ( $\text{R} = \text{Me}, \text{Ph}$ ) were also obtained from the reactions using  $\text{R}_3\text{GeCl}$  in place of  $\text{R}_3\text{SiCl}$ .<sup>147,148a</sup> Silylation of coordinated  $\text{N}_2$  by silyl halide has also been observed in the reaction of Mo complex  $[\text{C}_6\text{F}_5\text{-N}_3\text{N}]\text{Mo}(\text{N}_2)$ -

Scheme 13



{Na(ether)<sub>x</sub>} and Pr<sub>3</sub>SiCl (vide supra).<sup>50</sup> Further study of the silylation of the N<sub>2</sub> ligand in **1** and **2** has led to the finding that the silylcobalt complex [Ph<sub>2</sub>MeSiCo(CO)<sub>4</sub>] can also serve as the silylating reagent and the silyldiazenido complexes of the type *trans*-[M(NNSiPh<sub>2</sub>Me)(P)<sub>4</sub>](μ-OC)Co(CO)<sub>3</sub>] have been isolated.<sup>149</sup> Upon protonation by water, MeOH, or HBr, the diazenido complex (M = W, P = 1/2 dppe) is converted into a cationic silylhydrazido(2-) complexes *trans*-[WX(NNHSiPh<sub>2</sub>Me)(dppe)<sub>2</sub>][Co(CO)<sub>4</sub>] (X = OH, OMe, Br).


Interestingly the reduction of the silyldiazenido complexes **38** with Na in THF under Ar gives a mixture of N<sub>1</sub> products such as NH<sub>3</sub>, HN(SiMe<sub>3</sub>)<sub>2</sub>, NaNH<sub>2</sub>, NaNHSiMe<sub>3</sub>, and NaN(SiMe<sub>3</sub>)<sub>2</sub>, along with some N<sub>2</sub> gas. Furthermore, the parent N<sub>2</sub> complexes **2** can be isolated from the reaction mixtures in moderate yields when the reduction is carried out under N<sub>2</sub>. In the presence of excess Me<sub>3</sub>SiCl under Ar, complexes **38** give N(SiMe<sub>3</sub>)<sub>3</sub> as a principal nitrogenous product. On the other hand, treatment of **2** with Na in the presence of Me<sub>3</sub>SiCl under Ar gives N(SiMe<sub>3</sub>)<sub>3</sub> in good yield.<sup>146,150</sup> These findings shown in Scheme 13 have led to the attempts in converting N<sub>2</sub> gas directly into N(SiMe<sub>3</sub>)<sub>3</sub> by using the Me<sub>3</sub>SiCl/Na system containing complexes **2** as the catalyst (eq 12). In consequence, the Mo complex **2a**



has been found to remarkably facilitate the expected transformation.<sup>150</sup> In an optimum run using the equimolar mixture of Me<sub>3</sub>SiCl and Na in the presence of 0.5 mol % of **2a**, N(SiMe<sub>3</sub>)<sub>3</sub> was obtained in 37% yield (ca. 24 mol/Mo atom) at 30 °C after 15 h. The major byproduct was Me<sub>3</sub>SiSiMe<sub>3</sub>, the formation of which was also accelerated by **2a**. Among the complexes studied, the catalytic activity decreases in the order **2a** >> **1a** >> **1b**, **2b**.

One possible mechanism for the formation of N(SiMe<sub>3</sub>)<sub>3</sub> along with Me<sub>3</sub>SiSiMe<sub>3</sub> is depicted in Scheme 14. As in the alkylation of coordinated N<sub>2</sub> (vide supra), the silyl radical Me<sub>3</sub>Si· may be generated by homolytic cleavage of Me<sub>3</sub>SiCl within the coordination sphere and then attack on the remaining N<sub>2</sub> ligand to form a silyldiazenido ligand. If the radical couples to each other, Me<sub>3</sub>SiSiMe<sub>3</sub> may be formed. However, we must await further investigation to elucidate the detailed mechanism for this catalytic N<sub>2</sub> fixation. Other chlorosilanes can also be used for the transformation of N<sub>2</sub> gas into the corresponding


Scheme 14



silylamines by the catalysis of **2a**, although the turnover was lower.<sup>147</sup>

## VI. Reactions of Coordinated N<sub>2</sub> with Lewis Acids or Organometallic Reagents


Besides the reactivities toward a variety of protic acids and organic electrophiles described above, reactions of coordinated N<sub>2</sub> with metal species have also been investigated. As expected from the nucleophilicity or basicity of the terminal N atom in an end-on N<sub>2</sub> ligand, the reactions with various Lewis acidic metal compounds result in the formation of the corresponding Lewis acid–base adducts. Thus the adducts with a main group acceptor, e.g. [M-(NNAiMe<sub>3</sub>)(N<sub>2</sub>)(dppe)<sub>2</sub>] and *trans*-[ReCl(NNAiMe<sub>3</sub>)(P)<sub>4</sub>] (P = PMe<sub>2</sub>Ph, 1/2 dppe), as well as those with a transition metal π-acceptor, e.g. [ReCl(PMe<sub>2</sub>Ph)<sub>4</sub>-(NNMCl<sub>5</sub>)] (M = Nb, Ta) and [{ReCl(PMe<sub>2</sub>Ph)<sub>4</sub>-(NN)}<sub>2</sub>MCl<sub>4</sub>] (M = Ti, Zr), are known.<sup>151</sup> It may be noteworthy that the reactions of **2b** with AlX<sub>3</sub> (X = Cl, Br) in the presence of py afford not the simple adducts but the ditungsten complexes with a bridging NN(AlX<sub>2</sub>)<sub>2</sub>NN unit, [{WX(PMe<sub>2</sub>Ph)<sub>3</sub>(py)}<sub>2</sub>{μ-NN(AlX<sub>2</sub>)<sub>2</sub>-NN}] (**39**).<sup>152</sup> A structurally similar dicobalt complex,



[{(Me<sub>3</sub>P)<sub>3</sub>Co}<sub>2</sub>(μ-NN(AlMe<sub>2</sub>)<sub>2</sub>NN)], was also prepared upon treatment of [(Me<sub>3</sub>P)<sub>3</sub>CoNNK] with Me<sub>2</sub>AlCl.<sup>153</sup> The transition metal heteronuclear complexes [{WX(PMe<sub>2</sub>Ph)<sub>4</sub>}(μ-NN)MCp<sub>2</sub>Cl] (M = Ti, X = Cl; M = Zr, Hf, X = I)<sup>59</sup> described already are also available in the related reactions of **2b** with Cp<sub>2</sub>TiCl<sub>2</sub> or Cp<sub>2</sub>MCl<sub>2</sub>/NaI (M = Zr, Hf).

The N<sub>2</sub> ligand in [CpMn(CO)<sub>2</sub>(N<sub>2</sub>)] (**40**) can exhibit quite interesting reactivity. Thus **40** reacts with MeLi and then with [Me<sub>3</sub>O][BF<sub>4</sub>] to give methyldiazene complex [CpMn(CO)<sub>2</sub>(MeN=NMe)]. The initial


## Scheme 15



product obtained from the reaction with MeLi has been formulated as  $[\text{CpMn}(\text{CO})_2(\text{MeN}=\text{N}^-\text{Li}^+)]$ , which may be formed by the nucleophilic attack of the Me carbanion on the inner N atom.<sup>154</sup> This reaction was reported as early as 1978, but it is still unique in that the coordinated  $\text{N}_2$  undergoes nucleophilic attack, forming the satisfactorily characterizable metal species. When treated with pressurized  $\text{N}_2$ , the methyldiazene complex is converted into **40** with concurrent liberation of the diazene.


Although **40** does not react with the metal hydrides such as MH (M = Li, Na, K),  $\text{LiAlH}_4$ , and  $\text{LiBEt}_3\text{H}$ ,<sup>154</sup> **2b** reacts with  $\text{NaAlH}_2(\text{OCH}_2\text{CH}_2\text{OMe})_2$  and the base distillation of the evaporated reaction mixture residue gives  $\text{NH}_3$  in moderate yield.<sup>155</sup> Reactions of **2** with a range of transition metal hydrides have also been investigated. As expected from the facile  $\text{NH}_3$  formation from the reactions with sulfuric acid, **2b** reacts with acidic transition metal carbonyl hydrides such as  $[\text{HCo}(\text{CO})_4]$ ,  $[\text{H}_2\text{Fe}(\text{CO})_4]$ , and  $[\text{HFeCo}_3(\text{CO})_{12}]$  to give  $\text{NH}_3$  in substantial yields after base distillation of the reaction mixture.<sup>83c,155,156</sup> Besides these acidic hydrides, the hydridic transition metal compound  $[\text{Cp}_2\text{ZrHCl}]$  also reacts with **2** to give  $\text{NH}_3$  under the similar conditions.<sup>155</sup> Although any intermediate stages are unknown for the whole reactions of **2** with metal hydrides, a series of hydrazido(2-) complexes,  $\text{trans-}[\text{W}(\text{OR})(\text{NNH}_2)(\text{dppe})_2][\text{A}]$  (A =  $\text{Co}(\text{CO})_4$ ,  $\text{FeCo}_3(\text{CO})_{12}$ ), were isolated from the reactions of **1b** with  $[\text{HCo}(\text{CO})_4]$  or  $[\text{HFeCo}_3(\text{CO})_{12}]$  in various ROH.<sup>83c</sup> Related silylation reaction of **1** and **2b** with  $[\text{Ph}_2\text{MeSiCo}(\text{CO})_4]$  giving the silyldiazenido complexes has been cited already.<sup>149</sup> More recently, formation of **26b** from the reaction of **1b** with 2 equiv of acidic Ru complex  $[\text{CpRu}(\text{dtfpfe})\text{H}_2][\text{BF}_4]$  (**41**) has been reported (Scheme 15).<sup>157</sup> Complex **41** exists in solution as a slowly interconverting mixture of the dihydrogen and dihydride species  $[\text{CpRu}(\text{dtfpfe})(\eta^2\text{-H}_2)]^+$  and  $[\text{CpRu}(\text{dtfpfe})(\text{H})_2]^+$ , the  $\text{p}K_a$  values of which are estimated to be 4.3 and 4.4, respectively. In this reaction the heterolytic cleavage of the dihydrogen ligand within **41** into  $\text{HBF}_4$  and  $[\text{CpRu}(\text{dtfpfe})\text{H}]$  is taking place. Formation of **26b** may be important toward development of the  $\text{N}_2$ -reducing system by  $\text{H}_2$ , including two different metal sites suitable for activating  $\text{N}_2$  and  $\text{H}_2$ , although at present **41** is not available from gaseous  $\text{H}_2$ .

Arylation of coordinated  $\text{N}_2$  is of significant interest due to its possible relevance to the synthesis of aromatic amines. Although alkylation of coordinated  $\text{N}_2$  proceeds smoothly by reaction with alkyl halides, arylation of coordinated  $\text{N}_2$  is hardly accessible, except for the reaction of aryl halides with the Mo tetrathioether complex **10** (vide supra)<sup>115</sup> and the reaction of 2,4-( $\text{O}_2\text{N}$ ) $\text{C}_6\text{H}_3\text{F}$  with in situ generated  $[\text{WX}(\text{NNH})(\text{dppe})_2]$  (X = F, Br,  $\text{CF}_3\text{COO}$ ) (eq 13).<sup>158</sup>



Recently it has been demonstrated that the anionic  $\text{N}_2$  complex  $\text{trans-}[\text{Bu}^n_4\text{N}][\text{W}(\text{N}_2)(\text{NCS})(\text{dppe})_2]$  (**42**) reacts with fluoroarene complexes such as  $[(\eta^6\text{-p-FC}_6\text{H}_4\text{COOMe})\text{Cr}(\text{CO})_3]$  (**43**) and  $[\text{CpRu}(\eta^6\text{-FC}_6\text{H}_4\text{R})][\text{PF}_6]$  (**44**, R = H, Me, OMe, COOMe) under ambient conditions to give a series of novel dinuclear  $\mu$ -aryldiazenido complexes  $\text{trans-}[\text{W}(\text{NCS})\{\text{N}=\text{N}[(\eta^6\text{-p-C}_6\text{H}_4\text{-COOMe})\text{Cr}(\text{CO})_3]\}(\text{dppe})_2]$  (**45**) and  $\text{trans-}[\text{W}(\text{NCS})\{\text{N}=\text{N}[(\eta^6\text{-p-C}_6\text{H}_4\text{R})\text{RuCp}]\}(\text{dppe})_2][\text{PF}_6]$  (**46**) (Scheme 16).<sup>159</sup> Activation of both the  $\text{N}_2$  ligand and the

## Scheme 16




fluoroarene is required here; neither the reaction of the parent neutral complex **1b** with **43** or **44a** (R = H) nor that of the anionic complex **42** with free  $\text{p-FC}_6\text{H}_4\text{COOMe}$  proceeds. It is also to be noted that, in contrast to the smooth reaction of **44a** with **42** to give **46**, the reaction of  $\text{C}_6\text{H}_5\text{F}$  bound to the  $\text{Cr}(\text{CO})_3$  moiety with **42** did not yield the  $\text{N}$ -arylated product, indicating that the  $\text{CpRu}^+$  is a more efficient activating group for fluoroarenes than  $\text{Cr}(\text{CO})_3$ . On the other hand, the reaction of  $[\text{CpFe}(\eta^6\text{-C}_6\text{H}_5\text{F})][\text{PF}_6]$  with **42** presumably results in the redox reaction between these two complexes, since one electron oxidized product  $[\text{W}(\text{N}_2)(\text{NCS})(\text{dppe})_2]$  has been isolated from the reaction mixture.


Instead of starting from isolated **42**, when **1** was treated with **43** in the presence of  $[\text{Bu}^n_4\text{N}]\text{X}$  (X = NCS, F) under irradiation by a W filament, the NCS



complex **45** and its F analogue are obtained, by way of the in situ generation of  $[\text{WX}(\text{N}_2)(\text{dppe})_2]^-$  (eq 14).



In related reactions, formations of *trans*- $[\text{WF}\{\text{N}=\text{N}-[\eta^6\text{-}p\text{-C}_6\text{H}_4\text{R}]\text{M}\}(\text{dppe})_2]$  (M =  $\text{Cr}(\text{CO})_3$ , R = H; M =  $\text{CpFe}^+\text{PF}_6^-$ , R = H, Me) were observed from the reactions of the diazenido complex  $[\text{WF}(\text{NNH})(\text{dppe})_2]$  with  $[(\eta^6\text{-}p\text{-FC}_6\text{H}_4\text{R})\text{M}]$  (eq 15).<sup>159</sup>



## VII. Conclusions

It is now apparent that molecular nitrogen is activated sufficiently through coordination to one or more transition metals to form N–H, N–C, and N–Si bonds under mild conditions. Extensive studies on the synthesis and reactivities of  $\text{N}_2$  complexes have disclosed that the reduction of  $\text{N}_2$  can be promoted not only by the metals relevant to biological  $\text{N}_2$ -fixing system such as Fe, Mo, and V but also by the more diversified transition metals. Results of these studies may provide some basis for understanding reduction mechanism operating in the biological  $\text{N}_2$ -fixing system.

Other important advances in the chemistry of  $\text{N}_2$  complexes are found in the discovery of numerous transformations of coordinated  $\text{N}_2$  into organonitrogen compounds through the C–N bond formation at coordinated  $\text{N}_2$  by reactions with organic compounds. Although the reaction systems demonstrated to date give the organonitrogen compounds mostly in less than stoichiometric amounts, further studies will lead to the development of the effective systems for the production of valuable nitrogen-containing organic compounds directly from molecular nitrogen as a nitrogen source.

## VIII. Abbreviations

acac	acetylacetonate
$[\text{Bu}^t\text{Me}_2\text{Si}-\text{N}_3\text{N}]^{3-}$	$(\text{Bu}^t\text{Me}_2\text{SiNCH}_2\text{CH}_2)_3\text{N}$ trianion
$[\text{C}_6\text{F}_5-\text{N}_3\text{N}]^{3-}$	$(\text{C}_6\text{F}_5\text{NCH}_2\text{CH}_2)_3\text{N}$ trianion
citr	citrate trianion

Cl <sub>4</sub> -cat	tetrachlorocatechololate dianion
Cp	$\eta^5\text{-C}_5\text{H}_5$
Cp*	$\eta^5\text{-C}_5\text{Me}_5$
Cp'	$\eta^5\text{-C}_5\text{Me}_4\text{Et}$
depe	$\text{Et}_2\text{PCH}_2\text{CH}_2\text{PEt}_2$
diglyme	$\text{MeOCH}_2\text{CH}_2\text{OCH}_2\text{CH}_2\text{OMe}$
dmpe	$\text{Me}_2\text{PCH}_2\text{CH}_2\text{PMe}_2$
DPB	diporphyrinatobiphenylene tetraanion
dpepp	$\text{PhP}(\text{CH}_2\text{CH}_2\text{PPh}_2)_2$
dppe	$\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2$
dppee	$\text{O}(\text{CH}_2\text{CH}_2\text{PPh}_2)_2$
dppm	$\text{Ph}_2\text{PCH}_2\text{PPh}_2$
dppp	$\text{Ph}_2\text{PCH}_2\text{CH}_2\text{CH}_2\text{PPh}_2$
dtfpe	$(p\text{-CF}_3\text{C}_6\text{H}_4)_2\text{PCH}_2\text{CH}_2\text{P}(\text{C}_6\text{H}_4\text{CF}_3\text{-}p)_2$
EXAFS	extended X-ray absorption fine structure
Im*	1- <i>tert</i> -butyl-5-phenylimidazole
LDA	$\text{LiNPr}^i_2$
$\text{Me}_8[16]\text{aneS}_4$	3,3,7,7,11,11,15,15-octamethyl-1,5,9,13-tetrahydrocyclohexadecane
NBS	<i>N</i> -bromosuccinimide
NMP	<i>N</i> -methylpyrrolidone
OEPG	octaethylporphyrinogen tetraanion
pic	4-picoline
py	pyridine
TMEDA	$\text{Me}_2\text{NCH}_2\text{CH}_2\text{NMe}_2$
TMP	tetramesitylporphyrin dianion
Ts	<i>p</i> -toluenesulfonyl

## IX. References

- (1) *Chem. Eng. NY* **1993**, March, 19.
- (2) (a) Rees, D. C.; Chan, M. K.; Kim, J. *Adv. Inorg. Chem.* **1994**, *40*, 89. (b) Kim, J.; Rees, D. C. *Science* **1992**, *257*, 1677. (c) Kim, J.; Rees, D. C. *Nature (London)* **1992**, *360*, 553. (d) Chan, M. K.; Kim, J.; Rees, D. C. *Science* **1993**, *260*, 792. (e) Kim, J.; Woo, D.; Rees, D. C. *Biochemistry* **1993**, *32*, 7104. (f) Kim, J.; Rees, D. C. *Ibid.* **1994**, *33*, 389.
- (3) (a) Bolin, J. T.; Campobasso, N.; Muchmore, S. W.; Morgan, T. V.; Mortenson, L. E. In *Molybdenum Enzymes, Cofactors, and Model Systems*; Stiefel, E. I., Coucouvanis, D., Newton, W. E., Eds.; American Chemical Society: Washington, DC, 1993; p 186. (b) Chen, J.; Christiansen, J.; Campobasso, N.; Bolin, J. T.; Tittsworth, R. C.; Hales, B. J.; Rehr, J. J.; Cramer, S. P. *Angew. Chem., Int. Ed. Engl.* **1993**, *32*, 1592.
- (4) Liu, H. I.; Filippini, A.; Gavini, N.; Burgess, B. K.; Hedman, B.; Cicco, A. D.; Natoli, C. R.; Hodgson, K. O. *J. Am. Chem. Soc.* **1994**, *116*, 2418.
- (5) Deng, H.; Hoffmann, R. *Angew. Chem., Int. Ed. Engl.* **1993**, *32*, 1062.
- (6) For example: (a) Burgess, B. K. *Chem. Rev.* **1990**, *90*, 1377. (b) Eady, R. R. *Adv. Inorg. Chem.* **1991**, *36*, 77. (c) Eady, R.; Leigh, G. J. *J. Chem. Soc., Dalton Trans.* **1994**, 2739. See also references therein.
- (7) Allen, A. D.; Senoff, C. V. *J. Chem. Soc., Chem. Commun.* **1965**, 621.
- (8) (a) Vol'pin, M. E.; Shur, V. B. *Dokl. Akad. Nauk. SSSR* **1964**, *156*, 1102. (b) Vol'pin, M. E. *Pure Appl. Chem.* **1972**, *30*, 607.
- (9) (a) Hidai, M.; Tominari, K.; Uchida, Y.; Misono, A. *J. Chem. Soc., Chem. Commun.* **1969**, 1392. (b) Hidai, M.; Tominari, K.; Uchida, Y. *J. Am. Chem. Soc.* **1972**, *94*, 110. (c) Uchida, T.; Uchida, Y.; Hidai, M.; Kodama, T. *Acta Crystallogr.* **1975**, *B31*, 1197.
- (10) Chatt, J.; Dilworth, J. R.; Richards, R. L. *Chem. Rev.* **1978**, *78*, 589.
- (11) Dilworth, J. R.; Richards, R. L. In *Comprehensive Organometallic Chemistry*; Wilkinson, G., Stone, F. A. G., Abel, E. W., Eds.; Pergamon: Oxford, U.K., 1982; Vol. 8; p 1073.
- (12) Henderson, R. A.; Leigh, G. J.; Pickett, C. J. *Adv. Inorg. Radiochem.* **1983**, *27*, 197.
- (13) Hidai, M. In *Molybdenum Enzymes*; Spiro, T. G., Ed.; Metal Ions in Biology Vol. 7; Wiley-Interscience: New York, 1985; p 285.
- (14) Hidai, M.; Mizobe, Y. In *Reactions of Coordinated Ligands*; Braterman, P. S., Ed.; Plenum: New York, 1989; Vol. 2, p 53.
- (15) Hidai, M.; Mizobe, Y., in ref 3a, p 346.
- (16) George, T. A.; DeBord, J. R. D., in ref 3a, p 363.
- (17) Leigh, G. J. *Acc. Chem. Res.* **1992**, *25*, 177.
- (18) Henderson, R. A. *Transition Met. Chem.* **1990**, *15*, 330.
- (19) Leigh, G. J. *J. Mol. Catal.* **1988**, *47*, 363.
- (20) Colquhoun, H. M. *Acc. Chem. Res.* **1984**, *17*, 23.
- (21) George, T. A. In *Homogeneous Catalysis with Metal Phosphine Complexes*; Pignolet, L. H., Ed.; Plenum: New York, 1983; p 405.

- (22) Edema, J. J. H.; Meetsma, A.; Gambarotta, S. *J. Am. Chem. Soc.* **1989**, *111*, 6878.
- (23) (a) Woitha, C.; Rehder, D. *Angew. Chem., Int. Ed. Engl.* **1990**, *29*, 1438. (b) Rehder, D.; Woitha, C.; Friebach, W.; Gailus, H. *J. Chem. Soc., Chem. Commun.* **1992**, 364. (c) Gailus, H.; Woitha, C.; Rehder, D. *J. Chem. Soc., Dalton Trans.* **1994**, 3471.
- (24) Buijink, J.-K. F.; Meetsma, A.; Teuben, J. H. *Organometallics* **1993**, *12*, 2004.
- (25) Song, J.-I.; Berno, P.; Gambarotta, S. *J. Am. Chem. Soc.* **1994**, *116*, 6927.
- (26) Berno, P.; Hao, S.; Minhas, R.; Gambarotta, S. *J. Am. Chem. Soc.* **1994**, *116*, 7417.
- (27) Ferguson, R.; Solani, E.; Floriani, C.; Shiesi-Villa, A.; Rizzoli, C. *Angew. Chem., Int. Ed. Engl.* **1993**, *32*, 396.
- (28) Evans, W. J.; Ulibarri, T. A.; Ziller, J. W. *J. Am. Chem. Soc.* **1988**, *110*, 6877.
- (29) Jubb, J.; Gambarotta, S. *J. Am. Chem. Soc.* **1994**, *116*, 4477.
- (30) Ho, J.; Drake, R. J.; Stephan, D. W. *J. Am. Chem. Soc.* **1993**, *115*, 3792.
- (31) (a) Chatt, J.; Pearman, A. J.; Richards, R. L. *J. Chem. Soc., Dalton Trans.* **1974**, 2074. (b) Donovan-Mtunzi, S.; Richards, R. L.; Mason, J. *Ibid.* **1984**, 469.
- (32) Mizobe, Y.; Ishida, T.; Egawa, Y.; Ochi, K.; Tanase, T.; Hidai, M. *J. Coord. Chem.* **1991**, *23*, 57.
- (33) Chatt, J.; Pearman, A. J.; Richards, R. L. *J. Chem. Soc., Dalton Trans.* **1977**, 2139.
- (34) Dadkhah, H.; Dilworth, J. R.; Faiman, K.; Kan, C. T.; Richards, R. L.; Hughes, D. L. *J. Chem. Soc., Dalton Trans.* **1985**, 1523.
- (35) (a) Carmona, E.; Marin, J. M.; Poveda, M. L.; Atwood, J. L.; Rogers, R. D. *J. Am. Chem. Soc.* **1983**, *105*, 3014. (b) Carmona, E.; Galindo, A.; Poveda, M. L.; Rogers, R. D. *Inorg. Chem.* **1985**, *24*, 4033.
- (36) Anderson, S. N.; Richards, R. L.; Hughes, D. L. *J. Chem. Soc., Dalton Trans.* **1986**, 245.
- (37) George, T. A.; DeBord, J. R. D.; Kaul, B. B.; Pickett, C. J.; Rose, D. J. *Inorg. Chem.* **1992**, *31*, 1295.
- (38) Anderson, S. N.; Hughes, D. L.; Richards, R. L. *J. Chem. Soc., Dalton Trans.* **1986**, 1591.
- (39) Yoshida, T.; Adachi, T.; Kaminaka, M.; Ueda, T. *J. Am. Chem. Soc.* **1988**, *110*, 4872.
- (40) Cruz-Garrits, D.; Gelover, S.; Torrens, H.; Leal, J.; Richards, R. L. *J. Chem. Soc., Dalton Trans.* **1988**, 2393.
- (41) Dilworth, J. R.; Hu, J.; Thompson, R. M.; Hughes, D. L. *J. Chem. Soc., Chem. Commun.* **1992**, 551.
- (42) Dilworth, J. R.; Henderson, R. A.; Hills, A.; Hughes, D. L.; Macdonald, C.; Stephens, A. N.; Walton, D. R. *J. Chem. Soc., Dalton Trans.* **1990**, 1077.
- (43) O'Regan, M. B.; Liu, A. H.; Finch, W. C.; Schrock, R. R.; Davis, W. M. *J. Am. Chem. Soc.* **1990**, *112*, 4331.
- (44) Schrock, R. R.; Wesolek, M.; Liu, A. H.; Wallace, K. C.; Dewan, J. C. *Inorg. Chem.* **1988**, *27*, 2050.
- (45) (a) Camenzind, M. J.; James, B. R.; Dolphin, D. *J. Chem. Soc., Chem. Commun.* **1986**, 1137. (b) Camenzind, M. J.; James, B. R.; Dolphin, D.; Sparapany, J. W.; Ibers, J. A. *Inorg. Chem.* **1988**, *27*, 3054.
- (46) (a) Collman, J. P.; Hutchison, J. E.; Lopez, M. A.; Guillard, R. *J. Am. Chem. Soc.* **1992**, *114*, 8066. (b) Collman, J. P.; Hutchison, J. E.; Ennis, M. S.; Lopez, M. A.; Guillard, R. *Ibid.* **1992**, *114*, 8074.
- (47) Joachim, J. E.; Apostolidis, C.; Kanellakopulos, B.; Maier, R.; Meyer, D.; Rebizant, J.; Ziegler, M. L. *J. Organomet. Chem.* **1993**, *445*, 137.
- (48) Reinaud, O. M.; Theopold, K. H. *J. Am. Chem. Soc.* **1994**, *116*, 6979.
- (49) Duchateau, R.; Gambarotta, S.; Beydoun, N.; Bensimon, C. *J. Am. Chem. Soc.* **1991**, *113*, 8986.
- (50) Kol, M.; Schrock, R. R.; Kempe, R.; Davis, W. M. *J. Am. Chem. Soc.* **1994**, *116*, 4382.
- (51) Shih, K. Y.; Schrock, R. R.; Kempe, R. *J. Am. Chem. Soc.* **1994**, *116*, 8804.
- (52) (a) Kaul, B. B.; Hayes, R. K.; George, T. A. *J. Am. Chem. Soc.* **1990**, *112*, 2002. (b) George, T. A.; Kaul, B. B. *Inorg. Chem.* **1990**, *29*, 4969.
- (53) DuBois, D. L. *Inorg. Chem.* **1984**, *23*, 2047.
- (54) Koide, M.; Kato, M.; Tsuchida, E.; Kurimura, Y. *Makromol. Chem.* **1981**, *182*, 283.
- (55) Kurimura, Y.; Ohta, F.; Gohda, J.; Nishide, H.; Tsuchida, E. *Makromol. Chem.* **1982**, *183*, 2889.
- (56) (a) Mercer, M. *J. Chem. Soc., Dalton Trans.* **1974**, 1637. (b) Cradwick, P. D.; Chatt, J.; Crabtree, R. H.; Richards, R. L. *J. Chem. Soc., Chem. Commun.* **1975**, 351.
- (57) Schrock, R. R.; Kolodziej, R. M.; Liu, A. H.; Davis, W. M.; Vale, M. G. *J. Am. Chem. Soc.* **1990**, *112*, 4338.
- (58) Glassman, T. E.; Liu, A. H.; Schrock, R. R. *Inorg. Chem.* **1991**, *30*, 4723.
- (59) Mizobe, Y.; Yokobayashi, Y.; Oshita, H.; Takahashi, T.; Hidai, M. *Organometallics* **1994**, *13*, 3764.
- (60) (a) Krüger, C.; Tsay, Y.-H. *Angew. Chem., Int. Ed. Engl.* **1973**, *12*, 998. (b) Jonas, K.; Brauer, D. J.; Krüger, C.; Roberts, P. J.; Tsay, Y.-H. *J. Am. Chem. Soc.* **1976**, *98*, 74.
- (61) (a) Fryzuk, M. D.; Haddad, T. S.; Rettig, S. J. *J. Am. Chem. Soc.* **1990**, *112*, 8185. (b) Fryzuk, M. D.; Haddad, T. S.; Mylvaganam, M.; McConville, D. H.; Rettig, S. J. *Ibid.* **1993**, *115*, 2782.
- (62) Cohen, J. D.; Mylvaganam, M.; Fryzuk, M. D.; Loehr, T. M. *J. Am. Chem. Soc.* **1994**, *116*, 9529.
- (63) Blomberg, M. R.; Siegbahn, P. E. *J. Am. Chem. Soc.* **1993**, *115*, 6908.
- (64) Pez, G. P.; Apgar, P.; Crissey, R. K. *J. Am. Chem. Soc.* **1982**, *104*, 482.
- (65) Cusanelli, A.; Sutton, D. *J. Chem. Soc., Chem. Commun.* **1989**, 1719.
- (66) Jeffery, J.; Lappart, M. F.; Riley, P. I. *J. Organomet. Chem.* **1979**, *181*, 25.
- (67) Schwab, P.; Mahr, N.; Wolf, J.; Werner, H. *Angew. Chem., Int. Ed. Engl.* **1993**, *32*, 1480.
- (68) Klahn, A. H.; Sutton, D. *Organometallics* **1989**, *8*, 198.
- (69) Chatt, J.; Dilworth, J. R.; Leigh, G. J. *J. Chem. Soc., Dalton Trans.* **1973**, 612.
- (70) Ware, D. C.; Taube, H. *Inorg. Chem.* **1991**, *30*, 4605.
- (71) Che, C.-M.; Lam, H.-W.; Tong, W.-F.; Lai, T.-F.; Lau, T.-C. *J. Chem. Soc., Chem. Commun.* **1989**, 1883.
- (72) Sellmann, D. *Angew. Chem.* **1971**, *83*, 1017.
- (73) Sellmann, D. *J. Organomet. Chem.* **1972**, *36*, C27.
- (74) Churchill, M. R.; Li, Y.-J.; Theopold, K. H.; Schrock, R. R. *Inorg. Chem.* **1984**, *23*, 4472.
- (75) Rocklage, S. M.; Schrock, R. R. *J. Am. Chem. Soc.* **1982**, *104*, 3077.
- (76) Hughes, D. L.; Mohammed, M. Y.; Pickett, C. J. *J. Chem. Soc., Chem. Commun.* **1988**, 1481.
- (77) Mohammed, M. Y.; Pickett, C. J. *J. Chem. Soc., Chem. Commun.* **1988**, 1119.
- (78) (a) Pickett, C. J.; Talarmin, J. *Nature (London)* **1985**, *317*, 652. (b) Pickett, C. J.; Ryder, K. S.; Talarmin, J. *J. Chem. Soc., Dalton Trans.* **1986**, 1453.
- (79) Al-Salih, T. I.; Pickett, C. J. *J. Chem. Soc., Dalton Trans.* **1985**, 1255.
- (80) Thorneley, R. N. F.; Eady, R. R.; Lowe, D. J. *Nature (London)* **1978**, *272*, 557.
- (81) (a) Chatt, J.; Pearman, A. J.; Richards, R. L. *Nature (London)* **1975**, *253*, 39. (b) *J. Chem. Soc., Dalton Trans.* **1977**, 1852.
- (82) Deeth, R. J.; Field, C. N. *J. Chem. Soc., Dalton Trans.* **1994**, 1943 and references therein.
- (83) (a) Heath, G. A.; Mason, R.; Thomas, K. M. *J. Am. Chem. Soc.* **1974**, *96*, 259. (b) Hidai, M.; Kodama, T.; Sato, M.; Harakawa, M.; Uchida, Y. *Inorg. Chem.* **1976**, *15*, 2694. (c) Nishihara, H.; Mori, T.; Tsurita, Y.; Nakano, K.; Saito, T.; Sasaki, Y. *J. Am. Chem. Soc.* **1982**, *104*, 4367. (d) George, T. A.; Kaul, B. B.; Chen, Q.; Zubieta, J. *Inorg. Chem.* **1993**, *32*, 1706. (e) Hanson, I. R.; Hughes, D. L. *J. Chem. Soc., Dalton Trans.* **1981**, 390. (f) Chatt, J.; Pearman, A. J.; Richards, R. L. *Ibid.* **1978**, 1766. (g) Abu Bakar, M.; Hughes, D. L.; Hussain, W.; Leigh, G. J.; Macdonald, C. J.; Mohd-Ali, H. *Ibid.* **1988**, 2545. (h) Barclay, J. E.; Hills, A.; Hughes, D. L.; Leigh, G. J.; Macdonald, C. J.; Abu Bakar, M.; Mohd-Ali, H. *Ibid.* **1990**, 2503.
- (84) Chatt, J.; Pearman, A. J.; Richards, R. L. *J. Chem. Soc., Dalton Trans.* **1976**, 1520.
- (85) Galindo, A.; Hills, A.; Hughes, D. L.; Richards, R. L.; Hughes, M.; Mason, J. *J. Chem. Soc., Dalton Trans.* **1990**, 283.
- (86) (a) Takahashi, T.; Mizobe, Y.; Sato, M.; Uchida, Y.; Hidai, M. *J. Am. Chem. Soc.* **1980**, *102*, 7461; (b) **1979**, *101*, 3405.
- (87) Chatt, J.; Fakley, M. E.; Hitchcock, P. B.; Richards, R. L.; Luong-Thi, N. T. *J. Chem. Soc., Dalton Trans.* **1982**, 345.
- (88) Baumann, J. A.; Bossard, G. E.; George, T. A.; Howell, D. B.; Koczon, L. M.; Lester, R. K.; Noddings, C. M. *Inorg. Chem.* **1985**, *24*, 3568.
- (89) George, T. A.; Kaul, B. B. *Inorg. Chem.* **1991**, *30*, 883.
- (90) Yamamoto, A.; Miura, Y.; Ito, T.; Chen, H.-L.; Iri, K.; Ozawa, F.; Miki, K.; Sei, T.; Tanaka, N.; Kasai, N. *Organometallics* **1983**, *2*, 1429.
- (91) (a) Hills, A.; Hughes, D. L.; Jimenez-Tenorio, M.; Leigh, G. J.; Rowley, A. T. *J. Chem. Soc., Dalton Trans.* **1993**, 3041. (b) Leigh, G. J.; Jimenez-Tenorio, M. *J. Am. Chem. Soc.* **1991**, *113*, 5862.
- (92) Komiya, S.; Akita, M.; Yoza, A.; Kasuga, N.; Fukuoka, A.; Kai, Y. *J. Chem. Soc., Chem. Commun.* **1993**, 787.
- (93) Jonas, K.; Krüger, C. *Angew. Chem., Int. Ed. Engl.* **1980**, *19*, 520.
- (94) (a) Sanner, R. D.; Manriquez, R. E.; Marsh, R. E.; Bercaw, J. E. *J. Am. Chem. Soc.* **1976**, *98*, 8351. (b) Manriquez, R. E.; Sanner, R. D.; Marsh, R. E.; Bercaw, J. E. *Ibid.* **1976**, *98*, 3042.
- (95) Leigh, G. J.; Prieto-Alcon, R.; Sanders, J. R. *J. Chem. Soc., Chem. Commun.* **1991**, 921.
- (96) (a) Henderson, R. A.; Morgan, S. H.; Stephens, A. N. *J. Chem. Soc., Dalton Trans.* **1990**, 1101. (b) Henderson, R. A.; Morgan, S. H. *Ibid.* **1990**, 1107.
- (97) Lane, J. D.; Henderson, R. A. *J. Chem. Soc., Dalton Trans.* **1987**, 197.
- (98) See, for example: Burgess, B. K. in ref 13, p 161.
- (99) Schrock, R. R.; Glassman, T. E.; Vale, M. G.; Kol, M. *J. Am. Chem. Soc.* **1993**, *115*, 1760.
- (100) Vale, M. G.; Schrock, R. R. *Inorg. Chem.* **1993**, *32*, 2767.

- (101) Kuwata, S.; Mizobe, Y.; Hidai, M. *Inorg. Chem.* **1994**, *33*, 3619.
- (102) Block, E.; Ofori-Okai, G.; Kang, H.; Zubieta, J. *J. Am. Chem. Soc.* **1992**, *114*, 758.
- (103) Coucouvanis, D.; Mosier, P. E.; Demadis, K. D.; Patton, S.; Malinak, S. M.; Kim, C. G.; Tyson, M. A. *J. Am. Chem. Soc.* **1993**, *115*, 12193.
- (104) Palermo, R. E.; Singh, R. E.; Bashkin, J. K.; Holm, R. H. *J. Am. Chem. Soc.* **1984**, *106*, 2600.
- (105) Mosier, P. E.; Kim, C. G.; Coucouvanis, D. *Inorg. Chem.* **1993**, *32*, 2620.
- (106) Demadis, K.; Coucouvanis, D. *Inorg. Chem.* **1994**, *33*, 4195.
- (107) Vol'pin, M. E. *J. Organomet. Chem.* **1980**, *200*, 319 and references therein.
- (108) (a) Chatt, J.; Diamantis, A. A.; Heath, G. A.; Hooper, N. E.; Leigh, G. J. *J. Chem. Soc., Dalton Trans.* **1977**, 688. (b) Chatt, J.; Heath, G. A.; Leigh, G. J. *J. Chem. Soc., Chem. Commun.* **1972**, 444.
- (109) Sato, M.; Kodama, T.; Hidai, M.; Uchida, Y. *J. Organomet. Chem.* **1978**, *152*, 239.
- (110) Bossard, G. E.; Busby, D. C.; Chang, M.; George, T. A.; Iske, S. D. A., Jr. *J. Am. Chem. Soc.* **1980**, *102*, 1001.
- (111) Chatt, J.; Head, R. A.; Leigh, G. J.; Pickett, C. J. *J. Chem. Soc., Dalton Trans.* **1978**, 1638.
- (112) Hussain, W.; Leigh, G. J.; Mohd-Ali, H.; Pickett, C. J. *J. Chem. Soc., Dalton Trans.* **1986**, 1473.
- (113) (a) Abu Bakar, M.; Hughes, D. L.; Leigh, G. J. *J. Chem. Soc., Dalton Trans.* **1988**, 2525. (b) Chatt, J.; Hussain, W.; Leigh, G. J.; Terreros, F. P. *J. Chem. Soc., Dalton Trans.* **1980**, 1408.
- (114) Ben-Shoshan, R.; Chatt, J.; Leigh, G. J.; Hussain, W. *J. Chem. Soc., Dalton Trans.* **1980**, 771.
- (115) Yoshida, T.; Adachi, T.; Ueda, T.; Kaminaka, M.; Sasaki, N.; Higuchi, T.; Aoshima, T.; Mega, I.; Mizobe, Y.; Hidai, M. *Angew. Chem., Int. Ed. Engl.* **1989**, *28*, 1040.
- (116) Iwanami, K.; Mizobe, Y.; Takahashi, T.; Kodama, T.; Uchida, Y.; Hidai, M. *Bull. Chem. Soc. Jpn.* **1981**, *54*, 1773.
- (117) Aoshima, T.; Mizobe, Y.; Hidai, M.; Tsuchiya, J. *J. Organomet. Chem.* **1992**, *423*, 39.
- (118) (a) Hidai, M.; Mizobe, Y.; Sato, M.; Kodama, T.; Uchida, Y. *J. Am. Chem. Soc.* **1978**, *100*, 5740. (b) Bevan, P. C.; Chatt, J.; Hidai, M.; Leigh, G. J. *J. Organomet. Chem.* **1978**, *160*, 165. (d) Mizobe, Y.; Uchida, Y.; Hidai, M. *Bull. Chem. Soc. Jpn.* **1980**, *53*, 1781. (e) Mizobe, Y.; Ono, R.; Uchida, Y.; Hidai, M.; Tezuka, M.; Moue, S.; Tsuchiya, A. *J. Organomet. Chem.* **1981**, *204*, 377. (f) Hidai, M.; Mizobe, Y.; Uchida, Y. *J. Am. Chem. Soc.* **1976**, *98*, 7824.
- (119) Hidai, M.; Aramaki, S.; Yoshida, K.; Kodama, T.; Takahashi, T.; Uchida, Y.; Mizobe, Y. *J. Am. Chem. Soc.* **1986**, *108*, 1562.
- (120) Recent review on the reactions of coordinated diazoalkanes: Mizobe, Y.; Ishii, Y.; Hidai, M. *Coord. Chem. Rev.* **1995**, *139*, 281.
- (121) (a) Ishii, Y.; Miyagi, H.; Jitsukuni, S.; Seino, H.; Harkness, B. S.; Hidai, M. *J. Am. Chem. Soc.* **1992**, *114*, 9890. (b) Ishii, Y.; Miyagi, H.; Hidai, M. *J. Chem. Soc., Chem. Commun.* **1990**, 1569. (c) Miyagi, H.; Ishii, Y.; Aoshima, T.; Mizobe, Y.; Hidai, M. *Chem. Lett.* **1991**, 611.
- (122) Colquhoun, H. M.; Crease, A. E.; Taylor, S. A.; Williams, D. J. *J. Chem. Soc., Dalton Trans.* **1988**, 2781.
- (123) Seino, H.; Ishii, Y.; Hidai, M. *Organometallics* **1994**, *13*, 364.
- (124) Aoshima, T.; Tamura, T.; Mizobe, Y.; Hidai, M. *J. Organomet. Chem.* **1992**, *435*, 85.
- (125) Harada, Y.; Aoshima, T.; Mizobe, Y.; Oshita, H.; Hidai, M. *Chem. Lett.* **1994**, 797.
- (126) Aoshima, T.; Tanase, T.; Mizobe, Y.; Yamamoto, Y.; Hidai, M. *J. Chem. Soc., Chem. Commun.* **1992**, 586.
- (127) Walsh, P. J.; Baranger, A. M.; Bergman, R. G. *J. Am. Chem. Soc.* **1992**, *114*, 1708.
- (128) Glueck, D. S.; Wu, J.; Hollander, F. J.; Bergman, R. G. *J. Am. Chem. Soc.* **1991**, *113*, 2041.
- (129) Oshita, H.; Mizobe, Y.; Hidai, M. *J. Organomet. Chem.* **1993**, *461*, 43.
- (130) Seino, H.; Ishii, Y.; Hidai, M. *J. Am. Chem. Soc.* **1994**, *116*, 7433.
- (131) Bevan, P. C.; Chatt, J.; Leigh, G. J.; Leelamani, E. G. *J. Organomet. Chem.* **1977**, *139*, C59.
- (132) (a) Hussain, W.; Leigh, G. J.; Pickett, C. J. *J. Chem. Soc., Chem. Commun.* **1982**, 747. (b) Henderson, R. A.; Leigh, G. J.; Pickett, C. J. *J. Chem. Soc., Dalton Trans.* **1989**, 425.
- (133) (a) Watakabe, A.; Takahashi, T.; Jin, D.-M.; Yokotake, I.; Uchida, Y.; Hidai, M. *J. Organomet. Chem.* **1983**, *254*, 75. (b) Hidai, M.; Kurano, M.; Mizobe, Y. *Bull. Chem. Soc. Jpn.* **1985**, *58*, 2719.
- (134) Pickett, C. J.; Leigh, G. J. *J. Chem. Soc., Chem. Commun.* **1981**, 1033.
- (135) Colquhoun, H. M.; Crease, A. E.; Taylor, S. A. *J. Chem. Soc., Chem. Commun.* **1983**, 1158.
- (136) Yamamoto, A.; Go, S.; Ookawa, M.; Takahashi, M.; Ikeda, S.; Keii, T. *Bull. Chem. Soc. Jpn.* **1972**, *45*, 3110.
- (137) Sobota, P.; Jezowska-Trzebiatowska, B.; Janas, Z. *J. Organomet. Chem.* **1976**, *118*, 253.
- (138) Mori, M.; Uozumi, Y.; Shibusaki, M. *J. Organomet. Chem.* **1990**, *395*, 225.
- (139) (a) Uozumi, Y.; Kawasaki, N.; Mori, E.; Mori, M.; Shibusaki, M. *J. Am. Chem. Soc.* **1989**, *111*, 3725. (b) Uozumi, Y.; Mori, E.; Mori, M.; Shibusaki, M. *J. Organomet. Chem.* **1990**, *399*, 93.
- (140) (a) Uozumi, Y.; Mori, M.; Shibusaki, M. *J. Chem. Soc., Chem. Commun.* **1991**, 81. (b) Mori, M.; Uozumi, Y.; Shibusaki, M. *Heterocycles* **1992**, *33*, 819.
- (141) (a) Kawaguchi, M.; Hamaoka, S.; Mori, M. *Tetrahedron Lett.* **1993**, *34*, 6907. (b) Mori, M.; Kawaguchi, M.; Hori, M.; Hamaoka, S. *Heterocycles* **1994**, *39*, 729.
- (142) Shiina, K. *J. Am. Chem. Soc.* **1972**, *94*, 9266.
- (143) Hori, M.; Mori, M. *J. Org. Chem.* **1995**, *60*, 1480.
- (144) Hughes, D. L.; Ibrahim, S. K.; Macdonald, C. J.; Mohd. Ali, H.; Pickett, C. J. *J. Chem. Soc., Chem. Commun.* **1992**, 1762.
- (145) (a) Hidai, M.; Komori, K.; Kodama, T.; Jin, D.-M.; Takahashi, T.; Sugiura, S.; Uchida, Y.; Mizobe, Y. *J. Organomet. Chem.* **1984**, *272*, 155. (b) Komori, K.; Kodama, T.; Jin, D.-M.; Takahashi, T.; Uchida, Y.; Hidai, M. *Chem. Lett.* **1983**, 465.
- (146) Komori, K.; Sugiura, S.; Mizobe, Y.; Yamada, M.; Hidai, M. *Bull. Chem. Soc. Jpn.* **1989**, *62*, 2953.
- (147) Oshita, H.; Mizobe, Y.; Hidai, M. *J. Organomet. Chem.* **1993**, *456*, 213.
- (148) (a) Oshita, H.; Mizobe, Y.; Hidai, M. *Chem. Lett.* **1990**, 1303; (b) *Organometallics* **1992**, *11*, 4116.
- (149) Street, A. C.; Mizobe, Y.; Gotoh, F.; Mega, I.; Oshita, H.; Hidai, M. *Chem. Lett.* **1991**, 383.
- (150) Komori, K.; Oshita, H.; Mizobe, Y.; Hidai, M. *J. Am. Chem. Soc.* **1989**, *111*, 1939.
- (151) Donovan-Mtunzi, S.; Richards, R. L.; Mason, J. *J. Chem. Soc., Dalton Trans.* **1984**, 2429; see also references therein.
- (152) Takahashi, T.; Kodama, T.; Watakabe, A.; Uchida, Y.; Hidai, M. *J. Am. Chem. Soc.* **1983**, *105*, 1680.
- (153) Klein, H.-F.; König, H.; Koppert, S.; Ellrich, K.; Riede, J. *Organometallics* **1987**, *6*, 1341.
- (154) Sellmann, D.; Weiss, W. *J. Organomet. Chem.* **1978**, *160*, 183.
- (155) Hidai, M.; Takahashi, T.; Yokotake, I.; Uchida, Y. *Chem. Lett.* **1980**, 645.
- (156) Nishihara, H.; Mori, T.; Saito, T.; Sasaki, Y. *Chem. Lett.* **1980**, 667.
- (157) Jia, G.; Morris, R. H.; Schweitzer, C. T. *Inorg. Chem.* **1991**, *30*, 593.
- (158) Colquhoun, H. M. *J. Chem. Res., (Synop.)* **1979**, 325.
- (159) (a) Ishii, Y.; Kawaguchi, M.; Ishino, Y.; Aoki, T.; Hidai, M. *Organometallics* **1994**, *13*, 5062. (b) Ishii, Y.; Ishino, Y.; Aoki, T.; Hidai, M. *J. Am. Chem. Soc.* **1992**, *114*, 5429.

CR940323B