

Comparison of the Nucleophilicities of Alkynes and Alkenes. Quantitative Determination of the Nucleophilicities of Alkynes toward Carbenium Ions

Herbert Mayr^{*a}, Jose L. Gonzalez^a, and Kerstin Lüdtke^b

Institut für Organische Chemie der Technischen Hochschule Darmstadt^a,
Petersenstraße 22, 64287 Darmstadt, Germany

Institut für Chemie der Medizinischen Universität zu Lübeck^b,
Ratzeburger Allee 160, 23538 Lübeck, Germany

Received August 4, 1993

Key Words: Alkynes / Alkenes / Nucleophilicity / Carbenium ions / Vinyl cations / Kinetics

Relative reactivities of alkenes and alkynes toward diarylmethyl cations have been determined by direct rate measurements and by competition experiments in dichloromethane. At -70°C alkynes are found to be one to two orders of magnitude less reactive than analogously substituted alkenes

(e.g. phenylacetylene/styrene), but the reactivity difference reduces strongly as the temperature is raised. The stereochemistry of the vinyl chlorides produced by addition of benzhydryl chlorides to alkynes is characterized.

Alkynes react much more slowly with halogens than alkenes and are, therefore, often considered to be weak nucleophiles^[1]. On the other hand, protons attack alkynes and structurally related alkenes with comparable rates^[2,3], and carbenium ions have also been reported to reveal similar reactivity toward phenylacetylene and styrene^[4]. An explanation for these phenomena has been presented by Modena et al.^[5]: Electrophiles with strong bridging tendency react faster with alkenes than with alkynes, while electrophiles with weak bridging tendency show similar reactivities toward alkenes and alkynes.

We have recently studied the kinetics of the reactions of diarylmethyl cations with alkenes^[6–10], allyl-element compounds^[11,12], enol ethers^[13], and arenes^[14]. On the basis of these data we have set up a reactivity scale including different types of π nucleophiles^[14].

In order to add alkynes to this scale, we have measured the rate of the reaction of the 4-methoxybenzhydryl cation with phenylacetylene in CH_2Cl_2 at -70°C . Surprisingly, a rate constant ($k_2 = 0.072 \text{ l mol}^{-1} \text{ s}^{-1}$) 150 times smaller than that of styrene^[8] has been found, while Modena's competition experiments with $\text{Ph}_2\text{CHCl}/\text{ZnCl}_2$ in CH_2Cl_2 indicate comparable reactivities for structurally analogous alkenes and alkynes ($k_{\text{styrene}}/k_{\text{phenylacetylene}} = 3.8$)^[4].

We have, therefore, reinvestigated the problem of relative reactivities of alkenes and alkynes, and we will show that the discrepancy between Modena's competition experiments and our direct rate measurements is predominantly due to the different temperature used for the two types of experiments.

Reaction Products

As previously described by Modena and coworkers^[15], the ZnCl_2 -catalyzed reaction of chlorodiphenylmethane (**1a**) with phenylacetylene (**2**) gives a mixture of (*E*)-**3a** and (*Z*)-**3a**, the ratio of which is controlled by the different steric shielding of the two sites of **3⁺** (Scheme 1). A similar behavior has now been found for the two benzhydryl chlorides **1b** and **1c**.


While 1-phenyl-1-propyne (**4**) and ToI_2CHCl (**1b**) react with predominant formation of the (*E*)-isomer of **5b**^[4], the analogous reaction with 1-hexyne (**6**) gives a 1:1 mixture of the two stereoisomeric adducts **7b**, possibly due to an activationless collapse of the intermediate α -alkyl-substituted vinyl cation with the negative counterion (Scheme 2).

Compounds **8–10**, the products of the addition of diarylchloromethanes to styrene, *trans*-1-phenyl-1-propene, and 2-chloropropene, which are needed as reference compounds for the evaluation of the competition experiments, have been synthesized according to literature procedures^[16,17].

Determination of Absolute Rates




Spectrophotometry has been the method of choice in previous kinetic investigations of the reactions of diarylmethyl cations with π nucleophiles, since the solutions of the diarylmethyl cations completely fade upon addition of the CC double bond-containing nucleophiles^[7,12]. When alkynes are added to the diarylcarbenium ion solutions, only a *change of color* is observed, however, and the absorptions at λ_{max} of the carbenium ions do not usually disappear completely during the reactions.

Scheme 1



R, R'	1, 3		
H, H	a	9	1
Me, Me	b	7	1
OMe, H	c	8	1


Scheme 2

**10b**Table 1. Rate constants for the reactions of diarylmethyl cations with alkynes in CH₂Cl₂ at -70°C

Ar ₂ CH ⁺	Alkyne	MCl _n	k ₂ / l mol ⁻¹ s ⁻¹
1a⁺	Ph-C≡CH	(2) TiCl ₄	3.9 × 10 ²
1b⁺	Ph-C≡CH	(2) TiCl ₄	3.8 ^[a]
	Ph-C≡CH	(2) BCl ₃	3.9
1c⁺	Ph-C≡CH	(2) TiCl ₄	7.2 × 10 ⁻²
1b⁺	Ph-C≡C-Me	(4) TiCl ₄	4.3 × 10 ¹
	Ph-C≡C-Me	(4) BCl ₃	4.6 × 10 ¹
1b⁺	nC ₄ H ₉ -C≡CH	(6) TiCl ₄	1.6 × 10 ⁻²

^[a] ΔH[‡] = 25.56 ± 0.97 kJ mol⁻¹, ΔS[‡] = -104.0 ± 4.2 J mol⁻¹ K⁻¹.

Scheme 3



Conductivity measurements indicate that the colored by-products are of ionic nature, but we have not been able to reveal their identity. Possible explanations such as acid-catalyzed dimerizations of alkynes with formation of cyclobutenylium ions or hydride abstractions from **3**, **5**, and **7** with formation of aryl-substituted allyl cations, could not be substantiated. These side reactions disturb the initial period of the reactions only slightly, and consistent rate constants have been obtained by monitoring the course of the reactions at a low degree of conversion (see Table 3). As in analogous reactions with alkenes^[7] and other vinyl derivatives^[12,18], exchange of the carbenium counterion does not significantly affect the reaction rates (Table 1).

Nevertheless, a comparison of the individual kinetic runs (Table 3) shows that the averaged k_2 values given in Table 1 are much less accurate than those previously determined for alkenes^[6-9] and allyl-element compounds^[12]. We have, therefore, considered it necessary to confirm these data by competition experiments which would provide an additional link to the previously reported nucleophilicities of CC double bond-containing systems.

Determination of Relative Rates

When carbenium ions are generated in situ in the presence of two nucleophiles, the relative reactivities κ can be derived from eq. (1)^[19] (Scheme 3).

Table 2. Relative reactivities of alkynes and alkenes toward diarylmethyl cations in CH_2Cl_2 ($\text{ZnCl}_2 \cdot \text{OEt}_2$, -70°C)

Ar_2CHCl	$\text{Nu}_x / \text{Nu}_y$	$T / ^\circ\text{C}$	k_x / k_y
1a	Ph-CH=CH ₂ /	-70	214
		Ph-C≡CH	-40
	Ph-C≡CH	-10	35.6
		20	28.1
1b	Ph-CH=CH ₂ /	-70	164
		Ph-C≡CH	-40
	Ph-C≡CH	-10	27.6
		20	15.7
1b	(E)-Ph-CH=CH-CH ₃ /	-70	9.45
	Ph-C≡C-CH ₃		
1b	Cl(CH ₃)C=CH ₂ /	-20	1.37
	n-C ₄ H ₉ -C≡CH		

$$\kappa = k_x/k_y = \frac{\lg[\text{Nu}_x]_0 - \lg([\text{Nu}_x]_0 - [\text{P}_x])}{\lg[\text{Nu}_y]_0 - \lg([\text{Nu}_y]_0 - [\text{P}_y])} \quad (1)$$

The concentrations of the products $[\text{P}_x]$ and $[\text{P}_y]$, which are needed for the calculation of κ according to eq. (1), are obtained by HPLC analysis of the product mixtures using an internal standard. On the assumption that **1** is quantitatively converted into P_x and P_y , eq. (1) can be transformed into eq. (2)^[20], which allows us to calculate κ from the product ratio $R = [\text{P}_x]/[\text{P}_y]$ and does not require the determination of absolute product yields by use of an internal standard (Table 2). Independence of the competition constant of the reactant ratios justifies this procedure.

$$\kappa = k_x/k_y = \lg\left(1 - \frac{[\text{I}]_0 R}{[\text{Nu}_x]_0(1+R)}\right) / \lg\left(1 - \frac{[\text{I}]_0}{[\text{Nu}_y]_0(1+R)}\right) \quad (2)$$

with $R = [\text{P}_x]/[\text{P}_y]$

Discussion

In a previous work, a good agreement between relative olefin reactivities derived from direct rate measurements and from competition experiments has been observed, even





Figure 1. Relative reactivities of styrene/phenylacetylene (**2**) [$\kappa = \lg(k_{\text{styrene}}/k_{\text{phenylacetylene}})$] as a function of temperature

Scheme 4. Reactivity scale for π nucleophiles toward (*p*-MeC₆H₄)₂CH⁺ at -70°C (based on absolute rate constants)

[a] Extrapolated value.

when different Lewis acids were employed for the ionization of the benzhydryl chlorides. We have now found a styrene/phenylacetylene (**2**) reactivity ratio of 277 (ToI_2CH^+ , -70°C) from directly measured rates (counterion BCl_4^- or TiCl_5^- , Table 1 and ref.^[9]) and of 164 (counterion ZnCl_3^-) from competition experiments. Under the same conditions *trans*-1-phenyl-1-propene has been found to be 22 times more reactive than 1-phenyl-1-propyne (**4**) on the basis of absolute rate constants (counterion BCl_4^- or TiCl_5^- , Table 1 and ref.^[9]), while competition experiments yield the smaller ratio of 9.5 (counterion ZnCl_3^-). Though the deviations between the two methods are unusually large for these examples, probably because of the problems connected with the determination of absolute rate constants for alkynes, one can now be sure that the order of magnitude determined for alkyne reactivities is correct. Rough agreement between competition experiments ($k_x/k_y = 1.37$ at -20°C)

Table 3. Kinetics of the reactions of diarylmethyl cations with alkynes

1	[1] ₀ / mol l ⁻¹	[alkyne] ₀ / mol l ⁻¹	Lewis acid	[Lewis acid] ₀ / mol l ⁻¹	T / °C	Quantity observed (up to % conversion) ^[a]	k ₂ / l mol ⁻¹ s ⁻¹ ^[b]
<i>Phenylacetylene (2)</i>							
1 a	4.6 × 10 ⁻⁵	1.1 × 10 ⁻³	TiCl ₄	3.7 × 10 ⁻²	-70	A (81) / C (54)	4.52 × 10 ² / 3.27 × 10 ²
	4.6 × 10 ⁻⁵	5.3 × 10 ⁻⁴	TiCl ₄	3.7 × 10 ⁻²	-70	A (79) / C (52)	4.42 × 10 ² / 3.19 × 10 ²
1 b	6.7 × 10 ⁻⁵	5.5 × 10 ⁻⁴	TiCl ₄	7.1 × 10 ⁻⁴	-70	A (62) / C (23)	3.93 / 3.71
	6.0 × 10 ⁻⁵	4.8 × 10 ⁻⁴	TiCl ₄	5.5 × 10 ⁻⁴	-70	A (62) / C (24)	3.86 / 3.60
	1.1 × 10 ⁻⁴	1.1 × 10 ⁻³	BCl ₃	1.3 × 10 ⁻²	-70	A (60) / C (24)	3.89 / 3.88
	6.0 × 10 ⁻⁵	5.5 × 10 ⁻⁴	BCl ₃	2.6 × 10 ⁻²	-70	A (59) / C (23)	3.77 / 3.73
	5.4 × 10 ⁻⁵	5.5 × 10 ⁻⁴	TiCl ₄	1.2 × 10 ⁻⁴	-40	A (25)	4.50 × 10 ¹
	5.4 × 10 ⁻⁵	4.3 × 10 ⁻⁴	TiCl ₄	1.2 × 10 ⁻⁴	-40	A (24)	4.46 × 10 ¹
	5.4 × 10 ⁻⁵	5.5 × 10 ⁻⁴	TiCl ₄	1.2 × 10 ⁻³	-10	A (26)	2.06 × 10 ²
	5.4 × 10 ⁻⁵	8.3 × 10 ⁻⁴	TiCl ₄	1.2 × 10 ⁻³	-10	A (25)	2.06 × 10 ²
	5.4 × 10 ⁻⁵	5.5 × 10 ⁻⁴	TiCl ₄	1.2 × 10 ⁻³	+20	A (47)	4.98 × 10 ²
	5.4 × 10 ⁻⁵	4.3 × 10 ⁻⁴	TiCl ₄	1.2 × 10 ⁻³	+20	A (45)	4.87 × 10 ²
1 c	8.0 × 10 ⁻⁵	5.2 × 10 ⁻³	TiCl ₄	8.0 × 10 ⁻⁴	-70	A (17) / C (13)	7.8 × 10 ⁻² / 6.4 × 10 ⁻²
	2.0 × 10 ⁻⁴	1.1 × 10 ⁻³	TiCl ₄	8.0 × 10 ⁻⁴	-70	A (17) / C (12)	7.1 × 10 ⁻² / 6.9 × 10 ⁻²
	9.8 × 10 ⁻⁵	1.0 × 10 ⁻³	BCl ₃	2.7 × 10 ⁻²	-70	A (57)	8.0 × 10 ⁻² ^[c]
<i>1-Phenyl-1-propyne (4)</i>							
1 b	6.7 × 10 ⁻⁵	5.7 × 10 ⁻⁴	TiCl ₄	6.9 × 10 ⁻⁴	-70	A (18) / C (19)	6.23 × 10 ¹ / 4.29 × 10 ¹
	6.7 × 10 ⁻⁵	7.5 × 10 ⁻⁴	TiCl ₄	6.9 × 10 ⁻⁴	-70	A (19) / C (19)	6.14 × 10 ¹ / 4.29 × 10 ¹
	6.7 × 10 ⁻⁵	5.7 × 10 ⁻⁴	BCl ₃	2.6 × 10 ⁻²	-70	A (32) / C (34)	6.16 × 10 ¹ / 4.62 × 10 ¹
	6.7 × 10 ⁻⁵	7.5 × 10 ⁻⁴	BCl ₃	2.6 × 10 ⁻²	-70	A (33) / C (34)	6.05 × 10 ¹ / 4.82 × 10 ¹
<i>1-Hexyne (6)</i>							
1 b	6.1 × 10 ⁻⁵	5.2 × 10 ⁻⁴	TiCl ₄	6.9 × 10 ⁻⁴	-70	A (7) / C (17)	1.43 × 10 ⁻² / 1.85 × 10 ⁻²
	6.0 × 10 ⁻⁵	5.2 × 10 ⁻⁴	TiCl ₄	7.4 × 10 ⁻⁴	-70	A (8) / C (22)	1.32 × 10 ⁻² / 2.2 × 10 ⁻²

^[a] A = determination of absorbance; C = determination of conductance. — ^[b] If two values are given, the first refers to absorbance, the second to conductance. — ^[c] Reaction performed in the presence of [TEBA] = 4.39 · 10⁻³ mol l⁻¹.

and the results of absolute rate determinations ($k_x/k_y = 1.7$ at -70°C) has also been found for the couple 2-chloro-1-propene/1-hexyne (**6**), when a preliminary rate constant for the reaction of **1b** with 2-chloro-1-propene ($2.7 \cdot 10^{-2}$ l mol⁻¹ s⁻¹, -70°C)^[21] is employed.

A further congruence of both methods is noticeable. While ΔS^\ddagger for the reaction of diarylcarbenium ions with styrene has been determined to be -122 ± 5 J mol⁻¹ K⁻¹, similar to other terminal vinyl derivatives^[8], the activation entropy for the reaction of **1b** with phenylacetylene (-104 J mol⁻¹ K⁻¹) has been found to be considerably less negative, indicating a less ordered transition state for carbenium ion additions to alkynes than to alkenes. On the other hand, ΔH^\ddagger is greater for additions to phenylacetylene (**2**) than to styrene with the consequence that the alkyne additions have a higher temperature gradient than the alkene additions. This effect is also observed in competition experiments at variable temperature (Table 2): While styrene is two orders of magnitude more reactive than phenylacetylene (**2**) at -70°C , this ratio reduces to one order of magnitude at 20°C , and one can extrapolate that a $T \geq 300^\circ\text{C}$ **2** would react faster with diarylmethyl cations than styrene. An Eyring plot of the relative rates of the reaction of styrene and **2** with **1b** (Figure 1) yields $\Delta\Delta H^\ddagger = 13.0$ kJ mol⁻¹ and $\Delta\Delta S^\ddagger = 21.6$ J mol⁻¹ K⁻¹, which agrees fairly well with the average $\Delta\Delta S^\ddagger$ of 18 J mol⁻¹ K⁻¹ derived from measurements of absolute rates.

It is now clear that different reaction temperatures are the main reason for the deviating alkene/alkyne reactivity

ratio observed by Modena (40°C , competition method) and by our group (-70°C , direct rate measurements). An explanation why the alkene/alkyne ratios determined in this work are still somewhat larger than the previously reported reactivity ratios^[4] is hard to find, as details of the evaluation of the previous experiments have not been given^[4]. Possibly it is due to the fact that the previous experiments^[4] have been performed under heterogeneous conditions.

Conclusion

A comparison of the rate constants for the reactions of the bis(4-methylphenyl)methyl cation (**1b**) with various π nucleophiles (Scheme 4) shows that at -70°C alkynes are somewhat less reactive than structurally analogous alkenes. As previously discussed by Modena^[4,5], the reactivity of alkynes toward carbenium ions is comparable to that of ordinary alkenes, and alkynes cannot generally be regarded as weak nucleophiles. The ranking in Scheme 4 is only slightly affected when relative rates from competition experiments are used as the basis. Since the relative reactivity of styrene and phenylacetylene changes only slightly, when different reference electrophiles are employed (Table 1 and ref.^[9]), one can conclude that the reactivity order in Scheme 4 also holds roughly for other carbon electrophiles. It has to be kept in mind, however, that the relative reactivities of styrene/phenylacetylene and *trans*-1-phenyl-1-propene/1-phenyl-1-propyne (**4**) decrease dramatically as the temperature is increased, implying that the nucleophilicity order in Scheme 4 is strongly affected by temperature.

We thank Ms. M. Deters and Ms. H. Freter for experimental assistance (HPLC separations), the *Deutsche Forschungsgemeinschaft* and the *Fonds der Chemischen Industrie* for financial support.

Experimental

NMR: XL 200 (Varian) and WM 300 (Bruker). – MS: 70–250 E (VG-Instruments). – UV: Beckmann Spectrophotometer UV 5240. – HPLC: Knauer HPLC pump 64 with Rheodyne 7125 valve, steel column (250 × 4 mm) filled with LiChrospher 100 RP 18 (5- μ m particles). – All alkynes and alkenes are commercially available. The diarylchloromethanes **1a–c** were prepared by treatment of the corresponding diarylmethanols with HCl^[22].

Preparation of the Addition Products

Chlorodiphenylmethane (1a) and Phenylacetylene (2): The reaction was carried out as described in ref.^[15] to give 62% of a mixture of (*E*)-**3a** and (*Z*)-**3a** (9:1) from which (*E*)-**3a** was isolated by fractional crystallization from ethanol: (*E*)-1-Chloro-1,3,3-triphenyl-1-propene [(*E*)-**3a**]: Colorless crystals with m.p. 65–66°C (ref.^[15] 65–66°C). – ¹H NMR (CDCl₃, 300 MHz): δ = 4.78 (d, *J* = 11.0 Hz, 1H, 3-H), 6.45 (d, *J* = 11.0 Hz, 1H, 2-H), 7.11–7.36 (m, 15H, aromatic H). – ¹³C NMR (CDCl₃): δ = 50.8 (d), 126.7 (d), 128.2 (d), 128.4 (d), 128.7 (d), 128.9 (d), 129.6 (d), 131.5 (d), 131.6 (s), 137.0 (s), 143.4 (s). – UV (methanol): λ_{\max} (lg ϵ) = 246 nm (3.962), 263 (sh), 269 (sh).

Pure (*Z*)-**3a**, which is needed for the HPLC analysis of the competition experiments (see below), was obtained by fractional recrystallization (ethanol) of a 10:1 mixture of (*Z*)-**3a** and (*E*)-**3a**. This mixture was prepared as described in ref.^[15] by treatment of 1,3,3-triphenyl-1-propanone with PCl₅ and subsequently with NEt₃. (*Z*)-1-Chloro-1,3,3-triphenyl-1-propene [(*Z*)-**3a**]: Colorless crystals with m.p. 77–79°C (ethanol). – ¹H NMR (CDCl₃, 300 MHz): δ = 5.42 (d, *J* = 9.5 Hz, 1H, 3-H), 6.60 (d, *J* = 9.5 Hz, 1H, 2-H), 7.11–7.61 (m, 15H, aromatic H). – ¹³C NMR (CDCl₃): δ = 50.8 (d), 126.6 (d), 128.3 (d), 128.6 (d), 129.6 (d), 133.4 (s), 137.9 (s), 143.0 (s). – UV (methanol): λ_{\max} (lg ϵ) = 253 nm (4.267), 264 (sh), 270 (sh), 291 (sh).

Chlorobis(4-methylphenyl)methane (1b) and Phenylacetylene (2): A solution of **1b** (1.37 g, 5.94 mmol) in CH₂Cl₂ (5 ml) was added dropwise to a cooled solution (–20°C) of **2** (613 mg, 6.00 mmol) and ZnCl₂(OEt₂)_{1.6} (1.20 mmol)^[23] in CH₂Cl₂ (20 ml). After 3.5 h the mixture was poured into 20 ml of conc. aqueous ammonia. The aqueous phase was extracted with 5 ml of CH₂Cl₂, and the combined organic phases were dried with MgSO₄ and concentrated to give 1.82 g (92%) of crude (*E,Z*)-**3b** (7:1, determined by ¹H-NMR analysis). Two crystallizations from ethanol gave a pure sample of the (*E*) isomer. (*E*)-1-Chloro-3,3-bis(4-methylphenyl)-1-phenyl-1-propene [(*E*)-**3b**]: Colorless needles with m.p. 81–83°C. – ¹H NMR (CDCl₃, 300 MHz): δ = 2.32 (s, 6H, CH₃), 4.70 (d, *J* = 11.0 Hz, 1H, 3-H), 6.41 (d, *J* = 11.0 Hz, 1H, 2-H), 6.99–7.35 (m, 13H, aromatic H). – ¹³C NMR (CDCl₃): δ = 21.0 (q), 49.9 (d), 128.0 (d), 128.3 (d), 128.6 (d), 128.8 (d), 129.3 (d), 131.0 (s), 131.7 (d), 136.1 (s), 137.0 (s), 140.5 (s). – UV (methanol): λ_{\max} (lg ϵ) = 222 nm (4.354), 246 (4.017, sh), 266 (sh), 275 (3.508, sh). – MS (70 eV), *m/z* (%): 334, 332 (4, 9) [M⁺], 317 (4), 297 (100), 296 (17), 281 (14), 204 (71), 165 (17), 105 (17). – C₂₃H₂₁Cl: calcd. 332.1332; found 332.1346 (MS).

(*Z*)-**3b** was not isolated as a pure compound and was detected in the 300-MHz ¹H-NMR spectrum by the resonances of 2-H (δ = 6.59, d, *J* = 9.0 Hz) and 3-H (δ = 5.36, d, *J* = 9.0 Hz).

Chloro(4-methoxyphenyl)phenylmethane (1c) and Phenylacetylene (2): A solution of **1c** (900 mg, 3.87 mmol) in CH₂Cl₂ (10 ml) was added dropwise to a solution of **2** (2.00 g, 19.6 mmol) and

ZnCl₂ (54 mg) in 20 ml of CH₂Cl₂. After 6 h (40°C) the mixture was poured into 20 ml of conc. aqueous ammonia. The aqueous phase was extracted with 5 ml of CH₂Cl₂, and the combined organic phases were dried with MgSO₄ and concentrated to give 800 mg (62%) of oily (*E,Z*)-**3c** (8:1, determined by ¹H-NMR analysis). (*E*)-1-Chloro-3-(4-methoxyphenyl)-1,3-diphenyl-1-propene: ¹H NMR (CDCl₃, 300 MHz): δ = 3.70 (s, 3H, OCH₃), 4.65 (d, *J* = 10.5 Hz, 1H, 3-H), 6.39 (d, *J* = 10.5 Hz, 1H, 2-H), 6.6–7.7 (m, 14H, aromatic H).

(*Z*)-**3c** was detected in the 300-MHz ¹H-NMR spectrum by the resonance of 3-H (δ = 5.18, d, *J* = 10.0 Hz).

Chlorobis(4-methylphenyl)methane (1b) and 1-Phenyl-1-propyne (4): A solution of ZnCl₂(OEt₂)_{1.6} (1.4 mmol) in 0.5 ml of CH₂Cl₂ was added to a cold (–70°C) solution of **1b** (1.50 g, 6.50 mmol) and **4** (1.85 g, 15.9 mmol) in CH₂Cl₂ (25 ml). After 15 h the deep-red solution was worked up as described above, and the mixture was purified by liquid chromatography (RP 18, methanol) to give (*E*)-**5b** (1.80 g, 80%) and 0.18 g of a second fraction containing (*Z*)-**5b** and a cyclic compound. (*E*)-1-Chloro-2-methyl-3,3-bis(4-methylphenyl)-1-phenyl-1-propene [(*E*)-**5b**]: Colorless needles with m.p. 205°C (ethanol). – ¹H NMR (CDCl₃, 200 MHz): δ = 1.89 (s, 3H, 2-CH₃), 2.34 (s, 6H, aryl-CH₃), 5.07 (s, 1H, 3-H), 6.93–7.32 (m, 13H, aromatic H). – ¹³C NMR: δ = 17.7 (q), 21.0 (q), 53.2 (d), 128.2 (d), 128.4 (d), 128.8 (d), 128.9 (d), 135.2 (s), 135.9 (s), 139.0 (s), 139.0 (s). – MS (70 eV), *m/z* (%): 348, 346 (7, 20) [M⁺], 311 (83), 310 (22), 295 (23), 219 (100), 195 (16). – C₂₄H₂₃Cl: calcd. 346.1488; found 346.1487 (MS).

Chlorobis(4-methylphenyl)methane (1b) and 1-Hexyne (6): At ambient temperature, a solution of **1b** (1.90 g, 8.23 mmol) in CH₂Cl₂ (5 ml) was added dropwise (5 min) to a solution of **6** (2.10 g, 25.6 mmol) and ZnCl₂(Et₂O)_{1.6} (5.0 mmol) in CH₂Cl₂ (10 ml). After 15 h, the dark-red mixture was worked up as described above, and an HPLC analysis indicated the production of (*E,Z*)-**7b** and of oligomers. Short-path distillation (180°C/6 · 10^{–4} Pa) and chromatographic purification (RP 18, methanol) yielded 0.80 g (31%) of a 1:1 mixture of (*E*)-**7b** and (*Z*)-**7b** which was not separated. (*E,Z*)-3-Chloro-1,1-bis(4-methylphenyl)-2-heptene [(*E,Z*)-**7b**]: ¹H NMR (CDCl₃, 200 MHz): δ = 0.88, 0.91 (2 t, *J* = 7.2 Hz, 6H, *E/Z*-7-H), 1.30–1.57 (m, 8H, *E/Z*-5-H, *E/Z*-6-H), 2.32 (s, 12H, aryl-CH₃), 2.37, 2.43 (2 t, *J* = 7 Hz, 4H, *E/Z*-4-H), 4.79 (d, *J* = 10.2 Hz, 1H, *E*-1-H), 5.15 (d, *J* = 9.6 Hz, 1H, *Z*-1-H), 5.93 (d, *J* = 9.6 Hz, 1H, *Z*-2-H), 6.06 (d, *J* = 10.2 Hz, 1H, *E*-2-H), 7.02–7.25 (m, 16H, aromatic H). – ¹³C NMR (CDCl₃): δ = 13.83, 13.92 (2 q, C-7), 21.00 (q, aryl-CH₃), 21.75, 21.98 (2 t, C-6), 29.44, 29.53 (2 t, C-5), 33.70, 39.28 (2 t, C-4), 49.04, 49.10 (2 d, C-1), 127.68, 130.20 (2 d, C-2), 127.93, 128.01, 129.14, 129.21 (4 d, C_o, C_m), 135.05, 135.30 (2 s, C-3), 135.83, 136.00 (2 s, C_p), 140.56, 140.68 (2 s, C_i). – C₂₁H₂₅Cl: calcd. 312.1645; found 312.1643 (MS).

1-Chloro-1,3,3-triphenylpropane (8a) was synthesized from **1a** and styrene as described in ref.^[16] – ¹³C NMR (CDCl₃): δ = 45.7 (t, C-2), 48.4 (d, C-3), 61.6 (d, C-1), 126.5 (d), 126.6 (d), 127.0 (d), 127.8 (d), 127.9 (d), 128.4 (d), 128.6 (d), 128.7 (d), 141.5 (s), 143.3 (s), 143.6 (s). – UV (methanol): λ_{\max} (lg ϵ) = 249 nm (2.694), 254 (2.782), 259 (2.848), 262 (sh), 265 (2.771), 269 (2.671).

1-Chloro-3,3-bis(4-methylphenyl)-1-phenylpropane (8b) was prepared from **1b** and styrene as reported in ref.^[17] where also analytical data are given.

(*1R**,*2S**)-1-Chloro-2-methyl-3,3-bis(4-methylphenyl)-1-phenylpropane (**9b**): For synthesis and analytical data see ref.^[24]

3,3-Dichloro-1,1-bis(4-methylphenyl)butane (10b): 2-Chloro-1-propene (1.10 g, 14.4 mmol) and compound **1b** (0.70 g, 3.0 mmol)

Table 4. Determination of relative reactivities by competition experiments^[a]

T/ °C	1 / mg	Nu _x / mg	Nu _y / mg	Standard / mg	Products / mmol or molar ratio	k _x / k _y	
	1a	Styrene	Phenylacetylene (2)		8a : E-3a : Z-3a ^[b]		
20	66.3	62.2	104.2	-	81 : 6 : 1	27.1	
20	65.6	63.5	154.8	-	60 : 6 : 1	28.3	
20	61.4	62.7	305.9	-	32 : 6 : 1	28.8	
-10	60.4	62.8	151.9	-	98 : 8 : 1	35.6	
-40	64.1	65.3	160.6	-	273 : 6 : 1	134.5	
-70	66.5	62.2	124.0	-	487 : 6 : 1	203.2	
-70	59.9	62.7	155.6	-	527 : 7 : 1	227.7	
-70	64.4	63.2	303.0	-	247 : 7 : 1	210.3	
	1b	Styrene	Phenylacetylene (2)		8b : E-3b - Z-3b ^[c]		
20	61.6	63.5	105.0	-	46 : 5 : 1	16.2	
20	85.4	63.5	172.1	-	27 : 6 : 1	14.3	
20	62.8	68.5	303.9	-	28 : 8 : 1	16.6	
-10	72.7	64.9	163.3	-	57 : 6 : 1	27.6	
-40	76.1	64.8	164.6	-	123 : 6 : 1	62.7	
-70	63.8	67.2	102.9	-	503 : 5 : 1	170.1	
-70	73.2	64.3	155.2	-	366 : 7 : 1	156.3	
-70	74.0	68.4	309.4	-	238 : 8 : 1	164.4	
	1b	<i>trans</i> -1-Phenyl- 1-propene	1-Phenyl-1-propyne	Methyl benzoate	9b	5b ^[d]	
-70	78.5	141	418	39.1	0.215	0.0707	9.99
-70	78.5	123	603	45.3	0.200	0.124	8.90
-70	78.5	109	1067	47.4	0.154	0.176	9.45
	1b	2-Chloro-1- propene	1-Hexyne (6)	1-Phenyldecane	10b	7b ^[e]	
-70	92.6	169	335	87	0.0698	0.100	1.29
-70	92.6	227	407	134	0.0771	0.0896	1.44
-70	92.6	182	370	120	0.0696	0.0979	1.36

^[a] Because of the greater molar absorption coefficient of the alkyne adducts, the nucleophile ratios have been selected in a way to give comparable peak areas for the various products. — ^[b] Separation on RP 18 phase with acetonitrile/water/methanol (20:20:60); retention times: **8a** (16.3 min), (*E*)-**3a** (18.5 min), (*Z*)-**3a** (20.7 min); detection at 254 nm. — ^[c] Separation on RP 18 phase with acetonitrile/water/methanol (20:15:65); retention times: **8b** (34.8 min), (*E*)-**3b** (40.3 min), (*Z*)-**3b** (45.3 min); detection at 254 nm. — ^[d] Separation on NO₂ phase with isoctane; retention times: **5b** (6.0 min), **9b** (7.4 min); detection at 266 nm. — ^[e] Separation on RP 18 phase with water/methanol (7:93); retention times: **10b** (5.6 min), **7b** (8.6 min); detection at 258 nm.

were combined with ZnCl₂(OEt₂)_{1.6} (1 mmol) in 25 ml of CH₂Cl₂ for 15 h at -70°C. Workup as described above gave 650 mg (71%) of colorless crystals of **10b** with m.p. 45–46°C. — ¹H NMR (CDCl₃, 200 MHz): δ = 1.85 (s, 3H, 4-H), 2.29 (s, 6H, aryl-CH₃), 3.11 (d, *J* = 6.4 Hz, 2H, 2-H), 4.36 (t, *J* = 6.4 Hz, 1H, 1-H), 7.07–7.26 (m, 8H, aryl-H). — ¹³C NMR (CDCl₃): δ = 20.9 (q), 37.7 (q), 48.2 (d), 55.1 (t), 90.8 (s), 127.5 (d), 129.3 (d), 135.9 (s), 141.4 (s). — IR (KBr): ν̄ = 3010 cm⁻¹, 2857, 1648, 1509, 1436, 1379, 1170, 1066, 1031, 857, 811, 771, 679, 601, 564. — MS (70 eV), *m/z* (%): 308, 306 (7, 9) [M⁺], 195 (100), 165 (12). — C₁₈H₂₀Cl₂: calcd. 306.0942; found 306.0934 (MS).

Determinations of absolute rate constants were carried out at the workstation described previously^[7] by using the concentrations listed in Table 3.

Competition Experiments: The diarylchloromethane **1a** or **1b** (0.3–0.4 mmol) and a pair of nucleophiles (alkene or alkyne each > 2 equivalents) were dissolved in dry CH₂Cl₂ (20 ml), and the solution was placed in a cryostat. After thermal equilibration, ZnCl₂(Et₂O)_{1.6} (0.06–0.08 mmol) was added, and the mixture was kept at the temperature given in Table 4 until **1a** or **1b** had been

consumed. The solution was washed with conc. aqueous ammonia (40 ml) to destroy the catalyst, then the standard was added and the solution subsequently dried with MgSO₄. After partial evaporation of the solvent, the mixture was analyzed by HPLC using the conditions given in the footnotes of Table 4.

- ^[1] T. W. G. Solomons, *Organic Chemistry*, Wiley, New York, 1992, p 372.
^[2] K. Yates, G. H. Schmid, T. W. Regulski, D. G. Garratt, H.-W. Leung, R. McDonald, *J. Am. Chem. Soc.* 1973, 95, 160.
^[3] G. Modena, F. Rivetti, G. Scorrano, U. Tonellato, *J. Am. Chem. Soc.* 1977, 99, 3392.
^[4] F. Marcuzzi, G. Melloni, G. Modena, *J. Org. Chem.* 1979, 44, 3022.
^[5] G. Melloni, G. Modena, U. Tonellato, *Acc. Chem. Res.* 1981, 14, 227.
^[6] R. Schneider, U. Grabis, H. Mayr, *Angew. Chem.* 1986, 98, 94–95; *Angew. Chem. Int. Ed. Engl.* 1986, 25, 89–90.
^[7] H. Mayr, R. Schneider, C. Schade, J. Bartl, R. Bederke, *J. Am. Chem. Soc.* 1990, 112, 4446–4454.
^[8] H. Mayr, R. Schneider, B. Irrgang, C. Schade, *J. Am. Chem. Soc.* 1990, 112, 4454–4459.
^[9] H. Mayr, R. Schneider, U. Grabis, *J. Am. Chem. Soc.* 1990, 112, 4460–4467.

- [10] H. Mayr, *Angew. Chem.* **1990**, *102*, 1415–1428; *Angew. Chem. Int. Ed. Engl.* **1990**, *29*, 1371–1384.
- [11] H. Mayr, G. Hagen, *J. Chem. Soc., Chem. Commun.* **1989**, 91–92.
- [12] G. Hagen, H. Mayr, *J. Am. Chem. Soc.* **1991**, *113*, 4954–4961.
- [13] J. Bartl, S. Steenken, H. Mayr, *J. Am. Chem. Soc.* **1991**, *113*, 7710–7716.
- [14] H. Mayr, J. Bartl, G. Hagen, *Angew. Chem.* **1992**, *104*, 1689–1691; *Angew. Chem. Int. Ed. Engl.* **1992**, *31*, 1613–1615.
- [15] R. Maroni, G. Melloni, G. Modena, *J. Chem. Soc., Perkin Trans. 1*, **1973**, 2491.
- [16] H. Mayr, W. Striepe, *J. Org. Chem.* **1983**, *48*, 1159–1165.
- [17] R. Pock, H. Mayr, *Chem. Ber.* **1986**, *119*, 2497–2509.
- [18] M. Patz, H. Mayr, *Tetrahedron Lett.* **1993**, *34*, 3393–3396.
- [19] R. Huisgen, *Methoden Org. Chem. (Houben Weyl) 4th Ed.*, Bd. 3/1, Thieme Stuttgart, **1955**, 144.
- [20] C. K. Ingold, F. R. Shaw, *J. Chem. Soc.* **1949**, 575–581.
- [21] M. Glaser, Ph. D. Thesis in preparation, Med. Univ. zu Lübeck.
- [22] R. Schneider, H. Mayr, P. H. Plesch, *Ber. Bunsenges. Phys. Chem.* **1987**, *91*, 1369–1374.
- [23] H. Mayr, W. Striepe, *J. Org. Chem.* **1985**, *50*, 2995–2998.
- [24] R. Pock, H. Mayr, M. Rubow, E. Wilhelm, *J. Am. Chem. Soc.* **1986**, *108*, 7767–7772.

[259/93]