

β -LACTAM AS SYNTHETIC INTERMEDIATE: SYNTHESIS OF LEUCINE-ENKEPHALIN

Naoto HATANAKA, Rumiko ABE, and Iwao OJIMA*


Sagami Chemical Research Center, Nishi-Onnuma 4-4-1, Sagamihara, Kanagawa 229

A method for oligopeptide synthesis via β -lactam intermediate has been successfully applied to the synthesis of a biologically active pentapeptide, leucine-enkephalin.

In preceding papers,¹⁻⁴ we have reported the entirely new method for peptide synthesis via β -lactam intermediate. The β -lactam method has been proved to have an advantage over conventional methods⁵ in view of increased solubility and superior chromatographic behavior on silica gel. Now, we have successfully applied the current method to the synthesis of a pentapeptide, leucine-enkephalin (Tyr-Gly-Gly-Phe-Leu) which possesses opioid activity.⁶

The β -lactam 1 is prepared by using our modified Bose's method⁷ from N-(p-benzyloxy-benzylidene)ethoxycarbonylmethylamine and azidoacetyl chloride in the presence of triethylamine in methylene chloride (84 %). The ester group in 1 was successfully saponified with 1 equivalent of 1 N sodium hydroxide in methanol to the free β -lactamcarboxylic acid 3 without touching the sensitive β -lactam ring (72 % after recrystallization from benzene-ethyl acetate). The β -lactam 2 was prepared by azidoacetylation (azidoacetyl chloride and N-methylmorpholine in chloroform) of the corresponding amino- β -lactam⁸ (72 % after recrystallization from ether). The azide group in 2 was reduced under 1 atm of hydrogen on 5 % Pd-C in methanol at room temperature to give the amine 4 (100 %).

SYNTHESIS OF LEUCINE-ENKEPHALIN



The acid 3 and the amine 4 were subjected to the coupling reaction by using DCC (dicyclohexylcarbodiimide) and HOBT (1-hydroxybenztriazole) in DMF (dimethylformamide). The coupling product 5 was purified on a silica gel column using ethyl acetate as eluent (98 % after purification).⁹

After the azide group of the bis- β -lactam 5 was reduced to the amino group under 1 atm of hydrogen on 10 % Pd-C in ethanol at room temperature and the amino group was protected as hydrochloride by adding 1 equivalent of 1N hydrochloric acid, the hydrogenolysis was carried out at 50°C to give a diastereomeric mixture of leucine-enkephalin t-butyl ester hydrochloride 6 (84 %).¹⁰

Now that we have established an efficient route to leucine-enkephalin via β -lactam intermediate, an investigation on the synthesis of optically pure enkephalin and its analogues is now actively under way.

References

1. I. Ojima, S. Suga, and R. Abe, Chem. Lett., 1980, 853.
2. I. Ojima, S. Suga, and R. Abe, Tetrahedron Lett., 21, 3907 (1980).
3. N. Hatanaka and I. Ojima, Chem. Lett., 1980, 231.
4. N. Hatanaka and I. Ojima, J. Chem. Soc., Chem. Commun., 1981, 346.
5. E. Gross and J. Meienhofer, "The Peptides", Vol. 1, Chapters 3, 4, 5 and 6, Academic Press, New York (1979).
6. J. Hughes, T. W. Smith, H. W. Kosterlitz, L. A. Fothergil, B. A. Morgan, and H. R. Morris, Nature, 258, 577 (1975).
7. e.g. A. K. Bose, M. S. Manhas, J. C. Chib, H. P. S. Chawla, and B. Dayal, J. Org. Chem., 39, 2877 (1974).
8. Azido- β -lactam (prepared from N-benzylidene[(*S*)-1-t-butoxycarbonyl-3-methyl]butylamine 7 and azidoacetyl chloride in the presence of triethylamine in methylene chloride) was reduced to the amino- β -lactam under 1 atm of hydrogen on 5 % Pd-C in methanol at room temperature (100 %). The Schiff base 7 was obtained by the condensation of (*L*)-leucine t-butyl ester with benzaldehyde.
9. IR(KBr) 3330, 2110, 1770, 1665, 1540 cm^{-1} . NMR(CDCl_3) δ 0.61-1.08(m, 6H), 1.36, 1.47(s, 9H), 1.50-2.35(m, 2H), 2.91-4.50(m, 6H), 4.87-5.64(m, 6H), 6.76-7.53(m, 14H). Found: C, 64.58; H, 6.37; N, 13.41. Calcd. for $\text{C}_{39}\text{H}_{45}\text{N}_7\text{O}_7$; C, 64.72; H, 6.27; N, 13.55.
10. IR(KBr) 3300, 1730, 1660, 1550 cm^{-1} . NMR(CD_3OD) δ 0.50-1.10(m, 6H), 1.45(s, 9H), 1.10-1.95(m, 2H), 2.70-4.50(m, 12H), 6.70-7.40(m, 9H). The product was also identified by comparing the HPLC chromatogram with that of authentically prepared HCl·H-(*D/L*)-Tyr-Gly-Gly-(*D/L*)-Phe-(*L*)-Leu-OBu^t. HPLC analysis was carried out using a column packed with TOYO SODA LS410K (ODS SIL) and MeOH-H₂O as eluent.

(Received July 6, 1981)