

SIMULTANEOUS EVOLUTION OF HYDROGEN AND OXYGEN BY WATER PHOTOLYSIS WITH
PRUSSIAN BLUE AND TRIS(2,2'-BIPYRIDYL)RUTHENIUM(II) COMPLEX

Masao KANEKO*, Naoki TAKABAYASHI, and Akira YAMADA

The Institute of Physical and Chemical Research, Hirosawa, Wako-shi 351

Simultaneous evolution of hydrogen and oxygen gases was achieved by visible light irradiation of water containing prussian blue and tris(2,2'-bipyridyl)ruthenium(II) complex. The gases evolved were analyzed by gas chromatography and mass spectroscopy. The mechanism for the water photolysis was also discussed.

Water photolysis to give simultaneous evolution of hydrogen and oxygen gases is one of the most attracting and important subjects of solar energy conversion. Most photochemical systems of water photolysis are utilizing semiconductors such as titanium dioxide¹⁻³⁾ and cadmium sulfide.⁴⁾ The only one system containing no semiconductor for simultaneous H₂ and O₂ evolution was reported by Grätzel et al., who used tris(2,2'-bipyridyl)ruthenium(II) complex (abbreviated to Ru(bpy)₃²⁺), methylviologen (MV²⁺), platinum colloid, and ruthenium dioxide sol protected by poly(styrene-co-maleic anhydride).⁵⁾ This photolysis system is complicated and sensitive to the preparative techniques of the reagents, which often makes the conversion not reproducible. The O₂ analysis data of that paper would be insufficient. It might occur that the poly(styrene-co-maleic anhydride), instead of water, works as a reducing agent for H₂ production, because the copolymer can reduce MV²⁺ under visible light irradiation to give its cation radical⁶⁾ which can reduce protons to H₂ on Pt catalyst. The present authors have found a simple system of water photolysis composed of prussian blue (ferrous potassium hexacyanoferrate(III); KFe[Fe(CN)₆], abbreviated to PB) and Ru(bpy)₃²⁺ to evolve hydrogen and oxygen gases simultaneously, and the results are described.

PB has a polynuclear structure with mixed valent iron ions, and forms a colloidal solution.⁷⁾ A solution of PB was prepared by mixing a 40 mM aqueous solu-

tion of potassium ferricyanide, $K_3[Fe(CN)_6]$, and an equimolar aqueous solution of ferrous chloride while stirring, and used within one day after preparation. The irradiation of a 10 ml aqueous solution containing $10 \mu\text{M Ru}(\text{bpy})_3^{2+}$, 5 mM PB, and 0.5 M KCl at pH 2 (HCl-KCl buffer) under argon at room temperature with visible light (cutoff filters; Toshiba VY-42 and IRQ-80, light intensity; 78 mW/cm^2) from a 500 W xenon lamp gave simultaneous H_2 and O_2 evolution. The gas chromatogram (Fig. 1) of the gas evolved after 15 h's irradiation showed the presence of both H_2 and O_2 . The O_2 content evolved by the photochemical reaction was calculated by subtracting the content of O_2 coming from the air. The H_2 and O_2 evolved were 0.27 ± 0.01 and $0.14 \pm 0.02 \mu\text{l}$, respectively. Fig. 2 shows the mass spectrum of the gas evolved by the irradiation of the solution of 90 % D_2O containing 2.0 % H_2^{18}O , $10 \mu\text{M Ru}(\text{bpy})_3^{2+}$, 5 mM PB and 0.5 M KCl under the same condition as the above. The peaks at $m/e = 3$ and 4 are due to DH^+ and D_2^+ , respectively, and that at $m/e = 34$ to $^{18}\text{O}^{16}\text{O}^+$. The ratio of the peak intensity of D_2^+/DH^+ was 4.3 (theoretical; 4.5), and that of $^{18}\text{O}^{16}\text{O}^+ / ^{16}\text{O}_2^+$ was 0.039 (theoretical; 0.040). These results indicate that water was decomposed to give hydrogen and oxygen. The irradiation of a 10 ml aqueous solution containing $1 \mu\text{M Ru}(\text{bpy})_3^{2+}$, 500 μM PB, and 0.5 M KCl at pH 2 for 90 h gave $1.3 \mu\text{l H}_2$ and $0.7 \mu\text{l O}_2$, showing that the turnover number of the Ru complex is 11 under this condition. After this reaction the Ru complex and the PB suffered no change.

The addition of KCl caused the increase in the yield of the photochemical reaction probably by stabilizing PB. The optimum concentration of the KCl added




Fig. 1 Gas chromatogram of H_2 and O_2 evolved by irradiation of aqueous solution containing $\text{Ru}(\text{bpy})_3^{2+}$, PB, and KCl at pH 2 under Ar. for 15 h. ----; O_2 from the air.




Fig. 2 Mass spectrum of the gas evolved by irradiation of 90 % D_2O solution containing 2.0 % H_2^{18}O , $\text{Ru}(\text{bpy})_3^{2+}$, PB and KCl at pH 2 under Ar. for 15 h.

was around 0.5 M. The addition of more KCl retarded the reaction, which would be due to the partial aggregation of PB. Continuous stirring of the reaction mixture was required for the water photolysis, since the aggregation of PB leads to the inactivation of the water photolysis system.

Table 1. Effect of wavelength of the incident light on the photolysis of water containing 10 μM $\text{Ru}(\text{bpy})_3^{2+}$, 5 mM PB and 0.5 M KCl at pH 2 (HCl-KCl buffer) under argon at room temperature. Solution; 10 ml.

Wavelength (nm) of the incident light	Reaction time (h)	Gas evolved (μl)	
		H_2	O_2
>400	15	0.27	0.14
>550	15	0	0
450* ^a	70	0	0
700* ^a	70	0	0
450 + 700* ^{a,b}	70	0.09	0.05

*a) Interference filter was used; Light intensities; 1.8 mW/cm^2 (450 nm) and 3.8 mW/cm^2 (700 nm). Solution; 2 ml.






*b) Two light sources were used.

The effect of the wavelength of the incident light on the water photolysis was shown in Table 1. The excitation of the Ru complex ($\lambda_{\text{max}} = 452 \text{ nm}$) was essential for the photolysis. The irradiation either by 450 or 700 nm monochromatic light did not induce water photolysis, and both the excitation of the Ru complex and the PB ($\lambda_{\text{max}} = 700 \text{ nm}$) were required. The quenching of the excited state of the $\text{Ru}(\text{bpy})_3^{2+}$ by PB was very efficient. The rate constant of the quenching at 30 °C was measured to be $3.50 \times 10^{10} \text{ (M}^{-1} \text{ s}^{-1})$ from the emission quenching and the lifetime of $\text{Ru}(\text{bpy})_3^{2+*}$. The Stern-Volmer constant (k_{SV}) obtained from the lifetime measurement was about one third of that obtained from emission quenching, indicating that a static quenching of the $\text{Ru}(\text{bpy})_3^{2+*}$ by PB occurs. There must be some interaction between the two species, which might play an important role for the photochemical process. The details will be reported elsewhere.

One of the interesting features of the present water photolysis system is that PB provides more than two kinds of functions, i.e., electron mediator and catalysts for H_2 and O_2 evolution. The photochemical reaction of a 10 ml aqueous solution of 10 mM EDTA, 20 μM $\text{Ru}(\text{bpy})_3^{2+}$, and 5 mM PB at pH 6 (KH_2PO_4 -NaOH buffer) for 1 h gave 0.09 μl H_2 . The same reaction of a blank solution without PB gave no H_2 . In that H_2 evolving system, the PB must work as a mediator to accept electron from $\text{Ru}(\text{bpy})_3^{2+*}$ and also as a H_2 evolving site. In order to study the effect of the excitation of PB, the following experiments were carried out. L-ascorbic acid reduces PB, but the reaction of both the reagents in the dark does not give H_2 . How-

ever, the irradiation of a solution containing 5 mM PB, 10 μ M L-ascorbic acid, and 0.5 M KCl at pH 2 gave a small amount of H₂ (ca. 0.03 μ l) in 2 h, while no H₂ was formed in the dark. These results suggest that a partially reduced PB can give H₂ under irradiation. Reaction of 100 μ M Ru(bpy)₃³⁺ (ClO₄⁻)₃ with 5 mM PB in water at pH 2 gave 4.5 μ l O₂ (yield; 74 %) in 1 h by water oxidation, while a blank solution without PB did not give O₂, indicating that PB acts as O₂ evolving site with the trivalent Ru complex as oxidant.

The reaction scheme can be considered as follows. The quenching of Ru(bpy)₃^{2+*} by PB would be oxidative, because the irradiation of the mixed solution of the Ru complex, PB and EDTA gave H₂ as described above (Eq.(1)). Since PB has a poly-



nuclear structure⁷⁾, it could behave as a kind of electron pool to give PB²⁻ (Eq. (3)). Two protons would be reduced by the PB²⁻ under excitation to give H₂ (Eq. (4)). On the other hand, the function of PB as a kind of charge pool must be important in the O₂ evolving reaction (Eq.(5)).

The quantum yield for the H₂ and O₂ evolution with both 450 and 700 nm monochromatic light irradiation was the order of 0,01 % based on the 450 nm light absorbed. Although the quantum yield is not high so far, the simultaneous H₂ and O₂ evolution by such the simple two components system would be surprising in itself. The roles of PB both as electron and charge pool would be the crucial point, which must be due to its polynuclear structure and to its mixed valence state.

References

- 1) A. Fujishima and K. Honda, *Nature*, 238, 37 (1972).
- 2) T. Kawai and T. Sakata, *Chem. Phys. Lett.*, 72, 87 (1980).
- 3) E. Borgarello, J. Kiwi, E. Pelizzetti, M. Visca, and M. Grätzel, *Nature*, 289, 158 (1981).
- 4) K. Kalyanasundaram, E. Borgarello, and M. Grätzel, *Helv. Chim. Acta*, 64, 362 (1981).
- 5) K. Kalyanasundaram and M. Grätzel, *Angew. Chem.*, 91, 759 (1979).
- 6) M. Kaneko, K. Hamanishi, and A. Yamada, *Preprints of 43rd Congress of Chem. Soc. Jpn.*, 1128 (1981).
- 7) H.J.Buser, D.Schwarzenbach, W.Petter, and A.Ludi, *Inorg.Chem.*, 16, 2704 (1977).

(Received June 18, 1982)