

DIPHOSPHORUS TETRAIODIDE AS A NEW MILD CONDENSING AGENT
FOR THE SYNTHESIS OF AMIDES


Hitomi SUZUKI,* Junko TSUJI, Yoshio HIROI, Naofumi SATO,
and Atsuhiko OSUKA

Department of Chemistry, Faculty of Science,
Ehime University, Bunkyo-cho, Matsuyama 790

In the presence of diphosphorus tetraiodide, a variety of weak carboxylic acids readily react with amines to form the corresponding amides in high yields.

Dehydrative condensation of weak carboxylic acids with amines is too slow to serve as a practical process for amide synthesis.¹⁾ Thus the activation of carboxyl function is necessary to facilitate the reaction through the intermediate formation of reactive esters, acid anhydrides, or acyl-onium compounds. The reagents introduced for such purposes include carbodiimide,²⁾ phosphines,³⁾ phosphonates,⁴⁾ boranes,⁵⁾ metal or pseudometal halides,⁶⁾ 2-halopyridinium salts,⁷⁾ activated acetylenes,⁸⁾ and azole derivatives.⁹⁾ However, many of these are not free from some drawbacks such as limited availability, by-product formation, inconvenient handling, high costs, competing side-reactions, or low stability.

Diphosphorus tetraiodide (P_2I_4) is an easily prepared and handled dehydrating agent,¹⁰⁾ which can effect the conversion of aldoximes,¹¹⁾ amides,¹²⁾ and nitroalkanes¹³⁾ into nitriles under mild conditions. On continuing our research on the reactivity of P_2I_4 , we report herein a mild coupling reaction of a variety of weak carboxylic acids with amines in the presence of P_2I_4 .



A general procedure is as follows; amine (1.0 mmol) and 2,6-lutidine (1.0 mmol) are added to a suspension of P_2I_4 (1.0 mmol) in dry carbon tetrachloride (10 ml) and the mixture is stirred for several minutes to form a white precipitate. A solution of carboxylic acid (1.0 mmol) in dichloromethane (10 ml) is then added with stirring. The precipitate goes into solution to give a yellow homogeneous mixture, which is heated under gentle reflux for several hours. After cooling, the mixture is quenched with 1M-hydrochloric acid, the organic phase is separated, washed with 5%-sodium

Table 1 Amides Obtained from the P_2I_4 -assisted Condensation of Carboxylic Acids with Amines

Carboxylic acid <u>1</u>	Amine <u>2</u>	Amide <u>3</u>	
R	R'	Mp ($^{\circ}C$)	Yield (%)
4- $CH_3O-C_6H_4$	C_6H_5	172-173	~ 100
do	$C_6H_5CH_2$	128-129	90
3,4- $(CH_3O)_2-C_6H_3$	C_6H_5	162-163	~ 100
do	$C_6H_5CH_2$	175-176	98
4- $CH_3-C_6H_4$	C_6H_5	149-150	80
do	2,4,6- $(CH_3)_3-C_6H_2$	197-198	~ 100
2,4,6- $(CH_3)_3-C_6H_2$	C_6H_5	166-167	~ 100
4-Cl-2,3,5,6- $(CH_3)_4-C_6$	do	235-236*	~ 100
4- $CH_3CO-C_6H_4$	do	188-189	81
4- $NO_2-C_6H_4$	do	216-217	~ 100
do	$C_6H_5CH_2$	139-140	80
$C_6H_5CH=CH_2$	C_6H_5	149-150	~ 100
$(CH_3)_3C$	do	133-134	~ 100
do	$C_6H_5CH_2$	83-84	~ 100
do	2,6- $(CH_3)_2-C_6H_3$	205-206*	~ 100
do	3,5- $(CH_3)_2-C_6H_3$	126-127*	~ 100
do	4- $NO_2-C_6H_4$	185-186*	85

* Sublimes.

carbonate solution, dried with sodium sulfate, and evaporated to leave the expected amide as a sole product, which is further recrystallized from methanol. Yields of products range from 85 to 100%. The reaction can be carried out at room temperature, but it usually takes much longer time for completion. As apparent from the Table, the present method is especially attractive for converting weak or hindered amines into amides.

The reagent may also be used for the synthesis of some peptides. Thus N-acetyl- and N-benzyloxycarbonyl-D,L-phenylalanyl-glycine ethyl esters were obtained in

46% and 53% yields, respectively from the corresponding N-protected D,L-phenylalanine and glycine ethyl esters.

The use of P_2I_4 as a coupling agent for the amide synthesis has some distinct advantages; commercial availability, ease of handling (crystalline solid with no fuming property), simple work-up (oxygen-containing phosphorus by-products are water-soluble), mild reaction conditions (neutral and non-aqueous media; temperature below ca. 50°C), compatibility with various functionalities, and high purity of products.

References

- 1) For a general survey, see: B.C. Challis and A.R. Butler, in "The Chemistry of the Amino Group", S.Patai, ed., Interscience, London, p. 277 (1968); "The Chemistry of Amides", J.Zabicky, ed., Interscience, London (1970); N. Kato, in "Shin-jikken Kagaku Koza", Vol. 14-II, Maruzen, Tokyo, p.1134 (1977).
- 2) *Dicyclohexylcarbodiimide (DCC)*: W.A. Bonner and P.I. McNamee, *J. Org. Chem.*, 26, 2554 (1961); for a recent review of carbodiimide chemistry, see: A. Williams and I.T. Ibrahim, *Chem. Rev.*, 81, 589 (1981).
- 3) *Triphenylphosphine/carbon tetrachloride*: L.E. Barstow and V.J. Hruby, *J. Org. Chem.*, 36, 1305 (1971).
Tris(dimethylamino)phosphine/carbon tetrachloride: B. Castro and J.R. Dormoy, *Bull. Soc. Chim. Fr.*, 1971, 3034.
Triphenylphosphine/cupric chloride/p,p'-dichlorodiphenyl disulfide: R. Matsueda, H. Maruyama, M. Ueki, and T. Mukaiyama, *Bull. Chem. Soc. Jpn.*, 44, 1373 (1971).
Triphenylphosphine ditriflate: J.B. Hendrickson and S.M. Schwartzman, *Tetrahedron Lett.*, 1975, 277.
- 4) *Diphenyl phosphonate/pyridine*: N. Yamazaki and F. Higashi, *Tetrahedron Lett.*, 1972, 5047.
Diphenyl phosphorocyanidate: S.I. Yamada, Y. Kasai, and T. Shioiri, *Tetrahedron Lett.*, 1973, 1595.
Diphenyl phosphorobromidate: A. Gorecka, M. Leplawy, J. Zabrocki, and A. Zwierzak, *Synthesis*, 1978, 475.
- 5) *Trimethyl borate*: A. Pelter, T.E. Levitt, and P. Nelson, *Tetrahedron*, 26, 1539 (1970).
Boron trifluoride etherate: J. Tani, T. Oine, and I. Inoue, *Synthesis*, 1975, 715.
Catecholborane: D.B. Collum, S.C. Chen, and B. Ganem, *J. Org. Chem.*, 43, 4393 (1978).
- 6) *Silicon tetrachloride*: T.H. Chan and L.T.L. Wong, *J. Org. Chem.*, 34, 2776 (1969).
Titanium tetrachloride: J.D. Wilson and H. Weingarten, *Can J. Chem.*, 48, 983 (1970).
- 7) *2-Iodopyridinium iodide*: J.K. Sutherland and D.A. Widdowson, *J. Chem. Soc.*, 1964, 4650.

- 2-Fluoropyridinium iodide*: T. Mukaiyama, Y. Aikawa, and S. Kobayashi, Chem. Lett., 1976, 57.
- 8) *Dimethylpropargylsulfonium bromide*: G.D. Appleyard and C.J.M. Stirling, J. Chem. Soc., C, 1969, 1904.
- 3-Dimethylamino-1-phenyl-2-propyn-1-one*: M. Neuenschwander, U. Lienhard, H. P. Fahrni, and B. Hurni, Helv. Chim. Acta, 61, 2428 (1978).
- 9) *6-Chloro-1-(p-chlorobenzenesulfonyloxy)benzotriazole*: M. Itoh, H. Nojima, J. Notani, D. Hasegawa, and K. Takai, Tetrahedron Lett., 1974, 3089.
- 2-Chloro-N-methylbenzothiazolium triflate*: F. Souto-Bachiller, G.S. Bates, and S. Masamune, J.C.S. Chem. Commun., 1976, 719.
- 10) N.G. Feshchenko and A.V. Kirsanov, Zh. Obshch. Khim., 30, 3041 (1960); H. Suzuki and T. Fuchita, Nippon Kagaku Kaishi, 1977, 1679. Now available from the Aldrich Chemical Co., and Fluka AG.
- 11) H. Suzuki, T. Fuchita, A. Iwasa, and T. Mishina, Synthesis, 1978, 905; J. N. Denis, R. Magnane, M. Van Eenoo, and A. Krief, Nouveau J. Chim., 3, 705 (1979).
- 12) H. Suzuki and N. Sato, Nippon Kagaku Kaishi, 1981, 121.
- 13) J.N. Denis and A. Krief, Tetrahedron Lett., 1979, 3995.

(Received January 13, 1983)