


considered to proceed through the following reaction path. In the syntheses of 1,1-dialkylethenes²⁾ and ketones,³⁾ the initial formation of IV was rationalized by the attack of electrophiles on II with concurrent migration of alkyl group from boron to carbon, as depicted in Scheme (path A). It

Table 1. Synthesis of 1-Methoxy-2-alkanones

Organoborane R ₃ B, R=	Yield of 1-methoxy-2-alkanones RCOCH ₂ OCH ₃ , (%) ^{a)}
Butyl	65
Isobutyl	67
<i>sec</i> -Butyl	47
Pentyl	62
Cyclopentyl	61
Hexyl	60

a) Glpc yield based on the organoborane used.

has already been reported that complete elimination of the α -methoxy moiety in such a vinylborate is difficult.⁴⁾ The addition of BF₃:Et₂O to II may cause elimination of the α -methoxy group to give the intermediate (V) (path B), and the sequential oxidation of the B-C bond in V may form the expected product (III). However, we could not obtain satisfactory results by the usual aqueous sodium hydroxide-hydrogen peroxide oxidation, because the boron trifluoride seems to be partly decomposed in such an aqueous solution to boric acid and hydrogen fluoride, which lead to side reactions of II.⁵⁾ Utilization of sodium methoxide in methanol free from water was found to be very effective for the preparation of 1-methoxy-2-alkanones.



References

- 1) C. N. Skold, *Synth. Commun.*, **6**, 119 (1976).
- 2) T. Yogo and A. Suzuki, *Chem. Lett.*, 1980, 591.
- 3) T. Yogo, J. Koshino, and A. Suzuki, *Chem. Lett.*, 1981, 1059.
- 4) N. Miyaura, T. Yoshinari, M. Itoh, and A. Suzuki, *Tetrahedron Lett.*, **21**, 537 (1980).
- 5) Under the usual oxidation conditions with aqueous hydroxide and 30% hydrogen peroxide, II (R=Bu) gave dibutyl(methoxymethyl)methanol as a by-product together with the expected methoxyketone.

(Received April 18, 1983)