

UNPRECEDENTED *GAMMA*-SELECTIVE MICHAEL ADDITION OF TIN(II) DIENOLATES
TO ACYCLIC α,β -UNSATURATED KETONESRodney W. STEVENS and Teruaki MUKAIYAMA
Department of Chemistry, Faculty of Science,
The University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113


Tin(II) dienolates were found to react exclusively at the *gamma* carbon with acyclic α,β -unsaturated ketones in the Michael sense. Either acyclic 1,7-diketones or 2-cyclohexenol derivatives could be obtained in moderate to excellent yields by appropriate choice of reaction parameters. Moreover, in the latter case, only a *single* cyclohexenol adduct was afforded.

It has been well established that enolate anions, notably lithium enolates, of α,β -unsaturated carbonyl compounds react predominantly at the *alpha* carbon.^{1,2)} Because, in certain cases, *gamma*-substituted products would be of synthetic value, several methods have been devised to direct *gamma* addition. Notably, since it was first reported from this laboratory that 1-trimethylsilyloxybutadiene reacts with acetal exclusively from the *gamma* position in the presence of Lewis acid catalyst,³⁾ a series of silicon-directing *gamma*-substitution reactions with a variety of electrophiles has been reported.⁴⁾

Surprisingly, the *gamma*-selective addition of dienolates to Michael acceptors⁵⁾ has virtually been unexplored even though such a reaction may be of important synthetic, as well as theoretical, value. Maruyama and coworkers⁶⁾ in their synthetic efforts directed toward the synthesis of pyranonaphthoquinones employed an α -silyl β,γ -unsaturated ester as a dienolate anion equivalent of crotonic acid for directing *gamma*-selective Michael addition. Kraus⁷⁾ has also reported that treatment of 1-trimethylsilyloxybutadiene with butyllithium followed by treatment with α,β -unsaturated carbonyl compound gave an annulation product, albeit in poor yield. This intriguing reaction was speculated to arise from reversible Michael addition in which the observed product represents the most stable one. However, a metal-assisted Diels-Alder reaction cannot be disregarded.

During the course of our studies on the scope and limitation of tin(II) enolate chemistry,⁸⁾ we deemed it worthy to investigate the reactivity and regiochemistry of tin(II) dienolates.⁹⁾ And, in this communication we wish to disclose our preliminary findings on the reaction of tin(II) dienolates with α,β -unsaturated ketones.

In the first place, to the tin(II) dienolate of 2,2-dimethyl-5-hexen-3-one, generated by standard procedure at -45 °C in dichloromethane, was added 1-phenyl-

Table 1. Selective preparation of trans-olefinic 1,7-diketones^{a)}

Entry	R	α, β -Unsaturated ketone	Yield of <u>2</u> / $\%$ ^{b)}
1		(E)-PhC(O)CH=CHCH ₃ (<u>3</u>)	83
2		MVK ^{c)}	75 ^{d)}
3	^t Bu	(E)-PhC(O)CH=CHPh	81
4		(E,E)-PhC(O)CH=CHCH=CHPh	71
5		cyclohexenone	-

6	ⁱ Pr	<u>3</u>	72
7		MVK ^{c)}	55 ^{d, e)}

8		<u>3</u>	60 ^{e)}
9	Et	MVK ^{c)}	38 ^{d, e)}

Molar ratio of Sn(OTf)₂: N-ethylpiperidine : 1 : α, β -unsaturated ketone = 1.1 : 1.2 : 1.0 : 0.9.

a) Only the trans-isomer ($J = 16$ Hz) was obtained. b) Isolated yield. All products gave satisfactory spectral data. c) Methyl vinyl ketone. d) Reaction was conducted at -78°C . e) Enolization was performed at -78°C .

2-buten-1-one (3). Aqueous work-up of the reaction mixture revealed that the trans-olefinic 1,7-diketone 2 ($J = 16$ Hz) had been afforded as the only isolable product in 83% yield (Table 1, entry 1). This result represents unprecedented Michael addition of the tin(II) dienolate exclusively from the *gamma* carbon; no *alpha* Michael addition product or 1,2-addition products were detected. Next, the reaction of a number of readily available α, β -unsaturated ketones with a series of tin(II) dienolates, derived from the corresponding β, γ -unsaturated ketones,¹⁰⁾ was examined and the results are summarized in Table 1.

As can be seen from the results recorded in Table 1, with the exception of cyclohexenone (entry 5), in all examples examined a smooth reaction proceeds to give exclusively the 1,7-diketone 2 in moderate to excellent yields via *gamma*-selective 1,4-conjugate addition of the tin(II) dienolate to the α, β -unsaturated ketone. Although not entirely clear at this stage, preliminary investigations support that product formation is purely kinetic and not a result of Michael addition reversibility.

Scrutinization of the present reaction pathway suggests that *gamma*-selective addition of the tin(II) dienolate to the α, β -unsaturated ketone should regenerate an intermediate tin(II) enolate. Hence, subsequent intramolecular aldol cyclization would provide a simple step-wise route to the corresponding cyclohexenol derivatives, Diels-Alder type adducts. As a test of this postulate, examination of the reaction of the tin(II) dienolate generated from 2,2-dimethyl-5-hexen-3-one with 1-phenyl-2-buten-1-one (3) under a variety of reaction conditions


preferential pseudoequatorial orientation of the alkyl substituent seems most likely. Furthermore, prolonged reaction time led to the formation of diene products (entries 6 and 8), necessarily derived from the corresponding cyclohexenols 4, in which elimination to the exo-cyclic double bond product was exclusive. This result can be explained by considering that elimination to the alternate cyclohexadiene adduct is restricted by the severe gauche interactions that would arise between the ring substituents.

Although this reaction could well be the Diels-Alder type reaction, accumulated experimental data suggests otherwise. We tentatively speculate that a step-wise mechanism is operative by which *gamma*-selective 1,4-conjugate addition to α,β -unsaturated ketone is followed by successive intramolecular aldol-type cyclization in a highly stereoselective manner.

In summary, tin(II) dienolates were found to undergo unprecedented *gamma*-selective 1,4-conjugate addition to α,β -unsaturated ketones. Appropriate choice of reaction parameters led to the formation of either trans-olefinic 1,7-diketones or cyclohexenol derivatives at will. Furthermore, it was speculated that the *single* cyclohexenol adduct yielded arose via a step-wise mechanism with remarkable stereocontrol.

References

- 1) P. E. Pfeffer and L. S. Silbert, *J. Org. Chem.*, **36**, 3290 (1971); M. W. Rathke and D. Sullivan, *Tetrahedron Lett.*, **1972**, 4249; P. E. Pfeffer, L. S. Silbert, and E. Kinsel, *ibid.*, **1973**, 1163; J. L. Herrmann, G. R. Kieczkowski, and R. H. Schlessinger, *ibid.*, **1973**, 2433; S. A. G. de Graaf, P. E. R. Oosterhoff, and A. van der Gen, *ibid.*, **1974**, 1653; I. Casinos and R. Mestres, *J. Chem. Soc., Perkins Trans. 1*, **1978**, 1651.
- 2) For some special cases of *gamma*-selective substitution see; J. T. Wolfe, G. B. Trimitsis, and D. R. Morris, *J. Org. Chem.*, **34**, 3263 (1969); G. A. Koppel, *Tetrahedron Lett.*, **1972**, 1507; M. Yoshimoto, N. Ishida, and T. Hiraoka; *ibid.*, **1973**, 39; B. M. Trost and L. S. Melvin, *J. Am. Chem. Soc.*, **98**, 1204 (1976); J. A. Katzenellenbogen and A. L. Crumrine, *ibid.*, **98**, 4925 (1976); P. T. Lansbury and R. W. Erwin, *Tetrahedron Lett.*, **1978**, 2673; A. B. Smith III and R. M. Scarborough, Jr., *ibid.*, **1978**, 4193; P. M. Savu and J. A. Katzenellenbogen, *J. Org. Chem.*, **46**, 239 (1980).
- 3) T. Mukaiyama and A. Ishida, *Chem. Lett.*, **1975**, 319 and 1201; T. Mukaiyama and A. Ishida, *ibid.*, **1977**, 467; T. Mukaiyama and A. Ishida, *Bull. Chem. Soc. Jpn.*, **51**, 2077 (1978).
- 4) For example; I. Fleming, J. Goldhill, and I. Paterson, *Tetrahedron Lett.*, **1979**, 3205 and 3209; I. Fleming and T. V. Lee, *ibid.*, **1981**, 705; P. Albaugh-Robertson and J. A. Katzenellenbogen, *ibid.*, **1982**, 723.
- 5) G. A. Kraus and B. Roth, *Tetrahedron Lett.*, **1977**, 3129.
- 6) Y. Naruta, H. Uno, and K. Maruyama, *Chem. Lett.*, **1982**, 609.
- 7) G. A. Kraus and H. Sugimoto, *Tetrahedron Lett.*, **1977**, 3929.
- 8) T. Mukaiyama, N. Iwasawa, R. W. Stevens, and T. Haga, *Tetrahedron*, **40**, 1381 (1984) and references cited therein.
- 9) Tin(II) dienolates were found to react exclusively at the *alpha* carbon with aldehydes and saturated ketones (unpublished results).
- 10) Initial attempts to generate the tin(II) dienolate directly from the corresponding α,β -unsaturated ketone (for example, 3) were always complicated by self-coupling reactions.
- 11) 1,7-Diketone (2) was isolated with trans geometry, however, the formation of the cyclic adduct (4) implies that (E)-isomer (9) and (Z)-isomer (10) are in equilibrium in the reaction mixture and that the cyclization takes place via (Z)-isomer (10).



(Received March 1, 1985)