


SYNTHESIS OF β,γ -UNSATURATED CARBOXYLIC ACID DERIVATIVES BY THE NOVEL $\text{Ni}(\text{CO})_4$ -INDUCED RING-OPENING CARBONYLATION REACTION OF 1,1-DIBROMO-2-CHLOROCYCLOPROPANES

Toshikazu HIRAO,* Shinichiro NAGATA, and Toshio AGAWA

Department of Applied Chemistry, Faculty of Engineering, Osaka University,
2-1 Yamadaoka, Suita, Osaka 565


1,1-Dibromo-2-chlorocyclopropanes underwent the $\text{Ni}(\text{CO})_4$ -induced ring-opening carbonylation reaction with alcohol or amine giving the β,γ -unsaturated carboxylic acid and dicarboxylic acid derivatives. Use of *N,N*-dimethyltrimethylsilylamine as an initial nucleophile in the presence of benzaldehyde led to a dienecarboxamide presumably via codensation of the nickel enolate intermediate.

The $\text{Ni}(\text{CO})_4$ -induced carbonylation reactions of gem-dibromocyclopropanes permit gem-functionalization leading to a variety of cyclopropanecarboxylic acid derivatives.^{1,2)} This unique transformation is considered to depend on the intermediacy of nickel enolate compounds. We describe herein a novel ring-opening carbonylation reaction, which is effective for a synthesis of β,γ -unsaturated carboxylic acid derivatives.




Treatment of the 1,1-dibromo-2-chlorocyclopropane 1³⁾ with 1-propanol in the presence of $\text{Ni}(\text{CO})_4$ in DMF at 70 °C gave the β,γ -unsaturated ester 2 (Q=O) and the corresponding diester 3 (Q=O) as major products. No α,β -unsaturated isomer was detected. The 2-chlorocyclopropanecarboxylate 4 (Q=O) derived by normal reductive carbonylation was produced in a small amount. Some results including the preparation of β,γ -unsaturated carboxamides and carbothiolate are listed in Table 1.

Without the addition of alcohol, the carbonylation reaction did not occur and 1 was recovered. This finding excludes the possibility that $\text{Ni}(\text{CO})_4$ attacks 1 at the first step. One of the plausible reaction paths is outlined in the following scheme. The nickel enolate complex 5 (Q=O) might intervene as reported previously.²⁾ Further displacement with the alkoxycarbonyl group (Z=COOR) results in the formation of 7. Decarboxylation of 3 or 7 is assumed to lead to the β,γ -unsaturated ester 2 since a longer reaction time increased the 2a/3a ratio (Table 1). Another route to 2 is explained by hydride⁴⁾ transfer which gives the nickel enolate intermediate 6.



The present method provides a versatile synthetic procedure for β,γ -unsaturated carboxylic acid derivatives from vinyl chlorides³⁾ based on carbonylation with one-carbon homologation. Its utility was also demonstrated by preparing a precursor of the key intermediate, 7-hydroxy-3-octenoic acid, for the pyrenophorin synthesis^{5,6)} as shown by the bottom example of Table 1; the β,γ -unsaturated ester 2d was produced exclusively.⁷⁾ The corresponding diester and cyclopropanecarboxylate were not formed, this being in sharp contrast to the results mentioned above.

With *N,N*-dimethyltrimethylsilylamine as an initial nucleophile,²⁾ 1a underwent the similar ring-opening carbonylation reaction. This outcome contributes



another C-C bond formation between a nickel enolate intermediate and an electrophile. The Ni(CO)_4 -induced reaction⁸⁾ of 1a with the silylamine and benzaldehyde was carried out leading to the dienecarboxamide 8a.⁹⁾ It remains uncertain which of the enolate 6¹⁰⁾ or 7¹¹⁾ participates in the condensation reaction. This method achieves the introduction of two functional groups into a cyclopropane with ring cleavage only by mixing the silylamine, 1, Ni(CO)_4 , and benzaldehyde together in one flask.





























Table 1. Preparation of Carboxylic Acid Derivatives by the Ni(CO)₄-Induced Carbonylation Reaction of 1^{a)}

<u>1</u>	RQH, equiv.	Isolated yield/ % (ratio)			
 <u>1a</u>	PrOH	 <u>2a</u>	 <u>3a</u>	 b)	
		1	31	(61:29:10)	
		2.2 ^{c)}	46	(35:61:4)	
	2.2	57	(49:45:6)		
<u>1a</u>	PrNH ₂	 2a	 3a	 b)	
		2.2	62	(37:39:24)	
 <u>1b</u>	PrOH	 2a	 3a	 b)	
		2.2	80	(43:40:17)	
<u>1b</u>	PrNH ₂	 2a	 3a	 b)	
		2.2	74	(43:38:19)	
 <u>1c</u>	PrOH	 2a	 3a	 b)	
		2.2	33	(52:48:1)	
		— e)	— f)		
<u>1c</u>	PhSH	 2a	 3a	 b)	
		2.2	19	(58:1:42)	
 <u>1d</u> ^{d)}	PrOH	 2d	 3d	 b)	
		2.2	45		<u>2d</u> ^{d, g)}

a) Reaction temperature, 70 °C; time, 6 h. Ni(CO)₄, 7 equiv. b) An endo and exo mixture. c) Reaction time, 3 h. d) A cis and trans mixture. e) 1-Propanol was not added. f) The starting compound 1c was recovered. g) The other products were not isolated.

This work was partially supported by the Grant-in-Aid for Scientific Research from the Ministry of Education, Science, and Culture.

References

- 1) T. Hirao, Y. Harano, Y. Yamana, Y. Ohshiro, and T. Agawa, *Tetrahedron Lett.*, 24, 1255 (1983).
- 2) T. Hirao, S. Nagata, Y. Yamana, and T. Agawa, *Tetrahedron Lett.*, in press.
- 3) The 1,1-dibromo-2-chlorocyclopropanes 1 were easily obtained by the addition of dibromocarbene to the corresponding vinyl chlorides.
- 4) P. W. Jolly, "Comprehensive Organometallic Chemistry," ed by G. Wilkinson, Pergamon Press (1982), Vol. 6, p. 37.
- 5) R. S. Mali, M. Pohmakotr, B. Weidmann, and D. Seebach, *Liebigs Ann. Chem.*, 1981, 2272.
- 6) Another synthetic route to pyrenophorin via 7-hydroxy-cis-3-octenoic acid was explored by use of reductive deconjugation with diethyl phosphonate as a key reaction; T. Hirao, Y. Fujihara, K. Kurokawa, Y. Ohshiro, and T. Agawa, to be published.
- 7) The starting 1,1-dibromo-2-chlorocyclopropane 1d was prepared in the following way. 1,3-Dichloropropene was treated with the anion of ethyl acetoacetate, followed by decarboxylation in DMSO in the presence of LiCl. Thus obtained 6-chloro-5-hexen-2-one was reduced to the corresponding alcohol with NaBH_4 . After protecting the hydroxyl group, addition of dibromocarbene gave 1d.
- 8) To a mixture of 1a (1.0 mmol), *N,N*-dimethyltrimethylsilylamine (3.0 mmol), and benzaldehyde (3.0 mmol) in DMF (2.4 mL) was added $\text{Ni}(\text{CO})_4$ (3.0 mmol) at room temperature. The resultant solution was stirred at 70 °C for 3 h. After removal of excess $\text{Ni}(\text{CO})_4$ under the reduced pressure, the mixture was diluted with ether and filtered on celite 545. The ethereal solution was concentrated and flash column chromatographed to give *N,N*-dimethyl-2-(1-cyclohexenyl)cinnamamide (8a) in 31% yield.
- 9) The geometry of the benzylidene moiety was not determined yet although only one isomer was produced.
- 10) E. Yoshisato and S. Tsutsumi, *J. Chem. Soc., Chem. Commun.*, 1968, 33; I. Rhee, I. Ryu, H. Omura, S. Murai, and N. Sonoda, *Chem. Lett.*, 1979, 1435.
- 11) J. H. Nelson, P. N. Howells, G. C. DeLullo, and G. L. Landen, *J. Org. Chem.*, 45, 1246 (1980). Decarboxylation of the adduct might lead to 8a.

(Received July 24, 1985)