

## SYNTHESIS OF AZULENO[1,2-b]PYRROLE AND AZULENO[1,2-b]FURAN


Kunihide FUJIMORI,<sup>\*</sup> Hiroyuki FUKAZAWA, Yasuhiko NEZU, Kameji YAMANE,  
Masafumi YASUNAMI,<sup>†</sup> and Kahei TAKASE<sup>†</sup>

Department of Chemistry, Faculty of Science, Shinshu University,  
Asahi, Matsumoto 390


<sup>†</sup> Department of Chemistry, Faculty of Science, Tohoku University,  
Aramaki-aza-Aoba, Sendai 980

New heterocyclic compounds, methyl 1,2-dihydroazuleno[1,2-b]-pyrrole-9-carboxylate, its [1,2-c] isomer, and the furan analogues are synthesized by the reaction of 3-methoxycarbonyl-2H-cyclohepta[b]furan-2-one with morpholino enamines of N-ethoxycarbonyl-3-oxopyrrolidine or 3-oxotetrahydrofuran. By the dehydrogenation of the resulting [1,2-b] isomers followed by demethoxycarbonylation, azuleno[1,2-b]pyrrole and azuleno[1,2-b]furan are obtained, respectively.


Among three possible isomers of azulenopyrroles (1a-3a) and azulenofurans (1b-3b), respectively, only 2-phenyl derivatives of azuleno[2,1-b]pyrrole (2a) and azuleno[2,1-b]furan (2b) are known.<sup>1)</sup> In this communication, we wish to report the synthesis of new azulenopyrroles, azuleno[1,2-b]- and azuleno[1,2-c]pyrrole, and the furan analogues by the application of the new azulene synthesis<sup>2)</sup> using enamines derived from N-ethoxycarbonyl-3-oxopyrrolidine or 3-oxotetrahydrofuran.

1

a: X= NH  
b: X= O

2


a: X= NH  
b: X= O

3

a: X= NH  
b: X= O


The condensation of N-ethoxycarbonyl-3-oxopyrrolidine or 3-oxotetrahydrofuran with morpholine gave isomeric mixtures of two enamines (5a/5'a and 5b/5'b, respectively). A suspension of 3-methoxycarbonyl-2H-cyclohepta[b]furan-2-one (4)<sup>3)</sup> in anhydrous ethanol containing 3.5 molar equiv. of the morpholino enamines (mixture of 5a and 5'a) was refluxed for 7 days under argon atmosphere. After the reaction mixture was worked up in the usual way, the products were isolated by chromatography on silica gel using benzene-dichloromethane (1:1) as solvent to give methyl 3-ethoxycarbonyl-1,2-dihydroazuleno[1,2-b]pyrrole-9-carboxylate (6a; green prisms, mp 169.5-171 °C) and methyl 2-ethoxycarbonyl-1,3-dihydroazuleno[1,2-c]pyrrole-9-carboxylate (7a; purple microplates, 193-194.5 °C) in 34 and 37% yields, respectively. In a similar manner, the reaction of 4 with the morpholino enamines of 3-oxotetrahydrofuran (mixture of 5b and 5'b) gave methyl 1,2-dihydroazuleno[1,2-b]furan-9-car-

boxylate (6b; bluish green prisms, mp 114-114.5 °C) and methyl 1,3-dihydroazuleno[1,2-c]furan-9-carboxylate (7b; purple microneedles, mp 139.5-140 °C) in 59 and 24% yields.



Dehydrogenation of 6a by treatment with DDQ (2,3-dichloro-5,6-dicyano-p-benzoquinone) gave methyl 3-ethoxycarbonylazuleno[1,2-b]pyrrole-9-carboxylate (8a; dark purple prisms, mp 127-128.5 °C) in a 99% yield. Similar dehydrogenation of 6b gave methyl azuleno[1,2-b]furan-9-carboxylate (8b; bluish green prisms, mp 109-109.5 °C) in an 86% yield.

When the azuleno[1,2-b]pyrrole derivative (8a) was treated with 100% phosphoric acid at 90 °C, only demethoxycarbonylation at 9-position occurred to give ethyl azuleno[1,2-b]pyrrole-3-carboxylate (9; green plates, mp 51-52 °C) in a 95% yield. Further, heating of 9 with 100% phosphoric acid at 180 °C for 50 min resulted in deethoxycarbonylation at 3-position to give azuleno[1,2-b]pyrrole (1a; green microneedles, mp 185-186 °C(dec)) in a 97% yield. On the other hand, treatment of 8b with 100% phosphoric acid at 90-95 °C yielded azuleno[1,2-b]furan (1b; bluish green needles, mp 63-64 °C) in a 46% yield.



Moreover, refluxing of 7a with active manganese dioxide in dry benzene for 24 h resulted in dehydrogenation to give azuleno-[1,2-c]pyrrole derivative (10; yellowish brown crystals, 13% yield) having an ortho-quinoid structure. However, dehydrogenation of 7b has not been successful so far.

The difference between the vicinal coupling constants,  $\Delta J = J_{7,8} - J_{4,5}$ , in  $^1\text{H}$  NMR of azuleno[1,2-*b*]pyrrole (1a) and azuleno[1,2-*b*]-furan (1b) are 1.4 and 0.8 Hz, respectively, indicating that 7-membered ring ex-

hibits some bond alternation in analogy with azulenothiophene.<sup>4)</sup> The degree of bond alternation decreases in order of azuleno[1,2-b]thiophene,<sup>4)</sup> azuleno[1,2-b]-pyrrole (1a), and azuleno[1,2-b]furan (1b).

Detailed studies in the physical properties and reactivity of these azulenes (1a, 1b and azulenothiophenes<sup>4)</sup>) are now in progress.




Fig. 1. Electronic spectra of 1a, 1b, 8b, 9, and 10 in MeOH.

Table 1. Spectral data of azulenopyrroles and azulenofurans

	$^1\text{H}$ NMR(90 MHz, $\text{CDCl}_3$ ) $\delta$ ppm, $J$ in Hz; $^{13}\text{C}$ NMR( $\text{CDCl}_3$ ) $\delta$ ppm; UV $\lambda_{\max}/\text{nm}(\log \epsilon)$
<u>1a</u>	$^1\text{H}$ NMR: 6.64(1H, dm, $J=2.9$ , H-1), 6.7-7.4(3H, m, H-5,6,7), 7.26(1H, s, H-9), 7.49(1H, dd, $J=2.9$ , 2.5, H-2), 8.12(1H, dm, $J=8.6$ , H-4), 8.23(1H, dm, $J=10.0$ , H-8), and 8.72(1H, bs, NH) $^{13}\text{C}$ NMR: 99.8, 106.5, 119.5, 120.7, 123.0, 126.0, 129.0, 131.6, 133.6, 135.6, 138.4, and 139.5 UV(MeOH): 279(4.35), 312(4.69), 357(3.29, sh), 376(3.46), 394(3.57), 415(3.52), 590(2.30), 642(2.46), 708(2.39), and 794(2.02)
<u>1b</u>	$^1\text{H}$ NMR: 6.81(1H, d, $J=2.0$ , H-1), 6.7-7.6(3H, m, H-5,6,7), 7.11(1H, s, H-9), 7.81(1H, d, $J=2.0$ , H-2), 8.22(1H, dm, $J=10.1$ , H-8), and 8.36(1H, dm, $J=9.3$ , H-4) $^{13}\text{C}$ NMR: 103.1, 105.2, 118.8, 120.9, 121.5, 128.3, 135.7, 137.4, 138.8, 139.1, 149.9, and 151.2 UV(MeOH): 296(4.72), 354(3.52), 371(3.69), 389(3.79), 619(2.41), 652(2.30), 677(2.38), 721(1.96), and 750(2.07)
<u>8a</u>	$^1\text{H}$ NMR: 1.43(3H, t, $J=7.2$ , $\text{NCOOCH}_2\text{CH}_3$ ), 3.97(3H, s, $\text{COOCH}_3$ ), 4.43(2H, q, $J=7.2$ ,

	NCOOCH <sub>2</sub> CH <sub>3</sub> ), 6.78(1H, d, J=3.6, H-1), 7.2-7.8(3H, m, H-5,6,7), 7.70(1H, d, J=3.6, H-2), 9.56(1H, dm, J=9.6, H-4 or 8), and 9.67(1H, dm, J=9.6, H-8 or 4)
13	<sup>13</sup> C NMR: 14.4, 50.8, 63.6, 103.7, 106.5, 126.2, 126.3, 126.7, 127.6, 132.4, 134.8, 136.7, 137.3, 143.0, 143.6, 150.5, and 165.9
	UV(MeOH): 213(4.46), 283(4.28), 287(4.24, sh), 320(4.66), 331(4.73), 370(3.80, sh), 387(3.95), 405(3.61, sh), 560(2.62), 598(2.59), and 670(2.07, sh)
8b	<sup>1</sup> H NMR: 3.97(3H, s, COOCH <sub>3</sub> ), 7.05(1H, d, J=2.0, H-1), 7.2-7.8(3H, m, H-5,6,7), 7.89(1H, d, J=2.0, H-2), 8.48(1H, dm, J=9.3, H-4), and 9.69(1H, dm, J=10.1, H-8)
	<sup>13</sup> C NMR: 51.0, 102.3, 106.8, 122.2, 125.8, 126.8, 129.9, 137.2, 137.7, 140.2, 141.8, 149.0, 152.2, and 165.6
	UV(MeOH): 216(4.37), 265(4.10), 315(4.84), 327(4.88), 368(3.90), 385(4.07), 403(3.70), 577(2.58), 609(2.50), 629(2.51), 667(2.17), and 693(2.13)
9	<sup>1</sup> H NMR: 1.40(3H, t, J=7.2, NCOOCH <sub>2</sub> CH <sub>3</sub> ), 4.44(2H, q, J=7.2, NCOOCH <sub>2</sub> CH <sub>3</sub> ), 6.66(1H, d, J=3.6, H-1), 6.8-7.6(3H, m, H-5,6,7), 7.21(1H, s, H-9), 7.78(1H, d, J=3.6, H-2), 8.25(1H, dm, J=9.6, H-8), and 9.58(1H, dm, J=9.6, H-4)
	<sup>13</sup> C NMR: 14.5, 63.4, 105.2, 105.7, 121.0, 122.2, 123.4, 128.3, 131.5, 133.2, 135.7, 136.8, 140.5, 142.5, and 150.8
	UV(MeOH): 235(4.08, sh), 277(4.46, sh), 302(4.71), 310(4.71), 340(3.49, sh), 355(3.64), 372(3.67), 390(3.65), 560(2.57, sh), 608(2.88), 661(2.83), and 730(2.42)
10	<sup>1</sup> H NMR: 1.48(3H, t, J=7.3, NCOOCH <sub>2</sub> CH <sub>3</sub> ), 3.94(3H, s, COOCH <sub>3</sub> ), 4.50(2H, q, J=7.3, NCOOCH <sub>2</sub> CH <sub>3</sub> ), 6.6-7.0(3H, m, H-5,6,7), 7.47(1H, d, J=1.8, H-1), 7.5-7.7(1H, m, H-4), 7.86(1H, d, J=1.8, H-3), and 8.6-8.9(1H, m, H-8)
	<sup>13</sup> C NMR: 14.4, 51.1, 64.0, 107.3, 109.3, 111.1, 127.5, 129.2, 130.2, 130.5, 132.3, 134.5, 134.7, 141.1, 151.0, 151.2, and 165.9

## References

- 1) T. Morita, T. Nakadate, and K. Takase, *Heterocycles*, 15, 835 (1981).
- 2) P. W. Yang, M. Yasunami, and K. Takase, *Tetrahedron Lett.*, 1971, 4275; K. Takase and M. Yasunami, *Yuki Gosei Kagaku Kyokai Shi*, 39, 1172 (1981).
- 3) The compound was synthesized by the reaction of tropolone tosylate with dimethyl malonate (M. Yasunami and K. Takase, unpublished results).
- 4) K. Fujimori, T. Fujita, K. Yamane, M. Yasunami, and K. Takase, *Chem. Lett.*, 1983, 1721.
- 5) Satisfactory elemetal analyses have been obtained for all new compounds described in this paper.

( Received March 3, 1986 )