

Insertion Reactions of Dimethylgermanone $[(CH_3)_2Ge=O]$ into Si-O and Si-Cl BondsKyung-Tae KANG,[†] Georges MANUEL,^{††} and William P. WEBER*


Department of Chemistry,

Donald P. and Katherine B. Loker Hydrocarbon Research Institute,
University of Southern California, Los Angeles, CA 90089-1661, U.S.A.[†] Department of Chemical Education, Pusan National University,
Pusan 607, Korea^{††} Laboratoire des Organométalliques, Université Paul-Sabatier,
31062, Toulouse, Cedex, France

Flash vacuum co-pyrolysis (FVP) of 3,3-dimethyl-6-oxa-3-germabicyclo[3,1,0]hexane with various trapping reagents has been carried out. Adducts which result from insertion of dimethylgermanone into Si-O and Si-Cl bonds have been outlined.

The reaction chemistry of germanones, reactive intermediates which possess a germanium oxygen double bond is limited.¹⁾


Pyrolysis of 3,3-dimethyl-6-oxa-3-germabicyclo[3,1,0]hexane (1) has been shown to yield butadiene (3) and hexamethylcyclotrigermoxane (4).²⁾ The formation of 4 has been suggested to result from the cyclotrimerization of dimethylgermanone (2). We report two new insertions reaction of 2 into Si-O and Si-Cl bonds. These reactions have been carried out by FVP of 1 with various trapping reagents in the gas phase at 580° and 10⁻⁴ mm pressure. For example, FVP of 1 and 2,2,5,5-tetramethyl-1-oxa-2,5-disilacyclopentane (5) gave 2,2,4,4,7,7-hexamethyl-1,3-dioxa-2-germa-4,7-disilacycloheptane (6) (40%) and 3. Properties of 6: ¹H NMR (CDCl₃) δ = 0.068 (s, 12H), 0.524 (s, 6H), 0.695 (s, 4H); ¹³C NMR δ = 0.535 (4C, CH₃Si), 3.30 (2C, CH₃Ge), 11.98 (2C, CH₂Si); IR (NaCl) ν = 1005 and 980 cm⁻¹ Si-O-Ge; mass spectrum (rel. intensities) m/e 267 (6.6%), 265 (23.7%), 263 (17.1%), 261 (12.2%) P-15⁺, 240 (3.9%), 239 (26.7%), 237 (100%), 235 (73.8%), 233 (52.2%), 149 (6.3%), 148 (5.8%), 147 (34.2%), 133 (23.4%), 131 (15.4%), 119 (20.1%), 117 (16.4%), 115 (14.4%), 103 (11.8%); high resolution mass spectrum: calcd. for C₇H₁₉O₂⁷⁴Ge²⁸Si₂ 265.0135; found 265.0135. The formation of 6 probably results from the insertion of 2 into one of the Si-O bonds of 5. While 5 is a strained heterocyclic siloxane this is not an essential for insertion of 2 into siloxane (Si-O-Si) bonds. The FVP of 1 and



1,3-diphenyltetramethyldisiloxane gave bis(phenyldimethylsiloxy)dimethylgermane (7) (15%) and 2. Properties of 7: ^1H NMR δ = 0.308 (s, 12H), 0.417 (s, 6H), 7.3-7.6 (m, 10H); mass spectrum (rel. intensities) 393 (10.5%), 391 (36.9%), 389 (26.1%), 387 (17.9%) P-15⁺, 245 (10.7%), 244 (7.5%), 243 (52.2%), 242 (16.9%), 241 (39.0%), 239 (29.7%), 183 (21.6%), 181 (100%), 179 (74.6%), 177 (58.9%); high resolution mass spectrum: calcd. for $\text{C}_{17}\text{H}_{25}\text{O}_2^{74}\text{Ge}^{28}\text{Si}_2$ 391.0605; found 391.0601.

On the other hand, FVP of 1 and phenyldimethylmethoxysilane (8) gave 3, 4 and recovered 6. Apparently, Si-O bonds of siloxanes are more reactive than Si-O bonds of alkoxy silanes towards insertion of 2. By comparison, dimethylsilylone $[(\text{CH}_3)_2\text{Si}=\text{O}]$ inserts into both siloxane and alkoxy silane Si-O bonds.^{3,4)}

2 also inserts into Si-Cl bonds. For example, FVP of 1 and t-butyldimethylchlorosilane gave t-butyldimethylsiloxydimethylchlorogermane (9) (23%) and 2. Properties of 9: ^1H NMR δ = 0.057 (s, 6H), 0.837 (s, 6H), 0.858 (s, 9H); ^{13}C NMR δ = -2.29 (2C, CH_3Si), 7.43 (2C, CH_3Ge), 18.81 (1C, C), 26.125 (3C, $(\text{CH}_3)_3$); IR ν = 975 cm^{-1} Si-O-Ge; mass spectrum (rel. intensities) m/e 217 (6.5%), 216 (4.8%), 215 (43.4%), 214 (17.7%), 213 (100%), 212 (25.5%), 211 (77.9%), 209 (48.5%) P-57⁺. High resolution mass spectrum: calcd for $\text{C}_4\text{H}_{12}\text{O}^{35}\text{Cl}^{74}\text{Ge}^{28}\text{Si}$ 212.9558; found 212.9555.



A description of the FVP experiment can be found in Ref. 5.

This work was supported by the U.S. Air Force Office of Scientific Research 86-0042. K.T.K. thanks the Korean Science and Engineering Foundation. G.M. and W.P.W. thank NATO 845/83.

References

- 1) Review π -bonded germanium: J. Satge, *Adv. Organomet. Chem.*, 21, 241 (1982).
- 2) G. Manuel, G. Bertrand, W.P. Weber, and S.A. Kazoura, *Organometallics*, 3, 1340 (1984).
- 3) G.M. Golino, R.D. Bush, P. On, and L.H. Sommer, *J. Am. Chem. Soc.*, 97, 1957 (1975).
- 4) T.H. Lane, and C.L. Frye, *J. Organomet. Chem.*, 172, 213 (1979).
- 5) S.A. Kazoura, and W.P. Weber, *J. Organomet. Chem.*, 271, 47 (1984).

(Received July 5, 1986)