

Additions and Corrections

CHEMISTRY LETTERS, pp. 833-834, 1995

**Rhodium and Ruthenium Complexes with Optically Active Chelate
Cyclopentadienyl-Phosphine Bidentate Ligands**

Yasutaka Kataoka, Yoshinori Saito, Koji Nagata, Kenichi Kitamura, Atsushi Shibahara, and Kazuhide Tani*
Department of Chemistry, Faculty of Engineering Science, Osaka University, Toyonaka, Osaka 560

Page 834, **References and Notes** 7. The chemical shift value of ^{31}P NMR was lacking of a minus sign. The correct **References and Notes** 7 is shown below.

7 **2b**: $[\alpha]_{\text{D}}^{23} -35^\circ$ (*c* 0.074, CHCl_3); ^1H NMR (CDCl_3) $\delta =$ 1.37 (s, 3H), 1.40 (s, 3H), 2.27 (d, $J = 6.18$ Hz, 2H), 2.86 (m, 2H), 3.27 (d, $J = 1.48$ Hz, 2H), 3.82-3.92 (m, 1H), 4.24 (m, 1H), 6.22 (s, 1H), 7.13-7.49 (m, 14H); ^{31}P NMR (CDCl_3) $\delta = -23.0$ (s). HRMS: calcd. for $\text{C}_{28}\text{H}_{30}\text{O}_2\text{P}$ 429.1983, found 429.1987.