

Efficient Electrochemical *N*-Glycosylation of Silylated Pyrimidines with Protected Arylthioriboses in the Presence of a Catalytic Amount of NBS or Br₂

Junzo Nokami,* Masahiro Osafune, Yuichiro Ito, Fumiaki Miyake, Shin-ichi Sumida, and Sigeru Torii*[†]
 Department of Applied Chemistry, Faculty of Engineering, Okayama University of Science, Ridai-cho, Okayama 700-0005
[†]The Institute of Creative Chemistry Co., 874-5 Musa, Okayama 701-2141

(Received July 2, 1999; CL-990583)

Electrolysis of silylated pyrimidines with protected arylthioribose, in the presence of a catalytic amount of NBS or Br₂ as a mediator in an undivided cell, gave the corresponding *N*-glycosides (nucleosides) in good yield.

The glycosylation reaction of various nucleophiles (glycosyl acceptor) with sugar derivatives (glycosyl donor) to give *C*-,¹ *O*-,² and *N*-glycosides is one of the most important reactions in organic syntheses. Many chemists devote much effort to discover a convenient glycosylation procedure by use of suitable promoters as well as glycosyl donors. A reaction via oxonium ion under neutral conditions (without using Lewis acid) is well recognized as a characteristic electroorganic reaction.³ Thus, an electrochemical reaction seems to be one of the most effective ways to promote glycosylation. Noyori,⁴ Amatore,⁵ Balavoine,⁶ and Yoshida⁷ have already succeeded in electrochemical glycosylation of several types of alcohol with various stable glycosyl donors such as arylglycosides,⁴ aryl- or alkylthioglycosides,⁵ arylthio- or arylseleno-glycosides⁶ and aryltelluroglycosides,⁷ using supporting electrolytes such as LiClO₄, Bu₄NBF₄, Bu₄NClO₄ to give the corresponding *O*-glycosides effectively.

N-Glycosides such as nucleosides are very attractive in medicinal chemistry. However, electrochemical *N*-glycosylation of nucleoside bases with glycosyl donors to give *N*-glycosides has not been explored. Therefore, we investigated electrochemical *N*-glycosylation of nucleoside bases in order to discover a useful method for the synthesis of nucleosides, which are very important from an economic and environmental view point.


Electrolysis was carried out by employing arylthioriboses 2-5 as stable glycosyl donors, and silylated pyrimidines, bis(trimethylsilyl)thymine **1a** and bis(trimethylsilyl)-5-fluorouracil **1b** as soluble glycosyl acceptors, in the presence of a catalytic amount of supporting electrolyte in an undivided cell. We discovered that when tetrabutylammonium bromide (Bu₄NBr) was used as a supporting electrolyte, the desired *N*-glycosylation took place to give nucleosides in a moderate yield. However, the electrolysis using common supporting electrolytes, such as lithium perchlorate (LiClO₄), tetraethylammonium perchlorate (Et₄NClO₄), and tetraethylammonium tosylate (Et₄NOTs), was unsuccessful.⁸

The result suggests that the bromonium ion, formed by the electrolysis of the supporting electrolyte Bu₄NBr, serves as an effective mediator for the formation of the oxonium ion from arylthioribose via oxidative desulfurization, which then reacts with **1** to give nucleosides. Consequently, we examined the electrochemical *N*-glycosylation using a catalytic amount of *N*-bromosuccinimide (NBS)⁹ as a mediator, which was found to be more effective than Bu₄NBr. The results are listed in Table 1 (entries 1-12).


Table 1. *N*-Glycosylation of silylated nucleoside base **1** with thioglycosides **2**, **3**, **4** and **5**^a

Entry	Donor (α/β) ^b	Acceptor	C.D. ^c	F/mol ^d	Electrolyte (10 mol%)	Product	Yield/% ^e (α/β) ^b
1	2p (2/3)	1a	25	3	NBS	6a	73 (1.9/1)
2	2p (2/3)	1a	25	3	Bu ₄ NBr	6a	65 (2.2/1)
3	2t (1/1)	1a	25	3	NBS	6a	86 (2.3/1)
4	2t (1/1)	1a	25	3	Bu ₄ NBr	6a	73 (2.2/1)
5	2t (1/1)	1b	5	1	NBS	6b	74 (2.7/1)
6	2t (1/1)	1b	25	3	Bu ₄ NBr	6b	53 (2.2/1)
7	3p (1/3)	1a	25	3	NBS	7a	79 (4.2/1)
8	3p (1/3)	1b	5	1.5	NBS	7b	82 (3.2/1)
9	3t (1/3)	1a	25	3	NBS	7a	76 (5.2/1)
10	3t (1/4)	1a	25	3	Bu ₄ NBr	7a	41 (1.9/1)
11	3t (1/4)	1b	5	1.5	NBS	7b	92 (1.8/1)
12	3t (2/5)	1b	25	3	Bu ₄ NBr	7b	43 (1.8/1)
13 ^f	3t (1/2)	1b	25	3	Br ₂	7b	73 (2.7/1)
14 ^f	4 (2/3)	1b	25	3	NBS	8b	65 (2.6/1)
15 ^f	4 (2/3)	1b	25	2	Br ₂	8b	68 (2.0/1)
16 ^f	5 (1/2)	1b	25	4.5	NBS	9b	68 (β)
17 ^f	5 (1/2)	1b	25	4	Br ₂	9b	85 (β)

^a All reactions were performed with donor **2**, **3**, **4** or **5** (1 mmol) and acceptor **1** (1.2 mmol), using platinum plate (1 cm²) electrodes, in 2 mL of dry propanenitrile at room temperature unless otherwise noted. ^b Determined by ¹H NMR (400 MHz) integration of the mixture. ^c Current density (mA/cm²). ^d Electricity. ^e Isolation by column chromatography on silica gel. ^f Performed at 10 °C.



By employing the present glycosylation method, α -isomers were preferentially obtained in both dibenzylated 2-deoxyribose derivatives **2** and tribenzylated ribose derivatives **3**. This selectivity has already been shown in the reaction using stoichiometric amount of NBS,⁹ in which the formation of an ion-pair with a counter ion (succinimide anion) at the sterically less hindered β -site (*trans* to the most interactive substituent) is favorable to generation of the oxonium ion intermediate **10**. In this reaction, the nucleophile should be introduced from the opposite α -site (*cis* to the most interactive substituent) to give the α -isomer as a major product, bearing a sterically disadvantageous configuration. On the other hand, the triacetylated ribose derivatives **5** gave the β -isomer selectively due to the formation of the intramolecularly stabilized oxonium ion **11** (entry 16). This is in contrast to the reaction of diacetylated 2-deoxyribose derivative giving the α -isomer selectively (entry 14). The electrogenerated bromonium ion-assisted *N*-glycosylation is a versatile tool for nucleoside synthesis, on functioning a mediatory system, as shown in Scheme 1.



This finding prompted us to investigate further a similar electrolysis with a catalytic amount of bromine (Br_2)¹⁰ as the mediator. The results are also shown in Table 1 (entry 13, 15, 17). It can be assumed that the actual activator in the *N*-glycosylation using thioglycoside and NBS⁸ is also an in situ generated bromine species.

In conclusion, the electrochemical *N*-glycosylation undergoes preferentially in the presence of a catalytic amount (10 mol%) of NBS or bromine as a mediator. The stereoselectivity (β/α) of this glycosylation was similar to that of chemical glycosylation using a stoichiometric amount of NBS. It is most likely that the activation of the thioglycosides does not take place directly on the surface of the anode but indirectly in the bulk solution by the electrochemical mediatory system.

Further investigation of the synthetic applications of this

electrochemical glycosylation is now in progress.

This study was financially supported by the Electric Technology Research Foundation of Chugoku Electric Power Co. Ltd.

References and Notes

- For example, A. O. Stewart, and R. M. Williams, *J. Am. Chem. Soc.*, **107**, 4289 (1985) and references cited therein.
- For a recent review on *O*-glycosylation, see: K. Toshima and K. Tatsuta, *Chem. Rev.*, **93**, 1503 (1993).
- T. Shono and Y. Matsumura, *J. Am. Chem. Soc.*, **91**, 2803 (1969); J. Nokami, M. Kawada, R. Okawara, S. Torii, and H. Tanaka, *Tetrahedron Lett.*, **1979**, 1045; T. Mandai, H. Irie, M. Kawada, and J. Otera, *Tetrahedron Lett.*, **25**, 2371 (1984); J. Yoshida and S. Isoe, *Chem. Lett.*, **1987**, 631; S. Torii, "Electroorganic Syntheses," Kodansha VCH, Tokyo (1985), p. 246.
- R. Noyori and I. Kurimoto, *J. Org. Chem.*, **51**, 4320 (1986).
- C. Amatore, A. Jutand, J.-M. Mallet, G. Meyer, and P. Sinay, *J. Chem. Soc., Chem. Commun.*, **1990**, 718; J.-M. Mallet, G. Meyer, F. Yvelin, A. Jutand, C. Amatore, and P. Sinay, *Carbohydr. Res.*, **244**, 237 (1993); C. Amatore, A. Jutand, G. Meyer, P. Bourhis, F. Machetto, J.-M. Mallet, P. Sinay, C. Tabour, and Y.-M. Zhang, *J. Appl. Electrochem.*, **24**, 725 (1994).
- G. Balavoine, A. Gref, J.-C. Fischer, and A. Lubineau, *Tetrahedron Lett.*, **31**, 5761 (1990); G. Balavoine, S. Bertheina, A. Gref, J.-C. Fischer, and A. Lubineau, *J. Carbohydr. Chem.*, **14**, 1217, 1237 (1995).
- S. Yamago, K. Kokubo, and J. Yoshida, *Chem. Lett.*, **1997**, 111.
- The substrates **2** and **3** were recovered after electrolysis (passing 1-3 F/mol of electricity) with these supporting electrolytes, although lithium perchlorate was very effective for electrochemical *O*-glycosylation (Ref. 4-7).
- A stoichiometric amount of NBS (1.1-1.2 equiv.) was used as a promoter of *O*- and *N*-glycosylations of phenylthioglycosides. *O*-glycosylation: K. C. Nicolaou, S. P. Seitz, and D. P. Papahatjis, *J. Am. Chem. Soc.*, **105**, 2430 (1983); K. Fujiwara, S. Amano, and A. Murai, *Chem. Lett.*, **1995**, 191; M. Nakata, T. Tamai, T. Kamino, M. Kinoshita, and K. Tatsuta, *Tetrahedron Lett.*, **35**, 3099 (1994); *N*-glycosylation: H. Sugimura, K. Osumi, T. Yamazaki, and T. Yamaya, *Tetrahedron Lett.*, **32**, 1813 (1991); K. Sujino and H. Sugimura, *Synthlett*, **1992**, 553; H. Sugimura, K. Sujino, and K. Osumi, *Tetrahedron Lett.*, **33**, 2515 (1992); H. Sugimura, I. Muramoto, T. Nakamura, and K. Osumi, *Chem. Lett.*, **1993**, 169; H. Sugimura, K. Osumi, Y. Kodaka, and K. Sujino, *J. Org. Chem.*, **59**, 7653 (1994); R. Young, S. Shaw-Ponter, G. W. Hardy, and G. Mills, *Tetrahedron Lett.*, **35**, 8687 (1994).
- A stoichiometric amount of bromine (Br_2) was also used as a promoter of *O*- and *N*-glycosylations of thioglycosides: *O*-glycosylation: J. O. Kihkberg, D. A. Leigh, and D. R. Bundle, *J. Org. Chem.*, **55**, 2860 (1990); *N*-glycosylation: S. Hanessian, K. Sato, T. J. Liak, N. Danh, and D. Dixit, *J. Am. Chem. Soc.*, **106**, 6114 (1984).