

Alternative Synthesis of (*Z*)-1-Aryl-1-(tributylstannyl)-2-(triethylgermyl)ethenes and the Unprecedented Germyl 1,2-Migration during the Destannylation of the Adducts

Taichi Nakano,* Yoshiya Senda, and Takashi Miyamoto


Department of Material Science and Technology, School of High-Technology for Human Welfare,
Tokai University, 317 Nishino, Numazu, Shizuoka 410-0395

(Received October 6, 2000; CL-000910)

A specific combination catalyst, Pd(dba)₂ and 4-ethyl-1-phospha-2,6,7-trioxabicyclo[2.2.2]octane, effectively catalyzed the addition of tributyl(triethylgermyl)stannane to aryl-acetylenes in tetrahydrofuran to give (*Z*)-1-aryl-2-(germyl)-1-(stannyl)ethenes in high yields. The (*Z*)-1-aryl-2-(germyl)-1-(stannyl)ethenes were subject to the unprecedented germyl 1,2-migration during the destannylation using HI / TBAI in toluene to produce 1-aryl-1-(germyl)ethenes in high yields.

Palladium-catalyzed silastannylation of acetylenes has been well studied and known to afford vicinal sila(stannyl)ethenes.¹ The addition of a Ge–Sn bond of a (germyl)stannane to a triple bond of nonterminal α,β -acetylenic esters catalyzed by Pd(PPh₃)₄ forms a mixture of vicinal germa(stannyl)ethenes with (*E*)- and (*Z*)-configurations.² In contrast, the addition to terminal alkynes involving phenylacetylene only forms the (*Z*)-germyl(stannyl)ethenes, in which a germyl group is combined with the terminal sp² carbon.³ However, the yields did not exceed 50%. We found that a specific combination catalyst, Pd(dba)₂ (dba = dibenzylideneacetone) and P(OCH₂)₃CeT (4-ethyl-1-phospha-2,6,7-trioxabicyclo[2.2.2]octane, L), effectively catalyzed the addition of tributyl(triethylgermyl)stannane **1** to arylacetylenes to produce (*Z*)-1-aryl-1-(tributylstannyl)-2-(triethylgermyl)ethenes in high yields and the unprecedented germyl 1,2-migration occurred during the destannylation of the (*Z*)-1-aryl-2-(triethylgermyl)-1-(stannyl)ethenes to form 1-aryl-1-(triethylgermyl)ethenes. We report these preliminary results.

The germastannylation of phenylacetylene using **1** in THF using the Pd(dba)₂ and L⁴ combination catalyst afforded (*Z*)-1-(tributylstannyl)-2-(triethylgermyl)-1-phenylethenes **2a**⁵ in 85% yield. The use of the THF solvent is essential in Scheme 1, in contrast to no reaction in benzene.



Scheme 1.

The NMR coupling constants for **2a** between tin and the vinylic proton, ³J_{Sn-H}, were –161.2 and –168.0 Hz for tin-117 and tin-119, respectively. These constants are typical for the structure with tin and the vinylic proton in *trans*-disposition.^{3,6}

Table 1. Selected NMR data^a for **2a–e**


product No.	δ(=CH) ^b	δ(C ₁) ^c	δ(C ₂) ^c	³ J _{SnH} ^d
2a	6.63	165.5	151.8	-161.2 -168.4
2b	6.62	164.3	150.3	-157.2 -164.8
2c	6.67	164.3	152.4	-154.8 -162.0
2d	6.69	164.0	158.9	-150.0 -156.8
2e	6.63	164.9	148.2	-162.4 -169.6

^aChemical shifts referenced to TMS in ppm in chloroform-*d*.

^bVinyl proton. ^cC₁ is the tin-bearing vinyl carbon, C₂ bears germanium. ^dMeasured three times in CDCl₃ and its mean is described in Hz. The former is for tin-117, the latter for 119.


Other reaction examples are summarized in Scheme 1 and selected NMR data for **2a–e** in Table 1.

Next, we examined the Kosugi-Migita-Stille coupling⁷ of **2a** with allyl bromide in order to obtain chemical confirmation for the structure. Thus, a DMF (*N,N*-dimethylformamide, 5 mL) solution of **2a** (1 mmol), the bromide (5.5 mmol), Pd(dba)₂ (0.01 mmol) and CuI (0.018 mmol) was stirred at 40 °C. The reaction smoothly proceeded and went to completion in 5 h to produce (*E*)-1-(triethylgermyl)-2-phenylpenta-1,4-diene **3a**⁸ in 86% yield (Scheme 2). The reaction also proceeded at room temperature to yield **3a** in 85% yield, although a longer reaction time (20 h) was needed.




The NMR spectrum of the **3a** clearly showed that vinyl proton on the sp² carbon bearing the germyl group appeared as a singlet (δ = 6.02 ppm), and supported the fact that the allyl group did combine with the sp² carbon bearing the phenyl group, not another sp² carbon. Results from Scheme 2 again showed that **2a** had the (*Z*)-configuration with the stannyl group connected to the sp²-carbon bearing an aryl group. We next examined a preliminary experiment for the destannylation of **2a**, aimed at the synthesis of a series of *trans*-1-aryl-2-(triethylgermyl)ethenes using HI / TBAI (tetrabutylammonium iodide). The method has been reported by Mori et al. as a selective sub-

stitution of the stannyl moiety in (Z)-1-(silyl)-2-(stannyl)alk-1-enes by hydrogen with retention of the configuration.^{1g}



Scheme 3.

However, strikingly enough, the destannylation of **2a** did not form the expected product, but only 1-(triethylgermyl)-1-phenylethene **4a**⁹ {chemical shifts of vinyl protons; δ 5.89 ppm (d, 1H, $J = 2.4$ Hz), 5.43 ppm (d, 1H, $J = 2.4$ Hz)} in an isolated yield of 74%, as shown in Scheme 3. The reaction was very clean, and no other products were formed except for tributylstannyl iodide. Under similar reaction conditions, **2b** also underwent a migration to form 1-(*p*-chlorophenyl)-1-(triethylgermyl)ethene **4b**¹⁰ in 80% yield. The **4a** and **4b** possibly form through the Markovnikov addition followed by the formation of a transient cyclic cation, **I-b**, followed by destannylation forming a double bond (the unprecedented germyl 1,2-migration), as shown in Scheme 4.



Scheme 4.

Under similar conditions, (Z)-1-(tributylstannyl)-2-(trimethylsilyl)-1-phenylethene prepared by the palladium-catalyzed addition of tributyl(trimethylsilyl)stannane to phenylacetylene did not produce a similar type of silyl 1,2-migration. The reaction gave 1-phenyl-2-(silyl)ethene. A full study for the present reactions is now underway.

Financial support from the Shizuoka Shougou Kenkyu Kikoh Foundation of Japan is gratefully acknowledged by one of the authors (T. N.). The authors are indebted to Ms. Fumiyo O-ikawa (Tokai University) for the mass measurements, especially the high resolution ones.

References and Notes

- 1 a) T. N. Mitchell, H. Killing, R. Dicke, and R. Wickenkamp, *J. Chem. Soc., Chem. Commun.*, **1985**, 354. b) B. L. Chenard, E. D. Laganis, F. Davidson, and T. V. RajanBabu, *J. Org. Chem.*, **50**, 3666 (1985). c) B. L. Chenard and C. M. Van Zyl, *J. Org. Chem.*, **51**, 3561

- (1986). d) B. L. Chenard, C. M. Van Zyl, and D. R. Sanderson, *Tetrahedron Lett.*, **27**, 2801 (1986). e) T. N. Mitchell, R. Wickenkamp, A. Amamria, R. Dicke, and U. Schneider, *J. Org. Chem.*, **52**, 4868 (1987). f) K. Ritter, *Synthesis*, **1989**, 218. g) M. Mori, N. Watanabe, N. Kaneta, and M. Shibasaki, *Chem. Lett.*, **1991**, 1651. h) M. Murakami, H. Amii, N. Takizawa, and Y. Ito, *Organometallics*, **12**, 4223 (1993).
- 2 E. Piers and R. T. Skerlj, *J. Chem. Soc., Chem. Commun.*, **1987**, 1025.
- 3 T. N. Mitchell, U. Schneider, and B. Fröhling, *J. Organomet. Chem.*, **384**, C53 (1990).
- 4 P(OCH₂)₃CET: For the use of the particular ligand in the Pd-catalyzed addition of Si-Si and Ge-Ge bonds to acetylenes in benzene solvent, see: (the former); H. Yamashita, M. Catellani, and M. Tanaka, *Chem. Lett.*, **1991**, 241. (the latter); a) T. Hayashi, H. Yamashita, T. Sakakura, Y. Uchimar, and M. Tanaka, *Chem. Lett.*, **1991**, 245. b) K. Mochida, C. Hodota, H. Yamashita, and M. Tanaka, *Chem. Lett.*, **1992**, 1635.
- 5 A THF (5 mL) solution of 1 mmol, 4 mmol, 0.05 mmol and 0.1 mmol of **1**, phenylacetylene, Pd(dba)₂ and L was stirred at 80 °C for 50h. Treating the resulting mixture with aqueous KF followed by purification by column chromatography (silica gel, hexane) gave spectroscopically pure **2a** in 85% yield. ¹H-NMR (CDCl₃) δ 7.24 (m, 2H), 7.13 (m, 2H), 6.98 (m, 1H), 6.63 (s, 1H), 1.4 (m, 6H), 1.26 (sep, 6H, $J = 7.2$ Hz), 1.07 (t, 9H, $J = 7.6$ Hz), 0.87 (m, 21H) ppm. Other reactions were carried out similar to the synthesis of **2a**. Reaction time; 19 h for **2b**, 30 h for **2c**, 40 h for **2d**, and 35 h for **2e**.
- 6 a) T. N. Mitchell, A. Amamria, H. Killing, and D. Rutschow, *J. Organomet. Chem.*, **241**, C45 (1983). b) T. N. Mitchell, A. Amamria, H. Killing, and D. Rutschow, *J. Organomet. Chem.*, **304**, 257 (1986).
- 7 a) M. Kosugi, Y. Shimizu, and T. Migita, *Chem. Lett.*, **1977**, 1423. b) D. Milstein and J. K. Stille, *J. Am. Chem. Soc.*, **100**, 3636 (1978).
- 8 Treating the resulting mixture with aqueous KF followed by purification by column chromatography (silica gel, hexane) gave spectroscopically pure **3a** in 86% yield. ¹H-NMR (CDCl₃) δ 7.41 (m, 2H), 7.29 (m, 2H), 7.22 (m, 1H), 6.02 (s, 1H), 5.77 (m, 1H), 5.01 (m, 2H), 3.33 (m, 2H), 1.07 (t, 9H, $J = 7.6$ Hz), 0.91 (q, 6H, $J = 7.6$ Hz) ppm.
- 9 To a mixture of **2a** (0.23 mmol), TBAI (0.23 mmol) and toluene (1 mL) was added dropwise hydroiodic acid (0.35 mL, 57%) at 0 °C, then stirred for 1 h. The resulting mixture was washed with aqueous sodium bicarbonate. Treating the organic layer with aqueous KF followed by purification by column chromatography (silica gel, hexane) gave spectroscopically pure **4a** in 74% yield. ¹H NMR (CDCl₃) δ 7.28 (m, 2H), 7.19 (m, 3H), 5.89 (d, 1H, $J = 2.4$ Hz), 5.43 (d, 1H, $J = 2.4$ Hz), 1.01 (t, 9H, $J = 7.6$ Hz), 0.88 (q, 6H, $J = 7.6$ Hz) ppm.
- 10 By a procedure similar to that for **4a**, spectroscopically pure **4b** was isolated in 80% yield. ¹H-NMR (CDCl₃) δ 7.26 (dd, 2H, $J = 6.4, 2.2$ Hz), 7.09 (dd, 2H, $J = 6.4, 2.2$ Hz), 5.86 (d, 1H, $J = 2.8$ Hz), 5.43 (d, 1H, $J = 2.8$ Hz), 1.00 (t, 9H, $J = 8.4$ Hz), 0.87 (q, 6H, $J = 8.4$ Hz) ppm.