

Reactivities of Triethylgermylborate in Methanol

Masato Nanjo,* Kazuhiko Matsudo, and Kunio Mochida*


Department of Chemistry, Faculty of Science, Gakushuin University, 1-5-1 Mejiro, Toshima-ku, Tokyo 171-8588

(Received August 8, 2001; CL-010762)

The reactivity of lithium (triethylgermyl)triphenylborate, prepared from unsolvated triethylgermyllithium and triphenylborane, with organic substrates in methanol was investigated. The germylborates reacted with organic halides and acyl halides to give the corresponding germyl-substituted compounds. From the reaction with carbonyl compounds, esters, carboxylic acid, alkenes, and alkynes, germylborates were completely recovered.

The group 14 element-alkali metal species as nucleophiles have been extensively studied in organic and organometallic syntheses, e.g., the introduction of group 14 elements into organic halides, metal halides, carbonyl compounds, etc.¹⁻³


Very recently, our group reported the first successful isolation of lithium (triethylgermyl)triphenylborate (**1**), prepared from unsolvated triethylgermyllithium⁴ and triphenylborane^{5,6}



The polymeric structure of germylborate **1** in the solid state was destroyed by addition of MeOH to give a dimeric structure of **1** with coordination of MeOH to the lithium atoms, $\{[\text{Et}_3\text{GeBPh}_3][\text{Li}(\text{MeOH})]\}_2$ (**2**). The molecular structures of **1** and **2** were confirmed by X-ray diffraction analysis. We report herein some reactions of **2** with organic substrates in MeOH.

At first, the stability of the MeOH-coordinated germylborate **2** was examined in excess of degassed MeOH and H₂O. The germylborate **2** was very stable in excess degassed MeOH and H₂O at room temperature for 1-4 days, and was completely recovered unreacted. The germylborate **2** did not change in degassed MeOH even at 50 °C for 10 h. The stability of **2** may be explained by its planar four-membered ring composed of two lithium and two oxygen atoms.⁵

The displacement of halides and other groups from organic substrates by metalate anions represents one of the most important routes for the formation of metal-carbon σ bonds. The reactions of **2** with simple alkyl halides and aromatic halides (RX) in MeOH solvent were carried out at room temperature for 4 days. Hydrolysis of the reaction mixture led to triethylgermyl-substituted products as major products together with some unidentified byproducts. All products were identified by comparing their NMR, GC-MS spectra, and retention times on GC with those of authentic samples. The results are summarized in Table 1.



The yields of the products increased in the order of RCl < RBr

< RI for the octyl halides and phenyl halides shown in Table 1. This order is consistent with the recognized reactivities of triethylgermyl-alkali metals in HMPA and the bond strength of the carbon-halogen bond.⁷ The rest of **2** remained unreacted. The reaction of **2** with benzyl bromide afforded mainly benzyltriethylgermane in low yield. Several unidentified byproducts were also detected by GC-MS spectra.


No formation of coupling products derived by electron-transfer in the reactions of **2** with organic halides were detected by GC.

Table 1. Reactions of **2** with organic halides in MeOH

RX	Conditions	Products, Yields/% ^a
C ₈ H ₁₇ Cl	r.t., 4 d	C ₈ H ₁₇ GeEt ₃ (17)
C ₈ H ₁₇ Br	r.t., 4 d	C ₈ H ₁₇ GeEt ₃ (31)
C ₈ H ₁₇ I	r.t., 4 d	C ₈ H ₁₇ GeEt ₃ (98)
PhCl	r.t., 4 d	PhGeEt ₃ (9)
PhBr	r.t., 4 d	PhGeEt ₃ (38)
PhI	r.t., 4 d	PhGeEt ₃ (90)
PhCH ₂ Br	r.t., 4 d	PhCH ₂ GeEt ₃ (40)

^a Yields were determined by GC and NMR.

With acyl chlorides, the germylborate **2** in MeOH at room temperature for 2 days gave the corresponding triethylgermyl ketones as the sole product in high yields.⁸



The results of the reactions of **2** with various acyl chlorides (RCOCl) in MeOH are summarized in Table 2.

Table 2. Reactions of **2** with acyl chlorides (RCOCl) in MeOH

RCOCl	Conditions	Products, Yields/% ^a
t-BuCOCl	r.t., 2 d	t-BuCOGeEt ₃ (> 95)
C ₆ H ₅ COCl	r.t., 2 d	C ₆ H ₅ COGeEt ₃ (> 95)
p-MeC ₆ H ₄ COCl	r.t., 2 d	p-MeC ₆ H ₄ COGeEt ₃ (> 95)
p-MeOC ₆ H ₄ COCl	r.t., 2 d	p-MeOC ₆ H ₄ COGeEt ₃ (>95)


^a Yields were determined by NMR.

The MeOH-coordinated germylborate **2** smoothly reacted with trimethylacetyl chloride and aryl chlorides in MeOH to give the corresponding germylketones, respectively, in high yields shown in Table 2. These germylketones are commonly yellow, which were purified by rapid column chromatographic separation (silica gel, hexane/ether) and were identified by NMR and GC-MS spectra.

Representative methods of preparation of acylgermanes hitherto reported are (a) germylation of an acyl anion equivalent based on dithiane compounds followed by hydrolysis,^{9,10} (b)

germylation of metalated enol ethers,¹¹ (c) direct carbonylation by germyl-lithiums,¹² (d) Pd-catalyzed germylation of acyl halides,¹³ (e) germylation of acyl chlorides by germyl-cuprates,^{14–16} (f) hydrolysis of α -chloro ethers.¹⁷ However, these reactions involve strongly reducing conditions, oxidation procedures, or severe reaction conditions. The germylborate **2** is a very useful reagent of germylation of acyl halides in MeOH solvent.

The reactions of the MeOH-coordinated germylborate as nucleophiles with carbonyl compounds, esters, carboxylic acid and carbon-carbon multiple bonds were also examined. The reaction of MeOH-coordinated germylborate **2** with 3-pentanone in MeOH did not occur at 50 °C for 6 h, and remained unreacted. To the MeOH solution of **2** was added benzophenone under similar reaction conditions. The color of the solution turned to yellow, but the germylborate **2** was mostly recovered. The observed color may arise from a charge-transfer interaction of the σ HOMO of germylborate and the LUMO of carbonyl compound.



The reaction of **2** with acetic acid in MeOH did not proceed at room temperature for 4 days and recovered unreacted. With methyl acetate the germylborate **2** in MeOH under similar reaction conditions did not change.

The germylborate **2** in MeOH proved to be unreactive toward carbon-carbon multiple bonds. With 2,3-dimethyl-2-butene and 6-dodecyne under similar reaction conditions, the germylborate **2** recovered unreacted from the MeOH solutions. These results observed in MeOH are in marked contrast to those in the reactions of triethylgermyllithium with carbonyl compounds, esters, alkenes, and alkynes in HMPA.² It is well known that the germyllithium in HMPA reacts with carbonyl compounds, esters, alkenes, and alkynes to give the corresponding germyl-substituted products.¹

We thank Mitsubishi Material Co. Ltd., for the gift of germanium tetrachloride. This work was partially supported by the Ministry of Education, Science, Culture and Sports.

Dedicated to Professor Hideki Sakurai on the occasion of his 70th birthday.

References and Notes

- D. D. Davies and C. E. Gray, *Organomet. Chem. Rev. A*, **6**, 283 (1970).
- For examples see a) D. A. Armitage, "Comprehensive Organometallic Chemistry," ed. by G. Wilkinson, F. G. A. Stone and E. W. Abel, Pergamon, New York (1982), Vol. 2, Chap. 9, pp 99–104. b) H. Sakurai, *Kagaku no Ryoiki*, **30**, 67 (1976). c) M. Fujita and T. Hiyama, *Yukigosei Kagaku Kyokaishi*, **42**, 293 (1984).
- For example, see a) P. Riviere, M. R-Baudet, and J. Satge, "Comprehensive Organometallic Chemistry," ed. by G. Wilkinson, F. G. A. Stone, and E. W. Abel, Pergamon, New York (1982), Vol. 2, Chap. 10, pp 468–473. b) M. Lesbre, P. Mazerolles, J. Satge, "Organic Compounds of Germanium," Interscience, New York (1971). c) K. Mochida, *Yukigosei Kagaku Kyokaishi*, **49**, 288 (1991).
- Unsolvated triethylgermyllithium could be prepared from triethylgermane, di-*tert*-butylmercury, and lithium metal in benzene
- K. Matsudo, M. Nanjo, and K. Mochida, 10th International Conference on the Coordination and Organometallic Chemistry of Germanium, Tin and Lead, Bordeaux (France), 2001 (July), Abstr., No. 2p13; M. Nanjo, K. Matsudo, and K. Mochida, submitted for publication.
- No structural information of the HMPA complex of lithium (triethylgermyl)triphenylborate has been reported; E. J. Bulten and J. G. Noltes, *J. Organomet. Chem.*, **29**, 409 (1971).
- K. Mochida and N. Matsushige, *J. Organomet. Chem.*, **229**, 11 (1982).
- The typical experimental procedure of the reaction of MeOH-coordinated germylborate **2** with benzoyl chloride in the MeOH solvent is as follows: to a stirred MeOH-coordinated lithium (triethylgermyl)triphenylborane (100 mg, 0.245 mmol) and degassed MeOH (1.5 mL) in a 25-mL Schlenk tube was added benzoyl chloride (140 mg, 1.0 mmol) at room temperature under Ar. The reaction mixture was stirred for 2 days. After evaporation of MeOH and unreacted benzoyl chloride, the resulting residue was purified by column chromatography (silica gel, hexane/ether).
- A. G. Brook, J. M. Duff, P. F. Jones, and N. R. Davies, *J. Am. Chem. Soc.*, **89**, 431 (1967).
- E. J. Corey, D. Seebach, and R. Freeman, *J. Am. Chem. Soc.*, **89**, 434 (1967).
- J. A. Sonderquist and A. Hassner, *J. Am. Chem. Soc.*, **102**, 1577 (1980).
- R. J. Cross and F. Glockling, *J. Organomet. Chem.*, **3**, 146 (1965).
- K. Yamamoto, A. Hayashi, S. Suzuki, and J. Tsuji, *Organometallics*, **6**, 974 (1987).
- E. Piers and R. Lemieux, *Organometallics*, **14**, 5011(1995).
- D. P. Curran, U. Diederichsen, and M. Palovich, *J. Am. Chem. Soc.*, **119**, 4797 (1997).
- U. Iserloh and D. P. Curran, *J. Org. Chem.*, **63**, 471 (1998).
- R. Johannesen and T. Benneche, *J. Chem. Soc., Perkin Trans. 1*, **2000**, 2677.