


Additions and Corrections

CHEMISTRY LETTERS, pp 14-15, 2002

The First Total Synthesis of (\pm)-Napyradiomycin A1

Kuniaki Tatsuta,* Yoshiki Tanaka, Masazumi Kojima, and Hiroshi Ikegami
*Department of Applied Chemistry, School of Science and Engineering, Waseda University,
 3-4-1 Ohkubo, Shinjuku-ku, Tokyo 169-8555*

Page 15, the caption of Scheme 1 is incomplete. This mistake was caused on the publishing process. Scheme 1 and its caption should appear as given below.



Scheme 1. a) $H_2SO_4/MeOH$, reflux, 72 h, quant. b) $TBSCl$, NaH/THF , 0 °C, 30 min, quant. c) Et_2NH , $Me_3Al/PhMe$, reflux, 10 h, quant. d) $s\text{-}BuLi$, $TMEDA$, DMF/THF , -78 °C, 1 h, quant. e) $PhSO_2Na/AcOH$, 80 °C, 12 h, 80%. f) $MOMCl$, $i\text{-}Pr_2NEt/DMF$, 0 °C, 1 h, 88%. g) $t\text{-}BuOLi/THF$, -78 °C – rt, 2 h. h) $MnO_2/CHCl_3$, rt, 30 min, 2 steps 64%. i) SO_2Cl_2/CH_2Cl_2 , -40 °C, 40 min, 77%. j) $n\text{-}BuLi$, CuI , $LiBr/THF$, -78 °C, 1 h, 55%. k) $NCS/1,2\text{-dimethoxyethane}$, rt, 40 min, quant. l) KPh_3BH/THF , -78 °C, 30 min, 61%. m) $(PyS)_2$, $n\text{-}Bu_3P/PhMe$, rt, 5 min, 76%. n) KPh_3BH/THF , 0 °C, 30 min, 72%. o) NCS , $KHMDS/THF$, -78 °C, 3 h, 81%. p) $PPTS/t\text{-}BuOH$, 90 °C, 18 h, 91%.