

Chlorotrimethylsilane as a Mild and Effective Source of Acid Catalyst in Reductive Benzylation

Minoru Izumi* and Koichi Fukase†

Department of Bioresources Chemistry, Faculty of Agriculture, Okayama University,
1-1-1 Tsushimanaka, Okayama 700-8530

†Department of Chemistry, Graduate School of Science, Osaka University,
1-1 Machikaneyama, Toyonaka, Osaka 560-0043

(Received February 2, 2005; CL-050148)

New reductive benzylation of hydroxy functions was effected by using the combination of benzaldehyde, TMSCl, and Et₃SiH. TMSCl first reacts with alcohols to form HCl, which is probably the ultimate catalyst for this reaction system. TMSCl can also trap water and thus effectively promote dehydration reactions when more than stoichiometric amount of TMSCl is used. Although excess TMSCl was required for the foregoing reactions, TMSCl could be readily removed by simple evaporation.

Chlorotrimethylsilane (TMSCl) has been used as an acid catalyst for esterification and ketal formation.^{1,2} We have also reported the new convenient Fischer glycosidation using TMSCl as acid catalyst.³ In these reactions, TMSCl first reacts with alcohols or carboxylic acids to form HCl, which is probably the ultimate catalyst for these reaction systems. TMSCl can also trap water and thus effectively promote dehydration reactions when more than stoichiometric amount of TMSCl is used.

We first investigated reductive benzylation of hydroxy function using benzaldehyde, TMSCl, and Et₃SiH. Nishizawa^{4a} reported reductive benzylation of hydroxy function, in which the hydroxy group was once trimethylsilylated and then treated with benzaldehyde, TMSOTf, and Et₃SiH.^{4,5} However, partial cleavage of O-TMS group prior to benzylation reaction decreased the overall yields in some cases: no benzylation proceeds on a free hydroxy group. We previously described a new one-pot procedure to improve Nishizawa's method for the protection of the hydroxy group of 3-hydroxy fatty acid esters.⁶ Direct benzylation was effected in high yields via in situ O-trimethylsilylation by using benzaldehyde, TMS₂O, TMSOTf, and Et₃SiH. This method was, however, not suitable for benzylation of glucose derivatives having many benzyl groups, since some benzyl groups already present were reductively cleaved during the reaction. We, therefore, investigated a combination of TMSCl and Et₃SiH for reductive benzylation via in situ O-trimethylsilylation under milder conditions. A possible reaction mechanism is shown in Figure 1. In this reaction, TMSCl promotes O-trimethylsilyla-




Figure 1. Possible mechanism of the present reductive benzylation.

tion and then the formation of the oxocarbenium ion intermediate, which is reduced with Et₃SiH to give the desired ether.

As shown in Table 1, 10 equiv. of TMSCl was required for smooth reductive benzylation of benzyl alcohol (**1**) (Entry 3). The 6-hydroxy group of glucose derivative **3** was benzylated under these conditions in 81% yield (Entry 4).⁷ The more hindered 4-hydroxy group of glucose derivative **4** was also smoothly benzylated in 77% yield (Entry 5). Addition of excess benzaldehyde did not improve the yields of the desired benzylated compound but increased the formation of dibenzyl ether **2**.

Reductive alkylation of **3** with 1-naphthaldehyde and 9-anthracenecarboldehyde gave trace amount of the alkylated product but most of the starting aldehydes were transformed to symmetrical ethers via reduction of the aldehydes and subsequent reductive alkylation of the resulting alcohols. These results suggest more reactive aldehydes than benzaldehyde cannot be used, since they might be consumed prior to alkylation. By contrast, less reactive *p*-nitrobenzaldehyde can be used for the present reductive alkylation. We previously reported the use of *p*-nitrobenzyl group for temporary protection of hydroxy functions.⁸

Table 1. Reductive benzylation using TMSCl and Et₃SiH

Entry	R-OH	PhCHO	TMSCl	R-OBn	Yield/%
1			1 eq		8
2			5 eq		83
3			10 eq		quant
4			10 eq		81%
5			10 eq		77%
6			10 eq		70%
7			10 eq		52%

PNB-OR = OR PBB-OR = OR

The *p*-nitrobenzyl group has been so far introduced to hydroxy functions by using *p*-nitrobenzylbromide-Ag₂O, *p*-nitrobenzylbromide-AgOTf-2,4,6-collidine, or *p*-nitrobenzyl triflate or by Nishizawa's method. The hydroxy group of **3** was conveniently *p*-nitrobenzylated by the present method in a comparable yield (Entry 6). *p*-Bromobenzylated of **3** by the present method did not give good yield, because of debromination under the reducing condition of Et₃SiH (Entry 7). This PBB (*p*-bromobenzyl) ethers of **7** should be converted to the more labile, substituted *p*-arylbenzyl ethers via a Pd-catalyzed Suzuki–Miyaura coupling without using phosphine ligands and be subsequently removed by acids or oxidants.⁹

p-Methoxybenzyl group was completely cleaved under the present reaction conditions. We previously reported that *p*-acylamino benzyl group was much more stable to Lewis acids than *p*-methoxybenzyl group but was cleaved by DDQ in a comparable rate.^{4d,10} *p*-Acylamino benzyl group, however, also smoothly cleaved with TMSCl–Et₃SiH. In other words, a combination of TMSCl–Et₃SiH is useful for selective cleavage of *p*-methoxybenzyl and *p*-acylamino benzyl groups.¹¹

In conclusion, we have developed a new convenient method for the preparation of ethers from carbonyl compounds and non-protected alcohols under non-basic and non-metal conditions. TMSCl works as both an acid catalyst and dehydrating agent. Although excess TMSCl was required for the foregoing reactions, TMSCl could be readily removed by simple evaporation.

The present work was financially supported in part by “Research for the Future” Program No. 97L00502 from the Japan Society for the Promotion of Science, by Special Coordination Funds of the Science and Technology Agency of the Japanese Government, and by Grants-in-Aid for Scientific Research No. 09044086 and No. 12640571 from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

References and Notes

- 1 M. A. Brook and T.-H. Chan, *Synthesis*, **1983**, 201.
- 2 T.-H. Chan, M. A. Brook, and T. Chaly, *Synthesis*, **1983**, 203. Esterification was basically effected by using 2.2 equiv. of TMSCl and alcohols as solvents at room temperature or under reflux in THF. We found that esterification of *N*-fluorenylmethoxycarbonyl (Fmoc) amino acids also proceeded smoothly in CH₂Cl₂ at room temperature by using 2 equiv. of alcohols and 5 equiv. of TMSCl against carboxylic acids to give the desired esters quantitatively.
- 3 M. Izumi, K. Fukase, and S. Kusumoto, *Biosci. Biotechnol. Biochem.*, **66**, 211 (2002).
- 4 a) S. Hatakeyama, H. Mori, K. Kitano, H. Yamada, and M. Nishizawa, *Tetrahedron Lett.*, **35**, 4367 (1994). b) Trityl perchlorate promoted reduction of carbonyl compounds with triethylsilane: J. Kato, N. Iwasawa, and T. Mukaiyama, *Chem. Lett.*, **1985**, 743. c) Trimethylsilyl iodide catalyzed reductive coupling of carbonyl compounds with trialkylsilanes: M. B. Sassaman, K. Kotian, G. K. S. Prakash, and G. A. Olah, *J. Org. Chem.*, **52**, 4314 (1987). d) Reductive benzylation via *O*-tetrahydropyranyl ether: T. Suzuki, K. Ohashi, and T. Oriyama, *Synthesis*, **1999**, 1561. e) Bismuth trichloride catalyzed efficient reductive etherification of carbonyl compounds with alcohols: M. Wada, S. Nagayama, K. Mizutani, R. Hiroi, and N. Miyoshi, *Chem. Lett.*, **2002**, 248.
- 5 Reductive *N*-benzylation has also reported. a) S.-C. Shim, K.-T. Huh, S.-S. Oh, and D.-H. Oh, *Bull. Korean Chem. Soc.*, **7**, 484 (1986). b) R. B. C. Pillai, *Indian J. Chem. Sect. B*, **35B**, 487 (1996).
- 6 K. Fukase, Y. Fukase, M. Oikawa, W.-C. Liu, Y. Suda, and S. Kusumoto, *Tetrahedron*, **54**, 4033 (1998).
- 7 A typical procedure for reductive benzylation was as follows: To a solution of methyl 2,3,4-*O*-tribenzyl- α -D-glucopyranoside (**3**, 232 mg, 0.5 mmol) in CH₂Cl₂ (2.0 mL) was added chlorotrimethylsilane (0.63 mL, 5.0 mmol) and benzaldehyde (0.05 mL, 0.5 mmol) and triethylsilane (0.16 mL, 1.0 mmol). The mixture stirred at room temperature for 3 h, and evaporated under reduced pressure. The residue was submitted to flash chromatography (silica gel, eluent: CHCl₃/MeOH, 10:1, v/v), to give a pure colorless product **5**. Yield 224 mg (81%). **5**: ¹H NMR (CDCl₃) δ 3.37 (3H, s, OCH₃), 3.55 (1H, dd, *J* = 9.5, 3.6 Hz, H-2), 3.59–3.65 (2H, m, H-4 and H-5), 3.69–3.75 (2H, m, H-6), 3.97 (1H, dd, *J* = 9.5, 9.2 Hz, H-3), 4.45–4.98 (8H, m, PhCH₂-x4), 4.61 (1H, d, *J* = 3.6 Hz, H-1), 7.11–7.37 (20H, m, PhCH₂-x4). **6**: ¹H NMR (CDCl₃) δ 3.37 (3H, s, OCH₃), 3.55 (1H, dd, *J* = 9.5, 3.6 Hz, H-2), 3.60–3.64 (2H, m, H-4 and H-5), 3.69–3.75 (2H, m, H-6), 3.97 (1H, dd, *J* = 9.5, 9.2 Hz, H-3), 4.45–4.98 (8H, m, PhCH₂-x4), 4.62 (1H, d, *J* = 3.6 Hz, H-1), 7.12–7.37 (20H, m, PhCH₂-x3), 7.63 (2H, d, *J* = 9.0 Hz, NO₂PhCH₂-), 8.27 (2H, d, *J* = 9.0 Hz, NO₂PhCH₂-). **7**: ¹H NMR (CDCl₃) δ 3.35 (3H, s, OCH₃), 3.54 (1H, dd, *J* = 9.5, 3.6 Hz, H-2), 3.59–3.64 (2H, m, H-4 and H-5), 3.69–3.74 (2H, m, H-6), 3.97 (1H, dd, *J* = 9.5, 9.2 Hz, H-3), 4.44–4.98 (8H, m, PhCH₂-x4), 4.61 (1H, d, *J* = 3.6 Hz, H-1), 7.12–7.37 (20H, m, PhCH₂-x3), 7.25 (2H, d, *J* = 9.0 Hz, BrPhCH₂-), 7.51 (2H, d, *J* = 9.0 Hz, BrPhCH₂-).
- 8 a) K. Fukase, H. Tanaka, S. Torii, and S. Kusumoto, *Tetrahedron Lett.*, **31**, 389 (1990). b) K. Fukase, T. Matsumoto, N. Ito, T. Yoshimura, S. Kotani, and S. Kusumoto, *Bull. Chem. Soc. Jpn.*, **65**, 2643 (1992). c) K. Fukase, T. Yoshimura, S. Kotani, and S. Kusumoto, *Bull. Chem. Soc. Jpn.*, **67**, 473 (1994).
- 9 X.-Y. Liu and P. H. Seeberger, *Chem. Commun.*, **2004**, 1708.
- 10 K. Fukase, T. Yoshimura, M. Hashida, and S. Kusumoto, *Tetrahedron Lett.*, **32**, 4019 (1991).
- 11 Reductive cleavage of 4-methoxybenzyl group with Et₃SiH-BF₃·Et₂O: Y. Morimoto, M. Iwahashi, K. Nishida, Y. Hayashi, and H. Shirahama, *Angew. Chem., Int. Ed. Engl.*, **35**, 904 (1996).