

Expanding the Genetic Code for Biological Studies

Qian Wang,¹ Angela R. Parrish,¹ and Lei Wang^{1,*}

¹The Jack H. Skirball Center for Chemical Biology and Proteomics, The Salk Institute for Biological Studies, 10010 North Torrey Pines Road, La Jolla, CA 92037

*Correspondence: lwang@salk.edu

DOI 10.1016/j.chembiol.2009.03.001

Using an orthogonal tRNA-synthetase pair, unnatural amino acids can be genetically encoded with high efficiency and fidelity, and over 40 unnatural amino acids have been site-specifically incorporated into proteins in *Escherichia coli*, yeast, or mammalian cells. Novel chemical or physical properties embodied in these amino acids enable new means for tailored manipulation of proteins. This review summarizes the methodology and recent progress in expanding this technology to eukaryotic cells. Applications of genetically encoded unnatural amino acids are highlighted with reports on labeling and modifying proteins, probing protein structure and function, identifying and regulating protein activity, and generating proteins with new properties. Genetic incorporation of unnatural amino acids provides a powerful method for investigating a wide variety of biological processes both *in vitro* and *in vivo*.

Introduction

The genetic code consists of 64 triplet codons specifying 20 canonical amino acids and 3 stop signals. This code is preserved in all three kingdoms of life with few reassignments (Knight et al., 2001). Although various amino acid derivatives have been identified in natural proteins, most of them are post-translational modifications of the canonical amino acids (Uy and Wold, 1977). Two additional amino acids, selenocysteine (Bock et al., 1991) and pyrrolysine (Srinivasan et al., 2002), are also found to be incorporated into proteins cotranslationally and are thus regarded as natural expansions of the genetic code (Ambrogelly et al., 2006). With these amino acids and with the help of posttranslational modifications and cofactors, proteins are able to carry out various functions for sustaining life.

Given the important roles of proteins in biology, it is desirable to be able to manipulate proteins at will for understanding protein structure-function relationships, investigating protein-involved biological processes, and generating proteins and organisms with new properties. Therefore, tremendous efforts have been made regarding the incorporation of unnatural amino acids into proteins to introduce new functional groups apart from those found in the canonical amino acids, through chemical or biosynthetic means (Wang and Schultz, 2004). Chemical methods use chemistry to directly modify or prepare proteins. Early efforts focused on derivatizing reactive side chains of the protein, such as the thiol group of cysteine and the ϵ -amino group of lysine (Kaiser et al., 1985). The advance of stepwise solid-phase peptide synthesis makes possible the complete synthesis of peptides and small proteins (usually < 100 residues) containing any synthetically accessible unnatural amino acid (Kent, 1988). The protein size limitation is subsequently overcome by semi-synthetic protein ligation methods, in which two or more peptide fragments are chemically ligated to make the full-length protein (Dawson et al., 1994; Muir, 2003). By contrast, biosynthetic methods harness the endogenous cellular machinery to translate the target protein, in which the unnatural amino acid is incorporated cotranslationally. Bacterial strains auxotrophic for a particular amino acid have been used to globally replace the amino acid with a close structural analog (Cohen and Cowie,

1957; Minks et al., 2000; Link et al., 2003). For the site-specific incorporation of unnatural amino acids, a suppressor tRNA is chemically acylated with an unnatural amino acid. When added into cell extracts or *Xenopus* oocytes that support protein translation, the acylated tRNA incorporates the attached unnatural amino acid in response to a stop or extended codon (Noren et al., 1989; Bain et al., 1989; Nowak et al., 1995). All of these methods have proved to be extremely valuable, but each has limitations in the lack of site selectivity, the *in vitro* nature of the method, or the low incorporation efficiency (Wang and Schultz, 2004).

The ability to genetically encode unnatural amino acids in the same manner as the canonical amino acids would enable specific changes to be precisely made in proteins directly *in vivo*, thus providing novel tools for understanding biology in molecular terms in the native settings. Here, we review the methodology for genetically incorporating unnatural amino acids into proteins in live cells, and focus on recent advances in expanding the method to eukaryotic cells. Representative applications are discussed to illustrate the utility and scope of this technology in studying various biological questions.

Methodology

Similar to canonical amino acids, the genetic encoding of an unnatural amino acid requires a dedicated set of components including a tRNA, a codon, and an aminoacyl-tRNA synthetase (hereafter referred to as synthetase) (Figure 1) (Wang and Schultz, 2002). This tRNA/codon/synthetase set, termed the orthogonal set, must not crosstalk with endogenous tRNA/codon/synthetase sets and must be functionally compatible with other components of the translation apparatus. Specifically, the orthogonal tRNA should not be recognized by any endogenous synthetase, and it should decode the orthogonal codon, which is not assigned to any canonical amino acid. The orthogonal synthetase should not charge any endogenous tRNA but the orthogonal tRNA, and it should charge the orthogonal tRNA with the unnatural amino acid only. The unnatural amino acid cannot be a substrate for any endogenous synthetase and must be available in the cytoplasm of the host cell. Once expressed in cells, the orthogonal synthetase charges the




Figure 1. A General Method for Genetically Encoding Unnatural Amino Acids in Live Cells

orthogonal tRNA with the desired unnatural amino acid, and the acylated tRNA incorporates the attached unnatural amino acid into proteins in response to the unique codon by utilizing the endogenous translational machinery. As all elements are genetically encodable, this method has the potential to be applied to almost all genetically tractable organisms.

Method Development in *Escherichia coli*

Escherichia coli was initially chosen as the host organism to develop this general method for genetically encoding unnatural amino acids in vivo, because its translational machinery had been extensively studied and genetic manipulation of *E. coli* is efficient and relatively straightforward. A unique codon is required to specify the unnatural amino acid, and the amber stop codon has been chosen for this purpose. There are three stop codons that do not code for amino acids but function as a signal for the termination of translation. By reassigning a “blank” stop codon to an unnatural amino acid any overlap and ambiguity with sense codons is avoided. The amber codon became the “blank” codon of choice in this approach because (i) it is the least used among the three stop codons in *E. coli*, and because (ii) amber suppressor tRNAs have been identified or engineered in *E. coli* to readthrough the amber codon with canonical amino acids, and they do not significantly affect the growth rates of *E. coli* cells (Normanly et al., 1990). Therefore, specifying an unnatural amino acid with the amber codon was expected to minimize perturbations to the genetic code and to the health of the host.

Endogenous tRNAs and synthetases recognize their cognate partners through delicate interactions between various regions

(Ibba and Soll, 2000). As there is already a pool of tRNA/synthetase pairs inside cells, it is challenging to generate a new tRNA/synthetase pair orthogonal to existing ones and functional in translation. A successful strategy is to import a tRNA/synthetase pair from a species in a different kingdom of life, followed by fine tuning with selections. This strategy is based on early reports showing that in vitro cross-species aminoacylation is often low (Kwok and Wong, 1980). The first orthogonal tRNA/synthetase pair in *E. coli* generated from archaeal bacteria was derived from the tRNA^{Tyr}/tyrosyl-tRNA synthetase (TyrRS) from *Methanococcus jannaschii* (Wang et al., 2000). The anticodon of the tRNA is mutated to CUA for recognizing the amber codon UAG. This *M. jannaschii* tRNA^{Tyr}_{CUA} (*Mj*tRNA^{Tyr}_{CUA}) and its cognate *Mj*TyrRS function efficiently in translation in *E. coli*, but there is a small degree of aminoacylation of this *Mj*tRNA^{Tyr}_{CUA} by endogenous *E. coli* synthetases. To make a completely orthogonal tRNA while maintaining its affinity for the cognate synthetase, a general strategy has been developed to evolve orthogonal tRNAs in *E. coli* (Figure 2A) (Wang and Schultz, 2001). A library of tRNA mutants is prepared by randomizing specific nucleotides, and the orthogonal tRNA is selected by using a combination of negative and positive selection. The negative selection removes nonorthogonal tRNAs from the library, and the subsequent positive selection removes nonfunctional tRNAs but allows only orthogonal tRNAs with high affinity for the cognate synthetase to pass. This approach was applied to the *Mj*tRNA^{Tyr}_{CUA}, and the mutant tRNA^{Tyr}_{CUA} that is orthogonal in *E. coli* and functions efficiently with *Mj*TyrRS to translate the amber codon was identified.




Figure 2. Double-Sieve Selection with a Mutant Library for Evolving Orthogonal tRNAs or Unnatural Amino Acid-Specific Synthetases

(A) Evolving orthogonal tRNAs in *E. coli*. The negative selection uses a toxic barnase gene containing two TAG stop codons at permissive sites. tRNAs nonorthogonal to *E. coli* synthetases will be charged with amino acids and suppress the TAG codons to produce a functional barnase, which kills the cell. The positive selection uses the β -lactamase gene containing a TAG codon at a permissive site. tRNAs charged by the cognate synthetase and functional in translation will suppress the TAG codon and express active β -lactamase, which enables cells to survive in ampicillin. Cells containing nonfunctional tRNAs will be killed by ampicillin.

(B) Evolving orthogonal synthetases to be specific for an unnatural amino acid in *E. coli*. The positive selection uses the chloramphenicol acetyl transferase (CAT) gene containing a permissive TAG stop codon in the presence of the unnatural amino acid. Mutant synthetases capable of charging the orthogonal $\text{mutRNA}_{\text{CUA}}^{\text{Tyr}}$ with the unnatural or any natural amino acid will express full-length CAT, enabling cells to survive in chloramphenicol (Cm). The negative selection uses the barnase gene containing permissive TAG codons. Synthetase mutants active toward natural amino acids will charge the $\text{mutRNA}_{\text{CUA}}^{\text{Tyr}}$ to suppress TAG codons in the barnase gene, leading to cell death. In the absence of the unnatural amino acid, synthetase mutants specific for the unnatural amino acid will survive the negative selection. To improve activity and specificity, initial synthetase gene hits can be recombined by using DNA shuffling and are subject to the next round of selection.

are then subjected to a negative selection, in which synthetase mutants charging a common amino acid are eliminated. In the end, only synthetase mutants specific for the unnatural amino acid will survive both selections. Using this approach, a mutant synthetase specific for the unnatural amino acid *o*-methyltyrosine (**1**, Figure 3) was identified from the *MjTyrRS* library. Together with the evolved orthogonal $\text{mutRNA}_{\text{CUA}}^{\text{Tyr}}$ described above, it enables the site-specific incorporation of *o*-methyltyrosine into proteins in response to the

The next step is to change the substrate specificity of the orthogonal *MjTyrRS* from tyrosine to a desired unnatural amino acid, which is also achieved by using two-step selections (Figure 2B). A mutant synthetase library ($>10^9$ members) is generated by randomizing five amino acid residues in the active site of the *MjTyrRS*. The selection of the residues to be randomized was enabled by the availability of the *Bacillus stearothermophilus* TyrRS crystal structure (Brick et al., 1989). The library is first positively selected in the presence of the unnatural amino acid, in which active synthetase mutants able to charge the orthogonal $\text{mutRNA}_{\text{CUA}}^{\text{Tyr}}$ with the unnatural amino acid or any common amino acid are identified. The active synthetases

are then subjected to a negative selection, in which synthetase mutants charging a common amino acid are eliminated. In the end, only synthetase mutants specific for the unnatural amino acid will survive both selections. Using this approach, a mutant synthetase specific for the unnatural amino acid *o*-methyltyrosine (**1**, Figure 3) was identified from the *MjTyrRS* library. Together with the evolved orthogonal $\text{mutRNA}_{\text{CUA}}^{\text{Tyr}}$ described above, it enables the site-specific incorporation of *o*-methyltyrosine into proteins in response to the

amber codon in *E. coli*. Western blot and mass spectrometric analyses of the expressed protein indicate that the translational fidelity for *o*-methyltyrosine is close to that of common amino acids (Wang et al., 2001). This work represents the first expansion of the genetic code to include an unnatural amino acid by using engineering approaches. Structurally similar to tyrosine and phenylalanine, *o*-methyltyrosine provides an excellent case to demonstrate the high translational fidelity that can be achieved by using this approach. Meanwhile, it was important to test whether this method could be generally used to genetically encode various unnatural amino acids, especially those significantly deviating




Figure 3. Structures for Unnatural Amino Acids Discussed in the Text

that it is possible to artificially alter the genetic code in eukaryotes as well. The general strategy developed to genetically encode unnatural amino acids in *E. coli* should also be applicable to eukaryotic organisms. A number of tRNA/synthetase pairs derived from *E. coli* have been shown to be orthogonal in eukaryotic cells (Drabkin et al., 1996; Kowal et al., 2001), again due to the low crosskingdom aminoacylation. In particular, the *E. coli* tRNA^{Tyr}_{CUA}/TyrRS pair is orthogonal in yeast (Edwards and Schimmel, 1990; Kwok and Wong, 1980). Moreover, the success in evolving the *Mj*TyrRS to incorporate a wide range of unnatural amino acids in *E. coli* guided use of the *E. coli* tRNA^{Tyr}_{CUA}/TyrRS pair in yeast.

To evolve synthetases specific for unnatural amino acids in yeast, a general selection scheme analogous to that used in *E. coli* was set up, which also uses positive and negative selections with a mutant synthetase library. A synthetase library (10^8 in size) is similarly constructed by randomizing five active site residues in the *E. coli* TyrRS that correspond to the five residues randomized in the *Mj*TyrRS

(Wang et al., 2001). Mutant synthetases are identified after several rounds of selection that, together with the *E. coli* tRNA^{Tyr}_{CUA}, incorporate a number of unnatural amino acids into proteins in yeast, albeit with rather low protein yields (~0.05 mg/L) (Chin et al., 2003).

Low incorporation efficiency results in reduced yield of the full-length protein containing the unnatural amino acid and in an increase of truncated protein products terminated at the introduced amber codon, thus creating difficulties with employing genetically encoded unnatural amino acids in yeast. To address this problem, new methods have been developed to efficiently express orthogonal prokaryotic tRNAs and to improve the target mRNA stability in yeast. Transcription and processing of tRNAs are different in prokaryotes and eukaryotes. A major distinction is that *E. coli* tRNAs are transcribed through promoters upstream of the tRNA gene, whereas eukaryotic tRNAs are transcribed through promoter elements within the tRNA known as the A- and B-box (Figure 4A) (Galli et al., 1981). The A- and B-box identity elements are conserved among eukaryotic tRNAs but are absent in many *E. coli* tRNAs. Creating the consensus A- and B-box sequences in *E. coli* tRNAs through mutation could cripple the tRNA (Sakamoto et al., 2002). The *E. coli* tRNA^{Tyr}_{CUA} has the B-box but no fully matched A-box. In the previously described system (Chin et al., 2003), only the tRNA structural gene was inserted in a plasmid, resulting in a basal level of tRNA expression likely driven by a cryptic promoter elsewhere (Chen et al., 2007).

from common amino acids in structure. Subsequent experiments showed that L-3-(2-naphthyl)alanine (2), which represents a significant structural perturbation relative to tyrosine, can also be genetically incorporated into proteins in *E. coli* with high efficiency and fidelity (Wang et al., 2002). In the following years, the principle and methods developed herein have been used to genetically incorporate more than 40 different unnatural amino acids in *E. coli* (Wang et al., 2006b). For more than 80% of unnatural amino acids incorporated, the orthogonal tRNA is the same mutRNA^{Tyr}_{CUA}, and the orthogonal synthetases are all evolved from the *Mj*TyrRS (Wang et al., 2001), suggesting the great flexibility of the active site of TyrRS to be modified to accommodate different substrates. Suppression efficiencies of most unnatural amino acids range from 25% to 75% of the wild-type protein, and translational fidelity is >99%. Yields of unnatural amino acid-containing proteins are in the range of several milligrams to tens of milligrams per liter of cell culture. Additional orthogonal tRNA/synthetase pairs have since been generated (Anderson and Schultz, 2003), and four-base codons have also been used to encode unnatural amino acids in *E. coli* (Anderson et al., 2004). In addition, a biosynthetic pathway for *p*-aminophenylalanine was created in *E. coli* so that the strain is able to produce and genetically incorporate *p*-aminophenylalanine autonomously (Mehl et al., 2003).

Transition to Yeast

There are examples of eukaryotic organisms having natural changes in their genetic codes (Knight et al., 2001), suggesting




Figure 4. General Methods for Efficient Expression of Prokaryotic tRNAs in Eukaryotic Cells by Using Special Pol III Promoters
(A) Gene elements for tRNA transcription in *E. coli* and eukaryotic cells.
(B) Expression of prokaryotic tRNAs in yeast by using an external SNR52 or RPR1 Pol III promoter, which contains the consensus A- and B-box sequences and is posttranscriptionally cleaved from the primary transcript.
(C) A type-3 Pol III promoter, such as the H1 promoter, with a defined transcription initiation site is used to express prokaryotic tRNAs in mammalian cells.

One method for increasing tRNA expression is to use the 5'- and 3'-flanking sequences of an endogenous yeast suppressor, tRNA^{Tyr}_{CUA} (SUP4) (Francis and RajBhandary, 1990; Drabkin et al., 1996; Kowal et al., 2001; Chen et al., 2007). The *E. coli* tRNA^{Tyr}_{CUA} without the -CCA trinucleotide is flanked by these sequences, and this cassette is repeated in tandem 3–6 times. A strong PGK1 RNA Polymerase II promoter is additionally placed upstream of the tandem arrangements (Chen et al., 2007). This combination gives an overall > 50 fold increase in tRNA levels compared to the previous tRNA alone scheme. Together with optimized expression of the synthetase, this approach improves the yield of unnatural amino acid-containing proteins to 6–8 mg/L yeast cells. A new method to efficiently express prokaryotic tRNAs in yeast involves an external promoter containing the consensus eukaryotic A- and B-box sequences (Figure 4B) (Wang and Wang, 2008). The promoter is placed upstream of the *E. coli* tRNA (without 3'-CCA) to drive the transcription of a primary RNA consisting of the promoter and the tRNA. The promoter is cleaved posttranscriptionally to yield the mature

tRNA. Two yeast Pol III promoters, the RPR1 promoter and the SNR52 promoter, have been shown to efficiently drive the expression of the *E. coli* tRNA^{Tyr}_{CUA} in yeast. The expressed *E. coli* tRNA^{Tyr}_{CUA} is 6- to 9-fold more active in translation than the same tRNA transcribed by using the SUP4 5'-flanking sequence. Interestingly, the increased activity is not a result of a higher tRNA transcription level, suggesting the importance of proper tRNA processing. The cleavage of the SNR52 or RPR1 promoter from the primary RNA transcript could directly generate the correct 5' end of the tRNA. This external promoter method has also been used to express *E. coli* tRNA^{Leu}_{CUA} in yeast, which functions efficiently in translation as well, suggesting that it can be a general method for functionally expressing different prokaryotic tRNAs in yeast.

An important factor for unnatural amino acid incorporation in eukaryotes is the stability of the target mRNA. Eukaryotic cells have an mRNA surveillance mechanism, nonsense-mediated mRNA decay (NMD), to identify mRNAs containing premature stop codons and target the mRNA for rapid degradation (Amrani et al., 2006). The amber stop codon is thus far the most frequently used codon to encode unnatural amino acids, but NMD could shorten the target mRNA lifetime, resulting in a lower protein yield. Inactivation of NMD would preserve the stability of the UAG-containing mRNA and thus enhance the incorporation efficiency. An NMD-deficient yeast strain was generated by knocking out the UPF1 gene, an essential component for NMD. The unnatural amino acid incorporation efficiency was indeed increased more than 2-fold in the *upf1Δ* strain compared to the wild-type yeast (Wang and Wang, 2008). NMD in yeast shows a polar effect of nonsense codon positions (Cao and Parker, 2003). The steady-state mRNA level is reduced by NMD more significantly when the nonsense codon is closer to the 5' end than to the 3' end of an mRNA. Consistently, the increase of unnatural amino acid incorporation efficiency in the *upf1Δ* strain correlates with the position of the UAG codon: more than a 2-fold increase is measured when the amber codon is within the N-terminal two thirds of the gene, whereas no significant increase is detected when it is within the C-terminal fourth of the coding region (Q.W. and L.W., unpublished data). By using the external SNR52 promoter and the NMD-deficient *upf1Δ* strain, the overall yields of purified unnatural amino acid-containing proteins have reached ~15 mg/L yeast cells, ~300-fold higher than the previous system and comparable to the yield in *E. coli*.

Transition to Mammalian Cells

Genetic incorporation of unnatural amino acids into proteins in mammalian cells face the same challenges as in yeast. Specifically, a major challenge is how to efficiently express prokaryotic orthogonal tRNAs and to ensure they are translationally competent in mammalian cells, due to the aforementioned difference in tRNA transcription and processing. In addition, another major challenge is how to evolve mutant orthogonal synthetases to be specific for an unnatural amino acid for use in mammalian cells. The latter arises because the transfection efficiency of mammalian cells is much lower than the transformation efficiency of *E. coli* or yeast, making it impractical to generate large synthetase mutant libraries inside mammalian cells. Moreover, survival-death selection in mammalian cells is not as effective as in *E. coli* and yeast. Therefore, it is difficult to evolve a specific

synthetase out of a large mutant synthetase library directly in mammalian cells.

The *E. coli* tRNA^{Tyr}_{CUA}/TyrRS pair is orthogonal in mammalian cells, but the *E. coli* tRNA^{Tyr}_{CUA} is not easily expressed in mammalian cells, presumably due to the lack of a matched A-box. Fortunately, the *B. stearotherophilus* tRNA^{Tyr}_{CUA} happens to contain consensus A- and B-box sequences, is not charged by mammalian synthetases, and works with the heterologous *E. coli* TyrRS in suppressing the TAG codon. The *B. stearotherophilus* tRNA^{Tyr}_{CUA} (lacking 3'-CCA) was linked to the 5'-flanking sequence of the human tRNA^{Tyr}, and nine tandem repeats of this gene arrangement were used for tRNA expression (Sakamoto et al., 2002). A small collection of designed active-site variants of the *E. coli* TyrRS was screened in vitro, and a mutant synthetase using 3-iodotyrosine (**3**) more effectively than Tyr is identified. This mutant synthetase is used with nine tandem *B. stearotherophilus* tRNA^{Tyr}_{CUA} to incorporate 3-iodotyrosine into proteins in Chinese hamster ovary (CHO) cells and HEK293 cells with ~95% fidelity (Sakamoto et al., 2002). In another attempt, mutations were introduced into the A-box region of the *B. subtilis* tryptophan opal suppressor tRNA (tRNA^{Trp}_{UCA}) to create a consensus A-box (Zhang et al., 2004a). The 3'-CCA-lacking mutant tRNA was expressed by using the 5'- and 3'-flanking sequences from the *Arabidopsis* tRNA^{Trp}. Together with a rationally designed *B. subtilis* TrpRS mutant, this tRNA incorporates 5-hydroxytryptophan (**4**) into the foldon protein in HEK293 cells in response to the opal stop codon with ~97% fidelity.

Encouraging as these results are, it remained a challenge to incorporate various unnatural amino acids in mammalian cells with high efficiency and fidelity. The above-described approaches may not be generally applicable to other tRNA/synthetase pairs or other unnatural amino acids. As for tRNA expression, most bacterial tRNAs do not have the consensus eukaryotic A- and B-box sequences, and heterologous pairing of the tRNA with synthetase often results in low activity. Since the A- and B-box lie at nucleotides involved in tertiary interactions that support the L-shaped tRNA structure, mutations to create such consensus sequences, although they may not dramatically decrease the activity of the *B. subtilis* tRNA^{Trp}_{UCA}, greatly impair the suppression ability of the *E. coli* tRNA^{Tyr}_{CUA} (Sakamoto et al., 2002). In addition, using tandem repeats to boost the tRNA expression level may eventually lead to lower expression, especially in stable cell lines, as repeating sequences are frequently associated with gene silencing (Hsieh and Fire, 2000).

A new method has been developed that involves the use of an external type-3 Pol III promoter to efficiently express functional prokaryotic tRNAs in mammalian cells (Figure 4C) (Wang et al., 2007b). A type-3 Pol III promoter has its promoter elements exclusively upstream of the coding sequence and does not require any intragenic elements (such as the A- and B-box) for transcription (Willis, 1993). Therefore, tRNAs with or without the consensus A- and B-box sequences are both expected to be transcribed by this type of promoter in mammalian cells. In addition, the transcription initiation site of some type-3 Pol III promoters, such as the H1 promoter (Myslinski et al., 2001) and the U6 small-nuclear RNA promoter (Paule and White, 2000), is well defined, which could be used to generate the correct 5' end of the tRNA without further posttranscriptional processing. Using a fluorescence-based translation assay,

tRNA expression elements including the promoter, the 5'- and 3'-flanking sequences, and the 3'-CCA trinucleotide are linked to the *E. coli* tRNA^{Tyr}_{CUA} in various combinations and are systematically tested for their ability to generate functional tRNAs in mammalian cells. The gene arrangement that yields tRNAs with the highest translational activity is the H1 promoter followed by the *E. coli* tRNA^{Tyr}_{CUA} (without 3'-CCA) and then by the 3'-flanking sequence of tRNA^{fMet}. *E. coli* tRNA^{Tyr}_{CUA} expressed by using this method is ~70-fold more active in translation than the one expressed with the 5'- and 3'-flanking sequences of human tRNA^{fMet}. This method has also been used to express different *E. coli* tRNAs successfully, and it is effective in various mammalian cells such as HEK293 cells, HeLa cells, and primary neurons.

For generating unnatural amino acid-specific synthetase mutants, it is difficult to predict a priori which active site residues need to be mutated. More often, mutations at multiple sites are required to achieve high substrate specificity, suggesting that small collections of synthetase mutants will likely fall short. Mutants thus generated may still recognize common amino acids, as is the case with the synthetase used to incorporate 3-iodotyrosine.

To circumvent the challenge of evolving unnatural amino acid-specific synthetases directly in mammalian cells, a transfer strategy is used. Since the *E. coli* tRNA^{Tyr}_{CUA}/TyrRS pair is orthogonal in mammalian cells as well as in yeast, and the translational machinery of yeast is homologous to that of higher eukaryotes, it should be possible to evolve the specificity of this synthetase in yeast and transfer the optimized tRNA/synthetase pairs for use in mammalian cells. In one report, mutant synthetases that were evolved in yeast from the *E. coli* TyrRS specific for different unnatural amino acids (Deiters et al., 2003; Chin et al., 2003), together with the *B. stearotherophilus* tRNA^{Tyr}_{CUA}, have been successfully used in CHO cells and HEK293 cells for incorporating the corresponding unnatural amino acids (Liu et al., 2007). A high translational fidelity for the unnatural amino acid is confirmed with mass spectrometric analysis, and proteins could be produced up to 1 μg per 2 × 10⁷ cells. In another report, mutant synthetases evolved in yeast from the *E. coli* TyrRS are used together with the cognate *E. coli* tRNA^{Tyr}_{CUA} expressed by the H1 promoter in mammalian cells (Wang et al., 2007b). Because the H1 promoter also enables the functional expression of the *E. coli* tRNA^{Leu}_{CUA}, mutant synthetases evolved in yeast from the *E. coli* LeuRS (Summerer et al., 2006) are also successfully transferred in mammalian cells to incorporate the fluorescent unnatural amino acid dansylalanine (Wang et al., 2007b). Incorporation efficiencies are in the range of 13%–41%, depending on the activity of the evolved synthetase. The incorporation fidelity is also high, as measured by a sensitive fluorescence assay based on the translation of GFP. Unnatural amino acids have been incorporated in HEK293 cells, HeLa cells, and even in primary neurons.

Applications

New chemical, physical, and biological properties can be introduced into proteins via the incorporation of unnatural amino acids, and thus unnatural amino acids have been exploited to study a wide range of biological problems involving proteins. In particular, the stability, specificity, and catalytic properties of multiple proteins have been extensively studied with unnatural amino acids incorporated with chemically acylated tRNAs,

revealing fundamental principles of protein structure and function (Cornish et al., 1995). Microinjection of chemically acylated tRNAs into *Xenopus* oocytes enables studies of integral membrane proteins with unnatural amino acids. Various ion channels and neurotransmitter receptors have been probed with unnatural amino acids by using electrophysiological techniques, yielding novel insights into the function role of conserved amino acid residues and the gating mechanism of this important family of proteins (Beene et al., 2003; Lummis, et al., 2005). Recent years have also seen elegant studies of signal transduction, selectivity of ion channels, and chromatin biology by using unnatural amino acids that are introduced with native chemical ligation or expressed protein ligation (Hahn et al., 2007; Valiyaveetil et al., 2006; McGinty et al., 2008). There are already many excellent reviews on different aspects of this subject (Cornish et al., 1995; Hohsaka and Sisido, 2002; Beene et al., 2003; Link et al., 2003; Wang and Schultz, 2004; Schwarzer and Cole, 2005; Pellois and Muir, 2006), so here we will focus on the applications of unnatural amino acids that are genetically encoded by using the orthogonal tRNA/synthetase method. Representative examples are summarized to illustrate the usefulness of this technology in modifying and investigating proteins, regulating protein functions, and generating new protein properties. We note that some of the unnatural amino acids discussed below have also been incorporated into proteins by using other approaches, details of which can be found in the reviews cited above.

Labeling and Modifying Proteins

Reactive side chains of canonical amino acids such as cysteine and lysine have long been used for the selective chemical modification of proteins, but the intrinsic selectivity is low unless the amino acid is absent or can be mutated in the target protein. A nonproteinogenic chemical group embodied by an unnatural amino acid can serve as a unique chemical handle for bio-orthogonal chemical reactions, through which a variety of reagents can be selectively appended to proteins in vitro. These reagents can be biophysical probes, tags, posttranslational modifications, and groups modifying protein stability or activity. Currently, site-specific labeling of a target protein via unnatural amino acids directly inside live cells is still challenging.

The keto group is not present in any of the canonical amino acids, and it reacts with hydrazides, alkoxyamines, and semicarbazides under aqueous, mild conditions to respectively produce hydrazone, oxime, and semicarbazone linkages that are stable under physiological conditions. The keto group has been genetically encoded in *E. coli* in the form of *p*-acetylphenylalanine (**5**) and is specifically labeled in vitro with fluorescein hydrazide and biotin hydrazide with greater than 90% yield (Wang et al., 2003). Similarly, fluorescent dyes are selectively appended to the *m*-acetylphenylalanine (**6**) introduced into the membrane protein LamB in *E. coli* (Zhang et al., 2003). *p*-Acetylphenylalanine and *p*-benzoylphenylalanine (**21**) have also been incorporated into rhodopsin, a transmembrane G protein-coupled receptor, in mammalian cells (Ye et al., 2008). The purified mutant rhodopsin is labeled in vitro with fluorescein hydrazide.

Natural glycoproteins are often present as a population of different glycoforms, which complicates glycan structure analysis and the study of glycosylation effects on protein structure and function. Generation of pure glycoprotein mimetics with

defined glycan structures will be valuable for the systematic understanding of glycan function and the development of improved glycoprotein therapeutics. To prepare homogenous glycoprotein mimetics, sugars are synthesized in the aminoxy form and are attached to *p*-acetylphenylalanine incorporated at defined sites in proteins (Figure 5A) (Liu et al., 2003). The attached sugar is subsequently elaborated by adding additional saccharides with glycosyltransferases to generate oligosaccharides with defined structures. Alternatively, an aminoxy-derivatized glycan can also be covalently coupled to the *p*-acetylphenylalanine-containing protein in one step.

PEGylation, the attachment of polyethylene glycol (PEG) to proteins, increases protein stability and solubility. In comparison with using common amino acids, PEGylation using a unique chemical handle of an unnatural amino acid increases the homogeneity of the final protein product and provides flexibility in choosing PEGylation sites for optimizing protein activity. PEG has been attached to *p*-acetylphenylalanine introduced in the human growth hormone to afford a protein that retained wild-type activity but had a considerably improved half-life in serum (P.G. Schultz, personal communication). This approach can be extended to other therapeutic proteins to generate purer and more stable protein drugs.

Azide and acetylene are two additional nonproteinogenic chemical groups that react with each other through a copper (I)-catalyzed [2+3] cycloaddition reaction. *p*-Azidophenylalanine (**7**) and *p*-propargyloxyphenylalanine (**8**) have been genetically incorporated into human superoxide dismutase-1 in *E. coli* and yeast (Deiters et al., 2003, 2004). Purified mutant proteins are labeled in vitro with fluorescent dyes or PEG derivatized with the complementary functional group (Figure 5B). The requirement of Cu⁺ as a catalyst may make it difficult to use this reaction inside live cells. The azide group can also react with phosphine derivatives through the Staudinger ligation. *p*-Azidophenylalanine has been incorporated into the Z-domain protein in *E. coli* or into peptides displayed on phage, and it can be labeled with fluorescein-derived phosphines (Figure 5B) (Tsao et al., 2005).

Phenylselenide undergoes oxidative elimination in hydrogen peroxide followed by a Michael addition with thiols (Figure 5C). Phenylselenocysteine (**9**) is genetically incorporated into GFP in *E. coli* (Wang et al., 2007a). After treating with hydrogen peroxide, the resultant dehydroalanine is selectively labeled in vitro with thiol-containing mannopyranose to generate an analog of glycosylated GFP, or with *n*-hexadecylmercaptan to generate a palmitoylated GFP with a nonhydrolyzable linkage. Similarly, phenylselenocysteine has been used to append analogs of methyl- or acetyl-lysine onto the purified *Xenopus* histone H3, important histone modifications that contribute to chromatin structure and accessibility (Guo et al., 2008). The H3 protein with an acetyl-lysine 9 mimic is deacetylated by a histone deacetylation complex and phosphorylated by Aurora B kinase, suggesting that such chemically labeled histones are likely functional in nucleosomes to facilitate the analysis of chromatin.

Besides using a chemical handle, changes can be directly incorporated into proteins in the form of unnatural amino acids, in particular, to mimic posttranslational modifications. These modifications are common in eukaryotic proteins playing important roles in protein stability, localization, and function, but they cannot be easily produced in bacterial cells. It is also difficult




Figure 5. Protein Labeling and Modification through Genetically Incorporated Chemical Handles

(A) A general method for preparing glycoprotein mimetics with defined glycan structure by using the keto group.

(B) Protein modification with PEG through a copper-catalyzed cycloaddition reaction (top) and with fluorescein through Staudinger ligation (bottom).

(C) Protein modification through phenylselenocysteine.

endogenous glycosyl modification in *E. coli*, genetic incorporation of glycosylated amino acids will help improve the production of proteins that require glycosylation for proper folding and function.

Phosphorylation of serine, threonine, and tyrosine by kinases and reversion by phosphatases are key switches in various signal transduction cascades. A phosphorylation mimic resistant to phosphatase activity can be used to identify the role of a specific phosphorylated residue, and to create stably activated proteins for functional assays. *p*-Carboxymethylphenylalanine (*p*CMF, **12**), a non-hydrolyzable analog of phosphotyrosine, was found to be capable of mimicking the phosphorylated state of Tyr (Xie et al., 2007b). Wild-type human STAT1, after being phosphorylated at Tyr701, forms a homodimer and strongly binds a DNA duplex. The mutant STAT1 with Tyr701 substituted with *p*CMF also binds the same DNA duplex tightly, suggesting that *p*CMF could replace phosphotyrosine in the generation of constitutively active phosphoproteins.

Acetylation is another important reversible protein modification. For

to prepare homogeneously modified proteins from eukaryotic cells. Unnatural amino acids with the properties of posttranslational modifications may allow for the generation of homogenous proteins with the desired modifications to uncover their functional roles and possibly to control protein function.

Glycosylation is a frequent modification of eukaryotic proteins. Homogeneous glycoprotein mimetics with defined glycan structures can be prepared by using the keto- or phenylselenide-containing unnatural amino acids, but it leaves a small unnatural linkage between the protein and the first sugar. This linkage does not prevent glycosyltransferases from attaching additional sugars (Liu et al., 2003), but whether it affects other glycoprotein functions awaits more research. To prepare a native glycosylated protein, mutant synthetases have been evolved that site-specifically incorporate β -*N*-acetylglucosamine-*O*-serine (**10**) or *N*-acetylgalactosamine- α -*O*-threonine (**11**) into proteins in *E. coli* (Zhang et al., 2004b; Xu et al., 2004). Since there is no

instance, acetylation of lysine residues in histone proteins controls the secondary structure of chromatin as well as gene expression levels. One method for generating acetyl- and methyl-lysine analog-tagged histones is described above and uses phenylselenocysteine. *N*^F-acetyllysine (**13**) is genetically incorporated into proteins in *E. coli* (Neumann et al., 2008). In the presence of a deacetylase inhibitor, manganese superoxide dismutase containing *N*^F-acetyllysine at residue Lys44 is produced; however, acetylation of Lys44 does not appear to affect the enzyme activity. In addition, *N*^F-acetyllysine and other lysine analogs are also incorporated into GRB2 in HEK293 cells (Mukai et al., 2008).

Probing Protein Structure and Function

The structure and function relationship of proteins can be studied with greater precision and accuracy when various biophysical probes are site-specifically incorporated into proteins. Genetically encoded biophysical probes further extend

the potential of these studies into live cells, the native environment of proteins.

NMR spectra from large proteins or complexes are complicated to interpret. Specific NMR labels introduced at defined locations in a protein would greatly reduce complexity and simplify signal assignment. Tri-fluoromethylphenylalanine (tfm-Phe, **14**) was incorporated at the binding interface of two obligate dimers, nitroreductase and histidinol dehydrogenase, both of which contain active sites at the dimer interface (Jackson et al., 2007). Substrate- or inhibitor-induced conformational changes are evident by monitoring the ^{19}F NMR chemical shifts. In another report, the binding of a small-molecule ligand to the thioesterase domain of fatty acid synthase was studied by NMR by using tfm-Phe, ^{13}C - and ^{15}N -labeled *o*-methyltyrosine (**15**), and ^{15}N -labeled *o*-nitrobenzyltyrosine (**16**) at 11 different sites (Cellitti et al., 2008). Comparing the spectra of different mutants and the conformational changes upon addition of the small molecule, the binding site of this molecule is mapped on the protein. It is also worthy to note that photo-decaging of **16** to regenerate ^{15}N -labeled tyrosine provides a useful method of isotopic labeling of selected tyrosines without structural change.

p-Cyanophenylalanine (*p*CNPhe, **17**) is a probe for infrared (IR) spectroscopy (Schultz et al., 2006). The stretching vibration of its nitrile group has strong absorption and a frequency (ν_{CN}) at $\sim 2200\text{ cm}^{-1}$, which falls in the transparent window of protein IR spectra. *p*CNPhe was incorporated into myoglobin at His64, a site close to the iron center of the heme group, to examine ligand-bound states of the heme group. When the Fe(III) ligand is changed from water to cyanide in ferric myoglobin, ν_{CN} shifts from 2248 cm^{-1} to 2236 cm^{-1} . In the ferrous myoglobin, the linear Fe(II)CO complex shows a ν_{CN} absorption at 2239 cm^{-1} , whereas the bent Fe(II)NO and Fe(II)O₂ complexes show absorption at 2230 cm^{-1} . These results demonstrate that the nitrile group is a sensitive probe for ligand binding and local electronic environment.

Fluorescent unnatural amino acids may complement the widely used fluorescent proteins in imaging protein expression, dynamics, and function. Their smaller size can minimize potential perturbations to target proteins, and their fluorophores can be designed to report different microenvironmental changes. Dansylalanine (**18**) contains the dansyl fluorophore, whose fluorescence intensity increases in hydrophobic surroundings. When incorporated on the surface of a β barrel in human superoxide dismutase, dansylalanine shows little change in fluorescence after protein denaturation; when placed at an internal site of the β barrel, its emission wavelength is red shifted and intensity is greatly decreased upon denaturation (Summerer et al., 2006). Therefore, fluorescence change can be used to infer when a residue is exposed to solvent during unfolding. L-(7-hydroxycoumarin-4-yl) ethylglycine (**19**) bears the coumarin fluorophore, which is sensitive to pH and polarity. It was genetically incorporated into myoglobin at two different helices, A or C (Wang et al., 2006a). At 3 M urea, coumarin placed in helices A and C shows a similar increase in fluorescence. However, at 2 M urea, fluorescence increases only when coumarin is incorporated in helix A, suggesting that helix C is not destabilized until urea concentrations reach above 3 M. This result is consistent with previous NMR data, and it suggests that coumarin fluorescence can

also report local protein unfolding. Another reporter, *p*-nitrophenylalanine (**20**) quenches the fluorescence of nearby tryptophans (Tsao et al., 2006). When it is incorporated in a leucine zipper protein, GCN4, the tryptophanyl fluorescence is quenched in a distance-dependent manner, making *p*-nitrophenylalanine a useful distance probe to monitor protein folding or conformational changes. Although still in its infancy, the use of genetically encoded fluorescent amino acids as real-time optical reporters and biosensors in live cells is promising.

Introducing heavy atoms into protein crystals is a critical step for phase determination in X-ray crystallography. A new method to introduce heavy atoms into proteins is to genetically incorporate *p*-iodophenylalanine (**3**) into proteins in *E. coli* or yeast. *p*-Iodophenylalanine was incorporated into the T4 lysozyme, and the mutant protein was crystallized (Xie et al., 2004). Diffraction data were collected with a laboratory CuK α X-ray source, and the structure was solved by using single-wavelength anomalous dispersion phasing. A single iodinated amino acid among 164 residues results in a strong anomalous signal, about 3% of the total intensities, which compares favorably with the level achieved with selenomethionine by using synchrotron beams. *p*-Iodophenylalanine causes little structural perturbation when substituted for Phe in the core of the T4 lysozyme. This approach ensures that the heavy atom iodine is quantitatively introduced at a specific site of the target protein. The strong anomalous signal, the possibility of incorporation at multiple sites and in different cell types, and the use of an in-house X-ray source should facilitate solving protein structures in a high-throughput manner.

The side chain size of unnatural amino acids can be conveniently altered at atomic precision to provide desired changes for probing the bulk effects of amino acids on protein structure and function. The fast inactivation mechanism of the voltage-gated K⁺ channel Kv1.4 was examined by using unnatural amino acids in HEK293 cells (Wang et al., 2007b). The classic ball-and-chain model for channel inactivation proposes that the N-terminal inactivation peptide forms a ball-like domain to occlude the channel exit for ions. In contrast, a new model hypothesizes that the inactivation peptide threads through a side portal and extends into the inner pore to block ion flow. To experimentally test this, Tyr19 in the inactivation peptide was initially mutated to Phe or Trp, which shows no difference in channel inactivation. However, mutation to *o*-methyltyrosine (**1**) results in a markedly slower inactivation, as does mutation to dansylalanine (**18**) (Figure 6). Modeling suggested that the diameter of the inactivation peptide, which is unchanged for Phe or Trp at this site but is larger for *o*-methyltyrosine and dansylalanine, is important for channel inactivation. This is likely due to the narrow width of the side portal in the channel, supporting the new model for channel inactivation.

Identifying and Regulating Protein Activity

The identification of protein interactions and the control of protein activities *in vivo* would ideally require methods that are noninvasive and easy to administer. Application of light is an attractive approach. Photoresponsive unnatural amino acids that can crosslink with nearby molecules, shed protecting groups, or change conformation have been developed and genetically incorporated into proteins. They enable the *in vivo* manipulation of proteins, which should have great potential in molecular and synthetic biology applications.




Figure 6. Unnatural Amino Acids to Probe the Inactivation Mechanism of Ion Channel Kv1.4

o-Methyltyrosine or dansylalanine extends the side chain length of tyrosine, which impedes the inactivation peptide from threading through the side portal of the ion channel and abolishes the fast inactivation, as shown in the current time curve.

the amino acid to the wild-type active form, generating abrupt or localized changes. For example, mutation of the active site Cys residue in the proapoptotic protease caspase 3 to *o*-nitrobenzylcysteine (**23**) in yeast leads to a catalytically inactive enzyme. UV illumination of the cell lysate converts ~40% of the caged caspase to the active enzyme (Wu et al., 2004). *o*-Nitrobenzyltyrosine (**24**) is incorporated into β -galactosidase at Tyr503 in *E. coli*, which effectively reduced the activity of this enzyme to 5% of the wild-type form (Deiters et al.,

Crosslinking has been an important technique for uncovering protein interactions in cells. Site-specific photocrosslinking enabled by genetically encoded unnatural amino acids may help to pinpoint transient or weak interactions, to identify which region is involved, and to distinguish direct from indirect interactions. *p*-Azidophenylalanine (**7**), *p*-benzoylphenylalanine (pBpA, **21**), and *p*-(3-trifluoromethyl-3*H*-diazirin-3-yl)phenylalanine (TfmdPhe, **22**) contain photocrosslinking side chains and have been genetically incorporated into proteins (Chin et al., 2002a, 2002b; Tippmann et al., 2007). pBpA is incorporated into the heat shock protein ClpB in *E. coli* at residue Tyr251, which is considered as the main substrate recognition residue (Schlieker et al., 2004). Biotinylated substrate peptides are shown to be crosslinked upon UV light exposure, but not if pBpA is incorporated elsewhere. pBpA is also incorporated into Ste2p, a GPCR that binds a short peptide pheromone α factor, at several sites on the extracellular loops in yeast (Huang et al., 2008). At two sites, biotinylated α factor is indeed photocaptured and identified by western blot. pBpA has also been incorporated into Grb2 (growth-factor-receptor-bound protein 2) in the ligand-binding pocket of the SH2 domain in mammalian cells. When activated epidermal growth factor receptor (EGFR) and pBpA containing Grb2 are coexpressed and treated with UV light, a larger-molecular weight band is detected, suggesting the binding of EGFR to Grb2 (Hino et al., 2005). Genetically encoded photocrosslinkers should provide high specificity and be compatible with a wide range of biological processes, making this strategy attractive for probing protein complexes and interactions in vivo.

Photocaging of critical residues can be harnessed to regulate protein activity by using light. A photoremovable protecting group is installed onto the suitable amino acid in the target protein, which masks the amino acid and renders the protein inactive. Photolysis releases the caging group and converts

(2006). After a 30 min exposure of bacterial cells to UV light, the enzyme regains activity to ~70% of wild-type levels. A photocaged serine, 4,5-dimethoxy-2-nitrobenzylserine (DMNB-Ser, **25**), is incorporated at phosphoserine sites in the transcription factor Pho4 to control its phosphorylation in yeast (Lemke et al., 2007). DMNB-Ser blocks phosphorylation until photolysed with 405 nm light. Serine is then regenerated and subsequently phosphorylated, triggering the nuclear export of Pho4 (Figure 7A). Photocaging prevents posttranslational modification of the selected residue until a desired moment, which is a more versatile method than mutating the residue to a null amino acid in examining signal transductions.

Photolysis of a caged residue is an irreversible process. Reversible modulation can be achieved with the photochromic azobenzene group, which undergoes a reversible *cis-trans* isomerization by using light with different wavelengths (Figure 7B). The resultant change in geometry and/or dipole of the compound can be used to regulate protein activity. *p*-Azophenylphenylalanine (AzoPhe, **26**) has been incorporated at Ile71 of the *E. coli* catabolite activator protein, a transcriptional activator (Bose et al., 2006). Its binding affinity for the promoter sequence decreases 4-fold after irradiation at 334 nm, which converts the predominant *trans* AzoPhe to the *cis* form. The isomerized *cis* AzoPhe is switched back to the *trans* state by irradiation at > 420 nm, after which the protein affinity for the promoter is completely recovered.

Generating New Protein Properties

Natural evolution affords numerous proteins for life needs, whereas directed protein evolution and rational engineering further increase the diversity of protein functions. Genetically encoding additional functional groups into proteins has the potential to generate totally novel and previously impossible protein properties, and initial progress is encouraging.

Metal ions bound in proteins participate in a large number of catalytic and electron transfer reactions in cells. Rationally




Figure 7. Regulation of Protein Activity with Photoresponsive Amino Acids

(A) Photocaged serine is used to control when the target serine is regenerated and phosphorylated. (B) The photoisomerizable *p*-azophenyl-phenylalanine is used to control the binding of CAP to its promoter, which is determined by a gel mobility shift assay.

retained in mice after exposure to the mutant TNF- α , and it protects mice against lipopolysaccharide-induced death. These striking observations have the potential for generating new therapies for cancer or protein-misfolding diseases.

Conclusions and Future Perspectives

Since the first unnatural amino acid was genetically encoded in live cells in 2001 (Wang et al., 2001), more than 40 unnatural amino acids have been incorporated into proteins in *E. coli*, and about a dozen have been incorporated into yeast and mammalian cells to date. It should be noted that this method currently cannot be used to incorporate unnatural amino acids that are incompatible with the ribosome (such as α -amino acids and β amino acids) or toxic to cells (such as an analog that is structurally so similar

engineering a metalloprotein is challenging, as it is difficult to predict and control the amino acid shells surrounding the metal ion. Multidentate, metal-chelating amino acids with coordinating atoms preorientated in the correct configuration may facilitate such an effort. Bipyridylalanine (BpyAla, **27**) has been shown to reversibly bind copper ions when genetically incorporated into the T4 lysozyme (Xie et al., 2007a). The Cu(II)-binding property is exploited to oxidatively cleave the phosphosugar moiety of the nucleic acid backbone. BpyAla is incorporated into the catabolite activator protein (CAP) at Lys26, a site close to the protein-DNA interface (Lee and Schultz, 2008). In the presence of Cu(II) and 3-mercaptopropionic acid, the mutant CAP, originally a transcription factor, cleaves double-stranded DNA at its consensus sequence with high specificity. This method may be generally applied to other DNA-binding proteins to map their preferred DNA sequences.

The distance probe *p*-nitrophenylalanine (**20**) has recently been rediscovered to stimulate potent immune responses for novel immunogenic applications. An animal usually will not mount a substantial immune response against a self-protein. This immune self-tolerance has been shown to be broken by introducing *p*-nitrophenylalanine into the antigen (Grünevald et al., 2008). Incorporation of this unnatural amino acid into tumor necrosis factor α (TNF- α) leads to a strong immune response directed at the unnatural mutant TNF- α in mice, even in the absence of adjuvant. The generated antibodies crossreact with native mouse TNF- α . The immune response is

to a common amino acid that the analog is charged by the cognate synthetase of the amino acid and misincorporated into the proteome) (Hartman et al., 2007). The novel functional groups embodied by these genetically encoded new amino acids will enable an increasing number of new experiments for biological research. The utility of many unnatural amino acids has been demonstrated in proof-of-principle experiments, and they are expected to be applied to various biological systems for addressing challenging questions and uncovering unknowns in the coming years. Unnatural amino acids that are fluorescent, photoresponsive, bio-orthogonally reactive, or able to mimic various protein modifications would be particularly useful in addressing cell biology questions. Further improvement of the unnatural amino acid incorporation efficiency and detailed studies on the physiological effects of their incorporation would be important for effective application, especially in mammalian cells. The introduction of this technology into multicellular organisms would open the field to biological problems concerning how cells interact with one another in a variety of settings, allowing aspects of development, neural connectivity, and cellular signaling to be studied, to name only a few. The development of random unnatural amino acid mutagenesis may enable the evolution of protein properties that have never existed in nature. Lastly, the genetic incorporation of unnatural amino acids has implications on various aspects of the origin and evolution of the genetic code itself, which remains to be explored.

ACKNOWLEDGMENTS

L.W. acknowledges support from the Searle Scholar program, the Beckman Young Investigator Program, the March of Dimes Foundation (5-FY08110), the California Institute for Regenerative Medicine (RN1-00577-1), and the Office of the Director of the National Institutes of Health (DP2OD004744).

REFERENCES

- Ambrogelly, A., Palioura, S., and Söll, D. (2006). Natural expansion of the genetic code. *Nat. Chem. Biol.* 3, 29–35.
- Amrani, N., Sachs, M.S., and Jacobson, A. (2006). Early nonsense: mRNA decay solves a translational problem. *Nat. Rev. Mol. Cell Biol.* 7, 415–425.
- Anderson, J.C., and Schultz, P.G. (2003). Adaptation of an orthogonal archaeal leucyl-tRNA and synthetase pair for four-base, amber, and opal suppression. *Biochemistry* 42, 9598–9608.
- Anderson, J.C., Wu, N., Santoro, S.W., Lakshman, V., King, D.S., and Schultz, P.G. (2004). An expanded genetic code with a functional quadruplet codon. *Proc. Natl. Acad. Sci. USA* 101, 7566–7571.
- Bain, J.D., Glabe, C.G., Dix, T.A., Chamberlin, A.R., and Diala, E.S. (1989). Biosynthetic site-specific incorporation of a non-natural amino acid into a polypeptide. *J. Am. Chem. Soc.* 111, 8013–8014.
- Beene, D.L., Dougherty, D.A., and Lester, H.A. (2003). Unnatural amino acid mutagenesis in mapping ion channel function. *Curr. Opin. Neurobiol.* 13, 264–270.
- Bock, A., Forchhammer, K., Heider, J., Leinfelder, W., Sawers, G., Veprek, B., and Zinoni, F. (1991). Selenocysteine: the 21st amino acid. *Mol. Microbiol.* 5, 515–520.
- Bose, M., Groff, D., Xie, J., Brustad, E., and Schultz, P.G. (2006). The incorporation of a photoisomerizable amino acid into proteins in *E. coli*. *J. Am. Chem. Soc.* 128, 388–389.
- Brick, P., Bhat, T.N., and Blow, D.M. (1989). Structure of tyrosyl-tRNA synthetase refined at 2.3 Å resolution. Interaction of the enzyme with the tyrosyl adenylate intermediate. *J. Mol. Biol.* 208, 83–98.
- Cao, D., and Parker, R. (2003). Computational modeling and experimental analysis of nonsense-mediated decay in yeast. *Cell* 113, 533–545.
- Cellitti, S.E., Jones, D.H., Lagpacan, L., Hao, X., Zhang, Q., Hu, H., Brittain, S.M., Brinker, A., Caldwell, J., Bursulaya, B., et al. (2008). In vivo incorporation of unnatural amino acids to probe structure, dynamics, and ligand binding in a large protein by nuclear magnetic resonance spectroscopy. *J. Am. Chem. Soc.* 130, 9268–9281.
- Chen, S., Schultz, P.G., and Brock, A. (2007). An improved system for the generation and analysis of mutant proteins containing unnatural amino acids in *Saccharomyces cerevisiae*. *J. Mol. Biol.* 371, 112–122.
- Chin, J.W., Martin, A.B., King, D.S., Wang, L., and Schultz, P.G. (2002a). Addition of a photocrosslinking amino acid to the genetic code of *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 99, 11020–11024.
- Chin, J.W., Santoro, S.W., Martin, A.B., King, D.S., Wang, L., and Schultz, P.G. (2002b). Addition of *p*-azido-L-phenylalanine to the genetic code of *Escherichia coli*. *J. Am. Chem. Soc.* 124, 9026–9027.
- Chin, J.W., Cropp, T.A., Anderson, J.C., Mukherji, M., Zhang, Z., and Schultz, P.G. (2003). An expanded eukaryotic genetic code. *Science* 301, 964–967.
- Cohen, G.N., and Cowie, D.B. (1957). Total replacement of methionine by selenomethionine in the proteins of *Escherichia coli*. *C. R. Hebd. Seances Acad. Sci.* 244, 680–683.
- Cornish, V.W., Mendel, D., and Schultz, P.G. (1995). Probing protein structure and function with an expanded genetic code. *Angew. Chem. Int. Ed. Engl.* 34, 621–633.
- Dawson, P.E., Muir, T.W., Clarklewis, I., and Kent, S.B.H. (1994). Synthesis of proteins by native chemical ligation. *Science* 266, 776–779.
- Deiters, A., Cropp, T.A., Mukherji, M., Chin, J.W., Anderson, J.C., and Schultz, P.G. (2003). Adding amino acids with novel reactivity to the genetic code of *Saccharomyces cerevisiae*. *J. Am. Chem. Soc.* 125, 11782–11783.
- Deiters, A., Cropp, T.A., Summerer, D., Mukherji, M., and Schultz, P.G. (2004). Site-specific PEGylation of proteins containing unnatural amino acids. *Bioorg. Med. Chem. Lett.* 14, 5743–5745.
- Deiters, A., Groff, D., Ryu, Y., Xie, J., and Schultz, P.G. (2006). A genetically encoded photocaged tyrosine. *Angew. Chem. Int. Ed. Engl.* 45, 2728–2731.
- Drabkin, H.J., Park, H.J., and Rajbhandary, U.L. (1996). Amber suppression in mammalian cells dependent upon expression of an *Escherichia coli* aminoacyl-tRNA synthetase gene. *Mol. Cell. Biol.* 16, 907–913.
- Edwards, H., and Schimmel, P. (1990). A bacterial amber suppressor in *Saccharomyces cerevisiae* is selectively recognized by a bacterial aminoacyl-tRNA synthetase. *Mol. Cell. Biol.* 10, 1633–1641.
- Francis, M.A., and Rajbhandary, U.L. (1990). Expression and function of a human initiator tRNA gene in the yeast *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 10, 4486–4494.
- Galli, G., Hofstetter, H., and Birnstiel, M.L. (1981). Two conserved sequence blocks within eukaryotic tRNA genes are major promoter elements. *Nature* 294, 626–631.
- Grünwald, J., Tsao, M.L., Perera, R., Dong, L., Niessen, F., Wen, B.G., Kubitz, D.M., Smider, V.V., Ruf, W., Nasoff, M., et al. (2008). Immunochemical termination of self-tolerance. *Proc. Natl. Acad. Sci. USA* 105, 11276–11280.
- Guo, J., Wang, J., Lee, J.S., and Schultz, P.G. (2008). Site-specific incorporation of methyl- and acetyl-lysine analogues into recombinant proteins. *Angew. Chem. Int. Ed. Engl.* 47, 6399–6401.
- Hahn, M.E., Pellois, J.P., Vila-Perello, M., and Muir, T.W. (2007). Tunable photoactivation of apost-translationally modified signaling protein and its unmodified counterpart in live cells. *ChemBioChem* 8, 2100–2105.
- Hartman, M.C., Josephson, K., Lin, C.W., and Szostak, J.W. (2007). An expanded set of amino acid analogs for the ribosomal translation of unnatural peptides. *PLoS ONE* 2, e972.
- Hino, N., Okazaki, Y., Kobayashi, T., Hayashi, A., Sakamoto, K., and Yokoyama, S. (2005). Protein photo-cross-linking in mammalian cells by site-specific incorporation of a photoreactive amino acid. *Nat. Methods* 2, 201–206.
- Hohsaka, T., and Sisido, M. (2002). Incorporation of non-natural amino acids into proteins. *Curr. Opin. Chem. Biol.* 6, 809–815.
- Hsieh, J., and Fire, A. (2000). Recognition and silencing of repeated DNA. *Annu. Rev. Genet.* 34, 187–204.
- Huang, L.Y., Umanah, G., Hauser, M., Son, C., Arshava, B., Naider, F., and Becker, J.M. (2008). Unnatural amino acid replacement in a yeast G protein-coupled receptor in its native environment. *Biochemistry* 47, 5638–5648.
- Ibba, M., and Soll, D. (2000). Aminoacyl-tRNA synthesis. *Annu. Rev. Biochem.* 69, 617–650.
- Jackson, J.C., Hammill, J.T., and Mehl, R.A. (2007). Site-specific incorporation of a (19)F-amino acid into proteins as an NMR probe for characterizing protein structure and reactivity. *J. Am. Chem. Soc.* 129, 1160–1166.
- Kaiser, E.T., Lawrence, D.S., and Rokita, S.E. (1985). The chemical modification of enzymatic specificity. *Annu. Rev. Biochem.* 54, 565–595.
- Kent, S.B. (1988). Chemical synthesis of peptides and proteins. *Annu. Rev. Biochem.* 57, 957–989.
- Knight, R.D., Freeland, S.J., and Landweber, L.F. (2001). Rewiring the keyboard: evolvability of the genetic code. *Nat. Rev. Genet.* 2, 49–58.
- Kowal, A.K., Kohrer, C., and Rajbhandary, U.L. (2001). Twenty-first aminoacyl-tRNA synthetase-suppressor tRNA pairs for possible use in site-specific incorporation of amino acid analogues into proteins in eukaryotes and in eubacteria. *Proc. Natl. Acad. Sci. USA* 98, 2268–2273.
- Kwok, Y., and Wong, J.T. (1980). Evolutionary relationship between *Halobacterium cutirubrum* and eukaryotes determined by use of aminoacyl-tRNA synthetases as phylogenetic probes. *Can. J. Biochem.* 58, 213–218.
- Lee, H.S., and Schultz, P.G. (2008). Biosynthesis of a site-specific DNA cleaving protein. *J. Am. Chem. Soc.* 130, 13194–13195.

- Lenke, E.A., Summerer, D., Geierstanger, B.H., Brittain, S.M., and Schultz, P.G. (2007). Control of protein phosphorylation with a genetically encoded photocaged amino acid. *Nat. Chem. Biol.* **3**, 769–772.
- Link, A.J., Mock, M.L., and Tirrell, D.A. (2003). Non-canonical amino acids in protein engineering. *Curr. Opin. Biotechnol.* **14**, 603–609.
- Liu, H., Wang, L., Brock, A., Wong, C.H., and Schultz, P.G. (2003). A method for the generation of glycoprotein mimetics. *J. Am. Chem. Soc.* **125**, 1702–1703.
- Liu, W., Brock, A., Chen, S., Chen, S., and Schultz, P. (2007). Genetic incorporation of unnatural amino acids into proteins in mammalian cells. *Nat. Methods* **4**, 239–244.
- Lumms, S.C., Beene, D.L., Lee, L.W., Lester, H.A., Broadhurst, R.W., and Dougherty, D.A. (2005). Cis-trans isomerization at a proline opens the pore of a neurotransmitter-gated ion channel. *Nature* **438**, 248–252.
- McGinty, R.K., Kim, J., Chatterjee, C., Roeder, R.G., and Muir, T.W. (2008). Chemically ubiquitylated histone H2B stimulates hDot1L-mediated intranucleosomal methylation. *Nature* **453**, 812–816.
- Mehl, R.A., Anderson, J.C., Santoro, S.W., Wang, L., Martin, A.B., King, D.S., Horn, D.M., and Schultz, P.G. (2003). Generation of a bacterium with a 21 amino acid genetic code. *J. Am. Chem. Soc.* **125**, 935–939.
- Minks, C., Alefelder, S., Moroder, L., Huber, R., and Budisa, N. (2000). Towards new protein engineering: in vivo building and folding of protein shuttles for drug delivery and targeting by the selective pressure incorporation (SPI) method. *Tetrahedron* **56**, 9431–9442.
- Muir, T.W. (2003). Semisynthesis of proteins by expressed protein ligation. *Annu. Rev. Biochem.* **72**, 249–289.
- Mukai, T., Kobayashi, T., Hino, N., Yanagisawa, T., Sakamoto, K., and Yokoyama, S. (2008). Adding L-lysine derivatives to the genetic code of mammalian cells with engineered pyrrolysyl-tRNA synthetases. *Biochem. Biophys. Res. Commun.* **371**, 818–822.
- Myslinski, E., Ame, J.C., Krol, A., and Carbon, P. (2001). An unusually compact external promoter for RNA polymerase III transcription of the human H1RNA gene. *Nucleic Acids Res.* **29**, 2502–2509.
- Neumann, H., Peak-Chew, S.Y., and Chin, J.W. (2008). Genetically encoding N(ε)-acetyllysine in recombinant proteins. *Nat. Chem. Biol.* **4**, 232–234.
- Noren, C.J., Anthony-Cahill, S.J., Griffith, M.C., and Schultz, P.G. (1989). A general method for site-specific incorporation of unnatural amino acids into proteins. *Science* **244**, 182–188.
- Normanly, J., Kleina, L.G., Masson, J.M., Abelson, J., and Miller, J.H. (1990). Construction of *Escherichia coli* amber suppressor tRNA genes. III. Determination of tRNA specificity. *J. Mol. Biol.* **213**, 719–726.
- Nowak, M.W., Kearney, P.C., Sampson, J.R., Saks, M.E., Labarca, C.G., Silverman, S.K., Zhong, W.G., Thorson, J., Abelson, J.N., Davidson, N., et al. (1995). Nicotinic receptor binding site probed with unnatural amino acid incorporation in intact cells. *Science* **268**, 439–442.
- Paule, M.R., and White, R.J. (2000). Survey and summary: transcription by RNA polymerases I and III. *Nucleic Acids Res.* **28**, 1283–1298.
- Pellois, J.P., and Muir, T.W. (2006). Semisynthetic proteins in mechanistic studies: using chemistry to go where nature can't. *Curr. Opin. Chem. Biol.* **10**, 487–491.
- Sakamoto, K., Hayashi, A., Sakamoto, A., Kiga, D., Nakayama, H., Soma, A., Kobayashi, T., Kitabatake, M., Takio, K., Saito, K., et al. (2002). Site-specific incorporation of an unnatural amino acid into proteins in mammalian cells. *Nucleic Acids Res.* **30**, 4692–4699.
- Schlieker, C., Weibezahn, J., Patzelt, H., Tessarz, P., Strub, C., Zeth, K., Erbse, A., Schneider-Mergener, J., Chin, J.W., Schultz, P.G., et al. (2004). Substrate recognition by the AAA+ chaperone ClpB. *Nat. Struct. Mol. Biol.* **11**, 607–615.
- Schultz, K.C., Supekova, L., Ryu, Y., Xie, J., Perera, R., and Schultz, P.G. (2006). A genetically encoded infrared probe. *J. Am. Chem. Soc.* **128**, 13984–13985.
- Schwarzer, D., and Cole, P.A. (2005). Protein semisynthesis and expressed protein ligation: chasing a protein's tail. *Curr. Opin. Chem. Biol.* **9**, 561–569.
- Srinivasan, G., James, C.M., and Krzycki, J.A. (2002). Pyrrolysine encoded by UAG in Archaea: charging of a UAG-decoding specialized tRNA. *Science* **296**, 1459–1462.
- Summerer, D., Chen, S., Wu, N., Deiters, A., Chin, J.W., and Schultz, P.G. (2006). A genetically encoded fluorescent amino acid. *Proc. Natl. Acad. Sci. USA* **103**, 9785–9789.
- Tippmann, E.M., Liu, W., Summerer, D., Mack, A.V., and Schultz, P.G. (2007). A genetically encoded diazirine photocrosslinker in *Escherichia coli*. *ChemBioChem* **8**, 2210–2214.
- Tsao, M.L., Tian, F., and Schultz, P.G. (2005). Selective Staudinger modification of proteins containing *p*-azidophenylalanine. *ChemBioChem* **6**, 2147–2149.
- Tsao, M.L., Summerer, D., Ryu, Y., and Schultz, P.G. (2006). The genetic incorporation of a distance probe into proteins in *Escherichia coli*. *J. Am. Chem. Soc.* **128**, 4572–4573.
- Uy, R., and Wold, F. (1977). Posttranslational covalent modification of proteins. *Science* **198**, 890–896.
- Valiyaveetil, F.I., Leonetti, M., Muir, T.W., and Mackinnon, R. (2006). Ion selectivity in a semisynthetic K⁺ channel locked in the conductive conformation. *Science* **314**, 1004–1007.
- Wang, J., Xie, J., and Schultz, P.G. (2006a). A genetically encoded fluorescent amino acid. *J. Am. Chem. Soc.* **128**, 8738–8739.
- Wang, J., Schiller, S.M., and Schultz, P.G. (2007a). A biosynthetic route to dehydroalanine-containing proteins. *Angew. Chem. Int. Ed. Engl.* **46**, 6849–6851.
- Wang, L., and Schultz, P.G. (2001). A general approach for the generation of orthogonal tRNAs. *Chem. Biol.* **8**, 883–890.
- Wang, L., and Schultz, P.G. (2002). Expanding the genetic code. *Chem. Commun.* 1–11.
- Wang, L., and Schultz, P.G. (2004). Expanding the genetic code. *Angew. Chem. Int. Ed. Engl.* **44**, 34–66.
- Wang, L., Magliery, T.J., Liu, D.R., and Schultz, P.G. (2000). A new functional suppressor tRNA/aminoacyl-tRNA synthetase pair for the in vivo incorporation of unnatural amino acids into proteins. *J. Am. Chem. Soc.* **122**, 5010–5011.
- Wang, L., Brock, A., Herberich, B., and Schultz, P.G. (2001). Expanding the genetic code of *Escherichia coli*. *Science* **292**, 498–500.
- Wang, L., Brock, A., and Schultz, P.G. (2002). Adding L-3-(2-Naphthyl)alanine to the genetic code of *E. coli*. *J. Am. Chem. Soc.* **124**, 1836–1837.
- Wang, L., Zhang, Z., Brock, A., and Schultz, P.G. (2003). Addition of the keto functional group to the genetic code of *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **100**, 56–61.
- Wang, L., Xie, J., and Schultz, P.G. (2006b). Expanding the genetic code. *Annu. Rev. Biophys. Biomol. Struct.* **35**, 225–249.
- Wang, Q., and Wang, L. (2008). New methods enabling efficient incorporation of unnatural amino acids in yeast. *J. Am. Chem. Soc.* **130**, 6066–6067.
- Wang, W., Takimoto, J., Louie, G.V., Baiga, T.J., Noel, J.P., Lee, K.F., Slesinger, P.A., and Wang, L. (2007b). Genetically encoding unnatural amino acids for cellular and neuronal studies. *Nat. Neurosci.* **10**, 1063–1072.
- Willis, I.M. (1993). RNA polymerase III. Genes, factors and transcriptional specificity. *Eur. J. Biochem.* **212**, 1–11.
- Wu, N., Deiters, A., Cropp, T.A., King, D., and Schultz, P.G. (2004). A genetically encoded photocaged amino acid. *J. Am. Chem. Soc.* **126**, 14306–14307.
- Xie, J., Wang, L., Wu, N., Brock, A., Spraggon, G., and Schultz, P.G. (2004). The site-specific incorporation of *p*-iodo-L-phenylalanine into proteins for structure determination. *Nat. Biotechnol.* **22**, 1297–1301.
- Xie, J., Liu, W., and Schultz, P.G. (2007a). A genetically encoded bidentate, metal-binding amino acid. *Angew. Chem. Int. Ed. Engl.* **46**, 9239–9242.

- Xie, J., Supekova, L., and Schultz, P.G. (2007b). A genetically encoded metabolically stable analogue of phosphotyrosine in *Escherichia coli*. *ACS Chem. Biol.* **2**, 474–478.
- Xu, R., Hanson, S.R., Zhang, Z., Yang, Y.Y., Schultz, P.G., and Wong, C.H. (2004). Site-specific incorporation of the mucin-type N-acetylgalactosamine- α -O-threonine into protein in *Escherichia coli*. *J. Am. Chem. Soc.* **126**, 15654–15655.
- Ye, S., Köhrer, C., Huber, T., Kazmi, M., Sachdev, P., Yan, E.C., Bhagat, A., Rajbhandary, U.L., and Sakmar, T.P. (2008). Site-specific incorporation of keto amino acids into functional G protein-coupled receptors using unnatural amino acid mutagenesis. *J. Biol. Chem.* **283**, 1525–1533.
- Zhang, Z., Smith, B.A., Wang, L., Brock, A., Cho, C., and Schultz, P.G. (2003). A new strategy for the site-specific modification of proteins in vivo. *Biochemistry* **42**, 6735–6746.
- Zhang, Z., Alfonta, L., Tian, F., Bursulaya, B., Uryu, S., King, D.S., and Schultz, P.G. (2004a). Selective incorporation of 5-hydroxytryptophan into proteins in mammalian cells. *Proc. Natl. Acad. Sci. USA* **101**, 8882–8887.
- Zhang, Z., Gildersleeve, J., Yang, Y.Y., Xu, R., Loo, J.A., Uryu, S., Wong, C.H., and Schultz, P.G. (2004b). A new strategy for the synthesis of glycoproteins. *Science* **303**, 371–373.