


This article was downloaded by:

On: 15 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Chemistry and Ecology

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713455114>

Origin and Distribution of Suspended Organic Matter As Inferred From Carbon Isotope Composition in A Mediterranean Semi-Enclosed Marine System

A. Mazzola^{ab}; G. Sarà^a; F. Venezia^c; M. Caruso^a; D. Catalano^a; S. Hauser^c

^a Laboratory of Marine Biology and Resources, Department of Animal Biology, University of Palermo, Palermo, Italy ^b Dipartimento di Biologia Animale, Laboratorio di Biologia e Risorse Marine, Palermo, Italy ^c Institute of Mineralogy, Petrography and Geochemistry, University of Palermo, Italy

To cite this Article Mazzola, A. , Sarà, G. , Venezia, F. , Caruso, M. , Catalano, D. and Hauser, S.(1999) 'Origin and Distribution of Suspended Organic Matter As Inferred From Carbon Isotope Composition in A Mediterranean Semi-Enclosed Marine System', *Chemistry and Ecology*, 16: 2, 215 – 238

To link to this Article: DOI: 10.1080/02757549908037647

URL: <http://dx.doi.org/10.1080/02757549908037647>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

ORIGIN AND DISTRIBUTION OF SUSPENDED ORGANIC MATTER AS INFERRED FROM CARBON ISOTOPE COMPOSITION IN A MEDITERRANEAN SEMI-ENCLOSED MARINE SYSTEM

A. MAZZOLA^{a,*}, G. SARÀ^a, F. VENEZIA^b, M. CARUSO^a,
D. CATALANO^a and S. HAUSER^b

^a *Laboratory of Marine Biology and Resources, Department of Animal Biology,
University of Palermo, Via Archirafi, 18, I-90123 Palermo, Italy;*

^b *Institute of Mineralogy, Petrography and Geochemistry, University of
Palermo, Via Archirafi, 36, I-90123, Italy*

(Received 14 December 1998; In final form 22 February 1999)

The origin and distribution of suspended organic matter, the trophic features and the stable carbon isotopic composition of particulate organic carbon (POC) were studied monthly in a Western Mediterranean semi-enclosed basin. Sampling stations were selected as a function of wind-exposure and the degree of vegetation cover and then compared with an adjacent unvegetated site. The predominant vegetation was seagrass (*Posidonia oceanica* and *Cymodocea nodosa*) and *Caulerpa prolifera*. Water samples were analyzed for total suspended matter (inorganic and organic fractions), photosynthetic pigments (chlorophyll-a and phaeopigments), dissolved organic carbon, particulate organic carbon and their isotopic composition. Temperature and salinity were also measured at the same sampling sites within range of Mediterranean limits. The suspended organic matter concentration was $1.77 \pm 1.55 \text{ mg l}^{-1}$; the chlorophyll-a concentration was low ($0.35 \pm 0.24 \mu\text{g l}^{-1}$); the dissolved organic carbon concentration was $2,140 \pm 2,010 \mu\text{g l}^{-1}$; the particulate organic carbon concentration was $212 \pm 106 \mu\text{g l}^{-1}$ and the isotopic composition was $18.77 \pm 2.51\%$. There were significant temporal differences except for phaeopigments, POC and its POC isotopic composition, and there were no spatial differences other than for $\delta^{13}\text{C}$. This picture highlighted a general seasonal trend and trophical features similar to adjacent sea.

Spatial differences in $\delta^{13}\text{C}$ showed that the source of suspended organic matter was different between stations as that between sources and wind-hydrodynamic constraints. In

*Address for correspondence: Laboratorio di Biologia e Risorse Marine, Dipartimento di Biologia Animale, Via Archirafi, 18, I-90123 Palermo, Italy. Tel./Fax: (+39) 91 6167497-6172009, e-mail: amazzola@mbox.unipa.it

adjacent open sea, the phytoplankton component was identified as being the main isotopic signal of POC. In the Stagnone, resuspension from exposed sites and sedimentation into sheltered sites control biological dynamics, depending on wind exposition. Consequently, the dominance of the first or the second process determines the typology and the development of vegetation, allowing it to recognize different sub-systems where the origin and distribution of particulate organic carbon could be different. A first sub-system was exposed to dominant winds and affected by sediment erosion which was from wind-induced turbulence, and had a main effect on limited submersed vegetation. The POC isotopic composition ($\delta^{13}\text{C} = -18.5\text{‰}$) reflected a contemporary contribution of macrophyte and microphytobenthos. The combined effects of resuspension-sedimentation-lateral drifting permitted the autotrophic unicellular biomass marker to predominate, compared to *Cymodocea* and *Caulerpa*. The second sub-system was represented by the vitality of the waters and by unpredictability which was due to wind pulsing. As a result, a seasonal regime predominate by the trend in the seagrass leaves component (*Posidonia oceanica* and *Cymodocea nodosa*). The POC isotopic composition followed the seagrass, showing the heaviest values (spring and summer) in peaks in leaf biomass, and the lightest values in autumn and winter when the biomass was low.

Keywords: Organic matter sources; carbon isotope ratio; DOC; POC; MED

1. INTRODUCTION

In recent years, several studies have demonstrated that carbon stable isotope measurements allowed us to recognize the sources of carbon (*e.g.*, seagrass, macroalgae, microphytobenthos, phytoplankton or detritus) that affect the pool of suspended organic matter and trophic dynamics in shallow seas (Fry, 1984; Fry and Sherr, 1984; Mann, 1988; Rau *et al.*, 1991; Michener and Shell, 1994).

In these conditions, changes in the input of suspended organic matter (also by hydrodynamics, tides, internal seiches), have a primary role in affecting the quality and quantity of food sources for consumers (Albertelli and Fabiano, 1990; Fegley *et al.*, 1992; Sarà and Mazzola, 1997; Sarà *et al.*, 1998; Sarà *et al.*, in press). The main consequence is often an unpredictable spatio-temporal variability in the trophic components of these systems.

The semi-arid shallow systems of the southern Mediterranean Sea are characterized by a low or non-existent continental inflow. The main energy source is by wind-induced waves which often produces highly frequent and persistent resuspension events in exposed sites (Painchaud *et al.*, 1996; Sarà *et al.*, in press) affecting the distribution of suspended organic matter by means of lateral drifting. In wind-sheltered sites, the distribution of suspended organic matter is strongly constrained by water-dynamics.

In the Mediterranean, this picture can be important in studies dealing with physical and chemical constraints which affect the organic matter exchanges between different compartments where the primary production from phytoplankton, macroalgae and seagrass generally exceeds consumption by herbivores (Newell, 1982). Organic matter is directly available to consumers but it is highly refractory (Pirc, 1989) and needs to be fragmented and processed by means of bacterial decomposition (Pirc and Wollenweber, 1988; Hansen *et al.*, 1992).

Few studies deal with South Mediterranean environments (Giani *et al.*, 1995; Mazzola and Sarà, 1995; Sarà *et al.*, 1995; Pusceddu *et al.*, 1996; Sarà *et al.*, in press), with respect to other shallow areas (lagoons, sounds or estuaries), showing great variability of their trophic state (Painchaud *et al.*, 1990; Navarro *et al.*, 1993; Clavier *et al.*, 1995; Painchaud *et al.*, 1995; Savenkoff *et al.*, 1995; Sarà *et al.*, in press). Information about the role of the main source of suspended organic matter through carbon isotopic composition are quite scarce (Faganeli *et al.*, 1988; Dauby, 1989; Faganeli *et al.*, 1989; Posedel and Faganeli, 1991; Tufano, 1991; Jennings *et al.*, 1997).

In this paper, the spatial and temporal changes in the origin of particulate organic carbon in the Stagnone of Marsala, a semi-enclosed marine system (Western Sicily, Mediterranean Sea), will be studied. The main aims are: (1) to highlight the origin and distribution of particular organic carbon as a function of different sources; (2) to verify the existence of environmental constraints that affect the sources and spatio-temporal distribution of particular organic carbon and (3) to check how the carbon isotope measurements can improve the dynamic model of the Stagnone of Marsala.

2. MATERIAL AND METHODS

2.1. Study Site

The Stagnone of Marsala (15 km², 37° 52' N, 12° 28' E) (Fig. 1) is a semi-enclosed basin and is separated from the open sea by a low calcareous ooze (Grande Island). The northern mouth has a muddy bottom, and is 450 m wide, and allows turbulent inputs of marine waters. The southern mouth is 1450 m wide and is open to the sea




FIGURE 1 The study site.

water inflow. Three islands (S. Maria, S. Pantaleo and La Scuola) and Strada Romana (Roman Road) act as mechanical obstacles to the water flow in the middle of the basin and generate turbulence. The

dominant seagrasses are *Cymodocea nodosa* and *Posidonia oceanica*, the former luxuriant in the southern and central area. Macroalgae, principally *Caulerpa prolifera*, occurred, often coupled with *Cymodocea* at the station mostly exposed to the wind (Sarà *et al.*, submitted). No land inputs are present. Water exchange is ensured by currents with speeds ranging from $5.10 \pm 1.54 \text{ cm s}^{-1}$ in the southern mouth and from $2.21 \pm 0.87 \text{ cm s}^{-1}$ in the northern area.

2.2. Sampling and Environmental Variables

Superficial water samples (0.5 m depth) were collected monthly from March 1996 to February 1997, using 10 l Niskin bottles, at 5 stations located along a north–south transect (Fig. 1). They were characterized by having a different vegetal cover (Tab. I) (as degree of coverage) proposed by Scilipoti (1997) and Scilipoti *et al.* (1997) and confirmed by remote sensing (Calvo *et al.*, 1996). To obtain isotopic data of the particular organic carbon ($\delta^{13}\text{C}_{(\text{POC})}$), total suspended matter (TSM), suspended pigments and particular organic carbon (POC) in 2000–4000 ml of water was filtered in a gentle vacuum using Whatman GF/F glass-fibre filters. The filtered water was analyzed for dissolved organic carbon (DOC). On each sampling date, water temperature and salinity were measured *in situ* using a Hydrolab multiprobe, and salinity values from the probe were tested monthly using silver nitrate titration.

TABLE I Features of sampling sectors at Stagnone of Marsala

Site	Substrate	Depth (m)	Cover density	Exposure	Vegetation
Station 1 Open-sea	Sand-muddy	5	Absence	Exposed	Unvegetated
Station 2	Muddy-sand	1.05	Average	Exposed	<i>Cymodocea nodosa</i>
Northern basin Station 3	Sandy	1.20	High	Sheltered	<i>Caulerpa prolifera</i> <i>Cymodocea nodosa</i>
Middle basin Station 4	Muddy-sand	1.10	Low	Partially Exposed	<i>Posidonia oceanica</i> <i>Cymodocea nodosa</i>
East Channel					<i>Caulerpa prolifera</i> Other macroalgae
Station 5	Sandy	1.00	High	Sheltered	<i>Cymodocea nodosa</i>
Southern basin					<i>Posidonia oceanica</i>

2.3. Suspended Matter Features

To determine TSM, Whatman GF/F filters (0.45 μm nominal pore size) were weighed after desiccation (60°C, 24 h) using a Sartorius A200 balance (accuracy $\pm 1 \mu\text{g}$). Suspended organic matter (OSM) concentrations were determined after loss on ignition (450°C, 4 h; Strickland and Parsons, 1969). Chlorophyll-a (CHL-a) analyses were carried out according to Lorenzen and Jeffrey's (1980) method by using a spectrophotometer Jasco (Mod. V-530). Phaeopigments (PHAEO) were determined after acidification with 0.1N hydrochloric acid. The sum of chlorophyll-a and phaeopigment was reported as chloroplastic pigments equivalents (CPE; Pfannkuche, 1993). Particulate organic carbon was determined with a CHN Elemental Analyzer (Perkin-Elmer Mod. 2400), using acetanilide at 925°C as a standard, after the removal of inorganic carbon (Iseki *et al.*, 1987). DOC measures were carried out with Shimadzu TOC (Mod. 5000) (Sugimura and Suzuki, 1988).

Water samples for determining the composition of isotopic carbon were treated with 1N hydrochloric acid, then rinsed and dried (60°C; 24 h). Plankton, macroalgae, seagrass isotopic analyses were performed as above (2N hydrochloric acid); macroalgae and seagrass were scraped for vegetal epiphytes. The organic carbon was converted to carbon dioxide by combustion over cupric oxide (550°C, overnight) in a Pyrex tube. Purified carbon dioxide was analyzed by a mass spectrometer (Delta S, Finnigan MAT) and the results, expressed in the usual $\delta\text{‰}$ units, were reported against the PDB international standard. The reproducibility of the $\delta^{13}\text{C}$ determination was $\pm 0.2\text{‰}$.

A Spearman-Rank correlation was performed to test correlation degree between measured variables (Sokal and Rohlf, 1981). Data were ordered by means of a principal components analyses (PCA; Flury, 1988) using station by month environmental matrixes. Statistica (Statsoft Inc.; rel. 5.1) was used to perform statistical analysis.

3. RESULTS

Table II summarizes main statistics of measured hydrological parameters. Temperature and salinity (Fig. 2) showed seasonal trends with a

TABLE II Statistics of the measured variables. See the text for acronyms

Variables	Mean	$\pm s.d.$	Min	Max
T(°C)	18.8	5.5	11.2	29.1
S (psu)	38.6	3.2	32.8	47.1
OSM (mg l ⁻¹)	1.8	1.6	0.1	10.1
ISM (mg l ⁻¹)	4.3	5.5	0.7	28.4
CHL-a ($\mu\text{g l}^{-1}$)	0.4	0.2	0.0	1.3
PHAEO ($\mu\text{g l}^{-1}$)	0.2	0.1	0.0	1.7
DOC (mg l ⁻¹)	2.1	2.0	0.0	8.5
POC ($\mu\text{g l}^{-1}$)	211.6	106.0	22.5	1,344
$\delta^{13}\text{C}$ (‰)	-18.8	2.5	-25.1	-13.7




FIGURE 2 Monthly trend of temperature (°C) and Salinity (psu) in the Stagnone. Standard deviations are reported.

significant peak in summer and lower values between November and March. No significant differences occurred between open sea and inner sectors but a qualitative positive gradient was observed along a north-south transect.

Organic (OSM) and inorganic (ISM) fractions of total suspended matter (TSM) were significantly correlated ($n = 60$; $r_s = 0.74$; $p < 0.05$) and showed a peak in July (Fig. 3a). Maximum values were reached in middle and eastern stations whilst the lowest values occurred in the southern basin (Fig. 3b), although no significant spatial differences were observed.






FIGURE 3 Monthly trend (a) and spatial pattern (b) of total suspended matter (TSM; mg l^{-1}) in the Stagnone. Standard deviations are reported.

The trend of phytoplankton biomass (as chlorophyll-a concentration) (Fig. 4a) was marked by two primary peaks (May and October) and by a secondary peak (December), lowest values being reached in February. Chlorophyll-a concentration reached maximum values in

the open-sea and in the northern and middle basin (Fig. 4b). Phaeopigments showed a primary peak in correspondence to chlorophyll-a in May; secondary peaks were observed in November and



(a)



(b)

FIGURE 4 Monthly trend (a) and spatial pattern (b) of chlorophyll-a and phaeopigments (CHL-a and Phaeo; $\mu\text{g l}^{-1}$) in the Stagnone. Standard deviations are reported.

January with some time-delay respective to chlorophyll-a (Fig. 4a). Spatial pattern was different to chlorophyll-a, reaching a maximum value in the northern basin and a minimum value in the middle sector (Fig. 4b).

Dissolved organic carbon showed two marked loading periods in spring and summer (Fig. 5a), both of which preceded the chlorophyll-a peaks. DOC maximum values were reached in the middle and southern sectors, whilst they were minimum in the open sea and in northern sectors (Fig. 5b).

Particulate organic carbon showed maximum values in late spring and in winter (Fig. 6) and minimum values in summer and autumn. The POC pattern was different to the DOC in the other stations, except for the highest values in middle sector. The $\delta^{13}\text{C}_{(\text{POC})}$ values ranged between -25.1‰ to -13.7‰ (Tab. II) with the highest in summer (Fig. 6). Significant differences in $\delta^{13}\text{C}_{(\text{POC})}$ were found between the open sea and sites within the Stagnone with higher values in the middle station, corresponding to a maximum in POC concentration (Fig. 7).

Isotopic analyses were carried out on some organic carbon sources: seagrass (*Posidonia* and *Cymodocea*; $\delta^{13}\text{C} = -7\text{‰}$), macroalgae




FIGURE 5. Monthly trend (a) and spatial pattern (b) of dissolved organic carbon (DOC; $\mu\text{g l}^{-1}$) in the Stagnone. Standard deviations are reported.





FIGURE 5 (Continued).

FIGURE 6 Monthly trend of particulate organic carbon (POC; µg l⁻¹) and δ¹³C_{POC} (‰) in the Stagnone. Standard deviations are reported.

(*Caulerpa*, *Ritipilea*, *Rodophicaceae*, *Laurencia*, *Halimeda* and *Cystoseira* sp; δ¹³C = -13.5‰) and plankton (phytoplankton and zooplankton; δ¹³C = -21/-22 ‰).




FIGURE 7 Spatial pattern of carbon stable isotope ratio of particulate organic carbon ($\delta^{13}\text{C}_{\text{POC}}$, ‰) vs. concentration of particulate organic carbon (POC, $\mu\text{g l}^{-1}$) in the Stagnone.

4. DISCUSSION

The Stagnone of Marsala is a Mediterranean semi-enclosed marine system, characterized by the absence of chemical–physical gradients (*i.e.*, temperature and salinity). The low chlorophyll-*a* concentrations (*i.e.*, $\text{CHL} - a < 1.0 \mu\text{g l}^{-1}$) were similar to these found in the adjacent sea. DOC mean concentration (2.14 mg l^{-1}) was lower than that of Egadi Islands (4.54 mg l^{-1} ; Sarà, in preparation), but higher in other Mediterranean values (1.5 mg l^{-1} ; Copin-Montegut and Avril, 1993) or in other oceanic or coastal values (Mantoura and Woodward, 1983; Sugimura and Suzuki, 1988; Guo *et al.*, 1995). The absence of any differences in DOC concentrations between the sea and the sound may show that the trophic background similar and consequently each local escape is probably due to physical constraints and/or to *in situ* production and decomposition processes.

Unlike “true” lagoons (Pusceddu *et al.*, 1996; Friligos *et al.*, 1989; Chassany de Casabianca *et al.*, 1995) or estuaries (Painchaud *et al.*, 1995; Savenkof *et al.*, 1995; Clavier *et al.*, 1995), where chlorophyll-*a* and DOC concentrations are higher than in the adjacent sea, and

where often in the whole of the year, the living fraction (mostly labile) represents most of the particular organic carbon, in the Stagnone, the carbon content of the living biomass rarely exceeds 10% of a non-living fraction. Relatively high DOC concentration, high photosynthetic rates ($2 \text{ mg C m}^{-3} \text{ h}^{-1}$; Magazzù, 1977) and a significant correlation between chlorophyll-a and POM concentrations (Sarà *et al.*, 1995) could show that suspended pigments could be derived from a living planktonic component with a role in the POC pool (Pusceddu *et al.*, 1997; Sarà *et al.*, in press). Nevertheless, Maimone *et al.* (1997) have documented dominance of tytoplanktonic cells (especially benthic diatoms) in the water column. The remaining part of POC is probably of detritic or heterotrophic nature (Sarà *et al.*, in press).

Stable carbon isotopes may discriminate among the main carbon sources affecting the POC pool. The isotopic signature of macroalgae and plankton was quite similar to that documented in other marine areas, whilst *Posidonia oceanica* and *Cymodocea nodosa* appear to be slightly enriched ($\sim 1.5\%$ to -7.5%) compared to the literature data (Dauby, 1989; Jennings *et al.*, 1997). Such enrichment could be determined by higher temperature and sun irradiance (Hemmings and Mateo, 1996) due to the shallowness of the studied area compared to other investigated areas (Tufano, 1991; Dauby, 1989; Jennings *et al.*, 1997). Seagrass values were higher (Tab. III), although within in the bibliographic range (Fry, 1984; Mann, 1988; McMillan, 1980). Different isotopic values occurred as a function of different trophic processes prevailing in the different sites.

Principal component analysis summarizes the trophic picture of the Stagnone (Fig. 8).

The open sea site was characterized by values and a seasonal fluctuation in POC concentration similar to those reported by Fabiano *et al.* (1996) for various sites in the south-western Mediterranean. It showed phytoplanktonic production periods in spring and autumn and more negative $\delta^{13}\text{C}_{(\text{POC})}$ values (-21% to -22%) which were similar to those reported in other areas or similar to that of batch-cultured phytoplankton (Gearing *et al.*, 1984; Faganeli *et al.*, 1989; Lee Van Dover *et al.*, 1992). It may be possible to identify the planktonic component as the major source of POC due to a greater depth and unvegetated bottom (Figs. 8, 9). Conversely, Campolmi *et al.* (1997, 1998) and Maimone *et al.* (1997) have documented the

TABLE III Comparison of $\delta^{13}\text{C}$ values of different carbon sources in different environments

Source	Values	Environment	References
Phytoplankton	-18/-24	Open sea	Fry, 1984
Phytoplankton	-22	Shallow	Tufano, 1991
Benthic unicellular algae	-10/-20	Estuary	Haines and Montague, 1979
Benthic diatoms	-15/-16	Estuary	Riera and Richard, 1996
Phytoplankton and diatoms	-20/23	Estuary	Gearing <i>et al.</i> , 1984
<i>Cymodocea nodosa</i>	-7.20	Shallow	This paper
<i>Cymodocea nodosa</i>	-9.30	Shallow	Hemmings and Mateo, 1996
<i>Posidonia oceanica</i>	-7.80	Shallow	This paper
Phanerogames detritus	-4.50	Shallow	This paper
<i>Posidonia detritus</i>	-11.20	Shallow	Tufano, 1991
<i>Posidonia oceanica</i>	-11.80	Shallow	Tufano, 1991
<i>Posidonia oceanica</i>	-11.90	Shallow	Hemmings and Mateo, 1996
Macroalgae	-17/-20	Estuary	Riera and Richard, 1996
Macroalgae	-13.5	Shallow	This paper
Sediment - open sea - Stagnone	-17.50	Shallow	Mazzola <i>et al.</i> , submitted
Sediment - exposed site - Stagnone	-15.00	Shallow	Mazzola <i>et al.</i> , submitted
Sediment - sheltered site - Stagnone	-11.00	Shallow	Mazzola <i>et al.</i> , submitted




FIGURE 8 Principal components analysis: point-sector by point-month qualitative two-dimensional ordination as a function of correlation with F1 axis (contributing to F1 axis [35%] = $\delta^{13}\text{C}_{\text{POC}}$ -86%; *Cymodocea nodosa* -85%; *Caulerpa prolifera* -45%) and F2 axis (contributing to F2 axis [26%] = [CPE] 62%; [POC] 50%; *Posidonia* -73%).




FIGURE 9 Open-sea - monthly trend of carbon stable isotope ratio in comparison to CPE concentration ($\mu\text{g l}^{-1}$).

dominance of planktonic (phyto and zooplankton) species which are typical of the Mediterranean open-sea, even if Campolmi (1998) had found a tytoplanktonic peak (MacIntyre *et al.*, 1996) which might produce the evident enrichment in isotopic composition in December, whilst the depletion found in January could be produced by a terrestrial contribution. The prevalence of planktonic organic carbon could indicate a greater availability of more labile organic matter in the water column (Mann, 1988; Fichez, 1991).

Unlike the open sea, shallowness and presence of submersed vegetation may affect the POC isotopic composition inside the Stagnone. The distribution of suspended organic matter was determined by hydrodynamic constraints (wind-waves energy and small tides) (Sarà *et al.*, submitted). Resuspension from exposed sites and sedimentation into sheltered sites controlled the biological dynamics present in the Stagnone (Pusceddu *et al.*, 1997; Pusceddu *et al.*, in press), depending on the wind direction and intensity. The dominance of the first or the second process determined the vegetation typology and the degree of cover, allowing us to divide the Stagnone ecosystem into different sub-systems (Sarà *et al.*, in press) where origin and distribution of particulate organic carbon can differ (Fig. 8).

The northern basin and eastern channel could represent the first sub-system which being exposed to dominant winds (Sarà *et al.*, in press), and is characterized by the pulsing fluctuations of the pool of organic matter. These sites could produce as an effect, a limited vegetation (Scilipoti *et al.*, 1997) as affected by a frequently alternating of resuspension and sedimentation events (Kullenberg, 1972, 1976; Levasseur *et al.*, 1983; Millet and Cecchi, 1992). The POC isotopic composition ($\delta^{13}\text{C} = -18.5\text{‰}$) might reflect a contemporary contribution of *Cymodocea nodosa*, *Caulerpa prolifera* and probably microphytobenthos (MacIntyre *et al.*, 1996) as carbon sources. In the northern basin, higher wind exposure could justify chlorophyll-a and phaeopigments peaks, but these peaks correspond to a different vegetal cover degree (high in May and low in January; Fig. 10a; see also Fig. 8), the lightest isotopic ratios (about -21‰) could reflect the signature of planktonic and microbenthic components. In this situation, lateral drifting can be a further factor that could remove *Cymodocea* and *Caulerpa* detritus for storing it. Furthermore, Pusceddu *et al.* (1997, 1998) have documented a positive gradient of sedimentary organic matter along the N-S axis of the Stagnone. The total effect of resuspension-sedimentation-lateral drifting could permit the autotrophic uni-




FIGURE 10a Northern basin - monthly trend of carbon stable isotope ratio in comparison to vegetation type (cover class; Cym = *Cymodocea nodosa*; Cau = *Caulerpa prolifera*) and CPE ($\mu\text{g l}^{-1}$).

cellular biomass marker to become predominant with respect to *Cymodocea* and *Caulerpa*.

This dynamic frame well fits with the eastern channel situation, even if it is characterized by less dominant-wind exposure. A weakening of the resuspension frequency could produce an increase in the sedimentation rate of fine material (Sarà *et al.*, in press). Lower intensity of lateral drifting could explain the greater quantity of suspended material. According to Peres and Picard (1964), the prevalence of a muddy bottom can support *Caulerpa prolifera* which, being more abundant, can affect the composition of isotopic POC. Even if chlorophyll-a concentrations were high (Fig. 10b; see also Fig. 8), $\delta^{13}\text{C}_{(\text{POC})}$ values are always heavier. Confirming the weakening in resuspension frequency, more positive values resulted shifted with respect to peaks in *C. prolifera* (March and June) meaning that this alga probably affected the composition of isotopic POC, especially when it is successively resuspended after its storage in the sedimentary detritus pool.

In this context and even if Gambi and Di Meglio (1996) have documented high toughness of *Caulerpa prolifera*, the identification




FIGURE 10b Eastern channel – monthly trend of carbon stable isotope ratio in comparison to vegetation type (cover class; Cym = *Cymodocea nodosa*; Cau = *Caulerpa prolifera*) and CPE ($\mu\text{g l}^{-1}$).

of components which are more easily degradable (plankton and macroalgae) as carbon sources could support a hypothesis that the organic matter available is mostly labile (Mann, 1988).

Middle area and southern basin represent the second sub-system which are constrained by different hydrodynamic processes controlling the POC origin. These sites are characterized by a vitality of waters (Sarà *et al.*, in press) permitting a wide presence of seagrass (*Posidonia oceanica* and *Cymodocea nodosa*). In the absence of "unpredictability" due to wind pulsing, seasonal regimes predominate and are correlated to the trend in seagrass leaf component. The main source of organic material in the POC compartment is the detritus from phanerogames. In particular, the middle site represents the area where the sedimentation-resuspension process has been inverted. This sheltered site was probably affected by hydrodynamics, which was produced by currents coming from both the northern and southern areas. Apart from two phytoplankton production periods (May and October), when $\delta^{13}\text{C}_{(\text{POC})}$ values decreased (Fig. 10c; see also Fig. 8), POC isotopic composition followed the seasonality of seagrass, showing heaviest values (spring and summer) in correspondence to peaks in leaf biomass and lightest values in autumn and winter when the biomass is low (Mazzella and Ott, 1984; Velimirov, 1987).




FIGURE 10c Middle basin - monthly trend of carbon stable isotope ratio in comparison to vegetation type (cover class; Cym = *Cymodocea nodosa*; POS = *Posidonia oceanica*) and CPE ($\mu\text{g l}^{-1}$).




FIGURE 10d Southern basin – monthly trend of carbon stable isotope ratio in comparison to vegetation type (cover class; Cym = *Cymodocea nodosa*; POS = *Posidonia oceanica*) and CPE ($\mu\text{g l}^{-1}$).

The southern area was also affected by inputs of sea water with tidal periodicity coming from the southern mouth (P. Palermo) (Sarà *et al.*, 1995; Campolmi *et al.*, 1997, 1998). The reduced phenomena of re-suspension (being the sheltered site) and tidal periodicity of water movements caused the dominance of *Posidonia* with respect to *Cymodocea*. Seasonal dynamics of *Posidonia* strongly affected the composition of isotopic POC except in winter and spring when plankton contribution can predominate. In December (Fig. 10d; see also Fig. 8), the POC composition is probably affected by input of terrigenous material (Fry, 1984) which can lower the composition of isotopic POC. On the other hand, the density of zooplankton community is very scanty (Campolmi, 1998).

Thus, the prevalence of organic carbon which is more resistant to processes of degradation as that derived from seagrass (Pirc and Wollemweber, 1988), and could support the hypothesis of the low availability of organic matter is mostly refractory (Mann, 1988).

5. CONCLUDING REMARKS

The quality and quantity of organic matter and the carbon isotopic composition permit the discrimination between the main carbon sources

and the main physical processes, which regulate their distribution. In this shallow environment, relationships among the main hydrodynamic processes (resuspension-sedimentation-lateral drifting) and carbon sources could define the main factors determining the quality of suspended organic matter. Resuspension events seems to increase the level of lability (and consequently the availability for consumers; cfr. Newell, 1982) of organic matter which is characterized by the lightest isotopic values according to Hopkinson (1985) who said that resuspension appears to control the relative amounts of organic carbon as well as the sites and rates of organic matter degradation in the benthos and water column. By contrast, the predominance of sedimentation processes could produce organic matter, which is mostly refractory (thereby lowering the availability for consumers), with heavier values.

Nevertheless, further information about other factors, including nitrogen and sulphur isotope ratios, is necessary for a more complete analysis of the relationships between biotic and abiotic components in this Mediterranean area.

Acknowledgements

The authors would like to thank Prof. Bryan Fry (US Department of Interior; Geological Survey) and Dr. R. H. Michener (Stable Isotope Laboratory, Boston University, USA) and their proof-reader (Helen Maine, Edinburgh UK) for their fundamental help in improving this manuscript. This work has been funded by the Ministero Università Ricerca Scientifica e Tecnologica (MURST, Italy) and by Ministero Politiche Agricole (Italy).

References

- Albertelli, G. and Fabiano, M. (1990) Caratteristiche trofiche in una stazione costiera in Mar Ligure. *Italian Oceanography and Limnology Ass.*, pp. 59–66.
- Calvo, S., Ciraolo, G., La Loggia, G., Malthus, T. J., Savona, E. and Tomasello, A. (1996) Monitoring *Posidonia oceanica* meadows in the Mediterranean Sea by means of airborne remote sensing techniques. *International Airborne Remote Sensing Conference Exhibition*, June 1996 San Francisco, USA. 3, 659–668.
- Campolmi, M. (1998) Zooplankton community in the Stagnone di Marsala (West MED). *Ph.D. Dissertation*, University of Messina, Italy. p. 205.
- Campolmi, M., Zagami, G. and Guglielmo, L. (1998) Zooplankton space-time distribution in a shallow coastal area (Stagnone di Marsala, Western Sicily). *Biologia Marina Mediterranea* (in press).

- Chassany de Casabianca, M. L., Boonne, M. L. and Semroud, R. (1995) Principal component analysis of relationship between physicochemical variables in a Mediterranean lagoon. *C.R. Hebd. Seances Acad. Sci. (III)*, Paris, **310**, 397–403.
- Clavier, J., Chardy, P. and Chevillon, C. (1995) Sedimentation of particulate matter in the south-west lagoon of New Caledonia: Spatial and temporal pattern. *Estuarine, Coastal Shelf Science*, **40**, 281–294.
- Copin-Montegut, G. and Avril, B. (1993) Vertical distribution and temporal variation of dissolved organic carbon in the North-western Mediterranean Sea. *Deep-Sea Research*, **40**, 1963–1972.
- Dauby, P. (1989) The stable isotope ratios in benthic food webs of the Gulf of Calvi, Corsica. *Continental Shelf Research*, **9**, 181–195.
- Fabiano, M., Povero, P., Accornero, A., Mistic, C. and Ossola, C. (1996) Sostanza organica e nutrienti nel Tirreno Sud Orientale (sistema Eolie e bacini di Cefalù e Gioia). In: Faranda, F. M. and Povero, P. (Eds.), *Caratterizzazione ambientale del sistema Eolia e Dei bacini limitrofi di Cefalù e Gioia*, (EOCUMM95). CoNISMa Data Rep., pp. 115–134.
- Faganeli, J., Malej, A., Pezdic, J. and Malacic, V. (1988) C:N:P ratios and stable C isotopic ratios as indicators of sources of organic matter in the Gulf of Trieste (Northern Adriatic). *Oceanologica Acta*, **11**, 377–382.
- Faganeli, J., Pezdic, J. and Fanuko, N. (1989) Chemical and isotopic composition of some phytoplankton species. *Bioloski Vestnik*, **37**, 9–18.
- Fegley, S. R., MacDonald, B. A. and Jacobsen, T. R. (1992) Short-term variation in the quantity and quality of seston available to benthic suspension feeders. *Estuarine, Coastal and Shelf Science*, **34**, 393–412.
- Fichez, R. (1991a) Composition and fate of organic matter in submarine cave sediments: implications for the biogeochemical cycles of organic carbon. *Oceanologica Acta*, **14**(4), 369–378.
- Flury, B. (1988) Common Principal Components and Related Multivariate Analysis. John Wiley and Sons, New York., p. 258.
- Friligos, N. (1989) Nutrients status in a eutrophic Mediterranean lagoon. *Vie Milieu*, **39**(2), 63–69.
- Fry, B. (1984) $^{13}\text{C}/^{12}\text{C}$ ratios and the trophic importance of algae in Florida *Syringodium filiforme* seagrass meadow. *Marine Biology*, **79**, 11–19.
- Fry, B. and Sherr, E. B. (1984) $\delta^{13}\text{C}$ measurements and indicators of carbon flow in marine and freshwater ecosystems. *Contr. Mar. Sci.*, (Univ. Texas), **27**, 13–47.
- Gambi, M. C. and Di Meglio, A. (1996) Observations of feeding behaviour and food choice in the herbivore polychaete. *XI Italian Limnology and Oceanography Symposium*, Sorrento, Italy 1994, pp. 779–791.
- Gearing, J. N., Gearing, P. J., Rudnick, D. T., Requejo, A. G. and Hutchins, M. J. (1984) Isotopic variability of organic carbon in a phytoplankton-based, temperate estuary. *Geochimica Cosmochimica Acta*, **48**, 1080–1089.
- Giani, M., Mecozzi, M., Gesumundo, C., Ciuffa, G., Sunseri, G. and Andaloro, F. (1995) Physico-chemical features and nutrients distribution in the Marsala lagoon (Italy). *Rapp. Comm. Int. Mer Médit.*, **34**, 81.
- Guo, L., Santschi, P. H. and Warnken, K. W. (1995) Dynamics of dissolved organic carbon (DOC) in oceanic environments. *Limnology and Oceanography*, **40**, 1392–1403.
- Haines, E. B. and Montague, C. L. (1979) Food sources of estuarine invertebrates analyzed using $^{13}\text{C}/^{12}\text{C}$ ratios. *Ecology*, **60**, 48–56.
- Hansen, J. A., Klump, D. W., Alongi, D. M., Dayton, P. K. and Riddle, M. J. (1992) Detrital pathways in a coral reef lagoon. II. Detritus deposition, benthic microbial biomass and production. *Marine Biology*, **113**, 363–372.
- Helsel, D. R. and Hirsch, R. M. (1992) Statistical Methods in Water Resources. Elsevier, p. 522.

- Hemmings, M. A. and Mateo, M. A. (1996) Stable carbon isotopes in seagrasses: variability in ratios and use in ecological studies. *Marine Ecology Progress Series*, **140**, 285–298.
- Hopkinson, C. S. Jr. (1985) Shallow water benthic and pelagic metabolism: evidence of heterotrophy in the nearshore Georgia Bight. *Marine Biology*, **87**, 19–32.
- Iseki, K., McDonald, R. W. and Carmack, E. (1987) Distribution of particulate matter in the south-eastern Beaufort Sea in late summer. *Polar Biology*, **1**, 35–46.
- Jennings, S., Renones, O., Morales-Nin, B., Polunin, N. V. C., Moranta, J. and Coll, J. (1997) Spatial variation in the ^{15}N and ^{13}C stable isotope composition of plants, invertebrates and fishes on Mediterranean reefs: implications for the study of trophic pathways. *Marine Ecology Progress Series*, **146**, 109–116.
- Kullemberg, G. (1972) Apparent horizontal diffusion in stratified vertical shear flow. *Tellus*, **24**, 17–28.
- Kullemberg, G. (1976) On vertical mixing and the energy transfer from the wind to the water. *Tellus*, **28**, 159–165.
- Lee Van Dover, C., Grassle, J. F., Fry, B., Garritt, R. H. and Staczak, V. R. (1992) Stable isotope evidence for entry of sewage-derived organic material into a deep-sea food web. *Nature*, **360**, 153–156.
- Levasseur, M., Therrhault, J. C. and Legendre, L. (1983) Tidal currents, wind and the morphology of phytoplankton spatial structure. *Journal Marine Research*, **42**, 655–672.
- Lorenzen, C. J. and Jeffrey, S. W. (1980) Determination of chlorophyll and phaeopigments spectrophotometric equations. *Limnology and Oceanography*, **12**, 343–346.
- MacIntyre, H. L., Geider, R. J. and Miller, D. C. (1996) Microphytobenthos: The ecological role of the “secret” garden of unvegetated, shallow water habitats. I. Distribution, abundance and primary production. *Estuaries*, **19**, 186–201.
- McMillan, C. P., Parker, P. L. and Fry, B. (1980) $^{13}\text{C}/^{12}\text{C}$ ratios in seagrasses. *Aquatic Botany*, **9**, 237–249.
- Magazzù, G. (1977) Usefulness of the Marsala lagoon for aquaculture I. Nutrients and primary production. *Rapp. Comm. Int. Mer. Medit.*, **24**, 81–82.
- Maimone, G., Puglisi, A., Campolmi, M., Sarà, G. and Mazzola, A. (1997) Seasonal changes in phytoplanktonic community as a function of chemical and physical features (Stagnone di Marsala, Western Sicily). *Biologia Marine Mediterranea* (in press).
- Mann, K. H. (1988) Production and use of detritus in various freshwater, estuarine and coastal marine ecosystems. *Limnology and Oceanography*, **33**, 910–930.
- Mantoura, R. F. C. and Woodward, E. M. S. (1983) Conservative behaviour of riverine dissolved organic carbon in the Severn estuary: chemical and geochemical implications. *Geochimica Cosmochimica Acta*, **47**, 1293–1309.
- Mazzella, L. and Ott, J. (1984) Seasonal changes in some features of *Posidonia oceanica* (L.) Delile leaves and epiphytes at different depth. In: Boudouresque, C. F., Jeudy, A., de Grissac and Oliver, J., I Intern. Workshop on *Posidonia oceanica* beds, **1**, 119–127. *GIS Posidonie, Marseilles, France*.
- Mazzella, L., Scipione, M. B. and Buia, C. (1989) Spatio-temporal distribution of algal and animal communities in *Posidonia oceanica* meadows. *P.S.Z.N.I.: Marine Ecology*, **10**, 107–130.
- Mazzola, A. and Sarà, G. (1995) Hydrological features of Mediterranean coastal lagoon (Stagnone di Marsala – Western Sicily): qualitative hydrodynamic model. *Naturalista Siciliano*, **19**, 229–277.
- Mazzola, A., Sarà, G. and Michener, R. H. (submitted). The use of carbon stable isotope to investigate the origin and distribution of suspended and sedimentary organic matter in a semi-enclosed Mediterranean marine system. Springer Verlag Special Issue to the National Symposium “Diversità e Cambiamento” held in Ischia (NA, Italy), pp. 11–14 November 1998.

- Michener, R. H. and Shell, D. M. (1994) Stable isotope ratios as tracers in marine aquatic food webs. In: Lajtha, K. and Michener, R. (Eds.), *Stable Isotopes in Ecology and Environmental Sciences*. Blackwell Science, pp. 138–157.
- Millet, B. and Cecchi, P. (1992) Wind-induced hydrodynamic control of the phytoplankton biomass in a lagoon ecosystem. *Limnology and Oceanography*, **37**, 140–146.
- Navarro, J. M., Clasing, E., Urrutia, G., Asencio, F., Stead, R. and Herrera, C. (1993) Biochemical composition and nutritive value of suspended particulate matter over a tidal flat of southern Chile. *Estuarine Coastal Shelf Science*, **37**, 59–73.
- Newell, R. C. (1982) The energetic of detritus utilization in coastal lagoon and in nearshore waters. *Oceanologica Acta, Proceedings International Symposium on Coastal Lagoons*, SCOR/IABO/UNESCO, Bordeaux, France, September 1981, pp. 347–355.
- Painchaud, J., Lafavre, D., Therriault, J.-C. and Legendre, L. (1995) Physical processes controlling bacterial distribution and variability in the upper St. Lawrence estuary. *Estuaries*, **18**, 433–444.
- Painchaud, J., Lafavre, D., Therriault, J.-C. and Legendre, L. (1996) Bacterial dynamics in the upper St. Lawrence estuary. *Limnology and Oceanography*, **41**, 1610–1618.
- Painchaud, J., Lafavre, D., Tremblay, G. H. and Therriault, J.-C. (1990) Analysis of the distribution of suspended particulate matter, bacteria, chlorophyll a and PO₄ in the Upper St. Lawrence Estuary, using a two dimensional box model. In: Michaelis, W. (Editor), *Coastal and Estuarine Studies*, Springer-Verlag, Berlin, **36**, 59–65.
- Pfannkuche, O. (1993) Benthic response to the sedimentation of particulate organic matter at the BIOTRANS station, 47°N, 20°W. *Deep-Sea Research*, **40**, 135–149.
- Peres, J. M. and Picard, J. (1964) Nouveau manuel de bionomie benthique de la mer Méditerranée. *Recueil Travaux Stat. Mar. Endoume.*, **31**, 5–138.
- Pirc, H. (1989) Seasonal changes in soluble carbohydrates, starch, and energy content in Mediterranean seagrasses. *P.S.Z.N.I., Marine Ecology*, **10**, 97–105.
- Pirc, H. and Wollemweber, B. (1988) Seasonal changes in nitrogen, free amino acids and C/N ratio in Mediterranean seagrasses. *P.S.Z.N.I., Marine Ecology*, **9**, 167–179.
- Posedel, N. and Faganeli, J. (1991) Nature and sedimentation of suspended particulate matter during density stratification in shallow coastal waters (Gulf of Trieste, northern Adriatic). *Marine Ecology Progress Series*, **77**, 135–145.
- Pusceddu, A. and Fabiano, M. (1994) Short-term changes in the biochemical composition of particulate organic matter in a Mediterranean lagoon (southern Sardinia): the role of wind and rain. *Italian Limnology and Oceanography Ass.*, pp. 571–581.
- Pusceddu, A., Sarà, G., Armeni, M., Fabiano, M. and Mazzola, A. (in press) Seasonal and spatial changes in sediment organic matter composition of a semi-enclosed marine system (W-Mediterranean Sea). *Hydrobiologia*. (in press)
- Pusceddu, A., Sarà, G., Mazzola, A. and Fabiano, M. (1997) Relationships between suspended and sediment organic matter in a semi-enclosed marine system: the Stagnone di Marsala (western Sicily). *Water, Air and Soil Pollution*, **99**, 343–352.
- Pusceddu, A., Serra, E., Sanna, O. and Fabiano, M. (1996) Seasonal fluctuations in the nutritional value of particulate organic matter in a lagoon. *Chemistry and Ecology*, **13**, 21–37.
- Rau, G. H., Sullivan, C. W. and Gordon, L. I. (1991) $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ variations in Weddel Sea particulate organic matter. *Marine Chemistry*, **35**, 355–369.
- Sarà, G., Leonardi, M. and Mazzola, A. (in press) Spatial and temporal changes of suspended matter in relation to wind and vegetation cover in a Mediterranean shallow coastal environment. *Chemistry and Ecology*. (in press).
- Sarà, G., Manganaro, A., Cortese, G., Pusceddu, A. and Mazzola, A. (1998) The relationship between food availability and the growth of *Mytilus galloprovincialis* (LMK, 1819) in the open-sea (South Mediterranean Sea). *Aquaculture*, **167**, 1–15.
- Sarà, G. and Mazzola, A. (1997) Effects of trophic and environmental conditions on the *Crassostrea gigas* growth in culture. *Aquaculture*, **153**, 81–91.

- Sarà, G., Pusceddu, A., Mazzola, A. and Fabiano, M. (1995) Variazioni nictemerali della composizione biochimica del materiale organico particellato nello Stagnone di Marsala (Sicilia Occidentale): osservazioni preliminari. *Biologia Marina Mediterranea*, **2**, 127–129.
- Savenkoff, C., Chanut, J. P., Vézina, A. F. and Gratton, Y. (1995) Distribution of biological activity in the St. Lawrence Estuary as determined by multivariate analysis. *Estuarine Coastal Marine Science*, **40**, 647–664.
- Scilipoti, D. (1997) Fish community in the Stagnone of Marsala: distribution and resources repartition as a function of different habitat complexity degrees. *Ph.D. Dissertation*. University of Messina (Italy), p. 256.
- Scilipoti, D., Franzoi, P., Sarà, G. and Mazzola, A. (1997) Fish community in the Stagnone di Marsala sound (western Sicily): relationship with chemical-physical parameters and vegetal cover. *9th International Congress European Ichthyologist (CEI19) "To Review the History and Present Status of the World's Fishfauna, by Multidisciplinary Approach"*, Trieste (Italy), August 1997.
- Sokal, R. R. and Rohlf, F. J. (1981) *Biometry*. W.H. Freeman & C., New York, p. 859.
- Strickland, J. D. H. and Parsons, T. R. (1972) A practical handbook of sea water analysis. *Bulletin Fishery Research Board Canada*, **167**, pp. 310.
- Sugimura, Y. and Suzuki, Y. (1988). A high-temperature catalytic oxidation method for the determination of nonvolatile dissolved organic carbon in sea water by direct injection of liquid sample. *Marine Chemistry*, **24**, 105–131.
- Tucci, S., Capello, M. and Ferrari, M. (1996) Distribuzione del materiale particolato nel Tirreno Sud orientale. In: Faranda, F. M. and Povero, P. (Editors), *Caratterizzazione Ambientale del Sistema Eolia e dei Bacini limitrofi di Cefalù e Gioia (EOCUMM95)*. CoNISMa Data Rep., pp. 135–140.
- Tufano, M. (1991) Il rapporto isotopico $^{13}\text{C}/^{12}\text{C}$ ($\delta^{13}\text{C}$) nello studio dell'ecosistema bentonico costiero (Mar Ligure). *Ph.D. Dissertation*, University of Genoa (Italy), p. 102.
- Velimirov, B. (1987) Organic matter derived from a seagrass meadow: origin, properties and quality of particles. *P.S.Z.N.I., Marine Ecology*, **8**, 143–173.