

ALKALOIDS OF RAUWOLFIA CANESCENS

A. S. Belikov

Khimiya Prirodnykh Soedinenii, Vol. 5, No. 1, p. 64, 1969

Rauwolfia canescens L. (Trinidad devilpepper) has been introduced by the Transcaucasian Zonal Experimental Station of the All-Union Scientific Research Institute for Medicinal Plants [1]. We have studied the roots and epigeal part of the plant grown in Kobuleti in 1963, 1964, and 1966. The total alkaloid contents were: in the roots 3.1–4.3% and in the epigeal part 1.9–2.4%. By thin-layer chromatography [silica gel with 15% of gypsum, in the chloroform–ethanol (9:1) system], 18 alkaloids have been found in the roots and 16 in the epigeal part.

The alkaloids were extracted from the roots with 5% acetic acid, the bases being precipitated with ammonia and extracted with chloroform. By chromatographing the ether-soluble part of the mixture of alkaloids on alumina by a known method [2], ajmalicine with mp 252° C (from aqueous methanol) and ajmaline with mp 205° C (from methanol) were isolated. The fraction of the polar bases was rechromatographed and chloroform containing 0.5% of ethanol eluted reserpine with a yield of 0.05%, having mp 256–257° C (from ethanol, decomp.). Sarpagine was obtained by Stoll and Hofmann's method [3], mp 304–306° C (from methanol, decomp.).

The alkaloids from the epigeal part of the plant were extracted in the same way as from the roots or were extracted in the form of the bases with benzene, xylene, or dichloroethane. They were then separated by the oxalate method [4]. From the fraction of ethanol-insoluble oxalates, we isolated  $\alpha$ -yohimbine (0.4% of the raw material) with mp 230° C (from ethanol, decomp.). By chromatographing the bases from the fraction of ethanol-soluble oxalates we obtained aricine with mp 189–190° C (from ethanol) and reserpiline, the oxalate of which melted at 239–240° C (decomp.).

The ajmaline, ajmalicine, aricine, reserpiline,  $\alpha$ -yohimbine, and sarpagine were identified by their melting points and UV and IR spectra [5], and reserpine by direct comparison with an authentic sample.

REFERENCES

1. I. M. Rabinovich and Z. I. Chuprova, Nauchnye izvestiya Kazakhskogo meditsinskogo instituta, Alma-Ata, no. 18, 194, 1961.
2. A. Hofmann, Helv. Chim. Acta, **38**, 536, 1955.
3. A. Stoll and A. Hofmann, Helv. Chim. Acta, **36**, 1143, 1953.
4. A. Stoll and A. Hofmann, Helv. Chim. Acta, **38**, 270, 1955.
5. Physical Data of Indole and Dihydroindole Alkaloids, 5-th ed., Eli Lilly, Indianapolis, 1962.

26 July 1968

All-Union Scientific-Research Institute for Medicinal Plants

UDC 547.944/945

ALKALOIDS OF SWERTIA MARGINATA, S. GRACILIFLORA, AND DIPSACUS AZUREUS

T. U. Rakhmatullaev, S. T. Akramov, and S. Yu. Yunusov

Khimiya Prirodnykh Soedinenii, Vol. 5, No. 1, pp. 64–65, 1969

We have studied the alkaloids of Swertia (Gentianaceae) and D. azureus (family Dipsacaceae) (table).

By chloroformic extraction from S. marginata, we obtain 0.30% of ethereal and 0.08% of chloroformic bases. When the combined ethereal bases were treated with acetone, gentiananine [1] was isolated. The mother liquor was passed through a column of alumina. Gentianine [2] was obtained from the ethereal eluates.

From the chloroformic alkaloids, treatment with acetone isolated a yellow base with mp 207–208° C (acetone), which proved to be identical with gentioflavine [3], given to us by S. Popov (Bulgaria) (mixed melting point and IR spectra).