

presence of substituents in positions 5 and 7 of ring A. Apart from these signals, in the weak-field region the spectrum of the substance differs from those of many known flavonoids having substituents in positions 4'; 3',4'; or 3',4',5' [6]. In addition to the doublets mentioned, the spectrum contained three signals of one proton unit each at 6.32, 6.72, and 7.16 ppm. Two of them must be assigned to ring B, i.e., 7.16 ppm (H-2'), 6.72 ppm (H-5'), and 6.32 ppm (H-3). Hence it follows that the substituents in ring B of the compound isolated may be present in positions 3',4',6'. With a different arrangement of the substituents the signals of 2' and 5' protons would not appear in the form of a singlet.

The results of UV spectroscopy confirm the presence of hydroxy groups in positions 3', 4',5,7.

The substituents in position 3' 4',5,6',7 are hydroxy groups, as is shown by the results of UV spectroscopy, the absence of the signals of protons of other substituents in the NMR spectrum, and the mass-spectral results.

The mass spectrum of substance (V) has the molecular peak M 302⁺ (94%) corresponding to the number of substituents in the molecule and, in addition, the peaks of fragments of ring A with m/e 151 (16%) and a strong peak with m/e 153 (100%) corresponding to a fragment of ring B with the heteroatom. The peak with m/e 150 (30%) also corresponds to ring B with three hydroxy groups. Thus, the substance isolated has the structure of 3',4',5,6',7-penta-hydroxyflavone, which has not been described in the literature, and we have called it hier-acin.

LITERATURE CITED

1. N. A. Kaloshina, Abstracts of lectures at the IInd Congress of Pharmacists of the Belorussian SSR [in Russian], Minsk (1970), p. 70.
2. M. Haag-Berrurier and P. Duquéniois, *Compt. Rend.*, **257**, No. 21, 3239 (1963).
3. V. L. Shelyuto, V. I. Glyzin, A. I. Ban'kovskii, and N. T. Bubon, *Khim. Prirodn. Soedin.*, **371** (1971).
4. S. A. Medvedeva and N. A. Tyukavkina, *Khim. Prirodn. Soedin.*, **676** (1972).
5. T. A. Geissman, *The Chemistry of Flavonoid Compounds*, Pergamon, New York (1962), p. 424.
6. T. J. Mabry, K. R. Markham, M. B. Thomas, *The Systematic Identification of Flavonoids*, Springer, New York (1970), p. 254.

FLAVONIDS OF *Galinsoga parviflora*

T. I. Plekhanova, V. A. Bandyukova, and G. A. Mikhailova

UDC 547.972

We have studied the chemical composition of the epigeal part of *Galinsoga parviflora* Cav., collected in the flowering period in the region of Pyatigorsk. In the leaves we have found saponins of the triterpene series, polyphenolic compounds, tanning substances, and inulin. The amount of "crude" saponins was 1.59% (on the air-dry weight of raw material), of tanning substances 2.4% (by a standard method [1]), and of flavonoids 2.4% [2]. To isolate the inulin, the raw material was extracted with water and the inulin was precipitated with ethanol; its amount was 10.7% (mean of three determinations).

By paper chromatography, no less than five substances of flavonoid nature were detected in the ethanolic extracts. Acid hydrolysis showed the presence of three aglycones, two of which were identified as apigenin and luteolin. To isolate the glycosides, 1 kg of air-dry leaves was exhaustively extracted with 70% ethanol on the boiling water bath, the ethanolic extracts were concentrated, and the residue was freed from lipophilic impurities and treated with boiling water. The aqueous solution obtained was washed with chloroform, and the flavonoids were extracted with ethyl acetate.

After the elimination of the ethyl acetate, the residue was dissolved in ethanol and the flavonoids were separated on a column of polyamide sorbent using increasing concentrations of

Pyatigorsk Pharmaceutical Institute. Translated from *Khimiya Prirodnikh Soedinenii*, No. 6, p. 862, November-December, 1977. Original article submitted June 2, 1977.

aqueous ethanol. In this way, substances (I) and (II) were isolated. Substance (I) formed light yellow acicular crystals with mp 254°C. UV spectrum λ_{\max} , nm; C₂H₅OH: 255, 266 sh., 350; C₂H₅ONa: 262, 300 sh., 396; AlCl₃: 274, 300 sh., 330, 432; AlCl₃ + HCl: 274, 294 sh., 358, 386; CH₃COOH: 258, 266 sh., 365 sh., 405; CH₃COONa + H₃BO₃: 260, 372. Acid hydrolysis yielded luteolin and D-glucose. The substance was identified as luteolin 7- β -D-glucopyranoside. Substance (II) formed pale yellow crystals with mp 178-180°C. UV spectrum, λ_{\max} , nm; C₂H₅OH: 268, 335; C₂H₅ONa: 269, 386; AlCl₃: 276, 300, 348, 386; AlCl₃ + HCl: 277, 299, 340, 382; CH₃COONa: 267, 355, 386. Acid hydrolysis yielded apigenin and D-glucose. The substance was identified as apigenin 7- β -D-glucoside.

We have detected caffeic acid in the same plant by paper chromatography.

LITERATURE CITED

1. State Pharmacopoeia of the USSR [in Russian], Xth Ed., Moscow (1968).
2. A. R. Guseva and M. N. Nestyuk, *Biokhimiya*, 18, No. 4, 480 (1953).

THE STRUCTURES OF ISOFLAVONE C-GLYCOSIDES FROM *Lupinus luteus*

G. G. Zapesochneya and N. A. Laman

UDC 547.972

Continuing a chemical study of the flavonoids of *Lupinus luteus* L. (European yellow lupine) [1], from its flowers we have isolated two isoflavone glycosides.

Compound (I). C₂₁H₂₀O₁₀ • H₂O, mp 185-189°C (aq. MeOH), $[\alpha]_D^{20} + 24^\circ$ (c 1; MeOH); λ_{\max} ,




Fig. 1. NMR spectra of the trimethylsilyl ethers of genistein 8-C- β -D-glucopyranoside (I) and of orobol 8-C- β -D-glucopyranoside (II) in CCl₄.

All-Union Scientific-Research Institute of Medicinal Plants, Moscow. V. F. Kuprevich Institute of Experimental Botany, Academy of Sciences of the Belorussian SSR, Minsk. Translated from *Khimiya Prirodnikh Soedinenii*, No. 6, pp. 862-864, November-December, 1977. Original article submitted July 11, 1977.