

This article was downloaded by:

On: 15 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Chemistry and Ecology

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713455114>

Mineralization of particulate organic matter derived from coral-reef organisms in reef sediments of the Gulf of Aqaba

Mohammed Rasheed^a; Christian Wild^{bcd}; Carin Jantzen^c; Mohammed Badran^a

^a Marine Science Station, University of Jordan and Yarmouk University, Aqaba, Jordan ^b Max Planck Institute for marine Microbiology, Bremen, Germany ^c GeoBioCenter, Ludwig-Maximilians University, Munich, Germany ^d Ocean Sciences Section, Intergovernmental Oceanographic Commission, Paris, France ^e Zentrum für Marine Tropenökologie, Bremen, Germany

To cite this Article Rasheed, Mohammed , Wild, Christian , Jantzen, Carin and Badran, Mohammed(2006) 'Mineralization of particulate organic matter derived from coral-reef organisms in reef sediments of the Gulf of Aqaba', *Chemistry and Ecology*, 22: 1, 13 – 20

To link to this Article: DOI: 10.1080/02757540500456823

URL: <http://dx.doi.org/10.1080/02757540500456823>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Mineralization of particulate organic matter derived from coral-reef organisms in reef sediments of the Gulf of Aqaba

MOHAMMED RASHEED*†, CHRISTIAN WILD‡§¶, CARIN JANTZEN|| and MOHAMMED BADRAN†

†Marine Science Station, University of Jordan and Yarmouk University, PO Box 195, Aqaba, Jordan

‡Max Planck Institute for marine Microbiology, Celsiusstr. 1, 28359 Bremen, Germany

§GeoBioCenter, Ludwig-Maximilians University, Richard Wagner Str. 10, 80333 Munich, Germany

¶Ocean Sciences Section, Intergovernmental Oceanographic Commission, UNESCO,
1 rue Miollis, 75015 Paris, France

||Zentrum für Marine Tropenökologie, Fahrenheitstr. 6, 28359 Bremen, Germany

(Received 11 July 2005; in final form 4 November 2005)

In situ and laboratory incubation experiments in a fringing reef in the Gulf of Aqaba were performed to study degradation rates of particulate organic matter in reef sediments. Coral mucus, clam eggs, and zooxanthellae were used as model particulate organic compounds for these experiments. Aerobic and anaerobic mineralization rates were calculated by dissolved inorganic carbon (DIC) and O₂ fluxes from the sediments under different particulate organic matter additions. Fast enhancement (approximately twofold) of O₂ and DIC fluxes were found with the addition of coral mucus and clam eggs compared with control incubations without addition. Most of the degradation is believed to have occurred anaerobically rather than aerobically (DIC:O₂ ratios were 4.3–28.1). Higher degradation rates of coral mucus and clam eggs were estimated in carbonate sediment than in silicate sediment (1.2–1.6-fold), which was attributed to the different physical and chemical properties of both sediments. Our study shows the significance of the reef sediment as a suitable site for microbial degradation of particulate organic material excreted from different reef community organisms. This may increase the regeneration of nutrients in the reef environment necessary to sustain high biological productivity.

Keywords: Coral reef; Nutrient; Sediment; Particulate organic matter; Mineralization

1. Introduction

Particulate organic matter released from different benthic organisms in the reef environment represents a key source of energy in the nutrient-poor reef environments [1–5]. Degradation of these matters in the reef environment leads to release of inorganic nutrients like ammonium and phosphate, which support autotrophic growth in the reef [3, 6–7]. Particulate organic

*Corresponding author. Email: m.rasheed@ju.edu.jo

matter may be mineralized in reef sediment [6, 7], in the reef community [8–10], or in reef biota such as reef-dwelling organisms [10–14].

Carbonate sands, the dominating fraction in reef environment, are usually porous and permeable, exhibiting a large surface area. They thus have many suitable sites for bacterial growth [6–7, 15]. In this type of sediment, as a result of interaction between bottom current and permeable carbonate sands, particulate organic matters may be filtered and mineralized through the benthic microbial food chain [1, 10, 16]. Sandy sediments in reef environments cover roughly half of the surface area [10]; therefore, degradation of particulate organic matter in the reef sediment may be considered as the most important process in sustaining primary productivity in these environments.

In this study, we investigated the degradation of coral mucus, clam eggs, and zooxanthellae in carbonate and silicate sediments located in a coral reef environment. Degradation rates have been previously assessed from fluxes of dissolved inorganic carbon (DIC) and oxygen, which can be calculated by laboratory or *in situ* chamber incubation of sediment with measurements of the temporal variation of DIC and O₂ in the water enclosed in the chamber [15, 17–19]. *In situ* incubations do produce more realistic results than the laboratory incubations which suffer from artefacts caused by the removal of the sediment core from its particulate environment and the transport to the laboratory, which may seriously disturb the sediment core. However, in the laboratory, the conditions of the experiment can be controlled exactly. For this reason, we conducted both laboratory and *in situ* experiments with stirred benthic chambers.

2. Materials and methods

2.1 Study site

The study was carried out in a fringing reef of the northern Gulf of Aqaba at about 3 m water depth. The *in situ* chamber experiments took place in a marine reserve close to the Marine Science Station (MSS) in Aqaba (latitude: 29° 27', longitude: 34° 58'). The incubations were conducted at two sediment sites in front of the MSS, the first site covered by carbonate sands (water depth ~2.5 m), and the second site covered by silicate sands (water depth ~2 m). The distance between the two sites was approximately 200 m, and both sites were located within a radius of 5 m adjacent to coral reef. The mean grain size of both sediments was 229 and 559 μm, the organic carbon content was 0.24 and 0.36%, and the permeability was $27 \pm 11 \times 10^{-12} \text{ m}^2$ and $116 \pm 3 \times 10^{-12} \text{ m}^2$ for the silicate and carbonate sands, respectively [6, 20].

2.2 *In situ* and laboratory incubations

Seven *in situ* experiments and one laboratory incubation experiment were performed in order to estimate the degradation rate of different particulate matter in two different types of reef sediments. Fluxes of DIC and O₂ from the sediments were calculated under different organic matter additions and different light conditions (table 1), and used to calculate the mineralization rate of these organic substances. The particulate organic materials were collected as described by Wild *et al.* [20]. A description of the *in situ* incubation chambers can be found in Huettel and Gust [21]. These were transparent cylindrical acrylic containers 19 cm diameter and 30 cm height. Flat plastic lids covered the chambers and were fixed by four stainless steel clips. The lid contained a sampling port with a syringe holder for water samples and another port to replace the sampled water. The water inside the chambers was stirred at 20 rpm by a

Table 1. Summary of *in situ* and laboratory experiments performed for this study.

| Experiment | Sediment used | Light condition | Process | No. of incubated cores and types of organic matter added |
|--------------|---------------|-----------------|--------------------|--|
| In situ A | Carbonate | Dark | Control | 4 control |
| In situ B | Carbonate | Dark | Control and spiked | 2 control 2 with 40 ml of coral mucus |
| In situ C | Carbonate | Dark | Control and spiked | 1 control 1 with 300 ml of coral mucus 1 with 300 ml of zooxanthellae 1 with 200 ml of clam egg |
| In situ D | Carbonate | Dark | Control and spiked | 1 control 1 with 300 ml of coral mucus 1 with 300 ml of zooxanthellae 1 with 200 ml of clam egg |
| In situ E | Carbonate | Dark | Control and spiked | 1 control 2 with 300 ml of coral mucus 1 with 200 ml of clam egg |
| In situ F | Silicate | Dark and light | Control | 4 control |
| In situ G | Carbonate | Dark and light | Control and spiked | 3 control 1 with 300 ml of coral mucus |
| Laboratory A | Carbonate | Dark | Control and spiked | 3 control 3 with 100 ml of coral mucus |
| Laboratory B | Silicate | Dark | Control and spiked | 3 control 3 with 100 ml of coral mucus |

15 cm-diameter flat rotating disk, connected to a DC motor with a gear box. The chambers were inserted gently into the sediments to a depth of about 10 cm and thus included a water column of approximately 20 cm height, corresponding to a water volume of 5.7 l. The lids were then fitted to the chambers and fixed by the clips. Chambers used for dark incubations were covered with tough opaque black plastic foils secured with rubber bands at the top and the bottom. At defined time intervals (figures 1 and 2), 100 ml of water was removed for DIC, and O₂ determination and replaced by ambient sea water. The total duration of the incubations was around 400 min. The sea water volume in the chambers was determined after removing the lid by measurement of the sea water height at four different points in the chamber. For the laboratory incubation, the same chambers were employed under identical conditions as described by Rasheed *et al.* [6, 15]. These were cylindrical acrylic chambers with an inner diameter of 9.5 cm and a height of 40 cm. The lengths of sediment cores were about 20 cm, and the depth of the water column above the sediments was approximately 20 cm. The sediments used for the laboratory experiments were collected from the two sites by diving to the appropriate location, and upon return to the laboratory the sediments were washed with distilled water and dried in an oven at 60 °C. Fluxes of solutes in both the laboratory and *in situ* incubations were evaluated from linear regressions of solute concentrations over time.

2.3 Analytical procedure

Oxygen concentrations in the sea water were analysed using the Winkler titration method [22]. Dissolved inorganic carbon (DIC) concentrations in the water samples were measured using a flow-injection system [23]. Calibration standards were always freshly prepared from NaHCO₃.




Figure 1. Time course of oxygen (μM) and DIC (mM) during different *in situ* incubations (table 1) for the control chambers and spiked chambers. C: control, Muc-300: with 300 ml of mucus added; Muc-40: with 40 ml of mucus added; Cegg: with 200 ml of clam eggs, Zoox: with 300 ml of zooxanthella. The error bars represent the standard deviation of the results from the *in situ* experiments A–E.




Figure 2. Time course of oxygen (μM) and DIC (mM) during different *in situ* incubations (experiments *in situ* F and *in situ* G) of control chambers in silicate and carbonate sediments. C: carbonate chambers; Si: silicate chambers.

3. Results

3.1 Degradation of coral mucus in carbonate sediments

Concentrations of dissolved inorganic carbon, as the end product of organic matter re-mineralization, and dissolved oxygen, as the terminal electron acceptor of the aerobic metabolism, were measured in the reef sands. In dark control chambers as well as in spiked chambers, the DIC concentration increased gradually over time (figure 1), whereas O₂ concentrations decreased gradually in all chambers. The DIC release and oxygen consumption were always higher in the chambers with coral mucus than in the control chambers. Also, the changes in the chambers with high amounts of mucus (300 ml) were greater than those in the chambers with a low amount (40 ml). However, the differences between different chambers were more pronounced in DIC than in O₂. Flux calculations showed that positive DIC fluxes were always higher than the negative O₂ fluxes (table 2).

3.2 Degradation of zooxanthellae and clam eggs in carbonate sediments

The DIC concentration increase and O₂ concentration decrease in the chambers with zooxanthellae addition were very low compared with the control (figure 1). This may indicate low degradation rates of zooxanthellae in carbonate sediments compared with the other organic substances. In the chambers with clam eggs, DIC concentrations increased steadily, and dissolved oxygen concentration decreased (figure 1). The flux results (table 2) were similar compared with coral mucus incubations, with DIC production being higher than oxygen consumption in most chambers.

Table 2. Fluxes of coral mucus and clam eggs (mg C ml⁻¹ substrate m⁻² d⁻¹) inferred from DIC and O₂ fluxes of different *in situ* and laboratory experiments.

| | Sediment | Addition | No. of replicate | DIC flux (mmol m ⁻² d ⁻¹) | O ₂ flux (mmol m ⁻² d ⁻¹) |
|--------------|-----------|---------------|------------------|--|---|
| In situ A | Carbonate | Control | 4 | 58.3 ± 2.9 | -20.4 ± 6.2 |
| In situ B | Carbonate | Control | 2 | 45.6 ± 7.2 | -27.1 ± 4.3 |
| | Carbonate | Mucus | 2 | 91.3 ± 0.1 | -12.5 ± 0.9 |
| In situ C | Carbonate | Control | 1 | 45.6 | -22.5 |
| | Carbonate | Mucus | 1 | 172.4 | -30.0 |
| | Carbonate | Zooxanthellae | 1 | 86.2 | -13.6 |
| | Carbonate | Clam eggs | 1 | 121.7 | -33.5 |
| In situ D | Carbonate | Control | 1 | 45.6 | -19.8 |
| | Carbonate | Mucus | 1 | 157.2 | -37.0 |
| | Carbonate | Zooxanthellae | 1 | 106.5 | -20.1 |
| | Carbonate | Clam eggs | 1 | 142.0 | -34.9 |
| In situ E | Silicate | Control | 1 | 30.4 | -18.3 |
| | Silicate | Mucus | 2 | 103.9 ± 10.8 | -25.5 ± 10.1 |
| | Silicate | Clam eggs | 1 | 116.6 | -20.6 |
| In situ F | Silicate | Dark | 2 | 38.0 ± 3.6 | -15.2 ± 1.0 |
| | Silicate | Light | 2 | -35.5 ± 7.2 | 15.6 ± 0.7 |
| In situ G | Carbonate | Dark | 2 | 88.7 ± 46.6 | -20.1 ± 2.6 |
| | Carbonate | Light | 2 | -38.0 ± 3.6 | 19.5 ± 6.9 |
| Laboratory A | Carbonate | Control | 3 | 29.0 ± 1.5 | -7.6 ± 1.6 |
| | Carbonate | Mucus | 3 | 48.0 ± 1.9 | -13.14 ± 2.1 |
| Laboratory B | Silicate | Control | 3 | 23.9 ± 2.8 | -6.6 ± 1.5 |
| | Silicate | Mucus | 3 | 42.8 ± 3.7 | -11.2 ± 1.7 |

3.3 Benthic metabolism in carbonate and silicate sediments under light and dark conditions

Degradation of organic matter found naturally in carbonate and silicate sediments has been investigated to study the different rates with different sediment types. *In situ* and laboratory incubations were performed for this purpose. The *in situ* incubation (figure 2) shows that DIC was produced in the dark chambers and consumed in the light chambers, whereas the opposite was found for O₂ (figure 2). Laboratory incubations (not shown) showed the same trend as during the *in situ* incubation. Flux calculations for both *in situ* and laboratory incubations also showed similar trends, as positive values were obtained for DIC in dark chambers and negative values in light chambers (table 2). Here, also, apart from one exception (light silicate chamber), flux calculations showed that DIC fluxes were always higher than O₂ fluxes irrespective of the direction of the flux. As for the sediment type used in the *in situ* and laboratory incubations, generally higher fluxes were obtained for carbonate sediments compared with silicate sediments (figure 2 and table 2).

4. Discussion

The degradation rate of different substrates in sediment can be inferred either from DIC production or from O₂ consumption. When the degradation of organic matter exclusively occurs aerobically, the ratio between DIC production and O₂ consumption should be 1 [24]. Herein (table 3, figure 3), the mineralization rates of coral mucus and clam eggs illustrate that the ratios between the rates inferred from DIC production and from O₂ consumption are always more than 1 (3.43–28.11). This indicates that the mineralization of the substrates used occurs partially anaerobically in addition to aerobic degradation. When the oxygen concentration is very low, DIC can be produced anaerobically using different electron acceptors such as SO₄²⁻ and NO₃⁻ [25–26]. This was confirmed by several authors; Canfield *et al.* [27] estimated a minor contribution of oxygen in organic-matter degradation (4–17%), and Rasheed *et al.* [15] found DIC:O₂ flux rates of 5–33. Also, Wild *et al.* studied the degradation rate of mucus in permeable sediments and found DIC production to be twice as high as O₂ consumption. However, DIC production might also be enhanced by the precipitation and dissolution of carbonate sediments used in the incubation experiments of this type of study [28–30].

Table 3. Mineralization rates of coral mucus and clam eggs (mg C ml⁻¹ substrate m⁻² d⁻¹) inferred from DIC and O₂ fluxes of different *in situ* and laboratory experiments.

| Experiment | Addition | DIC | O ₂ | Ratio |
|--------------|-----------|-------|----------------|-------|
| In situ B | Mucus | 13.71 | -4.38 | |
| In situ C | Mucus | 5.07 | 0.4 | 12.78 |
| | Clam eggs | 4.56 | 0.88 | 5.21 |
| In situ D | Mucus | 4.46 | 0.92 | 4.85 |
| | Clam eggs | 5.78 | 1.21 | 4.78 |
| In situ E | Mucus | 2.94 | 0.38 | 7.66 |
| | Clam eggs | 5.17 | 0.18 | 28.11 |
| Laboratory A | Mucus | 2.28 | 1.11 | 3.43 |
| Laboratory B | Mucus | 2.27 | 0.92 | 4.11 |

Note: The calculations were performed by subtracting the control flux (table 2) from the average flux values of each substrate and then multiplying by the atomic weight of carbon or oxygen and dividing by the volume of each substrate in ml. The negative value indicates a higher chamber O₂ control consumption than the chamber with substrate.




Figure 3. Mineralization rates of coral mucus and clam eggs (mg C ml^{-1} substrate m^{-2} d^{-1}) inferred from DIC and O_2 fluxes of different *in situ* and laboratory experiments. Error bars represent the standard deviation of the *in situ* experiments C, D, E and the laboratory experiments A and B.

Minor differences existed between the mineralization rate of coral mucus estimated in the laboratory experiments and the *in situ* experiments (table 3, figure 3). These can be attributed to differences in the experimental conditions [6]. However, mineralization rates of coral mucus and eggs estimated from *in situ* or laboratory experiments in carbonate sediments were always higher than those in silicate sediments (~ 1.2 – 1.6 -fold). This was also reported by Rasheed *et al.* [6], who found a 1.5–3.8-fold higher degradation rate for *Spirulina* in carbonate sediments than in silicate sediments. The authors attributed this to the different mineralogical structure, larger carbonate surface area, higher permeability, and porosity of carbonate sand. In addition, higher numbers of fauna and flora live in the carbonate sands compared with the silicate sediments [7]. Besides high bacteria found in carbonate sand; cell number: 10^9 – 10^{10} cm^{-3} as compared with silicate sediments [5] may also facilitate degradation processes aerobically and anaerobically.

This investigation clearly shows the importance of sediment as a major incubator of organic-matter degradation. These particulate organic materials are produced by different organisms living in the coral-reef community. A significant fraction of these organic products is reworked by the coral reef itself [3]. The fraction that reaches the sediment does not stay there inert [4–5, 31] but becomes partially mineralized to inorganic nutrients such as ammonium and phosphate, which support a high level of biological productivity in the nutrient-deficient coral-reef environment.

Acknowledgements

Thanks are due to Marine Science Station staff in Aqaba-Jordan for their help in the laboratory and in the field. This work has been funded by the Deanship of Academic Research in Yarmouk University, Higher Council of Science and Technology, Jordan and Max Planck Society, Germany.

References

- [1] K. Grange. Mutualism between the Antipatharian *Antipathes fiordensis* and the Ophiuroid *Astrobrachion constrictum* in New Zealand fjords. *Hydrobiologia*, **216**, 297–303 (1991).
- [2] B. Rinkevich, Z. Wolodarsky, Y. Loya. Coral crab association—a compact domain of a multilevel trophic system. *Hydrobiologia*, **216**, 279–284 (1991).

- [3] C. Richter, M. Wunsch, M. Rasheed, I. Kotter, M.I. Badran. Endoscopic exploration of Red Sea coral reefs reveals dense populations of cavity-dwelling sponges. *Nature*, **413**, 726–730 (2001).
- [4] C. Wild, M. Huettel, A. Kluefer, S.G. Kremb, M. Rasheed, J.B. Bøslash rgenen. Coral mucus functions as an energy carrier and particle trap in the reef ecosystem. *Nature*, **428**, 66–70 (2004a).
- [5] C. Wild, M. Rasheed, U. Werner, U. Franke, R. Johnstone. Degradation and mineralization of coral mucus in reef environments. *Mar. Ecol. Prog. Ser.*, **267**, 159–171 (2004b).
- [6] M. Rasheed, M.I. Badran, M. Huettel. Influence of sediment permeability and mineral composition on organic matter degradation in three sediments from the Gulf of Aqaba, Red Sea. *Est. Coast. Shelf Sci.*, **57**, 369–384 (2003a).
- [7] M. Rasheed, M. IBadran, M. Huettel. Particulate matter filtration and seasonal nutrient dynamics in permeable carbonate and silicate sands of the Gulf of Aqaba, Red Sea. *Coral Reefs*, **22**, 167–177 (2003b).
- [8] C. Richter, M. Wunsch. Cavity-dwelling suspension feeders in coral reefs—a new link in reef trophodynamics. *Mar. Ecol. Prog. Ser.*, **188**, 105–116 (1999).
- [9] G.J. Gast, S. Wiegmann, E. Wieringa, F.C. van Duyl, R.P.M. Bak. Bacteria in coral reef water types: removal of cells, stimulation of growth and mineralization. *Mar. Ecol. Prog. Ser.*, **167**, 37–45 (1998).
- [10] M. Rasheed, M.I. Badran, C. Richter, M. Huettel. Effect of reef framework and bottom sediment on nutrient enrichment in a coral reef of the Gulf of Aqaba, Red Sea. *Mar. Ecol. Prog. Ser.*, **239**, 277–285 (2002).
- [11] R.A.I. Hatcher, B.G. Hatcher. Seasonal and spatial variation in dissolved inorganic nitrogen in One Tree Reef lagoon. *Proc. 4th Int. Coral Reef Symp.*, **1**, 419–424 (1981).
- [12] J.C. Andrews, H.R. Müller. Space–time variability of nutrients in a lagoonal patch reef. *Limnol. Oceanogr.*, **28**, 215–227 (1983).
- [13] G.W. Tribble, F.J. Sansone, Y.-H. Li, S.V. Smith, R.W. Buddemeier. Material fluxes from a reef framework. *Proc. Sixth Int. Coral Reef Symp.*, **2**, 577–582 (1989).
- [14] S.T. Larned. Nitrogen- versus phosphorus-limited growth and sources of nutrients for coral reef macroalgae. *Mar. Biol.*, **132**, 409–421 (1998).
- [15] M. Rasheed, C. Wild, U. Franke, M. Huettel. Benthic photosynthesis and oxygen consumption in permeable carbonate sediments at Heron Island, Great Barrier Reef, Australia. *Est. Coast Shelf Sci.*, **59**, 139–150 (2004).
- [16] M. Huettel, W. Ziebis, S. Forster. Flow-induced uptake of particulate matter in permeable sediments. *Limnol. Oceanogr.*, **41**, 309–322 (1996).
- [17] E. Kristensen. Seasonal variations in benthic community metabolism and nitrogen dynamics in a shallow, organic-poor danish lagoon. *Est. Coastal Shelf Sci.*, **36**, 565–586 (1993).
- [18] D.J. DeMaster, O. Ragueneau, C.A. Nittrouer. Preservation efficiencies and accumulation rates for biogenic silica and organic C, N, and P in high-latitude sediments: The Ross Sea. *J. Geophys. Res. Oceans.*, **101**, 18501–18518 (1996).
- [19] G.J. Nicholson, A.R. Longmore, W.M. Berelson. Nutrient fluxes measured by two types of benthic chamber. *Mar. Freshwat. Res.*, **50**, 567–572 (1999).
- [20] C. Wild, M. Rasheed, C. Jantzen, S. Kremb, P. Cook, U. Struck, M. Huettel, A. Boitius. Benthic metabolism and degradation of particulate organic matter in carbonate and silicate reef sands of the northern Red Sea. *Mar. Ecol. Prog. Ser.* **298**, 69–78 (2005).
- [21] M. Huettel, G. Gust. Solute release mechanisms from confined sediment cores in stirred benthic chambers and flume flows. *Mar. Ecol. Prog. Ser.*, **82**, 187–197 (1992).
- [22] L.W. Winkler. The determination of dissolved oxygen in water. *Ber Deutsch Chem. Ges.*, **21**, 2843–2857 (1888).
- [23] P.O.J. Hall, R.C. Aller. Rapid, small-volume, flow injection analysis for ΣCO_2 and NH_4^+ in marine and freshwaters. *Limnol. Oceanogr.*, **37**, 1113–1119 (1992).
- [24] E. Kristensen. Organic matter diagenesis at the oxic/anoxic interface in coastal marine sediments, with emphasis on the role of burrowing animals. *Hydrobiolog.*, **426**, 1–24 (2000).
- [25] B.B. Jorgensen. The sulfur cycle of a coastal marine sediment (Limfjorden, Demark). *Limnol. Oceanogr.*, **22**, 814–832 (1977).
- [26] B. Thamdrup, R.N. Glud, J.W. Hansen. Manganese oxidation and *in situ* manganese fluxes from a coastal sediment. *Geochim. Cosmochim. Acta*, **58**, 2563–2570 (1994).
- [27] D.E. Canfield, B. Thamdrup, J.W. Hansen. The anaerobic degradation of organic matter in Danish coastal sediments: Iron reduction, manganese reduction, and sulfate reduction. *Geochim. Cosmochim. Acta*, **57**, 3867–3883 (1993).
- [28] M.A. Green, R.C. Aller, J.Y. Aller. Carbonate dissolution and temporal abundances of foraminifera in Long-Island Sound sediments. *Limnol. Oceanogr.*, **38**, 331–345 (1993).
- [29] C.H.R. Heip, N.K. Goosen, P.M.J. Herman, J. Kromkamp, J.J. Middelburg, K. Soetaert. Production and consumption of biological particles in temperate tidal estuaries. *Oceanogr. Mar. Biol. Ann. Rev.*, **33**, pp. 1–149 (1995).
- [30] P. Lopez, M. Vidal, X. Lluch, J.A. Morgui. Sediment metabolism in a transitional continental marine area—the Albufera-Fo-Majorca (Balearic-Islands, Spain). *Mar. Freshwat. Res.*, **46**, 45–53 (1995).
- [31] E. Wolanski, E. Drew, K.M. Abel, J. O'Brien. Tidal jets, nutrient upwelling and their influence on the productivity of the alga *Halimeda* in the Ribbon Reefs, Great Barrier Reef. *Est. Coast Shelf Sci.*, **26**, 169–201 (1988).