

8-ALKYL AND 8-CYCLOALKYL-10-PIPERAZINODIBENZO[*b,f*]THIEPINS*

V. VALENTA, J. METYŠOVÁ, Z. ŠEDIVÝ and M. PROTIVA

Research Institute of Pharmacy and Biochemistry, 130 00 Prague 3

Received June 1st, 1973


Proceeding from 4-ethyl, 4-isopropyl, 4-n-butyl, 4-n-octyl, 4-cyclopropyl and 4-cyclopentyl thiophenols, the corresponding 8-substituted dibenzo [*b,f*] thiepin-10(11*H*)-ones *XIV* were synthesized *via* the intermediates *IX–XIII*. The ketones *XIV* were further converted to the 10-piperazino derivatives of the 10,11-dihydro series *I–VI* and further to enamines *VII*. Starting from the cyclopropyl acid *IXe*, the cyclopropyl analogue (*XXVII*) of the neuroleptic chlorprothixene was synthesized in three steps. The piperazine derivatives *Ia, Ib, Ic, IIa, IIb* and *VIIb* were found to be very effective neuroleptics with a high degree of central depressant and cataleptic activity.

In the present pharmacological research of 8-substitution derivatives of 10-(4-methylpiperazino)-10,11-dihydrodibenzo[*b,f*] thiepin which are generally neuroleptically active, relatively little attention has been devoted so far to alkyls as 8-substituents. The synthesis of the 8-methyl derivative (*I*, $R^1 = \text{CH}_3$) and of the 8-tert-butyl derivative (*I*, $R^1 = \text{C}(\text{CH}_3)_3$) (ref.¹) has been described, the first of these having been found to be a four-fold more potent central depressant than chlorpromazine, the second of these being slightly weaker than chlorpromazine. At the same time, in the phenothiazine series of neuroleptics with a similarly localized substituent of alkyl type, only the ethyl derivative² "ethylisobutrazine" found practical application while in the thioxanthene series of 2-alkyl derivatives of prothixene the isopropyl derivative was found to be most effective here³. For this reason we thought it useful to extend the information on the 8-alkyl derivatives of this series and we started to work with the 8-ethyl, 8-isopropyl, 8-n-butyl and 8-n-octyl derivatives *Ia–Id*. The study included two representatives of the series of 8-cycloalkyl derivatives, *viz.* the 8-cyclopropyl derivative *Ie* and the 8-cyclopentyl derivative *If*. In a number of cases, the principal N-methyl derivatives *I* were supplemented with further N-substitution analogues *II–VI* and further with the 10-unsaturated analogues, the enamines *VII*.


During synthesis of *I–VII* we generally adhered to the usual scheme¹, using the corresponding 4-substituted thiophenols as the starting compounds. The thiophenols were condensed in the first step with 2-iodobenzoic acid⁴ in the presence of potassium

* Part LXX in the series Neurotropic and Psychotropic Agents; Part LXIX: This Journal 39, 617 (1974).

hydroxide to the corresponding 2-(4-subst. phenylthio)benzoic acids *IX* (method *A*). The second step consisted in the reduction of these acids with sodium bis(2-metho-




- I*, R = CH₃
II, R = (CH₂)₃OH
III, R = COOC₂H₅
IV, R = H
V, R = (CH₂)₃OCO(CH₂)₆CH₃
VI, R = (CH₂)₃OCO(CH₂)₁₀CH₃




- VII*, R = N(CH₃)
VIII, R = H

In all formulae: *a*, R¹ = CH₂CH₃; *b*, R¹ = CH(CH₃)₂; *c*, R¹ = (CH₂)₃CH₃;
d, R¹ = (CH₂)₇CH₃; *e*, R¹ = ; *f*, R¹ =


xyethoxy)dihydroaluminat⁵ to the 2-(4-subst. phenylthio)benzyl alcohols *X* (method *B*). In a further step, these alcohols were converted by treatment with thionyl chloride (method *C*) to the 2-(4-subst. phenylthio)benzyl chlorides *XI* which were refluxed with aqueous-ethanolic potassium cyanide (method *D*) giving the nitriles *XII*. Alkaline hydrolysis of these nitriles (method *E*) then yielded the 2-(4-subst. phenylthio) phenylacetic acids (*XIII*). Cyclization to the 8-subst. dibenzo[*b,f*]thiepin-10(11*H*)-ones (*XIV*) was usually carried out by heating with polyphosphoric acid (method *F*). Ketones were reduced with sodium borohydride in aqueous ethanol (method *G*) to alcohols *XV* which were treated with hydrogen chloride in benzene (method *H*) to yield the chlorides *XVI*. Substitution reactions of these chlorides with 1-methylpiperazine, 1-(3-hydroxypropyl)piperazine⁶, or with 1-ethoxycarbonylpiperazine⁷ (method *J*) resulted in the piperazine derivatives *I–III*. The substitution reactions were accompanied by elimination reactions giving rise to 2-substituted dibenzo[*b,f*]thiepins (*VIII*) which were in most cases isolated and characterized. Alkaline hydrolysis of carbamates *III* (method *K*) yielded secondary amines *IV*. With a view to the possibility of obtaining a protracted effect by esterification of alcohols of type *II* with fatty acids containing a longer chain⁸, reaction of alcohols *II* with capryloyl⁹ and lauroyl chlorides¹⁰ in benzene (method *L*) was used for the preparation of esters *V* and *VI*. Reactions of ketones *XIV* with 1-methylpiperazine and titanium tetrachloride in benzene (method *M*) led to enamines *VII* (ref.^{11–13}). The experimental section shows only examples of preparations using the above general methods. All the compounds prepared are included together with the usual experimental data in Table I.



- IX, R = COOH
 X, R = CH₂OH
 XI, R = CH₂Cl
 XII, R = CH₂CN
 XIII, R = CH₂COOH



- XIV, R = =O
 XV, R = -OH
 XVI, R = -Cl



- XVII, R = H
 XVIII, R =

In the ethyl series (*a*) we proceeded from the described 2-(4-ethylphenylthio)benzoic acid³ (*IXa*) and used the above general reactions. Only in the case of cyclization of *XIIIa* with polyphosphoric acid to the ketone *XIVa* a modified method was used, consisting in the presence of toluene as solvent for the organic reaction components (method *F-1*). In the isopropyl series (*b*) the starting compound was 4-isopropylthiophenol¹ which had been obtained by reduction of 4-isopropylbenzenesulfonyl chloride¹ with phosphorus and iodine in boiling acetic acid (method¹⁴). This was then³ transformed to acid *IXb*. The ketone *XIVb* was obtained in approximately the same yield with method *F* at 150°C as with method *F-1*. Likewise, in the *n*-butyl series (*c*) the starting 4-*n*-butylthiophenol¹ was obtained by reduction of 4-(*n*-butyl)benzenesulfonyl chloride¹ with lithium aluminium hydride in ether. 2-(4-*n*-Butylphenylthio)benzoic acid (*IXc*) has been reported previously³. Ketone *XIVc* has been obtained by the *F-1* method. The other compounds of this series were prepared by application of general methods.

Synthesis of the *n*-octyl derivative *Id* was motivated differently from the preparation of other compounds of this series. Preliminary tests of the ethyl derivative *Ia* and of the *n*-butyl derivative *Ic* showed that with extension of the alkyl chain in position 8 the central activity decreased. On the other hand, the antimicrobial activity rose (Table III), this raising the interest in compounds with a longer alkyl in the given position. In the *n*-octyl series (*d*) the preparation of the starting 4-*n*-octylthiophenol has been described in the literature¹⁵ but here we used also the novel procedure, *viz.* reduction of 4-(*n*-octyl)benzenesulfonyl chloride¹⁶ with phosphorus and iodine in acetic acid. The work in this series was rather complicated by the highly lipophilic character of all the intermediates which decreased the yields and complicated the isolation. For reducing the acid *IXd* to the alcohol *Xd* we used lithium aluminium hydride in ether¹. Cyclization of acid *XIII d* was carried out by method *F* at 160 to 170°C; the ketone *XIV d* was characterized as crystalline 2,4-dinitrophenylhydrazone.

The attempt at preparing the analogous *n*-dodecyl derivative was motivated similarly to the work in the *n*-octyl series but the work was not completed because of the extremely lipophilic character of the intermediates. *n*-Dodecylbenzene¹⁷ was chlorosulfonated and the crude 4-(*n*-dodecyl) benzenesulfonyl chloride was reduced with phosphorus and iodine in acetic acid to the novel 4-*n*-dodecylthiophenol (*XVII*) which was converted in a poor yield by method *A* to the acid *XVIII*.

TABLE I

Dibenzo[*b,f*]thiepins (*I—VIII*) and Intermediates of Their Synthesis (*IX—XVI*)

Compound ^{a,b}	Method (yield %)	B. p., °C/Torr or m. p., °C (solvent)	Formula (m. w.)	Calculated/Found			
				% C	% H	% S	% N (Cl)
<i>IXd</i>	<i>A</i> (90)	111—112 ^c (ethanol)	C ₂₁ H ₂₆ O ₂ S (342.5)	73.64 73.39	7.65 7.82	9.36 9.41	—
<i>XVIII</i>	<i>A</i> (25)	90.5—91.5 ^d (ethanol)	C ₂₅ H ₃₄ O ₂ S (398.6)	75.33 75.36	8.60 8.70	8.04 7.91	—
<i>IXe</i>	<i>A</i> ^e (80)	192—194 (aqueous ethanol)	C ₁₆ H ₁₄ O ₂ S (270.3)	71.08 71.25	5.22 5.21	11.86 11.94	—
<i>IXf</i>	<i>A</i> (77)	208—209 ^f (aqueous ethanol)	C ₁₈ H ₁₈ O ₂ S (298.4)	72.45 72.20	6.08 6.11	10.75 10.94	—
<i>Xa</i>	<i>B</i> (89)	160—163/1.7 ^g	C ₁₅ H ₁₆ OS (244.3)	73.73 73.87	6.60 6.65	13.12 12.70	—
<i>Xb</i>	<i>B</i> ^e (94)	156—160/1.5	C ₁₆ H ₁₈ OS (258.4)	74.37 74.58	7.02 7.12	12.41 12.32	—
<i>Xc</i>	<i>B</i> (92)	175—180/0.5	C ₁₇ H ₂₀ OS (272.4)	74.95 74.81	7.40 7.52	11.77 11.50	—
<i>Xd</i>	<i>e</i>	195/0.4	C ₂₁ H ₂₈ OS (328.5)	76.78 76.56	8.59 8.80	9.76 9.60	—
<i>Xf</i>	<i>B</i> (82)	212—216/2 ^h	C ₁₈ H ₂₀ OS (284.4)	76.01 75.70	7.09 6.96	11.27 10.97	—
<i>XIa</i>	<i>C</i> —	170/2.5	C ₁₅ H ₁₅ ClS (262.8)	68.55 68.47	5.75 5.79	12.20 12.03	13.49 13.80
<i>XIb</i>	<i>C</i> (87)	163—165/2	C ₁₆ H ₁₇ ClS (276.8)	69.42 69.32	6.19 6.16	11.58 11.62	12.81 13.07
<i>XIc</i>	<i>C</i> —	133—138/2	C ₁₇ H ₁₉ ClS (290.8)	70.20 70.07	6.58 6.57	11.03 10.92	12.19 12.21
<i>XId</i>	<i>C</i> —	190—193/0.5	C ₂₁ H ₂₇ ClS (346.9)	—	—	9.24 9.44	—
<i>XIf</i>	<i>C</i> ^e (80)	180—182/1	C ₁₈ H ₁₉ ClS (302.9)	71.38 71.71	6.32 6.38	10.59 10.50	11.71 11.51
<i>XIIa</i>	<i>D</i> (62)	185—188/2.5	C ₁₆ H ₁₅ NS (253.4)	—	—	12.66 12.00	5.53 5.22
<i>XIIb</i>	<i>D</i> ^e (86)	158—162/1	C ₁₇ H ₁₇ NS (267.4)	76.36 76.74	6.40 6.53	11.99 11.68	5.24 4.86
<i>XIIc</i>	<i>D</i> (84)	195—200/2	C ₁₈ H ₁₉ NS (281.4)	76.82 76.81	6.81 6.87	11.39 11.25	4.98 4.41

TABLE I
(Continued)

Compound ^{a,b}	Method (yield %)	B. p., °C/Torr or m. p., °C (solvent)	Formula (m. w.)	Calculated/Found			
				% C	% H	% S	% N (Cl)
<i>XII f</i>	<i>D</i> (80)	190—195/2.5 ⁱ	C ₁₉ H ₁₉ NS (293.4)	77.79 77.60	6.53 6.49	10.91 11.04	4.77 4.81
<i>XIII a</i>	<i>E</i> ^e (83)	96—99 (aqueous ethanol)	C ₁₆ H ₁₆ O ₂ S (272.4)	70.55 70.70	5.92 6.03	11.77 11.96	—
<i>XIII b</i>	<i>E</i> (86)	111—113 ^j (aqueous ethanol)	C ₁₇ H ₁₈ O ₂ S (286.4)	71.29 71.34	6.33 6.35	11.20 11.13	—
<i>XIII c</i>	<i>E</i> (92)	84—85 (aqueous ethanol)	C ₁₈ H ₂₀ O ₂ S (300.4)	71.96 72.23	6.71 6.78	10.67 10.70	—
<i>XIII d</i>	<i>E</i>	68—68.5 ^k (aqueous ethanol)	C ₂₂ H ₂₈ O ₂ S (356.5)	74.11 74.67	7.92 8.02	8.99 9.12	—
<i>XIII e</i>	^e	122—125.5 (aqueous ethanol)	C ₁₇ H ₁₆ O ₂ S (284.4)	71.81 71.57	5.67 5.71	11.27 11.58	—
<i>XIII f</i>	<i>E</i> (92)	106—107 ^m (aqueous ethanol)	C ₁₉ H ₂₀ O ₂ S (312.4)	73.04 73.16	6.45 6.45	10.26 9.97	—
<i>XIV a</i>	<i>F-1</i> (90)	190/2.5 ⁿ	C ₁₆ H ₁₄ OS (254.3)	75.55 75.09	5.55 5.61	12.61 12.41	—
<i>XIV b</i>	<i>F</i> ^e (76)	167—170/1.5	C ₁₇ H ₁₆ OS (268.4)	76.08 76.40	6.01 6.05	11.95 11.96	—
<i>XIV c</i>	<i>F-1</i> ^e (82)	190—195/2	C ₁₈ H ₁₈ OS (282.4)	76.55 76.55	6.42 6.50	11.35 11.00	—
<i>XIV d-DFH</i>	<i>F</i> —	177.5—178.5 ^o (ethanol-acetone)	C ₂₈ H ₃₀ N ₄ O ₄ S (518.6)	64.84 64.56	5.83 6.01	6.18 6.43	10.80 10.90
<i>XIV e</i>	^e	76.5—77 (cyclohexane)	C ₁₇ H ₁₄ OS (266.4)	76.65 76.76	5.30 5.50	12.04 12.12	—
<i>XIV e-DFH</i>		244.5—245.5 ^p (dimethyl- formamide-ethanol)	C ₂₃ H ₁₈ N ₄ O ₄ S (446.5)	61.86 61.85	4.06 4.25	7.18 7.35	12.55 12.82
<i>XIV f</i>	<i>F-1</i> (60)	212—215/2 ^q	C ₁₉ H ₁₈ OS (294.4)	77.51 77.72	6.16 6.41	10.89 10.72	—
<i>XV a</i>	<i>G</i> (92)	87—88 ^r (benzene- light petroleum)	C ₁₆ H ₁₆ OS (256.4)	74.96 74.95	6.29 6.22	12.51 12.43	—
<i>XV b</i>	<i>G</i> (86)	87—88 ^s (light petroleum)	C ₁₇ H ₁₈ OS (270.4)	75.51 75.37	6.71 6.80	11.86 11.71	—

TABLE I
 (Continued)

Compound ^{a,b}	Method (yield %)	B. p., °C/Torr or m. p., °C (solvent)	Formula (m. w.)	Calculated/Found			
				% C	% H	% S	% N (Cl)
<i>XVc</i>	<i>G</i> (82)	62—63 (benzene— light petroleum)	$C_{18}H_{20}OS$ (284.4)	76.01 76.16	7.09 7.08	11.27 11.28	—
<i>XVe</i>	<i>G</i> (98)	105—107 ^f (cyclohexane)	$C_{17}H_{16}OS$ (268.4)	76.08 75.91	6.01 6.22	11.95 11.73	—
<i>XVf</i>	<i>G</i> (95)	77—81 ^h (light petroleum)	$C_{19}H_{20}OS$ (296.4)	76.98 76.77	6.80 6.77	10.82 11.00	—
<i>XVIIb</i>	<i>H</i> (93)	66—68 ^p (light petroleum)	$C_{17}H_{17}ClS$ (288.8)	70.69 70.82	5.93 6.07	11.10 11.28	12.28 12.48
<i>XVIIe</i>	<i>H</i> ^e (97)	76.5—77 (cyclohexane)	$C_{17}H_{15}ClS$ (286.8)	71.18 70.52	5.27 5.43	11.18 10.95	12.36 12.13
<i>Ia</i>	<i>J</i> (72)	96—98 (acetone)	$C_{21}H_{26}N_2S$ (338.5)	74.51 74.22	7.74 7.90	9.47 9.62	8.28 8.14
<i>Ia-MS</i>	—	190—192 (ethanol)	$C_{22}H_{30}N_2O_3S_2$ (434.6)	60.80 59.91	6.96 7.00	14.76 14.92	6.44 6.35
<i>Ib</i>	<i>J</i> ^e (91)	118—120 (benzene— light petroleum)	$C_{22}H_{28}N_2S$ (352.5)	74.95 74.45	8.00 8.11	9.09 9.23	7.95 8.22
<i>Ib-2MS</i>	—	203—206 (ethanol)	$C_{24}H_{36}N_2O_6S_3$ (544.7)	52.91 52.62	6.66 6.70	17.66 17.52	5.14 5.01
<i>Ic-MS</i> ^w	<i>J</i> (72)	171—173 ^x (ethanol—ether)	$C_{24}H_{35}N_2O_3.5S_2$ (471.7)	61.11 61.15	7.48 7.39	13.59 13.53	5.94 5.56
<i>Id-M</i>	<i>J</i> (51)	127—130 ^y (ethanol—ether)	$C_{31}H_{42}N_2O_4S$ (538.7)	69.11 68.87	7.85 7.84	5.95 6.44	5.21 5.07
<i>Id-2HM</i>	—	151—152.5 (ethanol—ether)	$C_{35}H_{46}N_2O_8S$ (654.8)	64.20 64.35	7.08 7.20	4.90 5.10	4.27 4.11
<i>Ie-M</i>	<i>J</i> (78)	168—170 ^z (ethanol)	$C_{26}H_{30}N_2O_4S$ (466.6)	66.93 67.08	6.48 6.50	6.87 7.04	6.00 5.90
<i>If-2MS</i> ^w	<i>J</i> (61)	203—205 (ethanol)	$C_{26}H_{39}N_2O_6.5S_3$ (579.8)	53.86 54.10	6.78 6.74	16.59 16.61	4.83 4.63
<i>Ila</i>	<i>J</i> (93)	71—73 ^{aa} (benzene— light petroleum)	$C_{23}H_{30}N_2OS$ (382.6)	72.21 72.11	7.90 7.96	8.38 8.00	7.32 7.07

TABLE I
(Continued)

Compound ^{a,b}	Method (yield %)	B. p., °C/Torr or m. p., °C (solvent)	Formula (m. w.)	Calculated/Found			
				% C	% H	% S	% N (Cl)
<i>Ila</i> -2MS ^{bb}	—	128—130 (ethanol-ether)	C ₂₅ H ₄₂ N ₂ O ₉ S ₃ (610·8)	49·16	6·93	15·75	4·59
				50·02	6·61	15·70	5·00
<i>Ila</i> -2HM ^w	—	153—155 (ethanol)	C ₃₁ H ₃₉ N ₂ O _{9·5} S ₃ (623·7)	59·63	6·30	5·14	4·49
				59·78	6·35	5·24	4·49
<i>Ilb</i> -2MS	<i>J</i> (94)	185—186 (ethanol)	C ₂₆ H ₄₀ N ₂ O ₇ S ₃ (588·8)	—	—	16·34	4·76
						16·20	4·83
<i>Ilc</i>	<i>J</i> (74)	92—94 ^{cc} (benzene- light petroleum)	C ₂₅ H ₃₄ N ₂ OS (410·6)	73·12	8·35	7·81	6·82
				73·10	8·39	7·85	6·46
<i>Ilc</i> -2HM	—	103—107 (ethanol)	C ₃₃ H ₄₂ N ₂ O ₉ S (642·7)	61·66	6·59	4·99	4·36
				61·77	6·53	5·57	4·23
<i>Ilf</i> -2MS ^w	<i>J</i> (82)	188—189 (ethanol)	C ₂₈ H ₄₃ N ₂ O _{7·5} S ₃ (623·8)	53·90	6·94	15·42	4·49
				54·07	7·14	15·65	4·45
<i>Illa</i> -M	<i>J</i> (70)	188—190 (ethanol)	C ₂₇ H ₃₂ N ₂ O ₆ S (512·5)	63·27	6·29	6·25	5·47
				63·15	6·32	6·52	5·47
<i>Illb</i> -M	<i>J</i> (68)	176—179 (ethanol)	C ₂₈ H ₃₄ N ₂ O ₆ S (526·6)	63·86	6·51	6·08	5·32
				63·81	6·53	6·00	5·32
<i>Illf</i> ^{dd}	<i>J</i> (76)	90—93 ^{ee} (benzene)	C ₂₈ H ₃₄ N ₂ O ₂ S (462·6)	72·70	7·41	6·92	6·06
				72·38	7·44	7·05	5·71
<i>Iva</i>	<i>K</i> ^e (84)	113—114 (acetone)	C ₂₀ H ₂₄ N ₂ S (324·4)	74·04	7·46	9·86	8·64
				73·86	7·49	9·91	8·38
<i>Iva</i> -M	—	159—161 (ethanol)	C ₂₄ H ₂₈ N ₂ O ₄ S (440·5)	65·43	6·40	7·28	6·36
				65·76	6·68	7·62	6·07
<i>Ivb</i>	<i>K</i> (93)	160—163 ^{ff} (benzene)	C ₂₁ H ₂₆ N ₂ S (338·5)	74·51	7·74	9·47	8·28
				74·43	7·85	9·44	7·98
<i>Ivf</i> -2HM	<i>K</i> (97)	143—146 (acetone)	C ₃₁ H ₃₆ N ₂ O ₈ S (596·7)	62·40	6·08	5·37	4·69
				62·18	6·10	5·57	4·67
<i>Va</i> -2HM	<i>L</i> ^c —	144—146 (acetone)	C ₃₉ H ₅₂ N ₂ O ₁₀ S (740·9)	63·22	7·07	4·33	3·78
				63·32	7·07	4·44	3·81
<i>Vb</i> -2HM	<i>L</i> —	146—148 (acetone)	C ₄₀ H ₅₄ N ₂ O ₁₀ S (754·9)	63·64	7·21	4·25	3·71
				63·43	7·40	4·69	3·39
<i>Vc</i> -2HM	<i>L</i> —	113—115 (acetone)	C ₄₁ H ₅₆ N ₂ O ₁₀ S (768·9)	64·04	7·34	4·17	3·64
				64·28	7·34	4·40	3·67

TABLE I
 (Continued)

Compound ^{a,b}	Method (yield %)	B. p., °C/Torr or m. p., °C (solvent)	Formula (m. w.)	Calculated/Found			
				% C	% H	% S	% N (Cl)
<i>Vla</i> -2HM	<i>L</i>	126—128	C ₄₃ H ₆₀ N ₂ O ₁₀ S	64.80	7.59	4.02	3.51
	—	(acetone)	(797.0)	64.63	7.67	4.14	3.46
<i>VIIIb</i> -MS ^w	<i>M</i>	210—212	C ₂₃ H ₃₁ N ₂ O _{3.5} S ₂	60.62	6.86	14.08	6.15
	—	(ethanol-ether)	(455.6)	60.88	6.92	14.58	6.07
<i>VIIc</i>	<i>M</i> ^e	140—141	C ₂₃ H ₂₈ N ₂ S	75.77	7.74	8.80	7.68
	—	(benzene)	(364.5)	76.13	7.91	9.06	7.91
<i>VIIc</i> -M	—	197—200	C ₂₇ H ₃₂ N ₂ O ₄ S	67.47	6.71	6.67	5.83
	—	(ethanol)	(480.6)	67.35	6.71	7.02	5.67
<i>VIIIf</i> -MS ^{gg}	<i>M</i>	204—208	C ₂₅ H ₃₄ N ₂ O ₄ S ₂	61.21	6.99	13.05	5.71
	—	(ethanol)	(490.5)	61.60	6.97	13.30	5.47
<i>VIIIa</i>	<i>J</i>	170—173/2	C ₁₆ H ₁₄ S	—	—	13.45	—
	—	—	(238.3)	—	—	12.86	—
<i>VIIIb</i>	<i>J</i> ^e	177—180/1.5	C ₁₇ H ₁₆ S	80.90	6.39	12.71	—
	—	—	(252.4)	80.52	6.52	12.56	—
<i>VIIIc</i>	<i>J</i>	150—160/1.7 ^{hh}	C ₁₈ H ₁₈ S	81.15	6.81	12.04	—
	—	—	(266.4)	80.77	7.07	11.71	—
<i>VIIIe</i>	<i>J</i>	72—73 ⁱⁱ	C ₁₇ H ₁₄ S	81.55	5.64	12.81	—
	—	(light petroleum)	(250.3)	81.39	5.93	12.51	—
<i>VIIIIf</i>	<i>J</i>	200/0.5 ^{jj}	C ₁₉ H ₁₈ S	81.96	6.52	11.52	—
	—	—	(278.4)	81.92	6.56	10.75	—

^a The intermediate nitrile *XIII*d, alcohol *XV*d and chlorides *XVI*a, *XVI*c, *XVI*d and *XVI*f could not be obtained in the crystalline form and were used in the crude state. They are not included in the table. ^b DFH 2,4-dinitrophenylhydrazone, MS methanesulfonate, M maleate, HM hydrogen maleate. ^c IR spectrum (KBr): 745 (4 vicinal aromatic C—H), 810 (2 vicinal aromatic C—H), 912 (COOH), 1250 (CO), 1550 and 1582 (Ar), 1670 cm⁻¹ (ArCOOH); NMR spectrum: δ 11.75 (bs, disappears after D₂O, 1 H, COOH), 8.18 (m, 1 H, aromatic 6-H), 7.54 (d, *J* = 9.0 Hz, 2 H, aromatic 2',6'-H₂), 7.27 (d, *J* = 9.0 Hz, 2 H, aromatic 3',5'-H₂), c. 7.21 (m, 2 H, aromatic 4,5-H₂), 6.80 (m, 1 H, aromatic 3-H), 2.65 (t, *J* = 7.0 Hz, 2 H, ArCH₂), c. 1.60 (m, 2 H, CH₂ in the vicinity of terminal methyl), 1.30 (bs, 10 H, remaining CH₂ groups of octyl), 0.88 (t, 3 H, CH₃). ^d UV spectrum: λ_{max} 222 nm (log ε 4.40), 256 nm (395), 276 nm (3.73), 317 nm (3.54); IR spectrum: 736 (4 vicinal aromatic C—H), 810 (2 vicinal aromatic C—H), 1562, 1585 (Ar), 1674 cm⁻¹ (ArCOOH). ^e See experimental section. ^f UV spectrum: λ_{max} 222 nm (log ε 4.41), 254 nm (4.01), 273 nm (3.76), 311 nm (3.58); IR spectrum (KBr): 740 (4 vicinal aromatic C—H), 810 (2 vicinal aromatic C—H), 920, 1254 (COOH), 1560, 1588 (Ar), 1730, 2560, 2640, 3100—3200 cm⁻¹ (COOH). ^g IR spectrum: 755 (4 vicinal aromatic C—H), 825 (2 vicinal aromatic C—H), 1030

(CH₂OH), 1 590 (Ar), 3 380 cm⁻¹ (OH). ^h NMR spectrum: δ 7.10–7.60 (m, 8 H, aromatic protons), 4.72 (bs, 2 H, ArCH₂O), 2.90 (m, 1 H, Ar—CH of cyclopentyl), 2.33 (bs, disappears after D₂O, 1 H, OH), 1.50–2.20 (m, 8 H, 4 CH₂ of cyclopentyl). ⁱ IR spectrum (film): 758 (4 vicinal aromatic C—H), 820 (2 vicinal aromatic C—H), 1 590 (Ar), 2 255 cm⁻¹ (R—CN). ^j IR spectrum: 758 (4 vicinal aromatic C—H), 828 (2 vicinal aromatic C—H), 928, 1 238, 1 710 and 3 100 cm⁻¹ (COOH). ^k IR spectrum: 760 (4 vicinal aromatic C—H), 820 (2 vicinal aromatic C—H), 940 (COOH), 1 240 (C—O), 1 700 cm⁻¹ (COOH); NMR spectrum: δ 10.40 (bs, disappears after D₂O, 1 H, COOH), 6.90–7.50 (m, 8 H, aromatic protons), 3.80 (s, 2 H, ArCH₂CO), 2.52 (t, *J* = 7.0 Hz, 2 H, ArCH₂ of octyl), c. 1.55 (m, 2 H, CH₂ in the vicinity of terminal methyl), 1.23 (bs, 10 H, remaining CH₂ groups of octyl), 0.85 (t, 3 H, CH₃). ^m IR spectrum: 755, 768 (4 vicinal aromatic C—H), 818, 828 (2 vicinal aromatic C—H), 945, 1 238 (COOH), 1 570, 1 593 (Ar), 1 710, 2 560, 2 730 and 3 150 cm⁻¹ (COOH). ⁿ UV spectrum: λ_{max} 241 nm (log ε 4.29); IR spectrum: 747, 766 (4 vicinal aromatic C—H), 830 (2 vicinal aromatic C—H), 880 (solitary aromatic C—H), 1 600 (Ar), 1 670 cm⁻¹ (Ar—CO). ^o UV spectrum: λ_{max} 254 nm (log ε 4.31), 356 nm (4.44); IR spectrum: 740, 760 (4 vicinal aromatic C—H), 840 (2 vicinal aromatic C—H), 909 (solitary aromatic C—H), 1 340, 1 518 (NO₂), 1 540, 1 595 (Ar), 1 620 (C=N), 3 310 cm⁻¹ (NH); NMR spectrum: δ 11.90 (bs, disappears after D₂O, 1 H, NH), 9.18 (d, *J* = 3.0 Hz, 1 H, aromatic proton between nitro groups), 8.35 (mcd, *J* = 9.0; 3.0 Hz, 1 H, another aromatic proton in the vicinity of NO₂), 8.02 (d, *J* = 9.0 Hz, 1 H, aromatic proton in *m*-position toward NO₂ groups), 7.00–7.80 (m, 7 H, remaining aromatic protons), 4.19 (s, 2 H, ArCH₂ in a ring), 2.58 (t, 2 H, ArCH₂ of octyl), c. 1.50 (m, 2 H, CH₂ adjacent to the terminal methyl), 1.24 (bs, 10 H, remaining CH₂ groups of octyl), 0.85 (t, 3 H, CH₃). ^p UV spectrum (ethanol): λ_{max} 228 nm (log ε 4.63), 251 nm (4.67), infl. 281 nm (4.37); IR spectrum: 760 (4 vicinal aromatic C—H), 842 and 833 (2 vicinal aromatic C—H), 860 (solitary aromatic C—H), 1 330 and 1 510 (NO₂), 1 590 (Ar), 1 610 (C=N), 3 310 cm⁻¹ (NH). ^q UV spectrum: λ_{max} 244 nm (log ε 4.26), 340 nm (3.47); IR spectrum (film): 748 (4 vicinal aromatic C—H), 828 (2 vicinal aromatic C—H), 875 (solitary aromatic C—H), 1 288 (CO), 1 594 (Ar), 1 673 cm⁻¹ (ArCO); NMR spectrum: δ 8.13 (d, *J* = 3.0 Hz, 1 H, aromatic 9-H), 7.00–7.80 (m, 6 H, remaining aromatic protons), 4.35 (s, 2 H, ArCH₂CO), 2.95 (m, 1 H, Ar—CH of cyclopentyl), 1.40–2.20 (m, 8 H, 4 CH₂ of cyclopentyl). ^r IR spectrum (KBr): 749 (4 vicinal aromatic C—H), 826 (2 vicinal aromatic C—H), 899 (solitary aromatic C—H), 1 052 (CHOH), 3 380 cm⁻¹ (OH). ^s NMR spectrum: δ 6.90–7.70 (m, 7 H, aromatic protons), 5.10–5.50 (m, dd after D₂O, 1 H, Ar—CH—O), 3.20–3.90 (dd, 2 H, ArCH₂ in a ring), 2.40–3.30 (m, 1 H, Ar—CH of isopropyl), 2.08 (bs, disappears after D₂O, 1 H, OH), 2.21 and 2.11 (2 s, 2 CH₃). ^t IR spectrum (KBr): 758 (4 vicinal aromatic C—H), 830 (2 vicinal aromatic C—H), 862 (solitary aromatic C—H), 1 060 (CHOH), 1 605 (Ar), 3 400 cm⁻¹ (OH); NMR spectrum: δ 7.05–7.70 (m, 6 H, aromatic protons in positions 1, 2, 3, 4, 6, 9), 6.90 (mcd, *J* = 9.0; 2.5 Hz, 1 H, aromatic 7-H), 5.33 (m, dd after D₂O, *J* = 8.0; 4.0 Hz, 1 H, Ar—CH—O), 3.76 and 3.32 (2 dd, *J* = 14.0; 4.0 and 14.0; 8.0 Hz, 2 H, ArCH₂ in a ring), 2.16 (d, disappears after D₂O, 1 H, OH), 1.85 (m, 1 H, Ar—CH of cyclopropyl), 0.50–1.10 (m, 4 H, 2 CH₂ of cyclopropyl). ^u IR spectrum: 751 (4 vicinal aromatic C—H), 820 (2 vicinal aromatic C—H), 900 (solitary aromatic C—H), 1 015 (CHOH), 3 240 cm⁻¹ (OH). ^v NMR spectrum: δ 6.90–7.70 (m, 7 H, aromatic protons), 5.80 (dd, *J* = 6.0; 8.0 Hz, 1 H, Ar—CH—Cl), 3.45–4.20 (m, 2 H, ArCH₂ in a ring), 2.50–3.10 (m, 1 H, Ar—CH of isopropyl), 1.17 (d, *J* = 7.0 Hz, 6 H, 2 CH₃). ^w Hemihydrate. ^x NMR spectrum (CD₃SOCD₃): δ 6.80–7.70 (m, 7 H, aromatic protons), 3.80–4.20 (m, 1 H, Ar—CH—N), 2.00–3.60 (m, 12 H, 2 ArCH₂ and 4 CH₂ of piperazine), 2.31 (s, 3 H, NCH₃), 1.35 (m, 4 H, 2 CH₂ of butyl), 0.82 (t, 3 H, C—CH₃). ^y NMR spectrum of base: δ 6.70–7.55 (m, 7 H, aromatic protons), 3.00–4.00 (m, 3 H, ArCH₂CHAr), 2.58 (m, 4 H, CH₂N¹CH₂ of piperazine), c. 2.50 (2 H, ArCH₂ of octyl), 2.45 (m, 4 H, CH₂N⁴CH₂ of piperazine), 2.24 (s, 3 H, NCH₃), c. 1.50 (m, 2 H, CH₂ in the vicinity



of the terminal methyl), 1.23 (bs, 10 H, remaining CH₂ groups of octyl), 0.85 (t, 3 H, C—CH₃)^z NMR spectrum of base: δ 6.50–7.10 (m, 7 H, aromatic protons), 3.00–4.00 (m, 3 H, ArCH₂CHAr), 2.60 (m, 4 H, CH₂N¹CH₂ of piperazine), 2.45 (m, 4 H, CH₂N⁴CH₂ of piperazine), 2.25 (s, 3 H, NCH₃), 1.80 (m, 1 H, Ar—CH of cyclopropyl), 0.50–1.10 (m, 4 H, 2 CH₂ of cyclopropyl). ^{aa} IR spectrum (KBr): 761 (4 vicinal aromatic C—H), 822 (2 vicinal aromatic C—H), 899 (solitary aromatic C—H), 1067 (C—OH), 2779, 2840 (NCH₂), 3215 and 3450 cm⁻¹ (OH); NMR spectrum: δ 6.75–7.60 (m, 7 H, aromatic protons), c. 4.00 (bs, 1 H, OH), 3.00–4.00 (m, 3 H, ArCH₂CHAr), 3.78 (t, $J = 7.0$ Hz, 2 H, CH₂O), c. 2.58 (m, 12 H, ArCH₂ and 5 NCH₂), 1.70 (m, 2 H, middle CH₂ of hydroxypropyl), 1.15 (t, $J = 8.0$ Hz, 3 H, C—CH₃). ^{bb} Dihydrate. ^{cc} IR spectrum (KBr): 760 (4 vicinal aromatic C—H), 828 (2 vicinal aromatic C—H), 888 (solitary aromatic C—H), 1070 and 1130 (CH₂OH), 2770 (NCH₂), 3190 cm⁻¹ (OH). ^{dd} Solute with 1/3 benzene molecule. ^{ee} NMR spectrum: δ 6.85–7.65 (m, 7 H, aromatic protons), 4.11 (q, 2 H, COOCH₂), 3.00–3.95 (m, 3 H, ArCH₂CHAr), 3.40 (m, 4 H, CH₂N⁴CH₂ of piperazine), 2.60 (m, 4 H, CH₂N¹CH₂ of piperazine), c. 2.00 (m, 1 H, Ar—CH of cyclopentyl), 1.68 (m, 8 H, 4 CH₂ of cyclopentyl), 1.24 (t, 3 H, C—CH₃). ^{ff} NMR spectrum: δ 7.56 (d, $J = 2.5$ Hz, 1 H, aromatic 9-H), 7.00–7.50 (m, 5 H, aromatic protons in positions 1, 2, 3, 4, 6), 6.94 (q, $J = 9.0$; 2.5 Hz, 1 H, aromatic 7-H), 3.00–4.00 (m, 3 H, ArCH₂CHAr), c. 3.00 (m, 1 H, Ar—CH of isopropyl), 2.76 and 2.64 (2 m, 8 H, 4 CH₂ of piperazine), 1.52 (s, 1 H, NH), 1.18 (d, $J = 7.0$ Hz, 6 H, 2 CH₃). ^{gg} Monohydrate. ^{hh} UV spectrum: λ_{\max} 260 nm (log ϵ 4.24), 285.5 nm (3.60); IR spectrum: 745 (4 vicinal aromatic C—H), 785 (*cis*-CH=CH), 820 (2 vicinal aromatic C—H), 888 (solitary aromatic C—H), 1590 cm⁻¹ (Ar); NMR spectrum: δ 7.05–7.60 (m, 7 H, aromatic protons), 7.05 (s, 2 H, ArCH=CHAr), 2.55 (t, $J = 7.0$ Hz, 2 H, ArCH₂), 1.45 (m, 4 H, C—CH₂CH₂—C of butyl), 0.88 (t, 3 H, CH₃). ⁱⁱ UV spectrum: λ_{\max} 222 nm (log ϵ 4.49), 262.5 nm (4.42), 297.5 nm (3.66); NMR spectrum: δ 6.80–7.50 (m, 7 H, aromatic protons), 6.90 (s, 2 H, ArCH=CHAr), 1.50–2.00 (m, 1 H, Ar—CH of cyclopropyl), 0.50–1.10 (m, 4 H, 2 CH₂ of cyclopropyl). ^{jj} UV spectrum: λ_{\max} 258 nm (log ϵ 4.22), 284 nm (3.70); IR spectrum (film): 750 (4 vicinal aromatic C—H), 788 (*cis*-CH=CH), 822 (2 vicinal aromatic C—H), 880 (solitary aromatic C—H), 1590 cm⁻¹ (Ar); NMR spectrum: δ 7.00–7.60 (m, 9 H, aromatic and olefinic protons), 2.92 (m, 1 H, Ar—CH of cyclopentyl), 1.69 (m, 8 H, 4 CH₂ of cyclopentyl).

Investigation in the cyclopropyl series (*e*) was attractive because of the similarity of cyclopropyl with isopropyl (the favourable effect of which on activity was confirmed in this series) as well as because of lack of reports on this substituent's effect on the aromatic ring on the neuroleptic activity in other series; the 2-cyclopropyl derivatives of the phenothiazine series, *e.g.* the cyclopropyl analogue of chlorpromazine, had been described in patent literature^{18,19} but were generally designated as coronary dilatants, spasmolytics and compounds with a central depressant activity. At the same time it was to be expected that the instability of the cyclopropane ring might make some of the synthetic steps rather difficult.


The starting compound used was cyclopropylbenzene which had been prepared according to Corbin and coworkers²⁰ by an intramolecular debromination of 1,3-dibromo-1-phenylpropane with the aid of a Zn—Cu-couple²¹. The authors carried out the reaction at 7–9°C. We observed while working in larger batches that under these conditions the reaction does not proceed with a low concentration of the dibromo derivative while after reaching higher concentrations it may

have the tendency to proceed too vigorously. It was found here that this difficulty can be overcome by working at 50°C. At the same time, the dibromo derivative must be added dropwise and very slowly to the mixture of the Zn-Cu-couple and dimethylformamide to maintain the low concentration of the starting compound. If it is added more rapidly an intermolecular debromination in the more reactive α -positions takes place as documented by the identification of *meso*-3,4-diphenylhexane (*XIX*) (ref.²²) in one of the experiments (debromination in the terminal positions probably takes place only during processing the reaction mixture).


Cyclopropylbenzene was converted by bromination at -70°C selectively and in a high yield to 4-cyclopropylbromobenzene²³ (see also ref.²⁴). The compound reacts with magnesium in ether smoothly to the Grignard reagent (ref.²⁵) which reacts with sulfur, giving rise to the novel 4-cyclopropylthiophenol (*XX*). Preparation of the analytically pure compound appeared to be difficult and hence the product was characterized by gentle oxidation to the crystalline di(4-cyclopropylphenyl) disulfide (*XXI*). Application of method *A* to the crude thiol *XX* yields readily the acid *IXe*. In view of the fact that during homologization of acid *IXe* to acid *XIIIe* by the usual procedure difficulties could be expected (particularly with method *C*), a modified procedure was preferred²⁶: the thiol *XX* was converted to the acid *XIIIe* by direct condensation with 2-(2-iodophenyl)acetic acid^{26,27}. Attempts at cyclization of *XIIIe* by the usual methods, *i.e.* with the aid of sulfuric acid, polyphosphoric acid, hydrofluoric acid or phosphorus pentoxide in toluene or xylene were not successful, apparently due to the instability of the cyclopropane ring in these media. On the other hand, cyclization proceeds relatively readily in trifluoroacetic anhydride²⁸⁻³⁰. Even in this case, the product formed is not homogeneous but chromatography on alumina permits the isolation of about 50% of the desired crystalline ketone in a pure state, the identity having been confirmed by analysis and spectra and finally by conversion to the 2,4-dinitrophenylhydrazone. In further steps of synthesis of *Ie* we used methods *G, H* and *J*. During treatment of alcohol *XVe* (method *H*) with hydrogen chloride no damage to the cyclopropane ring was observed.



XX, R = H




XXI, R = S-

XXII, R = SO₂ClXXIII, R = SO₂NH₂

XXIV, R = SH

In the cyclopentyl series (*f*) the starting compound was cyclopentylbenzene, obtained best by a Friedel-Crafts reaction of benzene with cyclopentanol³¹. The usual chlorosulfonation yielded the novel 4-cyclopentylbenzenesulfonyl chloride (*XXII*) which was converted for characterization to the sulfonamide *XXIII*. Reduction of the sulfonyl chloride with lithium aluminium hydride yielded 4-cyclopentylthiophenol (*XXIV*) which was processed further by application of general procedures.

The above 2-(4-cyclopropylphenylthio)benzoic acid (*Ixe*) was used for the synthesis of the hitherto unknown cyclopropyl analogue of the neuroleptic chlorprothixene³². Cyclization with the aid of trifluoroacetic anhydride yielded 2-cyclopropylthioxanthone (*XXV*) which reacted with 3-dimethylaminopropylmagnesium chloride in tetrahydrofuran to the tertiary alcohol *XXVI*. Its dehydration with boiling 20% sulfuric acid gave rise to the olefin *XXVII*, probably as a mixture of geometric isomers. Repeated crystallization of hydrogen maleate yielded a probably homogeneous compound but its configuration at the double bond cannot be determined.



Most of the prepared piperazine derivatives *I*, *II*, *IV* and *VII* and the thioxanthene derivative *XXVII* were tested pharmacologically with a view to the assumed central depressant and neuroleptic activity (for methods see ref.³³) in the form of salts, the values shown referring to bases. The results obtained are shown in Table II which includes as standards octoclothepein (*I*, $R^1 = \text{Cl}$) i.e. 8-chloro-10-(4-methylpiperazino)-10,11-dihydrodibenzo[*b,f*]thiepin (for parenteral administration as methanesulfonate⁵, for oral application as maleate³⁴), and further chlorprothixene, i.e. *cis*-2-chloro-9-(3-dimethylaminopropylidene)thioxanthene³⁵ (*p.o.* as hydrochloride, parenterally as methanesulfonate³²). Besides tests of acute toxicity for mice, the compounds were evaluated in the rotating-rod test (effect on motor coordination) in mice and for assessing the neuroleptic activity their cataleptic effect in rats was examined. The results are shown in the table as the usual mean lethal doses LD_{50} as well as in the form of the mean effective doses (ED_{50}), in mg/kg throughout.

Table II shows that ethyl, isopropyl and cyclopropyl are suitable 8-substituents from the point of view of central depressant and cataleptic activity in the series of neuroleptic 10-piperazinodibenzo[*b,f*]thiepins. Highly effective are especially *N*-methyl derivatives *I*, *N*-(3-hydroxypropyl) derivatives *II* and enamines *VII*. In the rotating rod test, none of the compounds exceeds the effect of octoclothepein even if some of them are almost equal to it (*Ia*, *Ie*, *Ila*). On the other hand, in the catalepsy test, it is surpassed by a number of compounds (*Ia*, *Ib*, *Ila*, *Iib*, *VIIb*) and others are almost equal to it (*Ie*, *IIf*). In contrast with octoclothepein, some compounds display a dissociation of the depressant from the cataleptic activity in favour of the latter which makes them to potential neuroleptics with decreased depressant activity. Of the highly effective compounds, this dissociation is best exhibited by the enamine of the isopropyl series (*VIIb*) which is 4 times weaker as a depressant but 14 times more

effective as a cataleptic than octoclothebin. Of the medium-potent compounds the hydroxyl derivative of the isopropyl series *I**b*** is 8 times weaker as a depressant and more than twice more effective as a cataleptic than octoclothebin. The cyclopropyl analogue of chlorprothixene (compound *XXVII*) retains some depressant activity but is completely ineffective cataleptically. It cannot be excluded, however, that the compound tested actually belongs to the practically inactive *trans* series.

Esters *Vb* and *Vc* in the form of di(hydrogen maleates) have been tested so far by methods of general pharmacological screening at the affiliated unit of this institute at Rosice n/L (Dr J. Němec). Both were applied *p.o.* Their LD₅₀ are greater than 1 g/kg and the compounds were applied *in vivo* in doses of 300 (*Vb*) or 200 (*Vc*) mg/kg. Both bring about signs of central depression, have cataleptic, antiamphetamine, and hypothermic activity and they potentiate thiopental narcosis. Further they display an antihistamine effect and, at higher doses, they depress the blood pressure and are antiinflammatory. In general, they thus behave also as neuroleptics.

TABLE II
Pharmacological Properties of Prepared Compounds

Compound	Acute toxicity ^a		Rotating rod ^a		Catalepsy ^a	
	LD ₅₀		ED ₅₀		ED ₅₀	
	<i>i.v.</i> ^b	<i>p.o.</i> ^b	<i>i.v.</i>	<i>p.o.</i>	<i>i.p.</i> ^b	<i>p.o.</i>
OCT ^c	46	78	0.06	2.2	2.4	4.3
<i>Ia</i>	54	—	0.088	—	0.72	—
<i>Ib</i>	28	—	0.18	—	1.0	—
<i>Ic</i>	—	250	—	9.6	—	17.0
<i>Ie</i>	—	86	—	2.5	—	4.8
<i>If</i>	47	—	0.8	—	4.2	—
<i>Ila</i>	47	—	0.074	—	0.56	—
<i>Ib</i>	29	—	0.45	—	1.0	—
<i>Ic</i>	—	275	—	12.0	—	38.0
<i>If</i>	37	—	0.6	—	3.2	—
<i>IVa</i>	—	160	—	3.7	—	9.6
<i>Vb</i>	35	—	0.25	—	0.17	—
<i>Vc</i>	—	200	—	7.2	—	8.2
<i>Vf</i>	—	—	0.9	—	—	—
CPTX ^d	38	217	0.11	5.2	2.6	23.0
<i>XXVII</i>	—	<200 ^e	—	11.0	—	>50.0 ^f

^a All values in mg/kg. ^b *i.v.* intravenously, *i.p.* intraperitoneally, *p.o.* per os. ^c Octoclothebin.

^d Chlorprothixene. ^e The dose shown was lethal for 4 out of 5 animals. ^f The dose shown caused catalepsy in 1 out of 10 rats.

TABLE III
Antimicrobial Activity of Prepared Compounds *in vitro*

Compound ^{a,b}	1 ^c	2	3	4	5	6	7	8	9	10
<i>Ia</i> -MS	12.5	12.5	12.5	12.5	—	12.5	62.3	62.3	125	125
<i>Ib</i> -2MS	12.5	12.5	12.5	12.5	—	3.1	125	62.3	125	125
<i>Ic</i> -MS	3.2	3.2	3.2	3.2	—	25	31.2	31.2	125	31.2
<i>Id</i> -M	12.5	12.5	12.5	12.5	50	25	125	125	—	—
<i>Ie</i> -M	25	25	25	25	—	12.5	125	125	—	—
<i>If</i> -2MS	6.2	6.2	6.2	6.2	—	12.5	62.3	62.3	125	125
<i>Ila</i> -MS	50	50	50	50	—	25	125	125	—	125
<i>Ilb</i> -MS	25	25	25	25	—	12.5	—	—	—	—
<i>Ilc</i> -2HM	3.2	3.2	3.2	3.2	—	25	62.3	62.3	125	125
<i>Ilf</i> -2MS	6.2	6.2	6.2	6.2	—	12.5	62.3	62.3	125	125
<i>IVa</i> -M	12.5	12.5	12.5	12.5	—	12.5	62.3	62.3	125	125
<i>VIIb</i> -MS	—	—	—	—	—	12.5	62.3	62.3	62.3	62.3
<i>VIIc</i> -M	3.2	3.2	3.2	3.2	—	25	62.3	62.3	125	62.3
<i>VIIlf</i> -MS	—	—	—	—	—	—	62.3	62.3	62.3	125
<i>XXVII</i> -M	12.5	12.5	12.5	12.5	—	6.2	125	125	—	—

^a Minimum inhibitory concentration in µg/ml. Unless a numerical value is given, the compound at 125 µg/ml causes no inhibition. ^b MS methanesulfonate, M maleate, HM hydrogen maleate.

^c *Streptococcus β-haemolyticus*, 2 *Streptococcus β-haemolyticus* WARD, 3 *Staphylococcus pyogenes aureus*, 4 *Staphylococcus pyogenes aureus* penicillin-resistant, 5 *Klebsiella pneumoniae*, 6 *Mycobacterium tuberculosis* H37Rv, 7 *Saccharomyces pastorianus*, 8 *Trichophyton mentagrophytes*, 9 *Candida albicans*, 10 *Aspergillus niger*.

In view of the antimicrobial activity of other derivatives of 10-piperazinodibenzo[*b,f*]thiepins³⁶ the compounds described here were tested at the bacteriological department of this institute (Dr A. Šimek, Dr J. Turinová) *in vitro* for their inhibitory activity toward 14 representative types of microorganisms. Table III shows the minimum found inhibitory concentrations in µg/ml. It should be noted that all the compounds up to a concentration of 125 µg/ml were inactive toward *Pseudomonas aeruginosa*, *Escherichia coli*, *Salmonella typhi abdominalis* and *Proteus vulgaris*. Table III shows the 8-*n*-butyl derivatives (*Ic*, *Ilc*, *VIIc*) and the 8-cyclopentyl derivatives (*If*, *Ilf*) to be relatively highly bacteriostatic, particularly toward cocci; they are more potent than the corresponding ethyl, isopropyl and cyclopropyl derivatives. This led to the preparation of the 8-*n*-octyl derivative *Id* but this was found again to be less effective. Hence the assumption of possible dissociation of central from antimicrobial activity with extending the alkyl chain in position 8 was not confirmed.

EXPERIMENTAL

The melting points of analytical preparations were determined in Kofler's block and are not corrected; the samples were dried for 8 h at a suitable temperature (100°C maximum) at oil-pump vacuum over P₂O₅. The UV spectra (in methanol unless stated otherwise) were recorded in a Unicam SP 700 spectrophotometer, the IR spectra (in Nujol unless stated

otherwise) in a Unicam SP 200 G or in an Infracan (Hilger and Watts) spectrophotometer, the NMR spectra (in CDCl_3 unless stated otherwise) in a ZKR 60 (Zeiss, Jena) spectrometer. The mass spectrum was obtained in a MS 902 (AEI) mass spectrometer.

4-Cyclopentylbenzenesulfonyl Chloride (XXII)

Chlorosulfonic acid (285 g) was added dropwise under stirring at 4–8°C over a period of 45 min to a solution of 90 g cyclopentylbenzene³¹ (b.p. 114–116°C/25 Torr) in 300 ml chloroform. The mixture was stirred for another hour at 10°C and then decomposed with ice and water. The product was isolated by extraction with chloroform. A total of 90 g (60%) crude sulfonyl chloride was obtained (residue), a sample of which was recrystallized for analysis from light petroleum; m.p. 52–54°C. IR spectrum: 838 (2 vicinal aromatic C—H), 1180 and 1385 (SO_2Cl), 1595 cm^{-1} (Ar). NMR spectrum: δ 8.00 (mcd, 2 H, aromatic protons in the vicinity of SO_2Cl), 7.50 (mcd, 2H, remaining aromatic protons), 2.85–3.35 (m, 1H, CH—Ar), 1.25–2.40 (m, 8H, CH_2 groups of cyclopentane). For $\text{C}_{11}\text{H}_{13}\text{ClO}_2\text{S}$ (244.7) calculated: 53.98% C, 5.35% H, 14.49% Cl, 13.10% S; found: 53.87% C, 5.32% H, 14.54% Cl, 12.92% S.

4-Cyclopentylbenzenesulfonamide (XXIII)

A mixture of 1.5 g sulfonyl chloride XXII and 15 ml concentrated aqueous ammonia was heated under stirring for 20 min to 65–70°C and then left for 2 h at room temperature. The precipitated product was filtered and recrystallized from aqueous ethanol; m.p. 124–125°C. NMR spectrum: δ 7.89 (mcd, 2H, $J = 9.0$ Hz, aromatic protons in the vicinity of SO_2NH_2), 7.38 (mcd, $J = 9.0$ Hz, 2H, remaining aromatic protons), 5.24 (s, 2H, SO_2NH_2), 3.00 (m, 1H, CH—Ar), 1.50–2.30 (m, 8H, 4 CH_2 of cyclopentane). For $\text{C}_{11}\text{H}_{15}\text{NO}_2\text{S}$ (225.3) calculated: 58.63% C, 6.71% H, 6.22% N, 14.23% S; found: 59.00% C, 6.74% H, 6.31% N, 14.49% S.

Cyclopropylbenzene

A. Crude 1-phenyl-1,3-dibromopropane²⁰ (69.5 g) was added dropwise under stirring at about 50°C over the period of 60 min to a mixture of 125 ml dimethylformamide and the Zn—Cu-couple (prepared from 32 g zinc and 100 ml 2% solution of cupric sulfate)²¹. The mixture was processed according to ref.²⁰ and a total of 23.9 g (84%) product was obtained: b.p. 58–62°C/7 Torr.

B. A double batch was carried out at 58–60°C and 1-phenyl-1,3-dibromopropane was added dropwise over 30 min. The mixture was decomposed with water²⁰, steam-distilled and the product was isolated by extraction with ether. After distilling off the cyclopropylbenzene a residue of 10.6 g was obtained, m.p. 90–92°C (cyclohexane). According to analysis and NMR spectrum we are dealing here with *meso*-3,4-diphenylhexane (XIX) with a reported²² m.p. of 92–93°C. NMR spectrum: δ 7.25 (m, 10H, aromatic protons), 2.55 (m, 2H, CH—CH), c. 1.30 (m, 4H, 2 CH_2), 0.48 (t, $J = 7.0$ Hz, 6H, 2 CH_3). For $\text{C}_{18}\text{H}_{22}$ (238.4) calculated: 90.70% C, 9.30% H; found: 90.70% C, 9.56% H.

4-Isopropylthiophenol

A solution of 700 g crude 4-isopropylbenzenesulfonyl chloride³ in 800 ml acetic acid was added dropwise over 1 h to a boiling mixture of 1320 ml acetic acid, 300 g red phosphorus and 17 g iodine and the mixture was refluxed under stirring for 3 h. After cooling, it was combined slowly with 400 ml water, the mixture was refluxed for 1 h and, after cooling, it was poured into 10 l icecold

water. The product was isolated by extraction with chloroform and redistilled; 424 g (87%), b.p. 100—103°C/12 Torr. For a product prepared differently a b.p. of 104°C/15 Torr was reported³.

4-n-Octylthiophenol

Similarly to the preceding case, 127 g crude 4-n-octylbenzenesulfonyl chloride¹⁶ was reduced with 24 g red phosphorus and 19 g iodine in 160 ml boiling acetic acid. Analogous treatment (extraction with benzene) led to 63.4 g (64%) product boiling at 124—130°C/0.3 Torr. Literature¹⁵ reports a b.p. of 136—137°C/2.3 Torr. for the compound prepared differently.

4-n-Dodecylthiophenol (XVII)

n-Dodecylbenzene¹⁷ (29.6 g; b.p. 186—190°C/15 Torr) was added dropwise under stirring over 45 min to 26 ml chlorosulfonic acid at 5—10°C, the mixture was stirred for 4 h at room temperature and then was left to stand overnight. After decomposition with ice and water the crude 4-(n-dodecyl)benzenesulfonyl chloride (35 g) was isolated by extraction with ether and evaporation of the extract. The total amount of the product was then reduced similarly to preceding cases with the aid of 6.4 g red phosphorus and 3.4 g iodine in 45 ml boiling acetic acid. Analogous treatment yielded 17.7 g (68%) product boiling at 177—181°C/0.7 Torr. For analysis, a redistilled middle fraction was used; b.p. 162—163°C/0.6 Torr. For $C_{18}H_{30}S$ (278.5) calculated: 77.63% C, 10.86% H, 11.51% S; found: 77.30% C, 10.83% H, 11.46% S.

4-n-Butylthiophenol

A solution of 23.2 g crude 4-n-butylbenzenesulfonyl chloride³ in 200 ml ether was added dropwise over 1 h to a suspension of 9.45 g $LiAlH_4$ in 100 ml ether and the mixture was refluxed for 2 h. After cooling, 150 ml water-saturated ether was added, followed with 400 ml 2.5M-HCl. After filtration, the ether phase was separated, the aqueous phase was extracted with ether, the ether solutions were dried with $MgSO_4$ and distilled; 9.0 g (55%), b.p. 130—135°C/30 Torr, or 120—125°C/20 Torr. For a compound prepared differently ref.³ reported a b.p. of 119°C/14 Torr.

4-Cyclopentylthiophenol (XXIV)

As in the preceding case, 194 g sulfonol chloride XXII was reduced with 75 g $LiAlH_4$ and the almost theoretical yield (130 g) of a crude residue was used for further work. Sample for analysis was distilled; b.p. 165°C/25 Torr. For $C_{11}H_{14}S$ (178.2) calculated 74.13% C, 7.92% H, 17.96% S; found: 74.14% C, 7.81% H, 17.61% S.

4-Cyclopropylthiophenol (XX)

A Grignard reagent was prepared from 6.4 g Mg and 49.4 g 4-cyclopropylbromobenzene²³ (b.p. 102°C/7 Torr) in 140 ml ether, using several drops of 1,2-dibromoethane for initiating the reaction. The preparation of the reagent took 9 h. After dilution with 80 ml ether the solution was combined over 4 h at room temperature with 6.5 g sulfur flowers. The suspension was stirred for 1 h at room temperature and then refluxed for 1 h. After standing overnight it was diluted with 100 ml ether and slowly poured into 280 ml water and 50 ml hydrochloric acid. Processing of the ether layer yielded 17.5 g (47%) product boiling at 110—116°C/12 Torr, which, even after redistillation, does not yield satisfactory analytical values. However, it was processed further without difficulties. For $C_9H_{10}S$ (150.2) calculated: 71.95% C, 6.71% H, 21.34% S; found: 72.73% C, 6.83% H, 20.38% S.

Di(4-cyclopropylphenyl)disulfide (*XXI*)

Thiol *XX* (1.5 g) was added to 15 ml 1M-NaOH and, under stirring at 5°C, this was followed with 1.1 ml 30% H₂O₂ in 9 ml water. The precipitate formed was filtered, washed with water and dried; 1.2 g (81%), m.p. 71–72.5°C (light petroleum). IR spectrum (KBr): 823 (2 vicinal aromatic C—H), 1498 and 1600 cm⁻¹ (Ar). NMR spectrum: δ 7.44 (d, $J = 9.0$ Hz, 4H, aromatic protons in the vicinity of S), 7.00 (d, $J = 9.0$ Hz, 4H, remaining aromatic protons), 1.45–2.10 (m, 2H, 2CH—Ar), 0.40–1.15 (m, 8H, 4CH₂ of cyclopropyls). For C₁₈H₁₈S₂ (298.5) calculated: 72.43% C, 6.07% H, 21.50% S; found: 72.59% C, 6.12% H, 21.45% S.

2-(4-Cyclopropylphenylthio)benzoic Acid (*IXe*) (Method *A*)

Thiol *XX* (7.5 g) was added to a solution of 10.8 g KOH in 110 ml water at 50°C, the mixture was stirred for a while and then 0.45 g "molecular" copper and 12.5 g 2-iodobenzoic acid⁴ was added and the mixture was refluxed under stirring for 7 h. It was filtered while hot with charcoal and the filtrate was cooled and acidified with 2.5M-HCl. The precipitated product was filtered, washed with water and recrystallized from aqueous ethanol; 10.6 g (80%), m.p. 192–194°C. IR spectrum (KBr): 734 (4 vicinal aromatic C—H), 815 (2 vicinal aromatic C—H), 1252 (C—O), 1672 cm⁻¹ (Ar—COOH). NMR spectrum: δ 12.00 (bs, disappears after D₂O, 1H, COOH), 8.16 (m, 1H, aromatic 6-H), 7.50 (d, $J = 9.0$ Hz, 2H, aromatic 2',6'-H₂), 7.13 (d, $J = 9.0$ Hz, 2H, aromatic 3', 5'-H₂), c. 7.10 (m, 2H, aromatic 4,5-H₂), 6.80 (m, 1H, aromatic 3-H), 1.90 (m, 1H, CH—Ar), 0.60–1.20 (m, 4H, 2CH₂ of cyclopropane). The analytical data are shown in Table I.

2-(4-Isopropylphenylthio)benzyl Alcohol (*Xb*) (Method *B*)

770 ml of a 70% benzene solution of sodium bis(2-methoxyethoxy)dihydroaluminat were added dropwise under stirring over 2 h to a solution of 362 g acid *IXb* (ref.³) in 2800 ml benzene. The temperature rose spontaneously to 40°C. The mixture was stirred for 3 h at room temperature and then decomposed under external cooling by adding dropwise 2000 ml 10% NaOH. The benzene phase was separated and the aqueous one extracted with more benzene. Usual treatment of the organic phases yielded 324 g (94%) product boiling at 170–170.5°C/1.5–2 Torr. IR spectrum: 752 (4 vicinal aromatic C—H), 822 (2 vicinal aromatic C—H), 1015 (CH₂OH), 1360 (isopropyl), 1590 (Ar), 3370 cm⁻¹ (OH). The analytical data are shown in Table I.

2-(4-n-Octylphenylthio)benzyl Alcohol (*Xd*)

Acid *IXd* (6.85 g) was added in parts under stirring to a suspension of 2.0 g LiAlH₄ in 40 ml ether and the mixture was refluxed for 6 h. It was cooled, decomposed by adding dropwise 7.5 ml water and 2 ml 20% NaOH, stirred for 30 min, the solid was filtered, the filtrate was dried with Na₂SO₄ and evaporated. A total of 5.8 g (88%) crude product was obtained, the sample of which was redistilled for analysis; b.p. 195°C/0.4 Torr. The analytical data are shown in Table I.

2-(4-Cyclopentylphenylthio)benzyl Chloride (*Xf*) (Method *C*)

SOCl₂ (18 g) was added dropwise under stirring at room temperature over a period of 15 min to a solution of 28.4 g alcohol *Xf* in 40 ml benzene and the mixture was refluxed for 30 min. After evaporation of excess SOCl₂ and benzene the product was distilled; 24.2 g (80%), b.p. 180–182°C/1 Torr or 210–215°C/3 Torr. The analytical data are shown in Table I.

2-(4-Isopropylphenylthio)phenylacetonitrile (*XIb*) (Method *D*)

A solution of 18.6 g chloride *XIb* in 35 ml ethanol was added to a solution of 9.1 g KCN in 13 ml water and the mixture was refluxed for 7 h. After cooling, it was diluted with 200 ml water and the product was isolated by extraction with ether; 15.5 g (86%), b.p. 158–162°C/1 Torr. NMR spectrum: δ 7.15–7.70 (m, 8H, aromatic protons), 3.84 (s, 2H, ArCH₂CN), 2.86 (d, $J = 7.0$ Hz, 1H, Ar—CH), 1.21 (d, $J = 7.0$ Hz, 6H, 2CH₃). The analytical data are shown in Table I.

2-(4-Ethylphenylthio)phenylacetic Acid (*XIIIa*) (Method *E*)

A mixture of 123 g nitrile *XIIa*, 850 ml ethanol and 136 g KOH in 125 ml water was refluxed for 5 h. After partial cooling, it was diluted with 1250 ml water (50°C), the mixture was filtered with charcoal and the filtrate made acid with excess dilute hydrochloric acid. The precipitated acid was filtered, washed with water and recrystallized from aqueous ethanol; 110 g (83%), m.p. 96–99°C. IR spectrum (KBr): 745 (4 vicinal aromatic C—H), 823 (2 vicinal aromatic C—H), 930, 1244, 1407, 1692 and 2600 cm⁻¹ (COOH). The analytical data are shown in Table I.

2-(4-Cyclopropylphenylthio)phenylacetic Acid (*XIIIe*)

Thiol *XX* (9.7 g) was added to a solution of 13.8 g KOH in 140 ml water at 40°C, the mixture was stirred for 5 min, 0.9 g "molecular" copper and 16.2 g 2-(2-iodophenyl)acetic acid^{26,27} was added and the mixture refluxed under stirring for 4 h. After standing overnight, 0.3 g copper was added and refluxing continued for 7 h. It was filtered while hot with charcoal and the filtrate was made acid with hydrochloric acid at 70°C to pH 1. The precipitated product was filtered, washed with water and dried; 16.8 g (96%), m.p. 122–122.5°C (aqueous ethanol). IR spectrum (CHCl₃): 820 (Ar—H), 1492 and 1590 (Ar), 928, 1235 and 1705 cm⁻¹ (COOH). NMR spectrum: δ 7.31 (s, 4H, aromatic protons in positions 3, 4, 5, 6), 7.25 and 7.00 (2d, $J = 9.0$ Hz, 4H, aromatic protons in positions 2', 3', 5', 6'), 3.80 (s, 2H, ArCH₂COO), 1.80 (m, 1H, Ar—CH), 0.50–1.15 (m, 4H, 2CH₂ of cyclopropane). The analytical data are shown in Table I.

8-Isopropylidibenzo[*b,f*]thiepin-10(11H)-one (*XIVb*) (Method *F*)

Acid *XIIIb* (14.3 g) was added at 100°C to 30 g polyphosphoric acid and the mixture was heated under stirring for 2 h to 150°C. After cooling, 60 ml benzene was added and the mixture decomposed under cooling with 60 ml water. The product was isolated by extraction with benzene, the extract was washed with 5% NaOH and water, dried with K₂CO₃ and evaporated. The crude product was distilled; 10.2 g (76%), b.p. 167–170°C/1.5 Torr. UV spectrum: λ_{\max} 242.5 nm (log ϵ 4.33), 335 nm (3.63), infl. 265 nm (4.07). IR spectrum (film): 748 (4 vicinal aromatic C—H), 838 (2 vicinal aromatic C—H), 1290 (C—O), 1360 (isopropyl), 1584 (Ar), 1675 cm⁻¹ (Ar—CO). The analytical data are shown in Table I.

8-n-Butyldibenzo[*b,f*]thiepin-10(11H)-one (*XIVc*) (Method *F-I*)

Acid *XIIIc* (60 g) was added to a hot mixture of 540 g polyphosphoric acid and 180 ml toluene and the mixture was refluxed under stirring for 2 h. After cooling, 500 ml benzene was added and decomposed with 750 ml ice-cold water. The organic phase was washed with water and 5% NaOH (acidification of the alkaline solution recovered 12.0 g acid *XIIIc*), dried with K₂CO₃ and processed by distillation; 37 g (82%), b.p. 190–195°C/2 Torr. UV spectrum: λ_{\max} 241 nm (log ϵ 4.29), 334 nm (3.55). IR spectrum: 745, 752 (4 vicinal aromatic C—H), 828 (2 vicinal aromatic C—H),

870 (solitary aromatic C—H), 1075 and 1288 (C—O), 1595 (Ar), 1670 cm^{-1} (Ar—CO). NMR spectrum: δ 8.06 (d, $J = 2.5$ Hz, 1H, aromatic 9-H), 7.00–7.30 (m, 6H, remaining aromatic protons), 4.34 (s, 2H, ArCH_2CO), 2.59 (t, $J = 7.0$ Hz, 2H, ArCH_2 of butyl), c. 1.45 (m, 4H, C— CH_2CH_2 —C of butyl), 0.88 (t, 3H, CH_3). The analytical data are shown in Table I.

8-Cyclopropyldibenzo[*b,f*]thiepin-10(11*H*)-one (XIVe)

A mixture of 80 ml trifluoroacetic anhydride and 13.2 g acid XIIIe was stirred for 4 h at room temperature. After standing overnight, 6 ml benzene were added and the mixture refluxed for 7 h, diluted with 25 ml benzene and decomposed with 1 l 12% NaHCO_3 . The mixture was extracted with benzene and the extract was evaporated. The residue was dissolved in benzene and chromatographed on a column of alumina (300 g, activity II). Elution with benzene yielded 6.8 g (55%) XIVe, m.p. 82–86°C; after recrystallization from cyclohexane, m.p. 85–86°C. UV spectrum: λ_{max} 242.5 nm ($\log \epsilon$ 4.39), 260 nm (4.15), 342 nm (3.60). IR spectrum (KBr): 755 (4 vicinal aromatic C—H), 842 (2 vicinal aromatic C—H), 878 (solitary aromatic C—H), 1475 and 1600 (Ar), 1670 cm^{-1} (Ar—CO). NMR spectrum: δ 8.06 (d, $J = 2.5$ Hz, 1H, aromatic 9-H), 7.63 (d, $J = 9.0$ Hz, 1H, aromatic 6-H), 7.21 (mcd, $J = 9.0$; 2.5 Hz, 1H, aromatic 7-H), 7.25–7.90 (m, 4H, remaining aromatic protons), 4.40 (s, 2H, ArCH_2CO), 1.88 (m, 1H, Ar—CH), 0.50–1.15 (m, 4H, 2CH_2 of cyclopropane). The analytical data are shown in Table I.

8-n-Butyl-10-hydroxy-10,11-dihydrodibenzo[*b,f*]thiepin (XVc) (Method G)

A solution of 11.4 g NaBH_4 with 0.65 ml 20% NaOH in 65 ml water was added dropwise over 1 h under stirring to a solution of 29.0 g ketone XIVc in 400 ml ethanol, the mixture was refluxed for 4 h, the ethanol was evaporated at reduced pressure, the residue was diluted with 400 ml water and the product was extracted with benzene. The extract was washed with water, 5% NaOH, dried and evaporated. The residue crystallized from a mixture of benzene and light petroleum; 24 g (82%), m.p. 62–63°C. IR spectrum (KBr): 743 (4 vicinal aromatic C—H), 820 (2 vicinal aromatic C—H), 867 and 882 (solitary aromatic C—H), 1047 (CHOH), 992, 1465 (Ar), 3280 and 3330 cm^{-1} (OH). NMR spectrum: δ 6.80–7.60 (m, 7H, aromatic protons), 5.25 (m, dd after D_2O , $J = 9.0$; 4.0 Hz, 1H, Ar—CH—O), 3.73 and 3.28 (2dd, $J = 14.0$; 4.0 Hz and 14.0; 4.0 Hz, 2H, ArCH_2 in a ring), 2.53 (t, $J = 7.0$ Hz, 2H, ArCH_2 of butyl), 2.15 (d, disappears after D_2O , 1H, OH), 1.42 (m, 4H, C— CH_2CH_2 —C of butyl), 0.88 (t, 3H, CH_3). The analytical data are shown in Table I.

8-Cyclopropyl-10-chloro-10,11-dihydrodibenzo[*b,f*]thiepin (XVie) (Method H)

Powder CaCl_2 (5 g) was added to a solution of 6.9 g alcohol XVe in 90 ml benzene and the suspension was saturated at room temperature with anhydrous HCl for 3 h. After standing overnight, it was filtered and the filtrate was evaporated at reduced pressure. The residue crystallized from cyclohexane; 7.10 g (97%), m.p. 76.5–77°C. NMR spectrum: δ 7.05–7.90 (m, 6H, aromatic protons in positions 1, 2, 3, 4, 6 and 9), 6.90 (mcd, $J = 9.0$; 2.5 Hz, 1H, aromatic proton in position 7), 5.85 (dd, $J = 9.0$; 4.0 Hz, 1H, Ar—CH—Cl), 4.02 and 3.65 (2dd, $J = 14.0$; 4.0 and 14.0; 9.0 Hz, 2H, ArCH_2 in the ring), 1.85 (m, 1H, Ar—CH of cyclopropane), 0.50–1.10 (m, 4H, 2CH_2 of cyclopropane). The analytical data are shown in Table I.

8-Isopropyl-10-(4-methylpiperazino)-10,11-dihydrodibenzo[*b,f*]thiepin (Ib) (Method J)

1-Methylpiperazine (7.5 g) was added to a solution of 7.2 g chloride XVib in 8 ml chloroform and the mixture was refluxed for 6 h. After evaporation of the chloroform the residue was dissolved

in 100 ml benzene, the solution was washed with 2.5M-NaOH and with water. The base was then transferred by shaking with 3 times 70 ml 1.25M-H₂SO₄ to the aqueous phase. Evaporation of the separated benzene phase and distillation of the residue yielded a small amount of 2-isopropylidibenzo[*b,f*]thiepin (*VIIIb*), b.p. 177–180°C/1.5 Torr. UV spectrum: λ_{\max} 221 nm (log ϵ 4.43), 261 nm (4.39), 297 nm (3.72). IR spectrum: 746 (4 vicinal aromatic C—H), 786 (*cis*-CH=CH), 826 (2 vicinal aromatic C—H), 880 (solitary aromatic C—H), 1061, 1553, 1589 cm⁻¹ (Ar). NMR spectrum: δ 7.00–7.60 (m, 7H, aromatic protons), 6.99 (s, 2H, olefinic CH=CH), 2.80 (m, 1H, Ar—CH of isopropyl), 1.18 (d, $J = 7.0$ Hz, 6H, 2CH₃). The analytical data are shown in Table I.

The acid aqueous phase was made alkaline with excess aqueous NH₄OH and the liberated base was isolated by extraction with benzene; 8.0 g (91%). It crystallized on standing, m.p. 118 to 120°C (benzene–light petroleum). NMR spectrum: δ 7.55 (d, $J = 2.5$ Hz, 1H, aromatic 9-H), 7.00–7.50 (m, 5H, aromatic protons in positions 1, 2, 3, 4, 6), 6.93 (mcd, 1H, aromatic 7-H), 3.00–4.00 (m, 3H, Ar—CH₂CH—Ar), c. 3.00 (obscured m, 1H, Ar—CH of isopropyl), 2.64 (m, 4H, CH₂N¹CH₂ of piperazine), 2.51 (m, 4H, CH₂N⁴CH₂ of piperazine), 2.30 (s, 3H, NCH₃), 1.18 (d, $J = 7.0$ Hz, 6H, 2CH₃ of isopropyl). Further data are in Table I.

8-Ethyl-10-piperazino-10,11-dihydrodibenzo[*b,f*]thiepin (*IVa*) (Method *K*)

A mixture of 12.0 g carbamate *IIIa*, 17 ml ethanol and 8.4 g KOH was refluxed for 3 h in a 120°C bath. It was then cooled, dissolved in 60 ml water and extracted with benzene. After washing and drying of the extract, the benzene was evaporated and the residue recrystallized from 12 ml ethanol; 8.2 g (84%), m.p. 113–114°C (acetone). NMR spectrum: δ 7.00–8.50 (m, 7H, aromatic protons), 2.95–4.00 (m, 6H, ArCH₂CHAr, ArCH₂ and NH), 2.30–2.95 (m, 8H, CH₂ groups of piperazine), 1.14 (t, 3H, C—CH₃ of ethyl). Further data are contained in Table I.

8-Ethyl-10-[4-(3-capryloyloxypropyl)piperazino]-10,11-dihydrodibenzo[*b,f*]thiepin (*Va*) (Method *L*)

Capryloyl chloride⁹ (10.5 g, b.p. 105°C/35 Torr) was added to a solution of 11.4 g aminoalcohol *IIa* in a mixture of 60 ml benzene and 20 ml chloroform, the mixture was stirred and left to stand for 4 days at room temperature. It was then heated for 1 h to 60°C, cooled, 100 ml water was added, made alkaline with ammonia and extracted with benzene. Processing of the extract yielded the crude base which was dissolved in 35 ml acetone and neutralized with a solution of 6.9 g maleic acid in 25 ml acetone. A total of 13.0 g (59%) crystalline di(hydrogen maleate) was obtained; m.p. 144–146°C (acetone). The analytical data are shown in Table I.

8-n-Butyl-10-(4-methylpiperazino)dibenzo[*b,f*]thiepin (*VIIc*) (Method *M*)

1-Methylpiperazine (15 g) was added to a solution of 8.5 g ketone *XIVc* in 70 ml benzene, followed over a period of 5 min with a solution of 3.0 g TiCl₄ in 20 ml benzene. The mixture was refluxed under stirring for 24 h. After cooling, it was decomposed with 100 ml water, the inorganic fraction was filtered and the filtrate processed by benzene extraction. Washing of the extract with water and evaporation yielded the crude base, which was recrystallized from benzene; 6.2 g (57%), m.p. 140–141°C. NMR spectrum: δ 7.00–7.65 (m, 7H, aromatic protons), 6.34 (s, 1H, Ar—CH=C), 3.00 and 2.55 (2 m, 10H, ArCH₂ of butyl and 4 CH₂ of piperazine), 2.36 (s, 3H, NCH₃), 1.40 (m, 4H, C—CH₂CH₂—C of butyl), 0.90 (t, 3H, CH₃ of butyl). Further data are included in Table I.

2-Cyclopropylthioxanthone (XXV)

A mixture of 11.4 g acid IXe and 100 g trifluoroacetic anhydride was stirred for 5 h at 35°C. After standing overnight, 120 ml of benzene was added and the mixture was stirred for 8 h at 35°C. After cooling, it was decomposed by pouring into 500 ml 10% NaHCO₃, 100 ml benzene was added, followed with 30 ml 5M-NaOH. After separation, acidification of the alkaline layer recovered 4.1 g acid IXe. The benzene layer was washed with water, dried with Na₂SO₄ and evaporated. A total of 3.8 g (57%) crude product melting at 95–98°C was obtained. For analysis it was recrystallized from a mixture of benzene and light petroleum, m.p. 104–106°C (change of the crystal modification at 95°C). The mass spectrum shows a molecular ion at $m/e = 252$ which corresponds to the assumption; the fragmentation is in agreement with the expected structure. For C₁₆H₁₂OS (252.3) calculated: 75.15% C, 4.80% H, 12.71% S; found: 76.07% C, 5.01% H, 12.80% S.

2-Cyclopropyl-9-(3-dimethylaminopropyl)thioxanthene-9-ol (XXVI)

Reaction of 0.8 g Mg and 4.06 g 3-dimethylaminopropyl chloride in 15 ml tetrahydrofuran produced a Grignard reagent, using a grain of iodine and several drops of 1,2-dibromoethane for initiating the reaction. The preparation of the reagent took 3 h. After cooling, a suspension of 4.0 g ketone XXV in 25 ml tetrahydrofuran was added and the mixture was left to stand for 3 days at room temperature. It was then decomposed with 100 ml 20% NH₄Cl and extracted with benzene. Processing of the extract yielded 4.7 g (87%) crude product which was recrystallized for analysis from a mixture of cyclohexane and light petroleum, m.p. 123–125.5°C. IR spectrum (CHCl₃): 770 (4 vicinal aromatic C—H), 820 (2 vicinal aromatic C—H), 900 (solitary aromatic C—H), 1020 (C—OH), 1596 (Ar), 2630 and 2700 cm⁻¹ (OH in a hydrogen bond at N). NMR spectrum: δ 8.02 (m, 1 H, aromatic 8-H), 7.75 (d, $J = 3.0$ Hz, 1 H, aromatic 1-H), 7.10–7.60 (m, 4 H, aromatic protons in positions 4, 5, 6, 7), 6.98 (mcd, $J = 9.0; 3.0$ Hz, 1 H, aromatic 3-H), 2.40 (s, 6 H, 2 N—CH₃), 1.60–2.40 (m, 5 H, Ar—CH of cyclopropyl and CH₂—C—CH₂N of the side chain), 0.50–1.40 (m, 6 H, 2 CH₂ of cyclopropyl and C—CH₂—C of the side chain). For C₂₁H₂₅NOS (339.5) calculated: 74.29% C, 7.42% H, 4.12% N, 9.46% S; found: 73.82% C, 7.69% H, 4.08% N, 9.23% S.

2-Cyclopropyl-9-(3-dimethylaminopropylidene)thioxanthene (XXVII)

A mixture of 125 ml water, 35 g H₂SO₄ and 8.25 g aminoalcohol XXVI was refluxed for 2 h, cooled, made alkaline with NH₄OH and the product was isolated by extraction with chloroform. The crude base (7.8 g) was converted by neutralization with maleic acid (2.9 g) in 2-propanol to the crystalline hydrogen maleate (9.2 g) which, after several recrystallizations from a mixture of 2-propanol and ether melted at 140–143°C. UV spectrum: λ_{\max} 231 nm (log ϵ 4.55), 271.5 nm (4.18), 329.5 nm (3.50). IR spectrum (KBr): 755 (4 vicinal aromatic C—H), 835 (2 vicinal aromatic C—H), 892 (solitary aromatic C—H), 1460 (Ar), 1528 and 1615 (COO⁻), 2460 cm⁻¹ (NH⁺). For C₂₅H₂₇NO₄S (437.5) calculated: 68.62% C, 6.22% H, 3.20% N, 7.33% S; found: 68.28% C, 6.52% H, 3.24% N, 7.31% S.

The authors are indebted to Drs B. Kakáč, E. Svátek, J. Holubek and Mrs P. Vejdělková of the physico-chemical department of this institute for registration and interpretation of IR, UV and NMR spectra. The mass spectrum was recorded by Dr M. Ryska, Institute of Macromolecular Chemistry, Czechoslovak Academy of Sciences, Prague. Technical cooperation with the preparative part of the work by Mrs M. Šebestíková is acknowledged. The analyses were done at the analytical department of this institute by Mr M. Čech, Mr K. Havel, Mrs V. Šmidová, Mrs J. Komancová, Mrs J. Hrdá, Mrs A. Slavíková and Mrs Z. Volková.

REFERENCES

1. Pelz K., Jirkovský I., Adlerová E., Metyšová J., Protiva M.: *This Journal* 33, 1895 (1968).
2. Schenker E., Herbst H.: *Fortschr. der Arzneimittelforsch.* 5, 324 (1963).
3. Pelz K., Protiva M.: *This Journal* 32, 2161 (1967).
4. Wachter W.: *Ber.* 26, 1744 (1893).
5. Jílek J. O., Šindelář K., Pomykáček J., Horešovský O., Pelz K., Svátek E., Kakáč B., Holubek J., Metyšová J., Protiva M.: *This Journal* 38, 115 (1973).
6. Zawisza T., Machoň Z., Kuczyński L.: *Acta Polon. Pharm.* 22 (6), 477 (1965).
7. Moore T. S., Boyle M., Thorn V. M.: *J. Chem. Soc.* 1929, 39.
8. Jílek J. O., Šindelář K., Dlabáč A., Kazdová E., Pomykáček J., Šedivý Z., Protiva M.: *This Journal* 38, 1190 (1973).
9. Fierz-David H. E., Kuster W.: *Helv. Chim. Acta* 22, 82 (1939).
10. Ralston A. W., Selby W. M.: *J. Am. Chem. Soc.* 61, 1019 (1939).
11. White W. A., Weingarten H.: *J. Org. Chem.* 32, 213 (1967).
12. Umio S., Ueda I., Sato Y., Maeno S. (Fujisawa Pharmaceutical Co., Ltd.): *Ger. Offen.* 1 801 523; *Chem. Abstr.* 71, 112 976 (1969).
13. Jílek J. O., Šindelář K., Metyšová J., Metyš J., Pomykáček J., Protiva M.: *This Journal* 35, 3721 (1970).
14. Wagner A. W.: *Chem. Ber.* 99, 375 (1966).
15. Kuliev A. M., Kuliev A. B., Mamedov F. N.: *Ž. Org. Chim.* 1 (10), 1787 (1965); *Chem. Abstr.* 64, 3392 (1966).
16. Cross J. M., Chiddix M. E. (General Aniline & Film Corp.): *U. S.* 2 740 814; *Chem. Abstr.* 50, 16 850 (1956).
17. Paquette R. G., Lingafelter E. C., Tartar H. V.: *J. Am. Chem. Soc.* 65, 686 (1943).
18. Boissier J. R., Malen Ch. (S. I. F. A.): *Fr.* 1 366 215 (*Appl.* 31. V. 1963); *Chem. Abstr.* 62, 570 (1965).
19. Boissier J. R., Malen Ch. (S. I. F. A.): *Fr.* 1 381 437 (*Appl.* 22. X. 1963); *Chem. Abstr.* 62, 10 444 (1965).
20. Corbin T. F., Hahn R. C., Schechter H.: *Org. Syn.* 44, 30 (1964).
21. Hennion G. F., Sheehan J. J.: *J. Am. Chem. Soc.* 71, 1964 (1949).
22. Späth E.: *Monatsh.* 34, 2010 (1913).
23. Levina P. J., Gembickij P. A., Treshova E. G.: *Ž. Obšč. Chim.* 33, 371 (1963).
24. Kurtz W., Fischer P., Effenberger F.: *Chem. Ber.* 106, 525 (1973).
25. Boissier J. R., Ratouis R., Dumont C., Derible P. H., Lavaux J. P.: *J. Med. Chem.* 13, 971 (1970).
26. Šindelář K., Metyšová J., Protiva M.: *This Journal* 37, 1734 (1972).
27. Taylor E. C., Kienzle F., Robey R. L., McKillop A., Hunt J. D.: *J. Am. Chem. Soc.* 93, 4845 (1971).
28. Bourne E. J., Stacey M., Tatlow J. C., Tedder J. M.: *J. Chem. Soc.* 1951, 718.
29. Ferrier R. J., Tedder J. M.: *J. Chem. Soc.* 1957, 1435.
30. Harfenist M.: *J. Org. Chem.* 28, 1834 (1963).
31. Hai P. V., Buu-Hoi N. P., Xuong N. D.: *J. Org. Chem.* 23, 39 (1958).
32. Jílek J. O., Rajšner M., Pomykáček J., Protiva M.: *Českoslov. farm.* 14, 294 (1965).
33. Šindelář K., Metyšová J., Protiva M.: *This Journal* 38, 2137 (1973).
34. Jílek J. O., Metyšová J., Pomykáček J., Protiva M.: *This Journal* 33, 1831 (1968).
35. Metyš J., Metyšová J., Votava Z.: *Českoslov. farm.* 15, 526 (1966).
36. Šimek A., Čapek A., Turinová J., Tůma J.: *Folia Microbiol.* 13, 134 (1968).

Translated by A. Kotyk.