

Reactivities of carbonyl sulfide (COS), carbon disulfide (CS₂) and carbon dioxide (CO₂) with transition metal complexes

Krishna K. Pandey

School of Chemistry, Devi Ahilya University Indore, Khandwa Road, Indore 452 001, India

Received 19 November 1993; in revised form 5 May 1994

Contents

Abstract	38
1. Introduction	39
2. Coordination chemistry of carbonyl sulfide	40
2.1. Transition metal–carbonyl sulfide bonding and energetics	40
2.1.1. End-on coordination (M– η^1 -OCS)	41
2.1.2. Side-on coordination (M– η^2 -COS)	41
2.1.3. Bridging carbonyl sulfide complexes	42
2.2. Preparation and structure	42
2.3. Reactions of carbonyl sulfide complexes	46
2.3.1. Carbonylation reaction	46
2.3.2. Reductive disproportionation reaction: dithiocarbonate complexes	47
2.3.3. Insertion reaction of carbonyl sulfide into M–X bonds	50
2.3.4. Formation of sulfide complexes	52
3. Coordination chemistry of carbon disulfide	52
3.1. Transition metal–carbon disulfide bonding and energetics	52
3.1.1. η^1 -end-on coordination	52
3.1.2. η^2 -side-on coordination	53
3.1.3. Bridging carbon disulfide complexes	53
3.2. Preparation and structure	54
3.2.1. Mononuclear metal–carbon disulfide complexes	54
3.2.2. Bridging carbon disulfide complexes	57
3.3. Reactions of carbon disulfide complexes	60
3.3.1. Thiocarbonylation reaction	60
3.3.2. Reductive dimerization of coordinated carbon disulfide	61
3.3.3. Insertion reaction of carbon disulfide	64
3.3.4. Alkylation reaction of carbon disulfide	71
3.3.5. Miscellaneous reactions	73
4. Coordination chemistry of carbon dioxide	75
4.1. Transition metal–carbon dioxide bonding and energetics	75
4.2. Preparation and structure	78
4.2.1. Mononuclear transition metal carbon dioxide complexes	78
4.2.2. Bimetallic and trimetallic carbon dioxide complexes	82

4.3. Spectroscopic studies	85
4.4. Insertion reactions of carbon dioxide complexes	85
4.4.1. Insertion into the M–H bond	85
4.4.2. Insertion into the M–C bond	89
4.4.3. Insertion into the M–N bond	89
4.4.4. Insertion into the M–O bond	90
4.4.5. Insertion into the M–Si bond	93
4.4.6. Insertion into the M–P bond	94
4.4.7. Oxidative coupling of carbon dioxide with coordinated organic substrate	94
4.4.8. Reduction of carbon dioxide	97
Acknowledgements	105
References	105

Abstract

Carbonyl sulfide, carbon disulfide and carbon dioxide interact with transition metal complexes to form complexes with a number of transition metals and undergo a variety of chemical transformations including insertion, dimerization, disproportionation, coupling and catalytic reactions. The coordination chemistry of these closely related molecules is very diverse. The metal-(η^2 -CS₂) complexes are much more stable than the metal-(η^2 -COS) and metal-(η^2 -CO₂) complexes. Elimination of sulfur from the metal-(η^2 -COS) complexes is more facile. Carbon dioxide undergoes coupling reactions with unsaturated organic compounds on transition metal centers to produce metallocycles. The reduction of carbon dioxide catalyzed by transition metal complexes, using different methods, is discussed.

Keywords: Reactivity; COS; CS₂; CO₂; Transition metal complexes

Abbreviations

alcn	HNCHCHCHO ⁻
bipy	2,2'-bipyridyl
<i>n</i> -Bu	<i>n</i> -butyl
<i>t</i> -Bu	tert-butyl
COD	cyclooctadiene
cp	cyclopentadienyl
Cy	cyclohexyl
dbu	diazabicyclo[5,4,0]undec-7-ene
diars	<i>o</i> -phenylenebisdimethylarsine
DMF	<i>N,N'</i> -dimethylformamide
dmpm	1,2-bis(dimethylphosphino)methane
dpam	1,2-bis(diphenylarsino)methane
dppe	1,2-bis(diphenylphosphino)ethane
dppm	bis(diphenylphosphino)methane
EDTA	ethylenediaminetetraacetate
EHMO	extended Hückel molecular orbital

en	ethylenediamine
Et	ethyl
HOMO	highest occupied molecular orbital
L	ligand
M	metal atom
Me	methyl
MeCN	acetonitrile
mes	2,4,6-Me ₃ C ₆ H ₂
NBD	norbornadiene
np ₃	tris(2-diphenylphosphinoethyl)amine, N(CH ₂ CH ₂ PPh ₂) ₃
pc	phthalocyanine
ph	phenyl
phen	1,10-phenanthroline
<i>n</i> -Pr	<i>n</i> -propyl
<i>i</i> -Pr	iso-propyl
py	pyridine
R	alkyl or aryl group
salen	bis(salicyldehydeethylenediimine)
sp	2-CH ₂ =CHC ₆ H ₄ PPh ₂
terpy	terpyridine
THF	tetrahydrofuran
TMED	<i>N,N,N',N'</i> -tetramethylethylenediamine
TPP	meso-tetraphenylporphyrin
triphos	1,1,1-tris(diphenylphosphinomethyl)ethane

1. Introduction

The interaction of heterocumulene molecules such as carbonyl sulfide (COS), carbon disulfide (CS₂) and carbon dioxide (CO₂), which are a potential source of C₁ chemistry, with transition metal complexes has been extensively studied in the past decade and has been shown to give rise to complexes with a number of transition metals and to produce a variety of chemical transformations including insertion, dimerization and disproportionation reactions. Since the first report by Baird and Wilkinson of the reaction of COS, which is structurally similar to CS₂ and CO₂, with a transition metal complex, this subject has received relatively little attention.

The important physical properties of the linear triatomic molecules COS, CS₂ and CO₂ are presented in Table 1 [1–24]. The values of ionization potential [2,11–18] and electron affinity [19–22] show that CS₂ is a relatively better σ -donor and a better π -acceptor than COS and CO₂. The bond dissociation energy of the COS molecule indicates that it can function as a good carbonylating agent in the presence of a suitable thiophilic agent. The carbon binding energy C_{1s} has been observed to shift 2.1 eV between COS and CS₂ and 2.3 eV between CO₂ and COS [25]. The shifts are due to the lower electronegativity of sulfur compared with oxygen.

Table 1
Selected properties of COS, CS₂ and CO₂

	COS	CS ₂	CO ₂	Reference
Point group	C _{∞v}	D _{∞h}	D _{∞v}	[1]
Ground state	1 _Σ ⁺	1 _{Σg} ⁺	1 _{Σg} ⁺	[2]
Boiling point (°C)	−50.2	46.3	−78.5	
LUMO	2π	2π _u	2π _u	[3]
HOMO	1π	1π _g	1π _g	[2]
Bond length (Å)	1.16 (C–O) 1.56 (C–S)	1.56 (C–S)	1.16 (C–O)	[4–9]
Bond energy (eV)	3.12 (OC–S) 6.81 (O–CS)	4.463	5.453	[4,10]
Ionization potential (eV)	11.2	10.09	13.78	[2,11–18]
Electron affinity (eV)	0.46	1.0	−0.6	[19–22]
IR data (cm ^{−1})	2062 859 521	1535 396	2396 677	[23,24]

The coordination chemistry of these closely related molecules is very diverse. The metal-(η^2 -CS₂) complexes are much more stable than the metal-(η^2 -COS) complexes. Sulfur abstraction from a metal-(η^2 -CS₂) complex is not a general procedure for the preparation of metal thiocarbonyl complex. In contrast, elimination of sulfur from metal-(η^2 -COS) complexes is more facile. Carbonyl sulfide chemistry has proven to be interesting with potential applications to carbonylation reactions. The weakening of the C=S bond of COS on coordination to a metal, due to $d\pi$ - π^* back donation, is apparently sufficient to promote elimination of sulfur. The stronger C=S bond of CS₂ is less susceptible to metal-promoted cleavage, but in the presence of a suitable sulfur acceptor can facilitate sulfur elimination.

Although CS₂ is structurally very similar to CO₂, its behavior towards metal centers is different. CS₂ is a very reactive molecule, while CO₂ is not.

A number of review articles, dealing with various aspects of the chemistry of COS [26,27], CS₂ [27–32] and CO₂ [33–53] have been written. Because of new developments, it is necessary to review recent research involving reactivities of COS, CS₂ and CO₂ with transition metal complexes. Carbon dioxide can be converted into useful organic compounds by coupling with an organic substrate using homogeneous transition metal catalysts [45–48,50,54–79]. These catalytic reactions of CO₂ with the organic substrate are not examined here. This article covers the chemistry of COS and CS₂ from 1982 to 1992, and the chemistry of CO₂ from 1983 to 1992.

2. Coordination chemistry of carbonyl sulfide

2.1. Transition metal–carbonyl sulfide bonding and energetics

General consideration of the electron donating and electron accepting properties of carbonyl sulfide shows three possible bonding modes with transition metals [26].

2.1.1. End-on coordination ($M-\eta^1\text{-OCS}$)

In this type of bonding, the transition metal behaves as a Lewis acid, accepting a lone pair of electrons from the 1π (HOMO) orbital of the COS into the corresponding metal vacant orbital.

2.1.2. Side-on coordination ($M-\eta^2\text{-COS}$)

In this bonding mode, the main bonding orbital is the overlap between the metal- $d\pi$ and the $\text{COS}-\pi^*$ orbitals. Thus, the main interaction is π -back donation from the metal atom to the COS ligand. Some contribution to the interaction between metal atom and COS arises from the COS ligand to metal σ -donation. Such complexes contain a three-membered ring.

In order to understand the nature and energetics of the interaction between metal and COS, ab initio molecular orbital calculations have been performed on the model systems $[\text{Fe}(\text{CO})_2(\text{PH}_3)_2(\eta^2\text{-COS})]$ with two possible side-on coordination modes (I) and (II) [80].

Bonding energy and Mulliken population analysis of the systems (I) and (II) are presented in Table 2. Results of the theoretical calculations suggest that COS prefers the $\eta^2\text{-C,S}$ coordination to the $\eta^2\text{-C,O}$ coordination, in agreement with the experimental evidence that no $\eta^2\text{-C,O}$ coordinated COS complex has so far been obtained.




Table 2

Binding energy and Mulliken population analysis of the systems (I) and (II) [80]

	System (I)	System (II)
Binding energy (kcal mol^{-1})	-35.8	-30.9
Atomic populations		
Fe s	6.40	6.39
Fe p	12.83	12.71
Fe d	7.17	7.18
COS	30.56	30.70
Net charges		
Fe	-0.40	-0.28
COS	-0.56	-0.70

The strength of the $d\pi(M)-\pi^*(COS)$ interaction increases as the overlap between $d\pi(M)$ and $\pi^*(COS)$ orbitals increases. The COS distortion on coordination lowers the $\pi^*(COS)$ orbital energy, reduces the gap between $d\pi(M)$ and $\pi^*(COS)$ and hence increases π -back donation. The π^* orbital energy of the distorted COS ligand is -0.0137 au for η^2-C,S coordination and -0.0099 au for η^2-C,O coordination, indicating stronger π -back donation in the η^2-C,S coordination mode. Mulliken population analysis reveals that the σ -donor and π -acceptor abilities of the η^2-C,S coordination mode are greater than those of the η^2-C,O coordination mode.

2.1.3. Bridging carbonyl sulfide complexes



In addition to the end-on coordination and side-on coordination modes, COS can function as a bridge between two metal atoms (III) and (IV).


2.2. Preparation and structure

Relatively few transition metal COS complexes have been reported in the literature. These complexes have been prepared using the free ligand which is readily available as a starting material. The most accessible technique for identifying metal–COS complexes is IR spectroscopy. The frequency range for the carbonyl stretching vibration in COS complexes is 1630 – 1765 cm^{-1} and the position of the in-ring $\nu(CMS)$ vibration, for $(M-\eta^2-COS)$ complexes lies in the regions 630 – 660 cm^{-1} and 800 – 860 cm^{-1} [26].


Titanium COS complex, $[H_4Ti(COS)]$, is prepared by the reaction of COS with a titanium hydride complex [81]. Reaction of carbonyl sulfide with the cyclooctadiene complex $[Ni(COD)(bipy)]$ results in the formation of the COS complex $[Ni(\eta^2-COS)(bipy)]$. Attempts to prepare $[Ni(\eta^2-COS)(PR_3)_2]$ ($R \equiv Ph$ or Cy) were unsuccessful [82]. The thermally labile COS complex $[(C_5Me_5)_2V(\eta^2-COS)]$ (V) is prepared by the reaction of $[(C_5Me_5)_2V]$ with COS [83]. Preparation of COS complexes of Ti, Ni, Pd and Pt have been reported [81,84]. Irradiation of $[Ni(S_2C_2O_2)(dppe)]$ in acetonitrile yields the carbonyl complex $[Ni(CO)_2(dppe)]$ and COS complex $[Ni(\eta^2-COS)(dppe)]$ [85].

Reaction of $[ReH(PMe_3)_5]$ with COS affords a white crystalline complex (VI) whose 1H NMR spectrum shows a singlet at δ 1.42, indicating a similar environment for the PMe_3 ligands, and a quintet at δ -3.10 due to $Re-H$. The IR spectrum






(V)



(VI)




(VII)

exhibits bands at 1935 cm^{-1} due to $\nu(\text{Re}-\text{H})$ and at 1791 cm^{-1} due to $\nu(\text{C}=\text{O})$ of the $\eta^2\text{-COS}$ ligand [86].


Exposing a toluene solution of $[\text{Ru}(\text{CO})_2(\text{PPh}_3)_3]$ to 1 atm COS for 5 min yields a cream white microcrystalline complex $[\text{Ru}(\text{CO})_2(\text{COS})(\text{PPh}_3)_2]$ (VII). It is stable in the solid state under a nitrogen atmosphere and decomposes readily in solution to afford $[\text{Ru}(\text{CO})_3(\text{PPh}_3)_2]$ and SPPH_3 . The IR spectrum of (VII) exhibits absorption bands at $1701, 1678\text{ cm}^{-1}$ due to $\nu(\text{CO})$ and at 638 cm^{-1} due to $\nu(\text{RuCS})$ which are characteristic of $\eta^2\text{-COS}$ through the $\text{C}=\text{S}$ bond. The presence of a singlet in the ^{31}P NMR spectrum of (VII) at 36.32(s) and two carbonyl stretching frequencies at 2012 and 1951 cm^{-1} due to terminal carbonyl ligands support the structure (VII) for the zerovalent ruthenium COS complex [87].

COS reacts with $[\text{Ru}(\text{CO})\text{Cl}(\text{CH}_3\text{COO})(\text{PPh}_3)_2]$ to give a bridged COS complex (VIII) whose IR spectrum shows a broad band at 850 cm^{-1} due to the COS ligand [88]. Ruthenocene reacts with COS to afford the COS complex $[(\eta^5\text{-C}_5\text{H}_5)\text{Ru}(\eta^2\text{-COS})]$ [89].

Reaction of COS with $[(\eta^2\text{-C}_5\text{H}_5)\text{Rh}(\text{PPh}_3)_2]$ gives the COS complexes (IX) and (X). The IR spectrum of (IX) shows absorption bands at 1720 cm^{-1} and 630 cm^{-1}




(VIII)



due to the coordinated COS group. In the ^1H NMR spectrum, the cyclopentadienyl resonance is a doublet of doublets ($\delta = 6.00$ ppm) with $J_{\text{P-H}} = 1.60$ Hz and $J_{\text{Rh-H}} = 0.66$ Hz. The electronic spectrum exhibits λ_{max} (ϵ_{max}) at 806 (77), 417 (3195) 283 nm ($14080 \text{ M}^{-1} \text{ cm}^{-1}$). The IR spectrum of (X) exhibits bands at 1750 cm^{-1} and 635 cm^{-1} due to the coordinated COS group. The proton NMR spectrum shows a multiplet, with the chemical shift of the most intense line at $\delta 7.51$ ppm due to the phenyl protons [90].

Reaction of COS with $\text{Rh}(\text{NO})\text{Cl}_3$ in ethanol affords the red complex (XI) which shows IR spectral bands at 1615 cm^{-1} due to $\nu(\text{NO})$ and at 1700 cm^{-1} due to $\nu(\text{CO})$ of the coordinated COS.


Reactions of COS with iridium complexes $[\text{Ir}(\text{CO})\text{Cl}(\text{PR}_3)_2]$ ($\text{PR}_3 \equiv \text{PMe}_2\text{Ph}$,




PMe_3 , PPh_3) have been investigated [91]. The COS complexes (XII) and (XIII) form at low temperatures, below -78°C . Attempts to prepare $[\text{Ir}(\text{CO})\text{Cl}(\text{COS})(\text{PPh}_3)_2]$ were unsuccessful. The complexes (XII) and (XIII), which decompose on warming to give COS, have IR spectra with absorption bands at 2036 cm^{-1} due to $\nu(\text{CO})$, at 1748 , 662 cm^{-1} due to COS ligand for the complex (XII), and at 2017 cm^{-1} due to $\nu(\text{CO})$, at 1730 , 660 cm^{-1} due to COS ligand and at 241 cm^{-1} due to $\nu(\text{Ir}-\text{Cl})$ for the complex (XIII). ^{31}P NMR spectra exhibit a singlet at $\delta -27.8(\text{s})$ for (XII) and at $\delta -32.1(\text{s})$ for (XIII). These spectral data are consistent with structures (XII) and (XIII).

A very unstable complex $[\text{Pd}(\text{COS})(\text{PPh}_3)_2]$ can be prepared from the reaction of COS and $\text{Pd}(\text{PPh}_3)_3$ or $\text{Pd}(\text{PPh}_3)_4$. The reaction of $\text{Pd}[\text{P}(p\text{-C}_6\text{H}_4\text{CH}_3)_3]_3$ with COS affords a relatively more stable complex (XIV) which exhibits IR spectral bands at 1739 and 630 cm^{-1} attributable to the bidentate COS ligand. ^{31}P NMR spectrum of (XIV) consists of an AB spin pattern ($\delta 22.2, 27.5$) consistent with (XIV) [92].

Reaction of COS with $\text{Cu}(\text{TPP})$ in toluene gives a pinkish-brown complex



(XII), $\text{PR}_3 = \text{PMe}_2\text{Ph}$
(XIII), $\text{PR}_3 = \text{PMe}_3$




[Cu(TPP)(COS)] whose IR spectrum shows a band at 1712 cm^{-1} due to the coordinated COS ligand [93].









2.3. Reactions of carbonyl sulfide complexes


2.3.1. Carbonylation reaction

The coordination chemistry of COS is dominated by reactions that involve cleavage of the C—S bond and form carbonyl complexes. A plausible explanation for the carbonylation reactions is the initial formation of η^2 -COS complex followed by sulfur elimination:









The carbonylation reactions, in which sulfur is eliminated by the phosphine ligand, are shown in the following equations.







The carbonylation reactions, in which the fate of the sulfur has not been determined, are presented in the following equations.





Some carbonylation reactions have also been reported in which sulfur remains in the complex either as a bridged ligand or associated with other ligands.



2.3.2. Reductive disproportionation reaction: dithiocarbonate complexes

Carbonyl sulfide affords a convenient route to metal dithiocarbonate complexes. In the formation of dithiocarbonate from COS, the COS undergoes a metal-promoted

two-electron reductive disproportionation reaction. Dithiocarbonates could arise from an electrophilic attack on the endocyclic sulfur atom by another COS molecule, followed by elimination of CO [92]:



Metal complexes with dithiocarbonate ligands show absorption bands in their IR spectra at 1680–1965, 1595–1615 and 830–865 cm^{-1} which are characteristics of a bidentate dithiocarbonate ligand [99,100]. Spectra data of the dithiocarbonate complexes are presented in Table 3 [101–104].

Reaction of COS with $[\text{Fe}(\text{CO})_2(\text{PPh}_3)_3]$ in a liquid-nitrogen-cooled pressure reactor and with $[\text{Ru}(\text{CO})_2(\text{PPh}_3)_3]$ in toluene solution result in the formation of dithiocarbonate complexes (XV) and (XVI). The IR and ^{31}P NMR spectra are consistent with the proposed structure with the presence of *cis*-carbonyl and *trans*-phosphine ligands.


The complexes $[\text{Pd}(\text{S}_2\text{CO})(\text{PR}_3)_2]$ ($\text{PR}_3 \equiv \text{PMePh}_2$, PMe_2Ph , PiPr_3) are prepared from the reaction of excess COS with $[\text{Pd}(\text{PR}_3)_4]$. The complex $[\text{Pd}(\text{S}_2\text{CO})(\text{PMe}_3)_2]$ is obtained by the displacement of phosphine ligand from $[\text{Pd}(\text{S}_2\text{CO})(\text{PR}_3)_2]$ with PMe_3 . The X-ray crystal structure of $[\text{Pd}(\text{S}_2\text{CO})(\text{PMe}_2\text{Ph})_2]$ shows approximate square planar coordination of the palladium with planar dithiocarbonate ligand (XVII) [101].

Platinum carbonyl sulfide complex $[\text{Pt}(\eta^2\text{-COS})(\text{PPh}_3)_2]$ reacts with PMe_3 or PMe_2Ph to afford dithiocarbonato-platinum(II) complexes $[\text{Pt}(\text{S}_2\text{CO})(\text{PR}_3)_2]$. Complex $[\text{Pt}(\text{S}_2\text{CO})(\text{PPh}_3)_2]$ is obtained by the reaction of excess COS with $[\text{Pt}(\text{PPh}_3)_4]$, $[\text{Pt}(\eta^2\text{-COS})(\text{PPh}_3)_2]$ or $[\text{Pt}_2\text{S}(\text{CO})(\text{PPh}_3)_3]$ [92].


Reactions of COS with *trans*- $[\text{M}(\text{C}_2\text{H}_4)_2(\text{PMe}_3)_4]$ ($\text{M} \equiv \text{Mo}$ or W) in diethylether give dithiocarbonate complexes $[\text{M}(\text{CO})_2(\text{S}_2\text{CO})(\text{PMe}_3)_3]$ ($\text{M} \equiv \text{Mo}$ (XVIII); W)

Table 3
Spectral data for dithiocarbonate complexes


Complex	IR frequencies (cm^{-1})		^{31}P NMR, δ (ppm)	Reference
	$\nu(\text{CO})$	$\nu(\text{CS})$		
$[\text{Mo}(\text{CO})_2(\text{S}_2\text{CO})(\text{PMe}_3)_3]$	1690, 1570	—	16.9 –18.0	[104]
$[\text{Fe}(\text{CO})_2(\text{S}_2\text{CO})(\text{PPh}_3)_2]$	1688, 1611	838	51.1(s)	[87]
$[\text{Ru}(\text{CO})_2(\text{S}_2\text{CO})(\text{PPh}_3)_2]$	1688, 1609	832	30.83(s)	[87]
$[\text{Pd}(\text{S}_2\text{CO})(\text{PMePh}_2)_2]$	1680, 1605	835	8.86(s)	[101]
$[\text{Pd}(\text{S}_2\text{CO})(\text{PMe}_2\text{Ph})_2]$	1680, 1610	835	–7.23(s)	[101]
$[\text{Pd}(\text{S}_2\text{CO})(\text{Pi-Pr}_3)_2]$	1695, 1605	855		[101]
$[\text{Pd}(\text{S}_2\text{CO})(\text{PMe}_3)_2]$	1690, 1595	860	–16.71(s)	[101]
$[\text{Pd}(\text{S}_2\text{CO})\{\text{P}(p\text{-C}_6\text{H}_4\text{Me})_3\}_2]$	1680, 1610	832	27.6(s)	[92]
$[\text{Pt}(\text{S}_2\text{CO})(\text{PMe}_2\text{Ph})_2]$	1680, 1615	835–850	–30.92(s)	[101]
$[\text{Pt}(\text{S}_2\text{CO})(\text{PMe}_3)_2]$	1695, 1605	855–865	–20.42(s)	[101]
$[\text{Pt}(\text{S}_2\text{CO})(\text{PPh}_3)_2]$	1688, 1615	835	16.6(s)	[92]



(XV), $\text{M} = \text{Fe}$; (XVI), $\text{M} = \text{Ru}$



(XVII)




(XVIII)

[102–105]. The complex (**XVIII**) has also been prepared by the reaction of COS with bis(nitrogen) complex *cis*-[Mo(N₂)₂(PMe₃)₄] or with bis(carbon dioxide) complex. The structure of (**XVIII**) has been determined by X-ray diffraction analysis and shows a pentagonal bipyramid coordination about the molybdenum with the axial positions occupied by one carbonyl and one PMe₃ ligand. Important bond distances and bond angles of dithiocarbonate complexes are given in Table 4.

2.3.3. Insertion reaction of carbonyl sulfide into M–X bonds

The insertion reactions of COS into metal–hydride, metal–carbon, metal–nitrogen and metal–oxygen bonds have been discussed by Pandey and Nigam [26]. In this article, recent developments in the insertion of COS into transition metal–hydride bonds is discussed.

Carbonyl sulfide insertion into a metal–hydride bond gives rise to monothioformate complexes:



The IR spectra of monothioformate complexes prove to be valuable in distinguishing between monodentate and bidentate bonding. The IR spectra of complexes containing a bidentate thioformate ligand show additional bands in the region 1340–1500 cm^{−1} which are assigned to ν(CO···M) (Table 5). The ¹H NMR chemical shifts for monodentate thioformate ligand occur at a lower field than the bidentate thioformate ligand. The ¹³C NMR spectra of monothioformate complexes exhibit a signal in the region δ 180–202 ppm.


Darensbourg and Rockicki reported the formation of monothioformate complexes [A][Cr(CO)₅(OSCH)] (A ≡ [Ph₂P=N=PPh₂] or Et₄N) by the exchange of COS for carbon dioxide in [A][Cr(CO)₅(O₂CH)] or by the direct reaction of COS with [Cr(CO)₅H][−] [106,107]. The monothioformate ligand is coordinated to the chromium in the monodentate manner (**XIX**). Bianchini and co-workers have reported the insertion of COS into Cu–H bonds in the reactions of COS with copper(I) tetrahydroborates [108–110]. COS reacts with the η²-BH₄ complexes

Table 4
Important bond distances (Å) and bond angles (deg) of dithiocarbonate complexes

Complex	M–S (Å)	S–C (Å)	C–O (Å)	S–M–S (deg)	S–C–S (deg)	Reference
[Mo(CO) ₂ (S ₂ CO)(PMe ₃) ₃]	2.577(7)	1.83(3)	1.236(12)	69.0(2)	111.0(5)	[104]
	2.533(7)	1.68(3)				
[Pd(S ₂ CO)(PMe ₂ Ph) ₂]	2.318(3)	1.736(9)	1.23(2)	75.4(1)	109.5(5)	[101]


Table 5
Spectroscopic data for monolithioformate complexes

Complex	IR frequencies (cm ⁻¹)				¹ H NMR, δ (ppm)	¹³ C NMR, δ (ppm)	Reference
	ν(CO)	ν(CO...M)	ν(CS)	ν(CH)			
[Cr(CO) ₅ (η ¹ -OCH)] ⁻						198.8	[106]
[(PPh ₃) ₂ Cu(η ² -SOCH)]	1632	1340	800	2800	10.12		[110]
	1580			2770			
[(PCy ₃) ₂ Cu(η ² -SOCH)]	1630	1350	805	2800	10.76		[110]
	1580			2760			
[(triphos)Cu(η ¹ -SOCH)]	1625		805	2755	12.44	201.3	[110]
	1570			2660			
[(np ₃)Cu(η ¹ -SOCH)]	1635			2760			[141]
	1570						
[(η ⁵ -C ₅ H ₅)Ru(η ¹ -SOCH)- (PPh ₃) ₂]	1670					181.8	[111]
	1630						[112]
[(η ⁵ -C ₅ H ₅)Ru(η ¹ -SOCH)- (AsPh ₃) ₂]	1660					181.64	[111]
[(η ⁵ -C ₅ H ₅)Ru(η ¹ -SOCH)- (SbPh ₃) ₂]	1658					179.9	[111]




(XIX)

[Cu(η²-BH₄)(PR₃)₂] (R ≡ Ph, Cy) to afford the η²-S,O-bonded monothioformates (**XX**, **XXI**) whereas the reaction of COS with the η¹-BH₄ complexes [(triphos)Cu(η²-BH₄)] and [(np₃)Cu(η²-BH₄)] results in the formation of the η¹-S-bonded monothioformate (**XXII**) and (**XXIII**).



(XX), R = Ph; (XXI), R = Cy



The IR spectra of (XX) and (XXI) exhibit absorption bands due to a monothioformate ligand bonding through sulfur in a bidentate manner. The IR spectra of (XXII) and (XXIII) are consistent with a monodentate thioformate ligand.

Carbonyl sulfide reacts with ruthenium hydride complexes $[(\eta^5\text{-C}_5\text{H}_5)\text{RuHL}_2]$ ($\text{L} \equiv \text{PPh}_3, \text{AsPh}_3, \text{SbPh}_3$) in methanol to give microcrystalline complexes (XXIV) [111,112].

2.3.4. Formation of sulfide complexes

The reactions of COS with metal complexes also results in the formation of sulfide complexes [26]. The trivalent uranium metallocene $[(\text{C}_5\text{H}_4\text{Me})_3\text{U}] \cdot \text{THF}$ reacts with COS to form $[(\text{C}_5\text{H}_4\text{Me})_3\text{U}]_2(\mu\text{-S})$ [113].



3. Coordination chemistry of carbon disulfide

3.1. Transition metal–carbon disulfide bonding and energetics

The coordination of CS_2 ligand to a transition metal (M) gives a wide variety of complexes which, apart from $\eta^1\text{-S}$ end-on coordination, all involve bonding to carbon.

3.1.1. η^1 -end-on coordination

In this type of bonding, CS_2 is bound to the transition metal through a sulfur atom with no change in the oxidation state of the metal. The CS_2 ligand behaves as a Lewis base, donating a lone pair of electrons from the $1\pi_g$ (HOMO) orbital to the corresponding metal vacant orbital [29,114–118].











3.1.2. η^2 -side-on coordination

The main interaction in η^2 -side-on coordination (XXV) is the π -back donation from the metal atom to the CS_2 ligand. Theoretical studies have led to a better understanding of the factors governing the bonding in CS_2 complexes [80,119–122]. The importance of the nature of the frontier molecular orbital has been emphasized. The best situation for the occurrence of the η^2 -side-on coordination mode is the presence of a metal- $d\pi$ orbital as HOMO and an empty $d\sigma$ orbital, pointing to the CS_2 ligand in the transition metal fragments. In the $\text{M}-(\eta^2\text{-CS}_2)$ complex, the HOMO is a combination of metal $d\pi$ and CS_2 $1b_2$ and $2a_1$ orbitals. The theoretical calculations show that the carbon atom exerts a trans-influence stronger than that exerted by the sulfur atom [80]. Mulliken population analysis shows negative charges on the uncoordinated sulfur atom which confirms its nucleophilic character.

3.1.3. Bridging carbon disulfide complexes

Carbon disulfide can also function as a bridge between two metal atoms and three metal atoms.





3.2. Preparation and structure

3.2.1. Mononuclear metal–carbon disulfide complexes

Transition metal CS_2 complexes are prepared by the reaction of CS_2 with transition metal complexes. The CS_2 ligand in $\text{M}-\eta^2\text{-CS}_2$ and $\text{M}-\eta^1\text{-CS}_2$ coordination modes acts as a better π -acceptor and a poor σ -donor. As a result, the net electron density shifts from metal ($d\pi$ orbital) to CS_2 (π^* orbital). This is expected to lead to a reduction in the C–S bond order of the CS_2 compared with the free CS_2 . In fact, the (CS) stretching frequency of a variety of π -bonded CS_2 complexes is appreciably lower than that of free CS_2 (Table 6).


The titanium– CS_2 complex $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\eta^2\text{-CS}_2)(\text{PMe}_3)]$ (XXXII) is prepared by the reaction of $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\text{PMe}_3)_2]$ with CS_2 in pentane at 0°C [123]. The reaction of $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\eta^3\text{-allyl})]$ with CS_2 yields $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\eta^2\text{-CS}_2)]$ (XXXIII) [124]. Complex $[(\eta^6\text{-C}_6\text{Et}_6)\text{Cr}(\text{CO})_2(\text{C}_4\text{H}_8)]$, obtained by UV irradiation of $[(\eta^6\text{-C}_6\text{Et}_6)\text{Cr}(\text{CO})_3]$ in the presence of cyclooctene, reacts with CS_2 to produce $[(\eta^6\text{-C}_6\text{Et}_6)\text{Cr}(\text{CO})_2(\eta^2\text{-CS}_2)]$ [125]. The complexes $[\text{A}]_2[\text{W}(\text{CO})_5(\eta^1\text{-CS}_2)]$ ($\text{A} \equiv \text{Li}$ or Na) may be prepared in solution by the reaction of CS_2 with $[\text{W}(\text{CO})_5]^-$ in THF at -78°C [126].

$\text{Na}[(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})_2]$ reacts with CS_2 at low temperature to give the $\eta^1\text{-CS}_2$ complex (XXXIV) [127]. Reaction of CS_2 with $[\text{M}(\text{CO})_2(\text{PPh}_3)_3]$ ($\text{M} \equiv \text{Fe}, \text{Ru}$) gives the $\eta^2\text{-CS}_2$ complexes (XXXV) [87].

Reaction of ruthenocene with CS_2 gives $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ru}(\eta^2\text{-CS}_2)]$ [89]. UV irradiation of a solution of $[(\eta^5\text{-C}_5\text{H}_5)\text{RuCl}(\text{PPh}_3)_2]$ in $\text{CH}_2\text{Cl}_2\text{-CS}_2$ yields $[(\eta^5\text{-C}_5\text{H}_5)\text{RuCl}(\eta^2\text{-CS}_2)(\text{PPh}_3)_2]$. Complex $[(\eta^5\text{-C}_5\text{H}_5)\text{RuCl}(\text{P}(\text{CH}_2\text{CH}_2\text{CN})_3)_2]$, owing to the lower basicity of the phosphine, does not react with CS_2 . Reaction of $[(\eta^5\text{-C}_5\text{H}_5)\text{Ru}(\text{CH}_3\text{CN})(\text{PPh}_3)_2]^+$, obtained by refluxing an acetonitrile solution of $[(\eta^5\text{-C}_5\text{H}_5)\text{RuCl}(\text{PPh}_3)_2]$ in the presence of HgCl_2 , with CS_2 results in the formation of $[(\eta^5\text{-C}_5\text{H}_5)\text{Ru}(\text{CS}_2)(\text{PPh}_3)_2](\text{HgCl}_3)$. Complex $[(\eta^5\text{-C}_5\text{H}_4\text{Me})\text{Ru}(\text{CS}_2)(\text{PPh}_3)_2](\text{BF}_4)$ is obtained by the reaction of $[(\eta^5\text{-C}_5\text{H}_5)\text{RuCl}(\text{PPh}_3)_2]$ and AgBF_4 in the presence of CS_2 [128]. CS_2 reacts with osmium nitrosyl complex $[\text{Os}(\text{NO})\text{Cl}(\text{PPh}_3)_3]$ to give (XXXVI) [129].

Table 6
IR frequencies of mononuclear carbon disulfide complexes







Complex	$\nu(\text{CS})$ (cm^{-1})	$\nu(\text{MCS})$ (cm^{-1})	Reference
$[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\eta^2\text{-CS}_2)(\text{PMe}_3)]$	1107		[123]
$[\text{Fe}(\text{CO})_2(\eta^2\text{-CS}_2)(\text{PPh}_3)_2]$	1142, 1155		[87]
$[\text{Ru}(\text{CO})_2(\eta^2\text{-CS}_2)(\text{PPh}_3)_2]$	1126	650	[87]
$[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ru}(\eta^2\text{-CS}_2)]$	1100		[89]
$[(\eta^5\text{-C}_5\text{H}_5)\text{RuCl}(\eta^2\text{-CS}_2)(\text{PPh}_3)_2]$	1173, 1000		[128]
$[(\eta^5\text{-C}_5\text{H}_5)\text{Ru}(\eta^2\text{-CS}_2)(\text{PPh}_3)_2][\text{HgCl}_3]$	1280	860	[128]
$[(\eta^5\text{-C}_5\text{H}_4\text{Me})\text{Ru}(\eta^2\text{-CS}_2)(\text{PPh}_3)_2][\text{BF}_4]$	1280	860	[128]
$[\text{Os}(\text{NO})\text{Cl}(\eta^2\text{-CS}_2)(\text{PPh}_3)_2]$	991		[129]
$[(\eta^5\text{-C}_5\text{H}_5)\text{Co}(\eta^2\text{-CS}_2)\{\text{P}(\text{OMe})_3\}]$	1165		[131]
$[(\text{triphos})\text{Ni}(\eta^2\text{-CS}_2)]$	1145	630	[132]
$[(\eta^5\text{-C}_5\text{H}_5)\text{Rh}(\eta^2\text{-CS}_2)(\text{PMe}_3)]$	1159		[133]
$[(\text{triphos})\text{Rh}(\eta^2\text{-CS}_2)]$	1160	640	[137]
$[(\text{triphos})\text{Rh}(\text{N}_3)(\eta^2\text{-CS}_2)]$	1150	630	[137]
$[(\text{triphos})\text{Ir}(\eta^2\text{-CS}_2)]$	1140	650	[137]
$[\text{Rh}(\text{NO})\text{Cl}_3(\eta^2\text{-CS}_2)]$	1140, 1000		[90]
$[\text{Pd}(\eta^2\text{-CS}_2)(\text{PMePh}_2)_2]$	1180		[140]
$[\text{Pd}(\eta^2\text{-CS}_2)(\text{dppe})]$	1173		[140]
$[\text{Pd}(\eta^2\text{-CS}_2)(\text{dpmb})]$	1173		[140]
$[\text{Pt}(\eta^2\text{-CS}_2)(\text{dppe})]$	1158		[140,141]
	1141	650	[142]
$[\text{Pt}(\eta^2\text{-CS}_2)(\text{dpmb})]$	1150		[140]
$[\text{Pt}(\eta^2\text{-CS}_2)(\text{P-}i\text{-Pr}_3)_2]$	1185		[140]



(XXXIII)

Complexes $[(\eta^5\text{-C}_5\text{H}_5)\text{Co}(\eta^2\text{-CS}_2)\text{L}]$ (XXXVII) are obtained, in low yield along with other products, by the reactions of CS_2 with $[(\eta^5\text{-C}_5\text{H}_5)\text{Co}(\text{CO})\text{L}]$ ($\text{L} \equiv \text{PEt}_3$, $\text{P-}n\text{-Pr}_3$, $\text{P-}n\text{-Bu}_3$, PMe_2Ph , PMePh_2 , PPh_3) [130]. $[(\text{np}_3)\text{CoH}]$ reacts with CS_2 to yield $[(\text{np}_3)\text{Co}(\eta^2\text{-CS}_2)]$ (XXXVIII) [13]. Kolb and Werner reported the formation of $[(\eta^5\text{-C}_5\text{H}_5)\text{Co}(\eta^2\text{-CS}_2)\{\text{P}(\text{OMe})_3\}]$ from the reaction of CS_2 with $[(\eta^5\text{-C}_5\text{H}_5)\text{Co}\{\text{P}(\text{OMe})_3\}_2]$ [132].

Rhodium CS_2 complexes $[(\eta^5\text{-C}_5\text{R}_5)\text{Rh}(\eta^2\text{-CS}_2)(\text{PMe})]$ ($\text{R} \equiv \text{H}$, Me) are formed when $[(\eta^5\text{-C}_5\text{R}_5)\text{Rh}(\text{PMe}_3)_2]$ reacts with CS_2 [133]. Bianchini and co-workers



have synthesized CS_2 complexes of rhodium and iridium containing tridentate phosphine ligands [134–137]. Reaction of CS_2 with $[(\text{np}_3)\text{RhH}]$ in the presence of NaBPh_4 yields $[(\text{np}_3)\text{Rh}(\eta^2\text{-CS}_2)](\text{BPh}_4)$. The molecular structure of $[(\text{np}_3)\text{Rh}(\eta^2\text{-CS}_2)](\text{BPh}_4)$ has been determined by X-ray diffraction analysis. $[(\text{triphos})\text{Rh}(\eta^2\text{-CS}_2) \cdot \text{C}_6\text{H}_6]$ is formed by the reaction of CS_2 with $[\text{RhCl}(\text{C}_2\text{H}_4)]_2$ in the presence of triphos. The reaction of $[\text{RhN}_3(\text{COD})]_2$ with CS_2 in the presence of triphos results in the formation of $[(\text{triphos})\text{RhN}_3(\eta^2\text{-CS}_2)]$. Complex



$[(\text{triphos})\text{IrCl}(\eta^2\text{-CS}_2)]$ can be obtained by the reaction of $[\text{IrCl}(\text{COD})]_2$ with CS_2 in the presence of triphos. Carbon disulfide reacts with $[\text{Rh}(\text{NO})\text{Cl}_3]$ in ethanol to produce $[\text{Rh}(\text{NO})\text{Cl}_2(\eta^2\text{-CS}_2)]$ [90]. Iridium complexes $[\text{Ir}(\text{COD})(\eta^2\text{-CS}_2)(\text{PR}_3)_2]$ ($\text{R} \equiv \text{Ph}, \text{Cy}$) can be obtained by the reaction of $[\text{Ir}(\text{COD})(\text{PR}_3)_2]$ with CS_2 in dichloromethane [138].



The nickel complex $[(\text{triphos})\text{Ni}(\text{S}_2\text{CSMe})][\text{BPh}_4]$ reacts with NaBH_4 to afford the CS_2 complex $[(\text{triphos})\text{Ni}(\eta^2\text{-CS}_2)]$ [96]. Hoffman and co-workers have reported a trimorpholinophosphine nickel(0) complex of CS_2 [139]. Palladium CS_2 complexes $[\text{Pd}(\eta^2\text{-CS}_2)(\text{PR}_3)_2]$ ($\text{PR}_3 \equiv \text{PMePh}_2, \text{P-i-Pr}_3$) are prepared by the reaction of CS_2 with $[\text{Pd}(\text{PMePh}_2)_4]$ and $[\text{Pd}(\text{P-i-Pr}_3)_2]$ complexes. Complexes $[\text{M}(\eta^2\text{-CS}_2)(\text{PR}_3)_2]$ ($\text{M} \equiv \text{Pd}, \text{Pt}$) react with bidentate phosphine ligands (dppe, dpmb) to give the chelate complexes $[\text{M}(\eta^2\text{-CS}_2)(\text{dppe})]$ (**XXXIX**) and $[\text{M}(\eta^2\text{-CS}_2)(\text{dpmb})]$ (**XL**) [94,140,141]. Walker and co-workers have also reported the formation of $[\text{Pt}(\eta^2\text{-CS}_2)(\text{dppe})]$ from the reaction of $[\text{M}(\eta^2\text{-CS}_2)(\text{PPh}_3)_2]$ [142,143]. Gregor has reported gas phase reactions of CS_2 with electron deficient carbonyl complexes $[\text{Cr}(\text{CO})_3]^-$, $[\text{Cr}(\text{CO})_4]^-$, $[\text{Mn}(\text{CO})_3]^-$, $[\text{Mn}(\text{CO})_4]^-$, $[\text{Fe}(\text{CO})_2]^-$, $[\text{Fe}(\text{CO})_3]^-$, $[\text{Ni}(\text{CO})_2]^-$ and $[\text{Ni}(\text{CO})_3]^-$ affording CS_2 complexes [144].

3.2.2. Bridging carbon disulfide complexes

A number of transition metal complexes containing bridged CS_2 has been reported. The IR spectra of complexes with bridged CS_2 ligand show a (CS) absorption in the region $900\text{--}1050\text{ cm}^{-1}$, substantially lower than the values ($1100\text{--}1175\text{ cm}^{-1}$) normally found for $\text{M-}\eta^2\text{-CS}_2$ complexes.

The complex anion $[(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})_2(\eta^1\text{-CS}_2)]^-$ which is unstable at room temperature and has not been isolated as solid, reacts with a variety of Lewis acids. A number of bimetallic and trimetallic CS_2 complexes, containing one or two other $(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})_2$, $\text{Cr}(\text{CO})_5$, $\text{Mn}(\text{CO})_5$, $\text{Re}(\text{CO})_5$ and $\text{W}(\text{CO})_5$ functionalities on the sulfur centers, has been obtained from $[(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})_2(\eta^1\text{-CS}_2)]^-$ [145–150]. $[(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})_2(\eta^1\text{-CS}_2)]^-$ reacts with $[(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})_2\text{I}]$ to afford (**XLI**). This bridged CS_2 complex $[(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{FeC}(\text{S})\text{SFe}(\text{CO})_2(\eta^5\text{-C}_5\text{H}_5)]$ (**XLI**) and the methylated complex $[(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})_2(\eta^1\text{-CS}(\text{SMe}))]$ are useful precursors for the synthesis of bimetallic and trimetallic CS_2 bridged complexes. Bimetallic complexes $[(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{FeC}(\text{SMe})\text{SM}'\text{Ln}]$ (**XLII**) ($\text{M}'\text{Ln} \equiv (\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})_2$, $\text{Cr}(\text{CO})_5$, $\text{Mn}(\text{CO})_5$, $\text{Re}(\text{CO})_5$ and $\text{W}(\text{CO})_5$) have been prepared by the reaction of











$[(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})_2(\eta^1\text{-CS}(\text{SMe}))]$ with electrophiles. The synthesis and characterization of complexes of the types **(XLIII)** and **(XLIV)** ($\text{M}'\text{Ln} \equiv (\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})_2$, $\text{W}(\text{CO})_5$) have also been reported.

The complex $[(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})_2(\eta^1\text{-CS}(\text{SMn}(\text{CO})_5))]$, spontaneously transforms on heating into **(XLV)**. Reaction of $[(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})_2(\eta^1\text{-CS}_2)]^-$ with bis-acetonitrile complex $[(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})(\text{CH}_3\text{CN})_2](\text{PF}_6)$ in THF produces **(XLVI)** ($\text{X} \equiv \text{O}$). The thiocarbonyl analog **(XLVI)** ($\text{X} \equiv \text{S}$) can be prepared from the reaction of $[(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})_2(\eta^1\text{-CS}_2)]^-$ with $[(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})(\text{CS})\text{I}]$ [150]. Busetto and co-workers [148] obtained **(XLVI)** ($\text{X} \equiv \text{O}$) by refluxing **(XLVII)** in dichloromethane. Photolysis of **(XLVII)** gives **(XLVI)** in 10%–17% yield [148,150].

Reaction of $[\text{Ni}(\text{COD})_2]$ with CS_2 in the presence of PPh_3 yields complex **(XLVIII)** ($\text{M} \equiv \text{Ni}$, $\text{R} \equiv \text{Ph}$) [151]. Bimetallic complexes of platinum and palladium **(XLVIII)** containing monodentate phosphine ligands have been synthesized by Farrar and co-workers [152,153].








The molybdenum complex $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Mo}_2(\text{CO})_5(\mu_2, \eta^3\text{-CS}_2)]$ contains a bridging CS_2 ligand bonded to one Mo atom through the carbon and one of the sulfur atoms and to the other Mo atom through the second sulfur atom [154].

Reaction of $[\text{Pt}(\eta^2\text{-CS}_2)(\text{PPh}_3)_2]$ with $[\text{Pt}(\eta^2\text{-C}_2\text{H}_4)(\text{PPh}_3)_2]$ results in the formation of (XLIX) [141,143].

The synthesis and characterization of bridged CS_2 complexes of platinum and palladium containing terminal or bridged diphosphine ligands have been a subject of interest. The complexes $[\text{M}(\eta^2\text{-CS}_2)(\text{dppe})]$ ($\text{M} \equiv \text{Pd, Pt}$) slowly dimerize in solution to produce bridging CS_2 complexes $[(\text{dppe})\text{M}(\mu\text{-CS}_2)\text{M}(\text{dppe})]$ (L) [141].

Reaction of $[\text{Pd}_2\text{Cl}_2(\text{dmpm})_2]$ with CS_2 results in the formation of a deep red-orange complex (LI) which may also be obtained by the reaction of $[\text{Pd}_2(\mu\text{-CO})\text{Cl}_2(\text{dmpm})_2]$ with CS_2 [155]. Complexes (LII) ($\text{R} \equiv \text{Ph, } p\text{-tolyl}$) are obtained by the reaction of $[\text{Pt}(\text{C}\equiv\text{CR})_2(\text{dppm})_2]$ with CS_2 in benzene [156].





Reaction of [Pt₂Cl₂(dppm)₂] with CS₂ leads to the formation of complex (LIII) [157,158].


3.3. Reactions of carbon disulfide complexes


3.3.1. Thiocarbonylation reaction

One of the most important routes for the synthesis of thiocarbonyl complexes is by the reaction of CS₂ with transition metal complexes. There are various paths for preparation of the thiocarbonyl ligand via the CS₂ ligand.

(a) Coordination of CS₂ to a metal center is followed by methylation and the methylated moiety undergoes hydride attack producing a thiocarbonyl complex.

(b) Coordination of CS₂ to a metal center is followed by sulfur elimination. There are three routes for the abstraction of sulfur from CS₂ by PR₃ producing S=PR₃: (i) an isolated η²-CS₂ complex is attacked by added phosphine ligand, (ii) coordinated phosphine ligand is displaced from a metal center and used as the abstracting molecule for the η²-CS₂ ligand, and (iii) a CS₂–PR₃ mixture is used.






(c) Reductive cleavage of CS₂ ligand produces sulfido and thiocarbonyl ligands. Some examples of thiocarbonylation reaction are given in Table 7 [159–169].

3.3.2. Reductive dimerization of coordinated carbon disulfide

3.3.2.1. Metal promoted head-to-head dimerization of carbon disulfide. Transition metal promoted reductive dimerization of two coordinated CS₂ molecules in the head-to-head coupling form the tetrathiooxalate ligand [135,165,170–174]. Maj and co-workers reported the first structurally characterized example of coordinated C₂S₄ (**LIV**) from the reaction of CS₂ with nickel pentamethylcyclopentadienyldicarbonyl [(η⁵-C₅Me₅)Ni(CO)₂]₂ [171]. Reaction of CS₂ with the iron cluster carbonyl [Fe₂(CO)₉] results in the formation of a tetranuclear complex [(CO)₆Fe₂(C₂S₄)Fe₂(CO)₆] in low yield, along with other products [165]. In this complex the central μ-C₂S₄ unit links, end-on-fashion, two Fe₂ fragments (**LV**). Synthesis and structural studies of [{(η⁵-C₅Me₅)Fe(CO)₂}(μ-C₂S₄)] and [{(η⁵-C₅Me₅)Co}₂(μ-C₂S₄)] have been reported [172]. Carbon–carbon bond formation by reductive head-to-head coupling of CS₂ is also activated by a titanium(II) complex. Reaction of [(η⁵-C₅H₅)Ti(CO)₂] with neat CS₂ under an inert atmosphere gives [{(η⁵-C₅H₅)Ti}₂(μ-C₂S₄)] (**LVI**) [173]. Complex (**LVI**), unlike the

Table 7
Thiocarbonylation reactions of carbon disulfide


Precursor complex	Isolated product	Reference
$[(np_3)CoH] + MeI + NaBH_4$	$[(np_3)Co(CS)]^+$	[131,159]
$[(np_3)CoH] + NaBPh_4$	$[(np_3)Co(CS)]^+$	[131]
$[(\eta^5-C_5H_5)Co(CO)_2] + L$ ($L \equiv PEt_3$, $P-n-Pr_3$, $P-n-Bu_3$, PMe_2Ph , PPh_3 , $t-BuNC$)	$[(\eta^5-C_5H_5)Co(CS)L] + \dots$	[130]
$[Fe(CO)_2(PPh_3)_3] + MeI + NH_4PF_6 + P-n-Bu_6$	$[Fe(CO)_2(CS)(PPh_3)_2]$	[160]
$[Fe(CO)_2((\eta^2-CS_2Me)L_2)]^+ + Na/Hg$ ($L \equiv PMe_3$, PMe_2Ph , $P-n-Bu_3$)	$[Fe(CO)_2(CS)L_2]$	[161]
$[(\eta^6-C_6Et_6)Cr(CO)_2(C_8H_{14})] + PPh_3$	$[(\eta^6-C_6Et_6)Cr(CO)_2(CS)]$	[125]
$[W(CO)_3(\eta^2-CS_2)(L)] + P-n-Bu_3$ ($L \equiv dmpe$, $dppe$)	$[W(CO)_3(CS)(L)]$	[162]
$[RuCl_2(PPh_3)_3] + PPh_3$	$[Ru(CS)Cl_2(H_2O)(PPh_3)_2]$	[163]
$[RuCl_2(PPh_3)_3]$	$[Ru(CS)Cl_2(PPh_3)_2(\mu-Cl)_2]$	[164]
$[Fe_3(CO)_{12}]$	$[Fe_3(CO)_8(CS)(S_2)] + \dots$	[165]
$[PtL_3]$ ($L \equiv t-BuN=P-N(SiMe_3)(t-Bu)$)	$[PtL(CS)]_2(\mu-S)_2$	[166]
$[M((\eta^2-CS_2)(dppe))] + [Pt(PPh_3)_4]$	$[(dppe)M(\mu-S)Pt(CS)(PPh_3)]$	[143]
$[Ru(CO)(TPP)(EtOH)]$	$[Ru(CO)(CS)(TPP)]$	[167]
$[Re_2X_4(\mu-dppm)_2]$ ($X \equiv Cl$, Br)	$[Re_2(\mu-S)(\mu-X)X_3(CS)(\mu-dppm)_2]$	[168]
$[Re_2Br_4(\mu-dpam)_2]$	$[Re_2(\mu-S)(\mu-Br)Br_3(CS)(\mu-dpam)_2]$	[169]
$[RuH(MeCOO)(PPh_3)_3]$	$[Ru(CS)(MeCOO)(PPh_3)_2(\mu-S)]$	[88]
$[Ru(MeCN)_2Cl_2(PPh_3) + PPh_3]$	$[Ru(CS)Cl(PPh_3)_2(\mu-Cl)_2]$	[80]
$[RuH(CO)Cl(AsPh_3)_3]$	$[RuH(CO)(CS)(AsPh_3)_2] + \dots$	[88]
$[Rh(NO)(PPh_3)_3]$	$[Rh(NO)(CS)(PPh_3)_3]$	[90]




(LIV)

C_2O_4 -bridged analog, is diamagnetic owing to much greater electron delocalization of the C_2S_4 -bridged ligand. The bridging C_2S_4 unit is coordinated to two centrosymmetrically equivalent $(\eta^5-C_5H_5)Ti$ moieties.

Bianchini and co-workers reported an extensive experimental and theoretical study of C_2S_4 -bridged bimetallic complexes of rhodium and iridium [135,174]. Complexes $[(triphos)M(\mu-C_2S_4)M(triphos)]X_2$ ($M \equiv Rh$, $X \equiv Cl^-$, BPh_4^- , BF_4^- , PF_6^- ; $M \equiv Ir$, $X \equiv Cl^-$, BPh_4^-) have been prepared from $[(triphos)RhCl(\eta^2-CS_2) \cdot C_6H_6]$, $[(triphos)RhN_3(\eta^2-CS_2)]$ and $[(triphos)IrCl(\eta^2-CS_2)]$. The complex $[(triphos)Rh(\mu-C_2S_4)Rh(triphos)](BPh_4)_2$ can also be prepared by the reaction of CS_2 with




(LV)




(LVI)

$[(\text{triphos})\text{Rh}(\text{COD})](\text{BPh}_4)$ or $[(\text{triphos})\text{Rh}(\text{C}_2\text{H}_4)_2](\text{BPh}_4)$. The neutral paramagnetic complex $[(\text{triphos})\text{Rh}(\mu\text{-C}_2\text{S}_4)\text{Rh}(\text{triphos})]$ is obtained by the reaction of $[(\text{triphos})\text{RhCl}(\eta^2\text{CS}_2)]$ with NaBH_4 or LiHBEt_3 . The molecular structure of $[(\text{triphos})\text{Rh}(\mu\text{-C}_2\text{S}_4)\text{Rh}(\text{triphos})](\text{BPh}_4)_2 \cdot \text{CH}_2\text{Cl}_2$ has been determined by X-ray diffraction analysis. The C_2S_4 unit is planar but the $\text{Rh}(\mu\text{-C}_2\text{S}_4)\text{Rh}$ unit is not completely planar, each Rh atom being $0.3 \text{ \AA}$ out of the C_2S_4 plane. The relative orientation of the bridge with respect to each terminal $(\text{triphos})\text{Rh}$ fragment is such that each metal is approximately square pyramidal (LVII). The C—C bond is shorter in these $\text{M}(\text{C}_2\text{S}_4)\text{M}$ complexes compared with the free tetrathiooxalate dianion (the C—C bond is $1.461(19) \text{ \AA}$) (Table 8).



(LVII)



(LVIII)

Table 8
Selected bond distances ($\text{\AA}$) of tetrathiooxalate complexes


Complex	C—C ($\text{\AA}$)	C—S ($\text{\AA}$)	Reference
$[(\eta^5\text{-Me}_5\text{C}_5)\text{Ni}(\mu\text{-C}_2\text{S}_4)\text{Ni}(\eta^5\text{-C}_5\text{Me}_5)]$	1.36(1)	1.718(2)	[171]
$[(\text{CO})_6\text{Fe}_2(\mu\text{-C}_2\text{S}_4)\text{Fe}_2(\text{CO})_6]$	1.332	1.775	[165]
$[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\mu\text{-C}_2\text{S}_4)\text{Ti}(\eta^5\text{-C}_5\text{H}_5)_2]$	1.41	1.736	[173]
$[(\text{triphos})\text{Rh}(\mu\text{-C}_2\text{S}_4)\text{Rh}(\text{triphos})][\text{BPh}_4]$	1.37(3)	1.76(2) 1.70(2)	[174]

3.3.2.2. Metal-promoted head-to-tail dimerization of carbon disulfide. The metal-promoted reductive dimerization of CS_2 in the head-to-tail coupling yields asymmetric heterometallocycles $\text{M}(\eta^2\text{-C}_2\text{S}_4)$ and $\text{M}(\eta^3\text{-C}_2\text{S}_4)$ [105,175–178]. The rhodium complex $[(\eta^5\text{-C}_5\text{H}_5)\text{Rh}(\text{C}_2\text{S}_4)(\text{PMe}_3)]$ has an $\eta^2\text{-C}_2\text{S}_4$ ligand with a five-membered RhSCSC heterometallocyclic ring (**LVIII**) [175]. *Trans*- $[\text{M}(\text{C}_2\text{H}_4)_2(\text{PMe}_3)_4]$ ($\text{M} \equiv \text{Mo}, \text{W}$) reacts with CS_2 in diethylether to afford black complexes. $[\text{M}(\text{C}_2\text{S}_4)(\text{C}_2\text{H}_4)_2(\text{PMe}_3)_3]$ [49,105,176]. The spectral studies (IR and NMR) are consistent with a head-to-tail reductive dimerization of CS_2 and formation of a heterometallocyclic unit $\text{M}(\text{SC}(\text{S})\text{SC}=\text{S})$. Complexes $[\text{Mo}(\text{C}_2\text{S}_4)(\text{C}_2\text{H}_4)(\text{PMe}_3)_3]$ (**LIX**), $[\text{Ni}(\text{C}_2\text{S}_4\text{PMe}_3)(\text{PMe}_3)]$ (**LX**) [177], and $[\text{Rh}_2(\text{C}_2\text{S}_4)(\text{CO})\text{Cl}_2\text{-}(\text{Ph}_2\text{PCH}_2\text{PPh}_2)]$ (**LXI**) [178] have an $\eta^3\text{-C}_2\text{S}_4$ ligand with bonding through two sulfur atoms and one carbon atom.


3.3.3. Insertion reaction of carbon disulfide

3.3.3.1. Insertion into the M–H bond. Insertion of CS_2 into an M-H bond results in the formation of dithioformato complexes.


Carbon disulfide reacts with cobalt hydride complexes $[\text{CoH}_3(\text{PPh}_3)_2]$ and $[\text{CoH}\{\text{P}(\text{OPh})_3\}_4]$ to give dithioformate complexes $[\text{Co}(\text{S}_2\text{CH})(\text{PPh}_3)_2]$ and $[\text{Co}(\text{S}_2\text{CH})\{\text{P}(\text{OPh})_3\}_2]$ respectively by the insertion of CS_2 into the Co-H bond [112].




(LIX)



(LX)




(LXI)




Bianchini and co-workers have reported insertion reactions of CS_2 into the copper-hydride bond [108,109,179–182]. The yellow complex $[(\text{PPh}_3)_2\text{Cu}(\mu\text{-S}_2\text{CSCH}_2\text{SCS}_2)\text{Cu}(\text{PPh}_3)_2]$ (**LXII**) is obtained from the reaction of excess CS_2 with $[\text{Cu}(\eta^2\text{-BH}_4)(\text{PPh}_3)_2]$. In complex (**LXII**) two $(\text{PPh}_3)_2\text{Cu}$ units are held by a bridging $\text{S}_2\text{CSCH}_2\text{SCS}_2^{2-}$ ligand. On addition of n-heptane, a red complex $[\text{Cu}(\eta^2\text{-S}_2\text{CH})(\text{PPh}_3)_2]$ is formed, which reacts with carbon disulfide to reproduce the yellow complex (**LXII**). The $\eta^1\text{-BH}_4$ complexes $[(\text{triphos})\text{Cu}(\eta^1\text{-BH}_4)]$ and $[(\text{np}_3)\text{Cu}(\eta^1\text{-BH}_4)]$ react with CS_2 to give the dithioformate complexes $[(\text{triphos})\text{Cu}(\eta^1\text{-S}_2\text{CH})]$ (**LXIII**) and $[(\text{np}_3)\text{Cu}(\eta^1\text{-S}_2\text{CH})]$ (**LXIV**) respectively [179].

The ruthenium dithioformate complex $[\text{Ru}(\text{CO})\text{Cl}(\text{S}_2\text{CH})(\text{PPh}_3)_2]$ is obtained from the reaction of CS_2 with $[\text{RuH}(\text{CO})\text{Cl}(\text{PPh}_3)(4\text{-vp})]$ (4-vp is 4-vinylpyridine) [183]. The IR absorption bands at 1235 cm^{-1} and 930 cm^{-1} are consistent with the presence of a bidentate dithioformate ligand (Table 9). Insertion of CS_2 into the Ru-H bond and the formation of dithioformate complexes $[(\eta^5\text{-C}_5\text{H}_5)\text{Ru}(\text{S}_2\text{CH})\text{L}_2]$ ($\text{L} \equiv \text{PPh}_3, \text{AsPh}_3, \text{SbPh}_3$) have been reported [111].


3.3.3.2. Insertion into the M-C bond. A number of insertion reactions of CS_2 into a metal-carbon bond have been studied [184–192]. Selected structural data of insertion products are summarized in Table 10. Campus and co-workers have



(LXII)



(LXIII)



(LXIV)

Table 9
IR and ^1H NMR spectral data for dithioformate complexes

Complex	IR frequencies (cm^{-1})			^1H NMR, δ (ppm)	Reference
	$\nu_{\text{as}}(\text{CS}_2)$	$\nu_{\text{s}}(\text{CS}_2)$	$\nu(\text{HCS})$		
$[\text{Co}(\text{S}_2\text{CH})(\text{PPh}_3)_2]$	910	740	1230	—	[112]
$[\text{Co}(\eta^2\text{-S}_2\text{CH})\{\text{P}(\text{OPh})_3\}_2]$	950	—	1240	—	[112]
$[\text{Cu}(\eta^2\text{-S}_2\text{CH})(\text{PPh}_3)_2]$	960	820	1235	11.27	[179]
$[(\text{triphos})\text{Cu}(\eta^1\text{-S}_2\text{CH})]$	1012	810	1245	—	[179]
$[(\text{np}_3)\text{Cu}(\eta^1\text{-S}_2\text{CH})]$	985	—	1235	11.26	[179]
$[\text{Ru}(\text{CO})\text{Cl}(\eta^2\text{-S}_2\text{CH})(\text{PPh}_3)_2]$	930	—	1235	11.88	[183]
$[(\eta^5\text{-C}_5\text{H}_5)\text{Ru}(\eta^2\text{-S}_2\text{CH})(\text{PPh}_3)(\text{AsPh}_3)]$	1020	—	—	11.6	[111]
$[(\eta^5\text{-C}_5\text{H}_5)\text{Ru}(\eta^2\text{-S}_2\text{CH})(\text{AsPh}_3)_2]$	1020	—	—	11.6	[111]


Table 10
Selected structural data of insertion products of carbon disulfide into the M—C bond

Complex	M—S (Å)	C—C (Å)	C—S (Å)	S—C—S (deg)	S—M—S (deg)	Reference
$[\text{Cu}_2(\text{S}_3\text{C-}o\text{-tolyl})(\text{S}_2\text{C-}o\text{-tolyl})$ (bipy)]	2.247(3) 2.201(2)	1.485(7)	1.680(4) 1.679(5)	128.2(3)	—	[186]
$[(\text{C}_5\text{Me}_5\text{CS}_2)\text{Cu}(\text{PPh}_3)_2]$	2.40(1) 2.381(9)	1.61(4)	1.68(3)	118(2)	75.2(3)	[187]
$[(\text{C}_5\text{Me}_5\text{CS}_2)\text{Au}(\text{P-i-Pr}_3)]$	2.317(3)	1.50	1.72(1) 1.65(1)	120.7(6)	—	[187]
$[\text{Ru}(\text{CO})\text{Cl}](\text{S}_2\text{C-CH=CHR})$ (PPh_3) $_2$]	2.52(2) 2.35(2)	1.56	1.65(4) 1.83(5)	—	72.4(7)	[188]


reported the insertion of CS_2 in copper–carbon bonds of arylcopper(I) [phenylcopper, *o*-tolylcopper and *p*-tolylcopper] complexes, in the presence of tertiary mono and bidentate phosphines [184–187] and in the presence of bidentate nitrogen-containing ligands such as 2,2'-bipyridine and 1,10-phenanthroline [188]. The structures of $[\text{Cu}(\text{S}_3\text{C-}o\text{-tolyl})(\text{phen})]$ (LXV), $[\{\text{Cu}(\text{S}_3\text{C-}o\text{-tolyl})\}_2(\text{phen})]$ (LXVI) and $[\text{Cu}_2(\text{S}_3\text{C-}o\text{-tolyl})(\text{S}_2\text{C-}o\text{-tolyl})(\text{bipy})]$ have been determined by X-ray diffraction analysis.

Otto and Werner reported the insertion of CS_2 into Cu—C, Ag—C and Au—C bonds. Reactions of CS_2 with $[(\eta^5\text{-C}_5\text{Me}_5)\text{Cu}(\text{PR}_3)]$ ($\text{PR}_3 \equiv \text{PPh}_3$, PMe_3 , P-i-Pr_3) and with $[(\eta^5\text{-C}_5\text{Me}_5)\text{Au}(\text{PR}_3)]$ ($\text{PR}_3 \equiv \text{PPh}_3$, P-i-Pr_3) result in the formation of dithiocarboxylato complexes $[(\text{C}_5\text{Me}_5\text{CS}_2)\text{Cu}(\text{PPh}_3)_2]$ (LXVII), $[(\text{C}_5\text{Me}_5\text{CS}_2)\text{Cu}(\text{PMe}_3)_2]$, $[(\text{C}_5\text{Me}_5\text{CS}_2)\text{Cu}(\text{P-i-Pr}_3)]$, $[(\text{C}_5\text{Me}_5\text{CS}_2)\text{Au}(\text{PPh}_3)_2]$ and $[(\text{C}_5\text{Me}_5\text{CS}_2)\text{Au}(\text{P-i-Pr}_3)]$ (LXVIII) by insertion of CS_2 into the M— C_5Me_5 bond. Complex $[(\text{C}_5\text{Me}_5\text{CS}_2)\text{Ag}(\text{PPh}_3)_2]$ (LXIX) can be obtained from the reaction of excess CS_2 with $[\text{AgCl}(\text{PPh}_3)]_4$ and $\text{C}_5\text{Me}_5\text{Li}$ [189].


Insertion of CS_2 into an Ru—C bond has been observed in the reactions of CS_2 with $[\text{Ru}(\text{CO})\text{Cl}(\text{HC}\equiv\text{CHR})(\text{PPh}_3)_2]$ ($\text{R} \equiv \text{Ph}$ or *t*-Bu). The insertion products




(LXV)




(LXVI)



(LXVII)




(LXVIII)




(LXIX)

$[\text{Ru}(\text{CO})\text{Cl}(\text{S}_2\text{C}-\text{CH}=\text{CHR})(\text{PPh}_3)_2]$ contain a bidentate alkenedithiocarboxylate ligand bonded through two sulfur atoms. The molecular structure of $[\text{Ru}(\text{CO})\text{Cl}(\text{S}_2\text{C}-\text{CH}=\text{CHR})(\text{PPh}_3)_2]$ (LXX) has been determined by X-ray crystallographic analysis [190]. Insertion of CS_2 into an $\text{Ru}-\text{C}$ bond has also been observed in the reaction of CS_2 with $[(\eta^5-\text{C}_5\text{H}_5)\text{Ru}(\text{C}_2\text{Ph})(\text{PPh}_3)_2]$ producing $[(\eta^5-\text{C}_5\text{H}_5)\text{Ru}(\eta^2-\text{S}_2\text{CC}\equiv\text{CPh})(\text{PPh}_3)]$ (LXXI) [191].


Reaction of CS_2 with $[\text{PdI}(\text{CH}_3)(\text{PMe}_3)_2]$ gives *trans*- $[\text{PdI}(\text{S}_2\text{CCH}_3)(\text{PMe}_3)_2]$ (LXXII) by insertion of CS_2 into the $\text{Pd}-\text{CH}_3$ bond [140].



(LXX)



(LXXI)



(LXXII)



3.3.3.3. *Insertion into the M–N bond.* Insertion of CS_2 into an $\text{M}–\text{NR}_2$ bond produces dithiocarbamate complexes $[\text{M}(\text{S}_2\text{CR}_2)]$ [193,194].

Complex $[\text{Pt}(\text{PhNO})(\text{PPh}_3)_2]$ reacts with CS_2 to give (LXXIII) by the insertion of CS_2 into the $\text{Pt}–\text{N}$ bond [195,196].

3.3.3.4. *Insertion into the M–P bond.* Insertion of CS_2 into an $\text{M}–\text{PR}_3$ bond produces phosphinodithiocarboxylate complexes $[\text{M}(\text{S}_2\text{CPR}_3)]$. Reaction of CS_2 with $[\text{RuH}(\text{CO})\text{Cl}(\text{PCy}_3)_3]$ gives a phosphinodithiocarboxylate complex (LXXIV), by insertion of CS_2 into the $\text{Ru}–\text{PCy}_3$ bond [197].

Insertion of CS_2 into an $\text{Rh}–\text{PPh}_3$ bond was observed in the reaction of CS_2 with $[\text{Rh}_2(\text{CO})_2(\text{PPh}_3)_4(\mu\text{-bipy})]$ and $[\text{Rh}(\text{CO})\text{L}(\text{PPh}_3)_2]\text{X}$ ($\text{L} \equiv \text{py}$, α -picoline; $\text{X} \equiv \text{ClO}_4$, NO_3). The insertion products are $[\text{Rh}_2(\text{CO})_2(\text{PPh}_3)_2(\text{S}_2\text{CPh}_3)_2(\mu\text{-bipy})]$ and $[\text{Rh}(\text{CO})\text{L}(\text{PPh}_3)(\text{S}_2\text{CPh}_3)]$ [198].

CS_2 inserts into the $\text{Pd}–\text{PR}_3$ bond in complexes $[\text{Pd}(\text{PR}_3)_4]$ ($\text{PR}_3 \equiv \text{PMe}_3$,







PMe_2Ph) to give binuclear complexes $[\text{Pd}(\text{S}_2\text{CPR}_3)(\text{PR}_3)]_2$. Complex $[\text{Pd}(\text{S}_2\text{CPMe}_3)(\text{PMe}_3)]_2$ can also be obtained from the reaction of $[\text{Pd}(\eta^2\text{-CS}_2)(\text{PPh}_3)_2]$ with PMe_3 [140]. Reactions of CS_2 with $[\text{PdI}(\text{COCH}_3)(\text{PMe}_3)_2]$ and $[\text{PdR}(\text{PMe}_3)_3](\text{BPh}_4)$ ($\text{R} \equiv \text{Me, COMe, Ph}$) give phosphinodithiocarboxylate complexes (LXXV) and (LXXVI) [140,199].

Carbon disulfide inserts into the Zr-P bond of $[(\eta^5\text{-C}_5\text{H}_5)_2\text{ZrCl}(\text{P}\{\text{SiMe}_3\}_2)]$ to give $[(\eta^5\text{-C}_5\text{H}_5)_2\text{ZrCl}\{\eta^2\text{-S}_2\text{CP}(\text{SiMe}_3)_2\}]$. The structure of the complex $[(\eta^5\text{-C}_5\text{H}_5)_2\text{ZrCl}(\eta^2\text{-S}_2\text{CP}(\text{SiMe}_3)_2)]$ is pyramidal with average Zr-S distance of 2.686(8) Å and angle S-Zr-S of $63.4(2)^\circ$ [200].

3.3.3.5. Insertion into the M-S bond. Insertion of CS_2 into M=S and $\text{M}(\eta^2\text{-S}_2)$ bonds gives trithiocarbonate and perthiocarbonate complexes:

Reaction of CS_2 with sulfidomolybdenum complexes leads to the insertion of CS_2 into Mo=S and $\text{Mo}(\eta^2\text{-S}_2)$ bonds and to the formation of CS_3^{2-} and CS_4^{2-} complexes respectively [201,202]. The $\text{Mo}-\eta^2\text{-CS}_4$ units dissociate CS_2 with the formation of the $\text{Mo}-\eta^2\text{-CS}_4$ units while the $\text{Mo}-\eta^2\text{-CS}_3$ units do not dissociate. Reaction of CS_2 with $[\text{Ph}_4\text{P}]_2[(\text{S})\text{Mo}(\text{S}_4)_2]$ in DMF results in the




formation of $[\text{Ph}_4\text{P}]_2[(\text{S})\text{Mo}(\text{CS}_4)_2]$ (LXXVII). Complex $[\text{Et}_4\text{N}]_2[\text{Mo}_2(\text{O})_2(\mu\text{-S})(\eta^2\text{-CS}_4)(\eta^2\text{-CS}_3)]$ (LXXVIII) is obtained from the reaction of $[\text{Et}_4\text{N}]_2[(\text{S})\text{Mo}(\mu\text{-S})\text{Mo}(\text{O})(\text{S}_4)]$. The synthesis and molecular structure of complex (LXXIX) has also been reported. The nickel complex $[\text{Ph}_4\text{P}]_2[\text{Ni}(\text{CS}_4)_2]$ can be obtained from the reaction of $[\text{Ph}_4\text{P}]_2[\text{Ni}(\text{S}_4)_2]$ [203].


3.3.3.6. Insertion into the M—Cl bond. Insertion of CS_2 into an Au—Cl bond has been reported in the reaction of gold(III) chloride $[\text{Au}_2\text{Cl}_6]$ with CS_2 . The insertion product $[\text{AuCl}_2(\eta^2\text{-S}_2\text{CCl})]$ contains a chlorodithioformate ligand (LXXX). The structure was determined by X-ray crystallography. The molecule is planar with S—Au—S and S—C—S angles of $75.0(2)^\circ$ and $114(1)^\circ$ respectively [204].

3.3.4. Alkylation reaction of carbon disulfide


The negative charges on the sulfur atoms of CS_2 ligand in $\eta^2\text{-CS}_2$ and $\eta^1\text{-CS}_2$ coordination modes show its nucleophilic character and therefore it undergoes alkylation with a variety of electrophiles. Reaction of $[\text{M}-\eta^2\text{-CS}_2]$ and





(LXXVII)



(LXXVIII)




(LXXIX)



$[M-\eta^1\text{-CS}_2]$ complexes with methyl iodide results in the formation of $[M-\eta^2\text{-CS(SMe)}]$, $[M-\eta^1\text{-CS(SMe)}]$ and $[M-\eta^1\text{-C(SMe)}_2]$ complexes. The CS stretching frequencies provide a useful diagnostic for determining the hapticity of the CS(SMe) ligand ($\nu(\text{CS}) \approx 1000\text{ cm}^{-1}$ for $\eta^1\text{-CS(SMe)}$ and $\nu(\text{CS}) \approx 1100\text{ cm}^{-1}$ for $\eta^2\text{-CS(SMe)}$ ligand).

$[\text{Os}(\text{NO})\text{Cl}(\text{CS}_2)(\text{PPh}_3)_2]$ reacts with methyl iodide in dichloromethane to afford the methylated complex (LXXXI). IR spectral studies ($\nu(\text{CS})$, 991 cm^{-1}) reveal that the dithiomethoxycarbonyl ligand binds in a monodentate manner. In ethanol, $[\text{Os}(\text{NO})\text{Cl}(\text{CS}_2)(\text{PPh}_3)_2]$ yields a cationic complex (LXXXII) containing a bidentate CS_2Me ligand which may be isolated by addition of a non-coordinating anion (PF_6^- or ClO_4^-).



The complex (LXXXII) can also be prepared by the reaction of trifluoromethylsulfo-nate and NH_4PF_6 with $[\text{Os}(\text{NO})\text{Cl}(\text{CS}_2)(\text{PPh}_3)_2]$ [129].

Complex $[\text{Fe}(\text{CO})_2(\eta^2\text{-CS}(\text{SMe})(\text{PPh}_3)_2)(\text{PF}_6)]$ is obtained by the reaction of MeI with $[\text{Fe}(\text{CO})_2(\eta^2\text{-CS}_2)(\text{PPh}_3)_2]$ in the presence of NH_4PF_6 [205]. A bimetallic complex (LXXXIII) is formed by treating $[\text{Fe}(\text{CO})_2(\eta^2\text{-CS}(\text{SMe})(\text{PPh}_3)_2)(\text{PF}_6)]$ with Na-Hg in THF.

The complex anion $[(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})_2(\eta^1\text{-CS}_2)]^-$ reacts with MeI or Me_3SiCl to afford dithiocarboxylate ester complexes (LXXXIV) ($\text{R} \equiv \text{Me}$ or SiMe_3) [127,150] which on further methylation with $\text{MeOSO}_2\text{CF}_2$ afford the dithiocarbene complex (LXXXV) containing triflate or PF_6^- counter anions.




Platinum CS_2 complexes $[\text{Pt}(\eta^2\text{-CS}_2)\text{L}_2]$ ($\text{L} \equiv \text{PPh}_3$, $1/2$ dppe) react with MeI to give $[\text{PtI}\{\eta^1\text{-C}(\text{SMe})_2\}\text{L}_2]^+ \text{I}^-$ (LXXXVI) [143].




3.3.5. Miscellaneous reactions

Carbon disulfide undergoes coupling reactions with ligands coordinated to a metal center. The thiocarbene complex $[\text{Cr}(\text{CO})_5\{\text{C}(\text{SEt})\text{Ph}\}]$ is isolated by successive treatment of CS_2 and $[\text{Et}_3\text{O}][\text{BF}_4]$ with the deprotonated aminocarbene complex $[\text{Cr}(\text{CO})_5\{\text{C}(\text{NH})\text{Ph}\}]^-$ [206].


No net incorporation of CS_2 occurs in the reaction of the secondary aminocarbene complex with CS_2 under the same conditions. The alkoxy carbene complex $[\text{Cr}(\text{CO})_5\{\text{C}(\text{OEt})\text{Me}\}]$ reacts with LiBu, CS_2 and $[\text{Et}_3\text{O}][\text{BF}_4]$ successively to produce (LXXXVII).


Reactions of deprotonated sulfur donor complexes of pentacarbonylchromium(0)





with carbon disulfide have been reported by Raubenheimer and co-workers [207,208].





Reaction of CS_2 with complex $[(\eta^5\text{-C}_5\text{H}_5)\text{Fe}\{\text{C}(\text{OMe})=\text{CH}_2\}]$ produces a thermo-labile complex which reacts with MeI to give $[(\eta^5\text{-C}_5\text{H}_5)\text{Fe}\{\text{C}(\text{OMe})=\text{CH}_2\text{C}(\text{S})\text{SMe}\}(\text{PMe}_3)_2]\text{I}$ [209].



Reaction of $[\text{Pt}(\eta^2\text{-CS}_2)(\text{PPh}_3)_2]$ with PMe_3 , PMe_2Ph and PMePh_2 results in the formation of trithiocarbonate complexes (LXXXVIII) [210].





Reduction of $[(\text{triphos})\text{Co}(\mu\text{-Cl})_2\text{Co}(\text{triphos})]^{2+}$ in THF solution under nitrogen atmosphere with sodium amalgam produces a deep red-brown solution which reacts with CS_2 to afford $[(\text{triphos})\text{Co}(\text{S}_2\text{CO})]$ (LXXXIX) [211]. Reaction of CS_2 with malonodinitrile in the presence of $[\text{IrH}(\text{PMe}_2\text{Ph})_4]$ yields $[\text{IrH}_2(\text{PMe}_2\text{Ph})_4]^+[\text{HS}(\text{SCC})(\text{CH}_2)_2]^-$ [212]. The coupling reaction of CS_2 in ruthenium complex $[(\eta^5\text{-C}_5\text{H}_5)\text{RuH}(\text{sp})]$ ($\text{sp} = 2\text{-CH}_2=\text{CHC}_6\text{H}_4\text{PPh}_2$) gives complex (XC) [213].

4. Coordination chemistry of carbon dioxide

4.1. Transition metal–carbon dioxide bonding and energetics




The coordination of CO_2 ligand to transition metal gives complexes of the following types: (a) η^1 -end-on coordination (XCI), (b) η^1 -C coordination (XCII),





(c) η^2 -side-on coordination (XCIII), and (d) bridging CO_2 complexes (XCIV-XCVI).

Molecular orbital studies of complexes containing a CO_2 ligand have led to a



better understanding of the nature of the coordinate bond, its electronic structure, conformational performance and reactivity. All three possible modes of coordination of the CO₂ molecule to a single metal atom, (a) η^1 -O end-on mode (XCI), (b) η^1 -C mode (XCII) and (c) η^2 -side-on mode (XCIII), have been discussed in molecular orbital studies [80,119–122,214–230]. The gross atomic charges of the CO₂ complexes are presented in Table 11.

The end-on η^1 -O coordination mode is preferred by the electrostatic interaction of the metal, having a considerable positive charge, with the negatively charged oxygen atom of the CO₂ ligand. No structural evidence for the end-on conformation has ever been observed, but it has been calculated theoretically for [Cu(CO₂)(PH₃)₂] [119] and [Cu(CO₂)] [221] and characterized spectroscopically for [Cu(CO₂)] in a matrix [231].

Both the η^1 -C coordination and η^2 -side-on coordination modes need strong charge transfer interactions from the metal to the CO₂ ligand and weak four-electron destabilization interactions [120,219]. Mulliken population analysis shows negative charges on the coordinated CO₂ and positive charges on the metal in transition metal CO₂ complexes. These results suggest that the CO₂ ligand removes an electron from the central metal atom as a result of strong metal to ligand charge transfer interactions [80,119,219–221,223,226].

The η^1 -C coordination requires charge transfer from metal $d\sigma$ to the CO₂ π^* orbitals and the electrons accepted by the η^1 -CO₂ ligand are distributed not only on the carbon atom but also on the oxygen atom. This coordination mode is most favored when the HOMO is mainly composed of a $d\sigma$ (d_z^2) orbital and when the metal is in a low oxidation state as found in the Rh(I) complex [RhCl(η^1 -CO₂)(diars)₂] [166], the Co(I) complex [Co(alcn)(η^1 -CO₂)][−] [162], and Ni(I) complexes

Table 11

Binding energy and net charges on metal and carbon dioxide for metal carbon dioxide complexes

Metal system	Binding energy (kcal mol ⁻¹)	Charge		Reference
		Metal	CO ₂	
<i>η¹-end-on coordination</i>				
[CuCO ₂]	-11.7	+0.72	-0.72	[221]
[Cu(CO ₂)(PH ₃) ₂] ⁻	-14.0	+0.85	+0.017	[119]
<i>η¹-C coordination</i>				
[Co(alcn) ₂ (CO ₂)]	-6.2	+0.40	-0.72	[219]
[RhCl(CO ₂)(AsH ₃) ₄]	-1.8	—	-0.53	[223]
[NiF(NH ₃) ₄ (CO ₂)]	-22.0	+0.687	-0.872	[226]
<i>η²-side-on coordination</i>				
[Mo(CO ₂) ₂ (PH ₃) ₄]	—	+1.31	-0.57	[220]
[Mo(CO ₂) ₂ (CNH)(PH ₃) ₃]	—	+1.68	-0.42	[220]
[Mo(C ₂ H ₄)(CO ₂) ₂ (PH ₃) ₃]	—	+0.09	-0.51	[220]
[Fe(CO) ₂ (CO ₂)(PH ₃) ₂]	-26.2	-0.38	-0.52	[80]
[Fe(CO ₂)(PH ₃) ₄]	-63.4	-0.24	-0.73	[80]
[Ni(CO ₂)(PH ₃) ₂]	-27.0	+0.569	-0.689	[119]

$[\text{NiF}(\text{NH}_3)_4(\eta^1\text{-CO}_2)]$ and $[\text{Ni}(\text{NH}_3)_5(\eta^1\text{-CO}_2)]^+$ [169]. Hoffman and co-workers, using the EHMO method, have proposed that for $\eta^1\text{-C}$ coordination, two conditions must be fulfilled: a closed shell configuration of metal fragment and the presence of a free coordination site [120].

The $\eta^2\text{-side-on}$ coordination requires charge transfer from metal $d\pi$ to CO_2 π^* orbitals. The best situation for $\eta^2\text{-side-on}$ coordination is the presence of a metal $d\pi$ as HOMO and an empty $d\sigma$ orbital pointing to the CO_2 ligand. The CO_2 distortion on coordination lowers the π^* orbital energy of CO_2 and hence increases π -back donation.

Influence ligand on the $\text{M}-\text{CO}_2$ moiety



Substitution of a ligand by a more efficient π -acceptor ligand reduces the electron density on the central metal atom and hence decreases the metal to CO_2 π -back donation as well as the binding energy of $\text{M}-\text{CO}_2$. A comparison of bonding energies of (XCVII) and (XCVIII) shows that the substitution of phosphines by carbonyls involves a decrease in the CO_2 binding energy. A relatively smaller negative charge on CO_2 in (XCVII) is in agreement with decreased π -back donation. Similarly, for complexes (XCIX) and (C), on substitution of a phosphine ligand by an isocyanide ligand, which is a better π -acceptor than the phosphine ligand, the molybdenum net charge increases from 1.31 to 1.68 and the CO_2 net charge changes from -0.57 to -0.42 .



4.2. Preparation and structure

4.2.1. Mononuclear transition metal carbon dioxide complexes

Transition metal CO_2 complexes are prepared by (a) reaction of CO_2 with transition metal complexes and (b) oxidation of transition metal carbonyls.



4.2.1.1. $\text{M}-\eta^1\text{-CO}_2$ complexes. The titanium complex $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\text{PMe}_3)_2]$ reacts with CO_2 to give the yellow $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\text{CO}_2)(\text{PMe}_3)]$ [123,232]. The results of detailed IR spectral studies of $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\text{CO}_2)(\text{PMe})]$ suggest $\eta^1\text{-CO}_2$ co-





ordination of CO₂ in the titanium complex (CI) [232]. The side-on coordination mode η^2 -CO₂ was proposed by Rausch and co-workers [123]. Complex Li₂[W(CO)₅(η^1 -CO₂)] is prepared in THF solution at -78°C by the reaction of CO₂ with Li₂[W(CO)₅]. An attempt to isolate a solid product was unsuccessful [126]. The IR and ¹³C NMR spectral studies are consistent with the formulation as a W- η^1 -CO₂ complex. The formation of complexes, containing η^1 -CO₂ ligand with alkali metal counter anions, of the type [A][(η^5 -C₅H₅)Fe(CO)₂(η^1 -CO₂)] (A⁺ $\equiv$ Li⁺, Na⁺, K⁺) [233,234], [A][(η^5 -C₅H₅)Fe(CO)(PPh₃)(η^1 -CO₂)] (A⁺ $\equiv$ Li⁺, K⁺) [235], [A][(η^5 -C₅H₅)Re(NO)(PPh₃)(η^1 -CO₂)] (A⁺ $\equiv$ Li⁺, K⁺) [236], [A][(η^5 -C₅H₅)Re(CO)(NO)(η^1 -CO₂)] (A⁺ $\equiv$ Na⁺, ET₃NH⁺) [237], [A][(η^5 -C₅H₅)Re(CO)(N₂Ph)(η^1 -CO₂)] (A⁺ $\equiv$ Li⁺, Na⁺) [238] has been observed.

Carbon dioxide complexes of Co(I) containing 14-membered tetraazamacrocycles (CII) have been prepared in acetonitrile solution by the addition of carbon dioxide to a solution of the Co(I) complexes, obtained by electrochemical reduction or sodium-amalgam reduction [239–241]. The solid perchlorate salt can be obtained from acetonitrile solution by rapid addition of THF [239]. [Co([14]diene)(η^1 -CO₂)] [ClO₄] shows C–O stretching frequencies at 1653, 1304 and 1222 cm^{−1} which are similar to those reported for [Co(n-pr-salen)(CO₂)(THF)][−] [242]. Electronic spectra, thermodynamic and reactivity studies carried out on the Co(I)- η^1 -CO₂ complexes were reported [241]. The



binding of CO_2 to Co(I) and Ni(I) tetraazamacrocyclic complexes has been determined using cyclic voltammetry or differential pulse polarography. The CO_2 binding constants showed a strong correlation with redox potential, solvent properties and ligand structure [243]. CO_2 binding by $[\text{Co(L)}]^+$ ($\text{L} \equiv \text{Me}_6[14]4,11\text{-diene}$) has been determined electrochemically. A K_{CO_2} value of $7 \times 10^4 \text{ M}^{-1}$ for CO_2 binding to $[\text{Co(L)}]^+$ has been obtained in 0.1 M tetrabutylammonium perchlorate in Me_2SO [244].

Herskovitz and co-workers reported the synthesis and X-ray crystal structure of $[\text{RhCl}(\text{CO}_2)(\text{diars})_2]$ where CO_2 is $\eta^1\text{-CO}_2$ bonded and CO_2 and Cl ligands are mutually trans (CIII) [245,246]. The $\eta^1\text{-CO}_2$ coordination is stabilized by additional $\text{C-H} \cdots \text{O}$ interactions with the ligand.

4.2.1.2. $\text{M}-\eta^2\text{-CO}_2$ complexes. The formation of transition metal CO_2 complexes as possible intermediates in the transition metal mediated oxidation of CO has been suggested [247–249]. The synthesis of CO_2 complexes from the reaction of transition metal carbonyls with molecular oxygen or air has been reported for the first time by Nicholas and co-workers [250]. Complexes $[(\eta^5\text{-C}_5\text{Me}_5)_2\text{Nb}(\eta^2\text{-CO}_2)(\text{R})]$ ($\text{R} \equiv \text{Me}, \text{CH}_2\text{CMe}_3, \text{CH}_2\text{Ph}, \text{CH}_2\text{SiMe}_3$) are prepared by the addition of air or O_2 into a solution of $[(\eta^5\text{-C}_5\text{Me}_5)_2\text{Nb}(\text{CO})(\text{R})]$ in toluene. The molecular structure of $[(\eta^5\text{-C}_5\text{Me}_5)_2\text{Nb}(\eta^2\text{-CO}_2)(\text{CH}_2\text{Ph})]$ was determined by X-ray crystallography. The structure of (CIV) is same as that of $[(\eta^5\text{-C}_5\text{H}_4\text{Me})_2\text{Nb}(\eta^2\text{-CO}_2)(\text{CH}_2\text{SiMe}_3)]$ (CV) [251] except for the longer Nb— CH_2R bond of (CIV) (2.337(4) Å vs. 2.282(11) Å for (CV)) (Table 12).

A molybdenum CO_2 complex $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Mo}(\eta^2\text{-CO}_2)]$ (CVI) is obtained from the reaction of $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Mo}(\text{PhC}\equiv\text{CPh})]$ [252]. Irradiation (more than 330 nm) of a THF solution of $[(\eta^5\text{-C}_5\text{H}_5)_2\text{MoH}_2]$ under 1 atm CO_2 at -10°C for 3 h results in the formation of (CVI) along with a dimer $[(\eta^5\text{-C}_5\text{H}_5)(\eta^5\text{-C}_5\text{H}_4)\text{Mo}]_2$ [253]. Complex (CVI) can also be prepared in low yield when a THF solution of $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Mo}(\text{CO})]$ is exposed to O_2 at 0°C [250].

The reaction of *cis*- $[\text{Mo}(\text{N}_2)_2(\text{PMe}_3)_4]$ with CO_2 under pressure (4–5 atm) gives bis(carbon dioxide) complex *trans*- $[\text{Mo}(\text{CO}_2)_2(\text{PMe}_3)_4]$ (CVII) which reacts with various isocyanides to afford CO_2 complexes *trans,mer*- $[\text{Mo}(\text{CO}_2)(\text{CNR})(\text{PMe}_3)_3]$ ($\text{R} \equiv \text{Me}, \text{i-Pr}, \text{t-Bu}, \text{Cy}$ and CH_2Ph) [102–105]. Molecular structures of








Table 12

X-ray structural data for mononuclear transition metal carbon dioxide complexes


Complex	M–C (Å)	M–O (1) (Å)	C–O (1) (Å)	C–O (2) (Å)	O (1)–C–O (2) (°)	Reference
K[Co(salen)(η^1 -CO ₂)]	1.99		1.22		132	[242]
[(η^5 -C ₅ H ₄ Me) ₂ Nb(η^2 -CO ₂)- (CH ₂ SiMe ₃)]	2.144(7)	2.172(4)	1.283(8)	1.216(8)	132.4(7)	[251]
[(η^5 -C ₅ H ₅) ₂ Mo(η^2 -CO ₂)]	2.112(11)	2.160(7)	1.288(14)	1.201(14)		[252]
[Mo(η^2 -CO ₂)(CN-i-Pr)(PMe ₃) ₃]	2.10(1)	2.146(7)	1.26(2)	1.22(2)	133(1)	[104]
	2.11(1)	2.147(7)	1.26(1)	1.22(1)	134(1)	
[Mo(η^2 -CO ₂)(CN-CH ₂ Ph)- (PMe ₃) ₃]	2.02(2)	2.14(1)	1.25(2)	1.28(3)	128(2)	[104]
	2.02(2)	2.16(1)	1.20(2)	1.28(3)	128(2)	
[Rh(η^1 -CO ₂)Cl(diars) ₂]	2.05(2)		1.25(2)		126(2)	[246]
			1.20(2)			



trans,mer-[Mo(CO₂)(CN-i-Pr)(PMe₃)₃] (CVIII) and *trans,mer*-[Mo(CO₂)(CNCH₂Ph)(PMe₃)₃] (CIX) were determined by X-ray diffraction. The complexes (CVIII) and (CIX) are isostructural with the molybdenum atom bonded to two *trans*, staggered CO₂ molecules. The overall molecular geometry about molybdenum is approximately octahedral.




(CVIII)



(CIX)

4.2.2. Bimetallic and trimetallic carbon dioxide complexes





The $M-\eta^1\text{-CO}_2$ complexes react with metal complexes to afford bimetallic $\mu_2(\eta^1\text{-C}; \eta^1\text{-O})$ and $\mu_2(\eta^1\text{-C}; \eta^2\text{-O})$ CO_2 complexes.



Geoffrey and co-workers reported the formation of bimetallic CO_2 complexes from the oxidation of transition metal carbonyl complexes with oxo complex [254]. The reaction of $[(\eta^5\text{-C}_5\text{H}_5)_2\text{M}'=\text{O}]$ ($\text{M}' \equiv \text{Mo}, \text{W}$) with carbonyl complexes $[(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})_3]^+$, $[(\eta^5\text{-C}_5\text{Me}_5)\text{Ru}(\text{CO})_3]^+$, $[(\eta^5\text{-C}_5\text{H}_4\text{Me})\text{Mn}(\text{NO})(\text{CO})_2]^+$ and $[(\eta^5\text{-C}_5\text{H}_5)\text{Re}(\text{NO})(\text{CO})_2]^+$ result in the formation of bimetallic CO_2 complexes. The structures of $\text{Fe}\{\mu_2(\eta^1\text{-C}, \eta^2\text{-O}, \text{O})\text{W}\}$ and $\text{Re}\{\mu_2(\eta^1\text{-C}, \eta^2\text{-O}, \text{O})\text{W}\}$ are shown as (CX) and (CXI) respectively. Reaction of $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Zr}(\text{CH}_3)\text{Cl}]$ with $[(\eta^5\text{-C}_5\text{H}_5)\text{Re}(\text{NO})(\text{CO})(\text{CO}_2\text{H})]$ result in the formation of the yellow complex $[(\eta^5\text{-C}_5\text{H}_5)(\text{NO})(\text{CO})\text{Re}-\text{C}(\text{O})\text{O}-\text{ZrCl}(\eta^5\text{-C}_5\text{H}_5)_2]$ (CXII) [255].

Reaction of metallocarboxylate $[(\eta^5\text{-C}_5\text{H}_5)\text{M}(\text{CO})_2(\text{CO}_2)]^-\text{Na}^+$ ($\text{M} \equiv \text{Fe}, \text{Ru}$) with zirconocene dichloride $[(\eta^5\text{-C}_5\text{H}_5)_2\text{ZrCl}_2]$ and titanocene dichloride $[(\eta^5\text{-C}_5\text{H}_5)_2\text{TiCl}_2]$ gives bridging CO_2 complexes $[(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{MCO}_2\text{M}'\text{Cl}(\eta^5\text{-C}_5\text{H}_5)_2]$ (CXIII) ($\text{M} \equiv \text{Fe}, \text{Ru}; \text{M}' \equiv \text{Zr}, \text{Ti}$) [256].



The nickel complex $[\{\text{Cy}_3\text{P}\}_2\text{Ni}\}_2(\mu\text{-CO}_2)]$ is obtained by the splitting of the CO_2




complex $[\text{Ni}(\eta^2\text{-CO}_2)(\text{PCy}_3)_2]$ [257]. The Vaska complex $[\text{Ir}(\text{CO})\text{Cl}(\text{PPh}_3)_2]$ reacts with osmium tetra-oxide in 4-*t*-BuC₅H₄N (L) to give the μ_2 -(η^1 -C, η^1 -O) bonded CO₂ complex (CXIV) [258].


A platinum carboxylic acid complex *trans*- $[\text{Pt}(\text{CO}_2\text{H})(\text{Ph})(\text{PEt}_3)_2]$, obtained from the reaction of CO₂ with *trans*- $[\text{Pt}(\text{OH})(\text{Ph})(\text{PEt}_3)_2]$, reacts with its precursor *trans*- $[\text{Pt}(\text{OH})(\text{Ph})(\text{PEt}_3)_2]$ to afford a bridging O₂ complex (CXV) which can also be obtained by the pyrolysis of *trans*- $[\text{Pt}(\text{CO}_2\text{H})(\text{Ph})(\text{PEt}_3)_2]$ [259]. The molecular structure of the complex $[\text{Pt}(\text{CH}_3)(\text{dppp})]_2(\mu\text{-CO}_2)$, prepared from $[\text{Pt}(\text{OH})(\text{CH}_3)(\text{dppp})]$ and $[\text{Pt}(\text{CO}_2\text{H})(\text{CH}_3)(\text{dppp})]$, has been determined by X-ray crystallography and shows the presence of a μ_2 -(η^1 -C, η^1 -O) CO₂ ligand.

The synthesis and characterization of bridging CO₂ complexes involving three or more metal centers (μ_3 -CO₂) have also been reported [260–266]. The complex $[\text{Re}(\text{OH})(\text{CO})_5]$ loses water at 20 °C in acetone to give the μ_3 CO₂ complex (CXVI) [264]. The yellow complex $[(\text{COD})_2\text{Rh}_2\text{OsH}_2(\mu_3\text{-CO}_2)(\text{PMe}_2\text{Ph})_2]$ (CXVII) is obtained by the reaction of $[(\text{COD})\text{RhH}_3\text{Os}(\text{PMe}_2\text{Ph})_3]$ with CO₂ in THF [265,266].





(CXVI)



(CXVII)

4.3. Spectroscopic studies

Transition metal CO₂ complexes exhibit characteristic CO₂ vibrations in the regions 1650–1620 cm⁻¹, 1300–1100 cm⁻¹, 900–800 cm⁻¹ and 600–400 cm⁻¹ for the M–η²-CO₂ coordination mode. The first two bands are assigned to ν_{as}(OCO) and ν_s(OCO) respectively. Comparison of the spectral data of the coordinated CO₂ ligand with the IR spectral data for free CO₂ [267] shows a decrease in ν_{as}(OCO) frequency on coordination to transition metal systems. A decrease in ν_{as}(OCO) on coordination to the metal and variation in ν(OCO) frequencies depend on the π-acceptor and σ-donor abilities of the CO₂ ligand. Mascetti and co-workers [231,232,268] have investigated the IR characterization of transition metal CO₂ complexes. The observations of isotropic shifts on the two ν_{as}(OCO) and ν_s(OCO) stretching modes and the splitting between the two CO stretching modes (ν_{as}(OCO)–ν_s(OCO)) give an indication of the CO₂ bonding mode in metal complexes. For the M–η¹-CO₂ coordination mode, the difference Δ (ν_{as}–ν_s) is approximately equal to 400 cm⁻¹ while for the M–η²-CO₂ coordination mode the difference Δ is above 500 cm⁻¹ (Table 13).

The ¹³C NMR spectrum of CO₂ exhibits absorption at δ 132 ppm. On coordination to transition metal systems, it shows downfield shifts (Table 14). The chemical shifts for the CO₂ carbon depend on (a) the extent of carbon hybridization, (b) charge delocalization and (c) the nature of the bond between the metal atom and CO₂. The higher chemical shifts δ of 260 ppm for [(η⁵-C₅H₅)(CO)₂-Fe=CO₂W(η⁵-C₅H₅)₂][PF₆] and δ 252 ppm for [(η⁵-C₅Me₅)(CO)₂-Fe=CO₂W(η⁵-C₅H₅)₂][BF₄] are in agreement with their formulation as dimetalated dioxocarbene ligands. ¹H NMR, ³¹P NMR and electronic spectral studies have been reported for a number of transition metal CO₂ complexes.

4.4. Insertion reactions of carbon dioxide complexes

4.4.1. Insertion into the M–H bond

The insertion of CO₂ into the M–H bond leads to formato complexes (CXVIII–CXIX) or a l-metalated formic acid complex (CXX). Most of the insertion reactions result in the formation of formato complexes.

Table 13

Important IR frequencies (cm^{-1}) of transition metal carbon dioxide complexes

Complex	$\nu(\text{CO}_2)$ (cm^{-1})	Reference
<i>M-η^1-CO₂ coordination</i>		
$[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\eta^1\text{-CO}_2)(\text{PMe}_3)]$	1671, 1280, 1215	[232]
$\text{Li}_2[\text{W}(\text{CO})_5(\eta^1\text{-CO}_2)]$	1450	[126]
$[\text{Co}([\text{14}] \text{diene})(\eta^1\text{-CO}_2)][\text{ClO}_4]$	1653, 1304, 1222	[239]
$\text{K}[(\text{Pr-salen})\text{Co}(\eta^1\text{-CO}_2)]$	1650, 1280, 1215	[242]
$[\text{RhCl}(\eta^1\text{-CO}_2)(\text{diars})_2]$	1610, 1210	[245]
<i>M-η^2-CO₂ coordination</i>		
$[(\eta^5\text{-C}_5\text{H}_4\text{Me})_2\text{Nb}(\eta^2\text{-CO}_2)(\text{CH}_2\text{Ph})]$	1738	[250]
$[(\eta^5\text{-C}_5\text{H}_5)_2\text{Mo}(\eta^2\text{-CO}_2)]$	1740	[252]
$[\text{Mo}(\eta^2\text{-CO}_2)(\text{PMe}_3)_4]$	1670, 1155, 1100	[104]
$[\text{Mo}(\eta^2\text{-CO}_2)(\text{CNMe})(\text{PMe}_3)_3]$	1660, 1150, 1100	[104]
$[\text{Mo}(\eta^2\text{-CO}_2)(\text{CN-i-Pr})(\text{PMe}_3)_3]$	1675, 1160, 1100	[104]
$[\text{Mo}(\eta^2\text{-CO}_2)(\text{CN-t-Bu})(\text{PMe}_3)_3]$	1680, 1155, 1100	[104]
$[\text{Mo}(\eta^2\text{-CO}_2)(\text{CNCy})(\text{PMe}_3)_3]$	1665, 1150, 1100	[104]
$[\text{Mo}(\eta^2\text{-CO}_2)(\text{CNCH}_2\text{Ph})(\text{PMe}_3)_3]$	1670, 1150, 1100	[104]
$[\text{Fe}(\eta^2\text{-CO}_2)(\text{PMe}_3)_4]$	1623, 1106	[268]
<i>Bridging carbon dioxide complexes</i>		
$[(\eta^5\text{-C}_5\text{H}_5)(\text{CO})(\text{NO})\text{Re}(\mu\text{-CO}_2)\text{ZrCl}(\eta^5\text{-C}_5\text{H}_5)_2]$	1348, 1288	[256]
$[(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Fe}(\mu\text{-CO}_2)\text{ZrCl}(\eta^5\text{-C}_5\text{H}_5)_2]$	1363, 1264	[256]
$[(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Fe}(\mu\text{-CO}_2)\text{TiCl}(\eta^5\text{-C}_5\text{H}_5)_2]$	1379, 1273	[256]
$[(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Ru}(\mu\text{-CO}_2)\text{ZrCl}(\eta^5\text{-C}_5\text{H}_5)_2]$	1344, 1285	[256]
$[(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Ru}(\mu\text{-CO}_2)\text{TiCl}(\eta^5\text{-C}_5\text{H}_5)_2]$	1349, 1284	[256]
$[(\text{COD})_2\text{Rh}_2\text{OsH}_2(\mu\text{-CO}_2)(\text{PMe}_2\text{Ph})_3]$	1365, 1260	[265]
$[(\text{PPh}_3)_2(\text{L})\text{ClIr}(\mu\text{-CO}_2)(\mu\text{-O})\text{Os}(\text{O})_2(\text{L})_2]$ ($\text{L} \equiv 4\text{-t-BuC}_5\text{H}_4\text{N}$)	1593, 1022	[258]
<i>trans</i> - $[\text{Pt}(\text{Ph})(\text{PEt}_3)_2]_2(\mu\text{-CO}_2)$	1495, 1290, 1190	[259]

The titanium formato complex $[(\eta^5\text{-C}_5\text{Me}_5)_2\text{Ti}(\text{O}_2\text{CH})]$, along with alkene, is formed in the reaction of $[(\eta^5\text{-C}_5\text{Me}_5)_2\text{TiR}]$ ($\text{R} \equiv \text{Et}$ or $n\text{-Pr}$) with CO_2 [269]. Lyons and co-workers reported the formation of a formato complex $[\text{MoH}(\text{O}_2\text{CH})(\text{PMe}_2)_4]$ from the reaction of CO_2 with $[\text{MoH}_2(\text{PMe}_3)_5]$ by the insertion of CO_2 into the Mo-H bond. The pentagonal bipyramidal structure (CXXI) was determined for $[\text{MoH}(\text{O}_2\text{CH})(\text{PMe}_3)_4]$ by X-ray diffraction [270]. The analogous formate complex $[\text{MoH}(\text{O}_2\text{CH})(\text{dppe})_2]$ (CXXII) is obtained by passing CO_2 into a irradiated yellow solution of $[\text{MoH}_4(\text{dppe})_2]$ in benzene [271]. The ^1H NMR spectra of (CXXI) and (CXXII) show signals at $\delta -8.66$ ppm and $\delta -6.47$ ppm due to $\text{Mo-O}_2\text{CH}$ proton respectively (Table 15). The complex $[\text{Mo}(\text{O}_2\text{CH})(\text{CO})_2(\text{PEt}_3)_2]$, obtained by the insertion of CO_2 into Mo-H bonds, contains both unidentate and bidentate formate ligands [272]. Reaction of CO_2 with $[\text{WH}_6(\text{PMe}_3)_3]$ results in the formation of complex $[\text{WH}_2(\eta^1\text{-O}_2\text{CH})(\eta^2\text{-O}_2\text{CH})(\text{PMe}_3)_3]$ containing both unidentate and bidentate formato ligands [273].

Carbon dioxide (at 50 atm) reacts with manganese dihydrides $[\text{Mn}_2(\mu\text{-H})_2(\text{CO})_6(\mu\text{-P-P})]$ ($\text{P-P} \equiv \text{dppm}$, tedp) to give hydridoformato complexes $[\text{Mn}_2(\mu\text{-H})(\text{O}_2\text{CH})(\text{CO})_6(\mu\text{-P-P})]$ (CXXIII) along with hydridohydroxy complexes

Table 14

¹³C NMR spectral data of transition metal carbon dioxide complexes

Complex	$\delta(\text{CO}_2)$ (ppm)	Solvent	Reference
$\text{O}=\text{C}=\text{O}$	132.0		
$[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\eta^1\text{-CO}_2)(\text{PMe}_3)]$	212.3(d)	CDCl_3	[123]
$[\text{Mo}(\eta^2\text{-CO}_2)(\text{PMe}_3)_4]$	206.1(q)	C_6D_6	[104]
$[\text{Mo}(\eta^2\text{-CO}_2)(\text{CNMe})(\text{PMe}_3)_3]$	201.1(m)	C_6D_6	[104]
$[\text{Mo}(\eta^2\text{-CO}_2)(\text{CN-i-Pr})(\text{PMe}_3)_3]$	201.4(m)	C_6D_6	[104]
$[\text{Mo}(\eta^2\text{-CO}_2)(\text{CN-t-Bu})(\text{PMe}_3)_3]$	201.2(td)	C_6D_6	[104]
$[\text{Mo}(\eta^2\text{-CO}_2)(\text{CNCy})(\text{PMe}_3)_3]$	201.4(td)	C_6D_6	[104]
$[\text{Mo}(\eta^2\text{-CO}_2)(\text{CNCH}_2\text{Ph})(\text{PMe}_3)_3]$	201.1(m)	C_6D_6	[104]
$[(\eta^5\text{-C}_5\text{H}_5)(\text{CO})(\text{NO})\text{Re}(\mu\text{-CO}_2)\text{ZrCl}(\eta^5\text{-C}_5\text{H}_5)_2]$	212.6	CdCl_2	[256]
$[(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Fe}(\mu\text{-CO}_2)\text{W}(\eta^5\text{-C}_5\text{H}_5)_2]^+$	260.0		[254]
$[(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Fe}(\mu\text{-CO}_2)\text{ZrCl}(\eta^5\text{-C}_5\text{H}_5)_2]$	206.2	C_6D_6	[256]
$[(\eta^5\text{-C}_5\text{Me}_5)(\text{CO})_2\text{Ru}(\mu\text{-CO}_2)\text{W}(\eta^5\text{-C}_5\text{H}_5)_2]^+$	252.0		[254]
$[(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Ru}(\mu\text{-CO}_2)\text{ZrCl}(\eta^5\text{-C}_5\text{H}_5)_2]$	213.2	C_6D_6	[256]
$[(\text{COD})_2\text{Rh}_2\text{OsH}_2(\mu\text{-CO}_2)(\text{PMe}_2\text{Ph})_3]$	193.0(d)	C_6D_6	[265]
<i>trans</i> -[Pt(Ph)(PEt ₃) ₂] ₂ ($\mu\text{-CO}_2$)	201.0(m)	CD_2Cl_2	[259]










Table 15

Spectral data of formate complexes

Complex	IR frequencies (cm ⁻¹)		¹ H NMR (ppm), $\delta(\text{O}_2\text{CH})$	Reference
	$\nu_{\text{as}}(\text{CO}_2)$	$\nu_{\text{s}}(\text{CO}_2)$		
[MoH(η^2 -O ₂ CH)(dppe) ₂]	1550	1360	8.83	[271]
[Mn ₂ (μ -H)(O ₂ CH)(CO) ₆ (μ -dppm)]	1570	1440		[274]
[Re(CO)(η^1 -O ₂ CH)(PMe ₃) ₄]	1634	1318	8.66	[86]
[Ru(NO)(O ₂ CH)(PPh ₃) ₂]	1610	1390		[112]
[(η^5 -C ₅ H ₅)Ru(O ₂ CH)(PPh ₃) ₂]	1640	1360		[112]
[NiH(O ₂ CH)(PCy ₃) ₂]	1619	1310	8.90	[276]
[Cu(η^2 -O ₂ CH)(PPh ₃) ₂]	1585	1350	8.55	[110]
[Cu(η^1 -O ₂ CH)(PPh ₃) ₃]	1610	1340	9.00	[110]
[Cu(η^1 -O ₂ CH)(PMePh ₂) ₃]	1620	1330	8.70	[110]
[Cu(η^2 -O ₂ CH)(PCy ₃) ₂]	1600	1330	8.60	[110]
[{(PPh ₃) ₃ Cu} ₂ (μ -O ₂ CH)][ClO ₄]	1585	1360	8.55	[110]
[(triphos)Cu(η^1 -O ₂ CH)]	1620	1320	—	[110]
[(np ₃)Cu(η^1 -O ₂ CH)]	1610	1330	8.70	[110]
[Cu(η^1 -O ₂ CH)(PPh ₃) ₂ (AsPh ₃)]	1610	1370		[112]
[Cu(η^1 -O ₂ CH)(PPh ₃) ₂ (SbPh ₃)]	1605	1370		[112]



(CXXIII)



(CXXIV)

[274]. Rhenium hydrido complexes reacts with CO₂ to give [Re(η^1 -OC(O)H] complexes [86,275]. Reactions of [ReH(PMe₃)₅] with CO₂ at 70 °C affords white crystals of *trans*-[Re(CO)(O₂CH)(PMe₃)₄] (CXXIV) containing a monodentate formate ligand [86].

Reaction of CO₂ with [RuH(NO)(PPh₃)₃] and [(η^5 -C₅H₅)RuH(PPh₃)₂] results in the formation of formato complexes [Ru(NO)(O₂CH)(PPh₃)₂] and [(η^5 -C₅H₅)Ru(O₂CH)(PPh₃)₂] respectively [112]. The synthesis and molecular structure of the complex [{CoL}₂(O₂CH)][ClO₄]₃, containing LCo—C(OH)O—CoL moiety (L $\equiv$ 5,7,7,12,14,14-hexamethyl-1,4,4,11-tetraazacyclotetradeca-4,11-diene) has been reported [239].

Carbon dioxide reacts with [NiH(CH₃)(PCy₃)₂] in benzene at -50 °C to give the orange brown complex [NiH(O₂CH)(PCy₃)₂] [276]. The platinum analog of

this complex $\text{trans-[PtH(O}_2\text{CH)(PCy}_3)_2]$ has been characterized by X-ray crystallography [277]. Complex $[\text{Ni(Ph)(O}_2\text{CH)(PCy}_3)_2]$ is obtained from the reaction of $[\text{NiH(Ph)(PCy}_3)_2]$ with CO_2 .

The formation of unidentate and bidentate formate ligands by the insertion of CO_2 into the Cu-H bond has been reported [108–110,112,278]. Reaction of CO_2 with $[\text{Cu}(\eta^2\text{-BH}_4)(\text{PPh}_3)_2]$ gives different insertion products under different conditions. Complex $[\text{Cu}(\eta^2\text{-O}_2\text{CH)(PPh}_3)_2]$ is formed when CO_2 is bubbled into a solution of $[\text{Cu}(\eta^2\text{-BH}_4)(\text{PPh}_3)_2]$ for 24 h. Passing CO_2 into a solution of $[\text{Cu}(\eta^2\text{-BM}_4)(\text{PPh}_3)_2]$ in the presence of one equivalent PPh_3 for 12 h gives complex $[\text{Cu}(\eta^1\text{-O}_2\text{CH)(PPh}_3)_3]$. In the presence of PPh_3 and NBu_4ClO_4 , $[\text{Cu}(\eta^2\text{-BH}_4)(\text{PPh}_3)_2]$ reacts with CO_2 to give $[(\text{Ph}_3\text{P})\text{Cu}(\mu\text{-O}_2\text{CH})\text{Cu}(\text{PPh}_3)_3][\text{ClO}_4]$ (CXXV) [110].

The molecular structure of $[\text{Cu}(\eta^2\text{-O}_2\text{CH)(PPh}_3)_2]$ has been determined by X-ray crystallography [110]. Reaction of CO_2 with $[\text{Cu}(\eta^2\text{-BH}_4)(\text{PCy}_3)_2]$ results in the formation of $[\text{Cu}(\eta^2\text{-O}_2\text{CH)(PCy}_3)_2]$. The reaction of CO_2 with copper complexes $[\text{LCu}(\text{BH}_4)]$ containing tridentate phosphine ligands L ($\text{L} \equiv \text{triphos, np}_3, \text{etp}_3$) and monodentate borohydrate ($\eta^1\text{-BH}_4$) ligand, give $\eta^1\text{-O}_2\text{CH}$ bonded formate complexes $[\text{LCu}(\text{O}_2\text{CH})]$. The coordination geometry about copper in $[(\text{triphos})\text{Cu}(\eta^1\text{-O}_2\text{CH})]$ (CXXVI) is distorted tetrahedral with three phosphorus atoms of the triphos ligand and an oxygen atom of the formate group [110].


4.4.2. Insertion into the M-C bond




The normal insertion of CO_2 into metal–carbon bonds yields carboxylato complexes while the abnormal insertion gives l-metalated formate ester.

Insertion reactions of carbon dioxide into a metal–carbon bond are summarized in Table 16 [279–298]. Reaction of nickelacyclobutane with CO_2 at room temperature affords the nickel carboxylate complex (CXXVII) [299]. Insertion of CO_2 into a Pd-allyl bond has been investigated by Behr and co-workers [300]. The insertion products react with HCl to give organic compounds.

4.4.3. Insertion into the M-N bond

Only a few insertion reactions of CO_2 into the M-N bond have been reported [301–304]. The interaction of CO_2 with a secondary amine in the presence of a transition metal system results in the formation of carbamate complexes [303].






The reaction of CO_2 with RZnNEt_2 ($\text{R} \equiv \text{Me, Et}$) yields $[\text{R}_2\text{Zn}_4(\text{O}_2\text{CNEt}_2)_6]$ by the insertion of CO_2 into the $\text{Zn}-\text{N}$ bond. The X-ray structure for $[\text{Me}_2\text{Zn}_4(\text{O}_2\text{CNEt}_2)_6]$ has been reported [304].

4.4.4. Insertion into the $\text{M}-\text{O}$ bond


The insertion of CO_2 into an $\text{M}-\text{OR}$ bond yields carbonate complexes.



Carbon dioxide insertion into the $\text{W}-\text{OR}$ bond of complexes $[\text{Et}_4\text{N}][\text{W}(\text{CO})_5(\text{OR})]$ ($\text{R} \equiv \text{Ph, C}_6\text{H}_4\text{Me-m}$) produces arylcarbonato complexes



Table 16
Insertion reaction of CO₂ into the M–C bond

Precursor complex	Isolated product	Reference
$[(\eta^5\text{-C}_5\text{H}_5)_2\text{TiMe}_2]$	$[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\text{O}_2\text{CMe})\text{Me}]$	[279]
$[(\eta^5\text{-C}_5\text{H}_5)\text{V}(\eta^1\text{-C}_3\text{H}_5)]$	$[(\eta^5\text{-C}_5\text{H}_5)\text{V}(\eta^1\text{-O}_2\text{CC}_3\text{H}_5)]$	[280]
$[\text{RM}(\text{CO})_5]^-$ (M = Cr, W; R = Me, Et)	$[(\text{RCO}_2)\text{M}(\text{CO})_5]^-$	[281–286]
$[\text{MeW}(\text{CO})_4\text{L}]^-$ (L = PMe ₃ , P(OMe) ₃)	$[(\text{MeCO}_2)\text{W}(\text{CO})_4\text{L}]^-$	[281–286]
$[\text{W}(\text{Me})(\text{CO})_2(\text{NO})\text{L}_2]$ (L = PPh ₃ , P(O-i-Pr) ₃)	$[\text{W}(\eta^2\text{-O}_2\text{CMe})(\text{CO})_2(\text{NO})\text{L}]$	[287,288]
$[\text{W}(\text{CO})_4(\eta^2\text{-CH}_2\text{CH}_2\text{PPh}_2)]^-$	$[\text{W}(\text{CO})_4\{\eta^2\text{-O}_2\text{C}(\text{O})\text{CH}_2\text{CH}_2\text{PPh}_2\}]^-$	[289]
$[\text{WH}(\text{Me}_2\text{PCH}_2)(\text{PMe}_3)_4]$	$[\text{W}(\eta^1\text{-Me}_2\text{PCH}_2)(\text{PMe}_3)_3]_2(\text{C}_3\text{H}_2\text{O}_6)$	[290]
$[\text{Mn}(\text{Ph})_2(\text{PCy}_3)_2]$	$[\text{Mn}(\eta^2\text{-O}_2\text{CPh})_2(\text{PCy}_3)_2]$	[291]
$[\text{Mn}(\text{CO})_4(\eta^2\text{-CH}_2\text{CH}_2\text{PPh}_2)]$	$[\text{Mn}(\text{CO})_4\{\eta^2\text{-O}_2\text{C}(\text{O})\text{CH}_2\text{CH}_2\text{PPh}_2\}]$	[292]
$[(\text{TDP})\text{Fe-Bu}]$	$[(\text{TPP})\text{Fe-OC}(\text{O})(\text{CH}_2)_3\text{CH}_3]$	[293]
$[\text{Rh}(\text{Ph})(\text{PPh}_3)_3]$	$[\text{Rh}\{\eta^1\text{-OC}(\text{O})\text{Ph}\}(\text{PPh}_3)_3]$	[286]
$[\text{Rh}(\text{Ph})(\text{PMe}_3)_3]$	$[\text{Rh}(\eta^2\text{-O}_2\text{CPh})(\text{PMe}_3)_2]$	[286]
$[\text{Rh}(\text{Me})(\text{PMe}_3)_3]$	$[\text{Rh}(\eta^2\text{-O}_2\text{CMe})(\text{PMe}_3)_2]$	[286]
$[\text{Rh}(\text{Ar})(\text{L})]$ (Ar = 4-MeC ₆ H ₄ , L = t-BuP(CH ₂ CH ₂ CH ₂ PPh ₂) ₂)	$[\text{Rh}\{\text{OC}(\text{O})\text{Ar}\}(\text{L})] + \dots$	[294]
$[\text{Ni}(\text{Et})_2(\text{bipy})]$	$[\text{Ni}(\text{Et})(\text{O}_2\text{CEt})(\text{bipy})] + [\text{Ni}(\text{O}_2\text{CEt})_2(\text{bipy})]$	[295]
$[\text{Ni}(\eta^2\text{-o-C}_6\text{H}_4\text{CH}_2)(\text{P-}n\text{-Bu}_3)_2]$	$[\text{Ni}\{\eta^2\text{-o-CH}_2\text{C}_6\text{H}_4\text{C}(\text{O})\text{O}\}(\text{P-}n\text{-Bu}_3)_2]$	[296]
$[\text{Ni}(\eta^2\text{-CH}_2\text{CMe}_2\text{-o-C}_6\text{H}_4)(\text{PMe}_3)_2]$	$[\text{Ni}\{\eta^2\text{-CH}_2\text{CMe}_2\text{-o-C}_6\text{H}_4\text{C}(\text{O})\text{O}\}(\text{PMe}_3)_2]$	[297]
$[(\text{phen})(\text{PPh}_3)\text{Cu-CH}(\text{CN})_2]$	$[(\text{Phen})(\text{PPh}_3)\text{Cu-OC}(\text{O})\text{CH}(\text{CN})_2]$	[298]
$[(\text{phen})(\text{PPh}_3)\text{Cu-CH}(\text{CN})(\text{COOEt})]$	$[(\text{Phen})(\text{PPh}_3)\text{Cu-OC}(\text{O})\text{CH}(\text{CN})(\text{COOEt})]$	[298]



$[\text{W}(\text{CO})_5\{\text{OC}(\text{O})\text{OR}\}]^-$. A THF solution of arylcarbonato complexes reacts with water to give the carbonato complex $[\text{Et}_4\text{N}]_2[\text{W}(\text{CO})_4(\text{CO}_3)] \cdot \text{H}_2\text{O}$ (CXXVIII) [305].


The formation of μ -carbonato-di- μ -hydroxybis(1,5-diamine-3-aza-pentane)cobalt (III) perchlorate from the interaction of CO₂ with $[(\text{dien})\text{Co}(\mu\text{-OH})_3\text{Co}(\text{dien})]^{3+}$ has been proposed [306]. The dioxygen complex $[\text{Rh}(\text{O}_2)(\text{S}_2\text{CNMe}_2)(\text{PPh}_3)_2]$ reacts with CO₂ to afford peroxycarbonato complex $[\text{Rh}(\text{CO}_4)(\text{S}_2\text{CNR}_2)(\text{PPh}_3)_2]$ which reacts with PPh₃ to give the yellow carbonato complex $[\text{Rh}(\text{CO}_3)(\text{S}_2\text{CNR}_2)(\text{PPh}_3)_2]$ (CXXIX) [307]. Reaction of CO₂ with $[(\eta^5\text{-C}_5\text{Me}_5)\text{IrH}(\text{OEt})(\text{PPh}_3)]$







leads to insertion of CO_2 into the $\text{Ir}-\text{OEt}$ bond and no insertion into the $\text{M}-\text{H}$ bond is observed. The insertion product is $[(\eta^5-\text{C}_5\text{Me}_5)(\text{PPh}_3)(\text{H})\text{Ir}-\text{O}-\text{C}(\text{O})-\text{OEt}]$ [308].

Insertion of CO_2 into a $\text{Pd}-\text{O}$ bond has been observed in the reaction of CO_2 with hydroxy bridged palladium complexes. Reactions of $[\text{Pd}(\mu\text{-OH})\text{L}_2][\text{BF}_4]_2$ ($\text{L} \equiv \text{PPh}_3$, PMePh_2 , $1/2 \text{ dppe}$, $1/2 \text{ dcpe}$) with CO_2 result in the formation of bis(bicarbonato) complexes $[\text{Pd}\{\text{OC}(\text{O})\text{OH}\}_2\text{L}_2]$. The structure of $[\text{Pd}\{\text{OC}(\text{O})\text{OH}\}_2(\text{dppe})]$ (CXXX) was determined by X-ray diffraction analysis. The geometry about the palladium atom is square planar [309]. Reaction of CO_2 with $[\text{Pt}(\text{OH})(\text{Ph})\text{L}_2]$ ($\text{L} \equiv \text{P}(\text{CH}_2\text{Ph})_3$) affords a dimeric carbonato complex (CXXXI) [310].

The $(\mu\text{-oxo})$ dichlorobis(1,10-phenanthroline)copper(II) reacts with CO_2 to afford the carbonate complex $[(\text{phen})\text{ClCu}(\mu,\eta^3\text{-CO}_3)\text{CuCl}(\text{phen})]$ [311]. The IR spectrum of this complex shows absorption bands at 1530, 1360, 820 and 750 cm^{-1} characteristic of the bridging tridentate carbonate ligand [312]. Photolysis of the μ -carbonato complex produces a μ -oxo complex by loss of CO_2 . Alkyl carbonato complexes of $\text{Zn}(\text{II})$, (CXXXII) and (CXXXIII) are obtained by the reaction of CO_2 with zinc(II)-tetraazacycloalkane complexes in the presence of base (ROH) in alcohol [313,314].





4.4.5. Insertion into the M–Si bond

Attempts to observe insertion of CO_2 into an M–Si bond were unsuccessful: complexes $[(\eta^5\text{-C}_5\text{Me}_5)\text{TaCl}_3(\text{SiMe}_3)]$, $[(\eta^5\text{-C}_5\text{H}_5)_2\text{ZrCl}(\text{SiMe}_3)]$ and $[\text{W}(\text{CO})_5(\text{SiMe}_3)]$ do not react with CO_2 [315–317]. Reaction of excess CO_2 with zirconium-silyl complex $[(\eta^5\text{-C}_5\text{Me}_5)(\eta^5\text{-C}_5\text{H}_5)\text{ZrCl}\{\text{Si}(\text{SiMe}_3)_3\}]$ gives an unidentified mixture of products [318].

Tilley and co-workers have successfully isolated the insertion product, for the first time, from the reaction of CO_2 with scandium-silyl complexes. $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Sc}(\text{SiR}_3)(\text{THF})]$ ($\text{R}_3 \equiv (\text{SiMe}_3)_3$, $t\text{-BuPh}_2$) react with CO_2 to afford

$[(\eta^5\text{-C}_5\text{H}_5)_2\text{Sc}(\mu\text{-O}_2\text{CSiR}_3)_2\text{Sc}(\eta^5\text{-C}_5\text{H}_5)_2]$ (CXXXIV). The structure of (CXXXIV) has been established by X-ray crystallography [319].

4.4.6. Insertion into the $M\text{—}P$ bond



Carbon dioxide undergoes insertion into metal— PR_2 bonds to give phosphinocarboxylate O_2CPR_2 ligands [320–322].






Carbon dioxide reacts rapidly with phosphino-amido complexes $[\text{M}(\text{NMe}_2)_4(\text{PR}_2)_2]$ to afford complexes $[\text{M}_2(\text{O}_2\text{CNMe}_2)_2(\text{NMe}_2)_2(\text{O}_2\text{CPR}_2)_2]$ ($\text{M} \equiv \text{Mo}$, $\text{R} \equiv \text{t-Bu}$ (CXXXV); $\text{M} \equiv \text{W}$, $\text{R} \equiv \text{t-Bu}$ (CXXXVI), Cy). The ligands O_2CNMe_2 and O_2CPR_2 are bound to metal centers in an η^2 fashion. In benzene solution, complex (CXXXV) decomposes to give $\text{Mo}_2[\text{O}_2\text{CP}(\text{t-Bu})_2]_4 \cdot 2\text{C}_6\text{H}_6$ (CXXXVII). The molecular structures of (CXXXVI) and (CXXXVII) have been determined by X-ray analysis [322].

The tungsten complex (CXXXVI), unlike its molybdenum analog, is inert with respect to decomposition in organic solvents. The coordination at each tungsten in complex (CXXXVI) is a distorted square planar WO_3 moiety. The $\text{W}\text{—}\text{W}$ distance of $2.290(1) \text{ \AA}$ is consistent with other $d^3\text{—}d^3$ ($\text{W} \equiv \text{W}$) $^{6+}$ units [322].

4.4.7. Oxidative coupling of carbon dioxide with coordinated organic substrate

Carbon dioxide undergoes coupling reactions with unsaturated hydrocarbons (alkenes, alkynes, dienes, conjugated dienes and cumulenes), imines and aldehydes on the transition metal centers to produce metallocycles (CXXXVIII–CXLIH).





Two main requirements, which play an important role in the coupling reactions of CO₂ and unsaturated hydrocarbons, are (a) high electron density at the metal center and (b) the unsaturated coordination of the transition metal [323–345].


The reaction of CO₂ with $[(\eta^5\text{-C}_5\text{Me}_5)_2\text{Ti}(\eta^2\text{-C}_2\text{H}_4)]$ in toluene at low temperature (-78°C) gives complex $[(\eta^5\text{-C}_5\text{Me}_5)_2\text{Ti}\{\eta^2\text{-CH}_2\text{CH}_2\text{C}(\text{O})\text{O}\}]$ [323]. In the reaction of CO₂ with allyltitanium complexes, the formation of complexes $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\eta^2\text{-O}_2\text{C}_4\text{H}_7)]$ has been suggested [324]. The coupling reaction between alkynytitanium complex $[(\eta^5\text{-C}_5\text{Me}_5)(\eta^5\text{-C}_5\text{H}_5)\text{Ti}(\text{PhC}\equiv\text{CPh})]$ and CO₂ to produce a five-membered metallacyclic complex $[(\eta^5\text{-C}_5\text{Me}_5)(\eta^5\text{-C}_5\text{H}_5)\text{Ti}\{\eta^2\text{-PhC}\equiv\text{CPhC}(\text{O})\text{O}\}]$ has been reported by Demerseman and co-workers [325]. Complex $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\text{HC}\equiv\text{CH})(\text{PMe}_3)]$ reacts with CO₂ to give $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}\{\eta^2\text{-CH}=\text{CHC}(\text{O})\text{O}\}]$ and PMe_3 [326]. Zirconium complexes $[(\eta^5\text{-C}_5\text{Me}_5)_2\text{Zr}(\eta^2, \eta^2\text{-C}_2\text{H}_6)]$ and $[(\eta^5\text{-C}_5\text{Me}_5)_2\text{Zr}(\text{CH}_2\text{CH}=\text{C}(\text{Me})\text{CH}_2)]$ react with CO₂ to give complexes $[(\eta^5\text{-C}_5\text{Me}_5)_2\text{Zr}\{\eta^3\text{-C}_3\text{H}_4\text{CH}_2\text{C}(\text{O})\text{O}\}]$ and $[(\eta^5\text{-C}_5\text{Me}_5)_2\text{Zr}\{\eta^2\text{-OC}(\text{O})\text{CH}_2\text{CH}=\text{CMeCH}_2\text{C}(\text{O})\text{O}\}]$ [327].

The vanadium complex $[\text{V}\{\eta^2\text{-t-BuNC}(\text{mes})_3\}]$ ($\text{mes} \equiv 2,4,6\text{-Me}_3\text{C}_6\text{H}_2$) reacts with CO₂ to give coupling product $[\text{V}\{\eta^2\text{-t-BuNC}(\text{mes})\}\{\eta^2\text{-t-BuNC}(\text{mes})\text{C}(\text{O})\text{O}\}_2]$ by insertion of CO₂ in two V–C bonds [328]. Coupling of CO₂ to the activated benzene in chromium complex $[\text{Cr}(\eta^4\text{-C}_6\text{H}_6)(\text{CO})_3]^{2-}$ yields an intermediate cyclohexanycarboxylate complex $[\text{Cr}(\eta^5\text{-C}_6\text{H}_6\text{CO})(\text{CO})_3]^{2-}$ which is oxidized and silylated to form $[\text{Cr}(\eta^6\text{-C}_6\text{H}_5\text{CO}_2\text{SiMe}_3)(\text{CO})_3]$ [329]. Reaction of CO₂, at room temperature and normal pressure, with molybdenum and tungsten complexes *trans*- $[\text{M}(\text{C}_2\text{H}_4)_2(\text{PMe}_3)_4]$ ($\text{M} \equiv \text{Mo}, \text{W}$) results in the formation of acrylate complexes $[\text{M}\{\mu\text{-CH}_2\text{C}=\text{CHC}(\text{O})\text{OH}\}(\text{C}_2\text{H}_4)(\text{PMe}_3)_2]$ (CXLIV) [330].


Complex $[\text{Mn}(\text{CO})_3\{\eta^3\text{-C}_3\text{H}_4\text{CH}_2\text{C}(\text{O})\text{O}\}]^-$ can be obtained from the reaction of $[\text{Mn}(\text{CO})_3(\eta^2, \eta^2\text{-C}_4\text{H}_6)]$ with CO₂ by the coupling of coordinated butadiene with CO₂ [331].

The coupling reaction of complex $[\text{Fe}(\text{C}_2\text{H}_4)_2(\text{PEt}_3)_2]$ with CO₂ yields $[\text{Fe}\{\eta^2\text{-CH}_2\text{CH}_2\text{C}(\text{O})\text{O}\}(\text{PEt}_3)_3]$ [332]. The iron complex $[\text{Fe}(\eta^2, \eta^2\text{-C}_4\text{H}_6)(\text{PMe}_3)_3]$ reacts with CO₂ to give the allylcarboxylato complex $[\text{Fe}\{\eta^3\text{-C}_3\text{H}_4\text{CH}_2\text{-C}(\text{O})\text{O}\}(\text{PMe}_3)_3]$ [333].

Carbon dioxide reacts with *mer*- $[\text{RhH}(\text{CH}_2\text{COMe})(\text{PMe}_3)_3\text{Cl}]$ to give



(CXLIV)



(CXLV)

$[\text{Rh}(\text{CH}_2\text{CHMeOCO}_2)(\text{PMe}_3)_3\text{Cl}]$ (CXLV) [334]. The formation of coupling product is reversible and the starting complex is regenerated by its dissolution in various solvents at 20 °C. Behr and co-workers have reported the coupling reaction between malonodinitrile and CO_2 on rhodium [335] and iridium [336] centers.

The coupling of CO_2 with alkynes [336–339], alkenes [340–345], conjugated dienes [346–353] or cumulenes [354] on nickel(0) centers, in the presence of chelating ligands such as 2,2'-bipyridyl TMED, 1,2-bis(dicyclohexylphosphino)ethane, diazabicyclo[5,4,0]undec-7-ene, have been reported. The coupling product $[(\text{dbu})_2\text{Ni}\{\eta^2\text{-CH}_2\text{CH}_2\text{C}(\text{O})\text{O}\}]$ (CXLVI), formed from ethylene and CO_2 at $\text{Ni}(\text{COD})_2$ in the presence of dbu, has been isolated and structurally characterized by Hoberg and co-workers [345].


4.4.8. Reduction of carbon dioxide

Recently, there has been a considerable upsurge in the study of the reduction of CO_2 catalyzed by transition metal complexes using different methods: (a) chemical reduction, (b) electrochemical reduction and (c) photochemical catalytic reduction. The reduction processes which have been observed are as follows: (a) reduction of CO_2 to formic acid or formate ion, (b) reductive disproportionation of CO_2 to CO and CO_3^{2-} , (c) reduction of CO_2 to oxalate, and (d) reduction of CO_2 to CO.

Transition metal complexes, which are powerful two-electron reductants, are promising substrates for reductive disproportionation of CO_2 .

4.4.8.1. Chemical reduction. The binuclear oxalate complex $[\{(\eta^5\text{-C}_5\text{H}_5)\text{Ti}\}_2(\mu\text{-C}_2\text{O}_4)]$ is prepared via a reductive dimerization of CO_2 [355]. Reaction of CO_2 with $\text{Na}_2[(\eta^5\text{-C}_5\text{H}_5)\text{V}(\text{CO})_3]$ results in reductive disproportionation of CO_2 and in the formation of $[(\eta^5\text{-C}_5\text{H}_5)\text{V}(\text{CO})_4]$ and Na_2CO_3 [356]. Reactions of complex $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Nb}(\eta^2\text{-CO}_2)(\text{CH}_2\text{SiMe}_3)]$ with several Lewis acids have been performed [357]. $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Nb}(\eta^2\text{-CO}_2)(\text{CH}_2\text{SiMe}_3)]$ reacts with ZnCl_2 to form an unstable carbonyl complex $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Nb}(\text{CO})_2(\text{CH}_2\text{SiMe}_3) \cdot \text{ZnCl}_2]$ which decomposes with CO loss to produce $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Nb}(\text{CH}_2\text{SiMe}_3)\text{O} \cdot \text{ZnCl}_2]_2$.


Reactions of CO_2 with $\text{A}_2[\text{M}(\text{CO})_5]$ ($\text{A} \equiv \text{K}$, $\text{M} \equiv \text{Cr}$, Mo ; $\text{A} \equiv \text{Li}$, Na , K , $\text{M} \equiv \text{W}$) give hexacarbonyls $[\text{M}(\text{CO})_6]$ and alkali metal carbonates by the disproportionation of CO_2 to CO and CO_3^{2-} . Cooper and co-workers suggested that the reaction



(CXLVI)



involves discrete CO_2 adducts (CXLVII) and (CXLVIII) containing $\text{M}-\eta^1\text{-CO}_2$ [356].


Nicholas and co-workers have reported the reactions of $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Mo}(\eta^2\text{-CO}_2)]$ with several electrophilic reagents and with transition metal hydrides which form carbonyl complexes via transfer of oxygen [358]. Reaction of $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Mo}(\eta^2\text{-CO}_2)]$ with R_3SiCl (or BCl_3 or HCl) gives $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Mo}(\text{CO})\text{Cl}]\text{Cl}$ and with $\text{Me}_3\text{SiOSO}_2\text{CF}_3$ in THF yields complex $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Mo}(\text{CO})(\text{THF})(\text{OSO}_2\text{CF}_3)]$. RCOCl reacts with $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Mo}(\eta^2\text{-CO}_2)]$ to afford $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Mo}(\text{CO})(\text{OC}(\text{O})\text{R})]\text{Cl}$. Reactions of $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Mo}(\eta^2\text{-CO}_2)]$ with $[\text{HCO}(\text{CO})_4]$ and $[\text{H}_2\text{Fe}(\text{CO})_4]$ produce cationic carbonyl complexes [359].




The complex $[(\eta^5\text{-C}_5\text{H}_5)_2\text{MoH}(\text{CO})][\text{CO}(\text{CO})_4]$ slowly converts to $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Mo}(\mu\text{-H})(\mu\text{-CO})\text{CO}(\text{CO})_3]$ (CXLIX) [359].

Reaction of CO_2 with *cis*- $[\text{Mo}(\text{N}_2)_2(\text{PMe}_3)_4]$ is very sensitive to the reaction conditions, in particular to the nature of the solvent used. If the reaction is performed in weakly coordinating solvents (Et_2O , THF, Me_2CO), the main products are either the red dimeric complex (CL) or the dark blue monomeric complex (CLI), obtained from the metal-induced reductive disproportionation of CO_2 to CO_3^{2-} and CO [360].






(CXLIX)




(CL)



(CLI)


Carbon dioxide reacts with the tungsten complex $[\text{WCl}_2(\text{PMePh}_2)_2]$ to produce the carbonyl complex $[\text{W}(\text{O})(\text{CO})\text{Cl}_2(\text{PMePh}_2)_2]$ [361]. The carbonato complex $[(\eta^5\text{-C}_5\text{H}_5)_2\text{W}(\eta^2\text{-CO}_3)]$ (CLII) can be obtained from the reaction of $[(\eta^5\text{-C}_5\text{H}_5)_2\text{WH}(\text{Ph})]$ with CO_2 in acetone in the presence of water [362].

Reactions of CO_2 with $\text{Na}_2[\text{M}'(\text{CO})_4]$ ($\text{M}' \equiv \text{Fe, Ru, Os}$) result in reductive disproportionation of CO_2 and in the formation of carbonyl metalates $[\text{M}'(\text{CO})_5]$ and Na_2CO_3 [356]. The reductive hydrogenation of CO_2 has also been examined in the presence of cobalt complexes [238–240,363] and ruthenium clusters [364,365].




(CLII)

The reaction of CO_2 with transition metal hydroxides results in the formation of formate complexes [366–368]. The complex $[\text{H}_2\text{Rh}\{\text{P}(\text{t-Bu})_2\text{CH}_2\text{C}_6\text{H}_3\text{CH}_2(\text{t-Bu})_2\}]$ reacts with CO_2 to produce a rhodium carbonyl complex by the reduction of CO_2 to CO and H_2O [369].



The iridium complex $[\text{Ir}_2(\mu\text{-CNR})_2(\text{CNR})_2(\text{dmpm})_2]$ ($\text{R} \equiv 2,6\text{-Me}_2\text{C}_6\text{H}_3$) reacts with CO_2 to give the CO_2 adduct $[\text{Ir}_2(\mu\text{-CN}(\text{CO}_2)\text{R})_2(\text{CNR})_2(\text{dmpm})_2]$ which loses two equivalent of CO_2 on heating. In dichloromethane, the CO_2 adduct decomposes to give bridged carbonyl complex $[\text{Ir}_2(\mu\text{-CO})(\mu\text{-H})\{\text{C}(\text{O})\text{NHR}\}_2(\text{CNR})_2(\text{dmpm})_2]$ by the reduction of CO_2 to CO [370]. Kubiak and co-workers have reported the mechanism of oxygen atom transfer from CO_2 to the methyl isocyanide ligand of a binuclear $\text{Ni}(0)$ complex [371].

The reaction of RSH ($\text{R} \equiv \text{H}$, alkyl, benzyl, phenyl and substituted phenyl) with the CO_2 complex $[\text{Ni}(\eta^2\text{-CO}_2)(\text{PCy}_3)_2]$ leads to the formation of a nickel carbonyl complex.



Reduction of CO_2 by $[\text{NiCl}(\text{PCy}_3)_2]$ and PhSH gives small amounts of the dicarbonyl complex $[\text{Ni}(\text{CO})_2(\text{PCy}_3)_2]$. All the $\text{Ni}(\text{II})$ complexes $[\text{NiCl}_2\text{L}_2]$ (L = monodentate phosphines, L_2 = bidentate phosphines) are inactive in the CO_2 reduction by thiols [372]. The formation of a nickel carbonate complex $[\text{Ni}_3(\text{N,N-Me}_2\text{en})_6(\text{CO}_3)(\text{H}_2\text{O})_4][\text{ClO}_4]_4$ arises from the reaction of CO_2 with the nickel(II) complex of N,N -dimethylethylenediamine, $[\text{Ni}(\text{N,N-Me}_2\text{en})][\text{ClO}_4]_2$ in aqueous ethanol [373]. Reaction of CO_2 with $[\text{Pt}(\text{C}_2\text{H}_4)(\text{PPh}_3)_2]$ results in the formation of the carbonate complex $[\text{Pt}(\eta^2\text{-O}_2\text{CO})(\text{PPh}_3)_2]$ and carbon monoxide [210].

Reductive hydrogenation of CO_2 gives formic acid in the presence of moisture or other proton source. The transition metal complex activates a water molecule with

the formation of a hydride complex which transforms the hydrogen to the CO₂ molecule coordinated to the metal. The hydrogenation of CO₂ to formic acid is catalyzed by complexes [RuH₂(PPh₃)₄] [374], [RuCl₃(EDTA)] [374] and [Rh(NBD)(PMe₂Ph)₃][BF₄] [375]. The hydrogenation of CO₂ catalyzed by soluble metal catalysts with added alcohols [376,377], alkyl halides [378] or secondary amines [379] to give formate esters or formamides is enhanced by the formation of H₂O or HX.

The reaction of H₂ with [Rh(NBD)(PMe₂Ph)₃][BF₄] gives rhodium dihydride complexes [Rh(H)₂(PMe₂Ph)₃(S)] (S ≡ H₂O, THF) which are active catalysts for the hydrogenation of CO₂ to formic acid. The formation of the monodentate formate complex [RhH(η¹-OC(O)H)(H₂O)(PMe₂Ph)₃]⁺ and bidentate formate complex [RhH(η²-O₂CH)(H₂O)(PMe₂Ph)₂]⁺ as unstable intermediate has been observed under catalytic conditions [375].

4.4.8.2. Electrocatalytic reaction. The electrocatalytic reaction of CO₂ in the presence of transition metal complexes has been extensively studied in the last decade [51]. The results are summarized in Table 17. In the electrocatalytic reactions, the coordination compound is either deposited onto the electrode surface or dissolved in the electrolyte. The formation of various reduction products depends mainly on the transition metal complex and solvent used.

Transition metal complexes which can reduce electrocatalytically CO₂ to various products are grouped into five categories: (a) transition metal tetraazamacrocyclic complexes [380–382], (b) phthalocyanine and porphyrin complexes [383–392], (c) metal clusters [393–395], (d) transition metal bipyridyl and polybipyridyl complexes [275,396–416], (e) phosphine complexes [417–420].

4.4.8.3. Photocatalytic reduction. A number of studies concerning photoreduction of CO₂ using transition metal complexes as photocatalyst have been reported [395–397,401–403,410,416,421–441]. This approach involves the photochemical generation of electron-rich transition metal species in homogeneous solution and these photogenerated species act as redox catalyst.


The oxo complex [(η⁵-C₅H₄Me)₂Nb(O)(CH₂SiMe₃)] can be isolated from the irradiated (λ = 250 nm) solution of [(η⁵-C₅H₄Me)₂Nb(η²-CO₂)(CH₂SiMe₃)] in THF at –20 °C [429]. Irradiation (at more than 330 nm) of a THF solution of [(η⁵-C₅H₅)₂MoH₂] at –10 °C under 1 atm CO₂ for 30 h yields molybdenum carbonyl complex [(η⁵-C₅H₅)₂Mo(CO)] and carbonate complex (CLIII) [430]. The photolysis of [Mo(η²-CO₂)(PMe₃)₄] in toluene solution at –20 °C results in oxygen transfer from CO₂ to phosphine and gives *cis*-[Mo(CO)₂(PMe₃)₄] and OPMe₃ as the major products along with lesser amounts of *mer*- and *fac*-[Mo(CO)₃(PMe₃)₃] and [Mo(CO)(PMe₃)₅] [431]. The adducts [W(CO)₅(PR₃)][–] (R ≡ Ph, *t*-Bu), obtained by the reaction of phosphine ligands with photogenerated [W(CO)₅][–], are powerful reducing agents. Irradiation of a CO₂ saturated solution containing Na₂[W₂(CO)₁₀] and phosphine ligand gives formate, carbon monoxide, carbonate and bicarbonate [432].

The generation of CO by photoreduction of CO₂ to CO via visible light irradiation



Table 17

Electrocatalytic reduction of carbon dioxide

Catalyst	Solvent	Products	Reference
$[\text{CoL}_3]^{2+}$	MeCN–H ₂ O	CO, H ₂	[380]
$[\text{CoL}_2]^{2+}$ (L ≡ 14 membered tetraazamacrocyclic ligands)	DMF–H ₂ O	CO, H ₂	[380]
$[\text{Ni}(\text{cyclam})]^{2+}$ (cyclam ≡ 1,4,8,11-tetraazacyclotetradecane)	H ₂ O	CO	[381,382]
$[\text{Ti}(\text{pc})]$	H ₂ O	CO	[390,391]
$[\text{Fe}(\text{pc})]$	H ₂ O	CO	[390,391]
$[\text{Co}(\text{pc})]$	H ₂ O	HCOO [−]	[383,385]
	H ₂ O	CO, H ₂	[384]
$[\text{Ni}(\text{pc})]$	H ₂ O	HCOO [−]	[383]
$[\text{Cu}(\text{pc})\text{Cl}]$	H ₂ O	CO	[391]
$[\text{Co}(\text{TMPP})]$	H ₂ O	CO, HCOOH	[387]
$[\text{Co}(\text{TPP})]$	CH ₂ Cl ₂	CO	[392]
$[\text{Pd}(\text{TPP})]$	CH ₂ Cl ₂	C ₂ O ₄ ^{2−}	[386]
$[\text{Fe}_4\text{S}_4(\text{SCH}_2\text{Ph})_4]^{2−}$	DMF	HCOO [−]	[393]
$[\text{Fe}_4\text{S}_4(\text{SXN}−)_4]^{2−}$ (X ≡ −COCM ₂ −, −COC ₆ H ₄ CH ₂ −)	DMF	HCOO [−]	[395]
$[\text{Re}(\text{CO})_3\text{Cl}(\text{bipy})]$	DMF–H ₂ O	CO	[396–398]
$[\text{Re}(\text{CO})_3\text{Cl}(\text{vbipy})]$ (vbipy ≡ 4-vinyl-4'-methyl-2,2'-bipyridine)	MeCN	CO, CO ₃ ^{2−}	[399,400]
$[\text{Re}(\text{CO})_3\text{Cl}(\text{bipy-py})]$ (bipy-py ≡ 4[4''-{N-pyrrolyl}n-butyl-4'-methyl-2,2'-bipyridine])	MeCN	CO	[407]
$[\text{Fe}(\text{vtpy})_2]^{2+}$ (vtpy ≡ 4'-vinyl-2,2':6',2''-terpyridine)	MeCN	CO	[411]
$[\text{Ru}(\text{CO})_2(\text{bipy})_2]^{2+}$	DMF	HCOO [−]	[401–403]
<i>cis</i> - $[\text{Ru}(\text{CH}_2\text{Ph})(\text{CO})(\text{bipy})_2]^+$	MeCN	CO	[400]
$[(\eta^6\text{-C}_6\text{H}_6)\text{Ru}(\text{bipy})\text{Cl}]^+$	MeCN	CO, HCOO [−]	[404]
$[\text{Ru}(\text{terpy})(\text{dppe})\text{Cl}]^+$	MeCN	CO, HCOO [−]	[404]
<i>cis</i> - $[\text{OsH}(\text{CO})(\text{bipy})_2]^+$	MeCN	CO	[400]
<i>cis</i> - $[\text{Os}(\text{Me})(\text{CO})(\text{bipy})_2]^+$	MeCN–H ₂ O	CO, HCOO [−]	[400]
<i>cis</i> - $[\text{Os}(\text{Ph})(\text{CO})(\text{bipy})_2]^+$	MeCN	CO	[400]
$[\text{Co}(\text{vtpy})_2]^{2+}$	MeCN	HCOOH	[411]
$[\text{Rh}(\text{COD})(\text{bipy})]^+$	MeCN	CO, HCOO [−]	[404]
<i>cis</i> - $[\text{Rh}(\text{bipy})_2(\text{CF}_3\text{SO}_3)_2]^+$	MeCN	CO, HCOO [−]	[404,408]
<i>cis</i> - $[\text{Ir}(\text{bipy})_2(\text{CF}_3\text{SO}_3)_2]^+$	MeCN	CO, HCOO [−]	[404,408]
$[\text{Ni}(\text{vtpy})_2]^{2+}$	MeCN	HCOOH	[411]
$[\text{Ni}(\text{bipy})_3]^{2+}$	DMF	CO	[415]
$[\text{RhCl}(\text{dppe})]$	MeCN	HCOO [−]	[417]
$[\text{Rh}(\text{CO})\text{Cl}(\text{PPh}_3)_2]$	DMF	CO	[419]
$[\text{Ir}(\text{CO})\text{Cl}(\text{PPh}_3)_2]$	DMF	CO	[419]
$[\text{Ni}(\text{MeCN})_4(\text{PPh}_3)_2]^{2+}$	DMF	CO	[415]
$[\text{Ni}_3(\mu\text{-CNMe})(\mu_3\text{-I})(\text{dppm})_3]^+$	THF	CO, CO ₃ ^{2−}	[420]
$[\text{Pd}(\text{triphos})(\text{PPh}_3)]$	MeCN	CO, H ₂	[418]
$[\text{Pd}(\text{triphos})(\text{PEt}_3)]$	MeCN	CO, H ₂	[418]
$[\text{Pd}(\text{triphos})\{\text{P}(\text{OMe})_3\}]$	MeCN	CO, H ₂	[418]



of solution containing $[\text{Re}(\text{CO})_3\text{X}(\text{bipy})]$ ($\text{X} \equiv \text{Cl}, \text{Br}$) and CO_2 in triethanolamine–dimethylformamide (TEOA–DMF) has been studied [396,398,433–435]. Similar irradiations, conducted in the presence of $[\text{Co}(\text{bipy})]^{2+}$, lead to CO and H_2 . The complex $[\text{Re}(\text{CO})_3\text{X}(\text{bipy})]$ absorbs visible light to yield the excited state $[\text{Re}(\text{CO})_3\text{X}(\text{bipy})^*]$ which may undergo either oxidative or reductive electron transfer quenching, depending on the redox potential of the quencher.



The $[\text{Re}(\text{CO})_3\text{X}(\text{bipy})]^-$ species should reduce CO_2 to CO [396,435]. The formation of CO depends on (a) the nature and concentration of the reducing agent, (b) the time of irradiation, and (c) the presence of additives (Table 18). Owing to labilization of the ligand species, the activity of the system decreases slowly. Addition of chloride or bromide anions to the corresponding $[\text{Re}(\text{CO})_3\text{X}(\text{bipy})]$ complex leads to an increase in stability, producing more CO [396].


Kitamura and Tazuke reported, for the first time, the photochemical reduction of

Table 18


Photochemical reduction of CO_2 to CO via visible light irradiation of solutions containing $[\text{Re}(\text{CO})_3\text{X}(\text{bipy})]$ and CO_2 in TEOA–DMF [396]

Complex	Additive	Irradiation time (h)	Volume of CO produced (ml)
$[\text{Re}(\text{CO})_3\text{Cl}(\text{bipy})]$	—	1	6.5
$[\text{Re}(\text{CO})_3\text{Cl}(\text{bipy})]$	—	2	9.7
$[\text{Re}(\text{CO})_3\text{Cl}(\text{bipy})]$	—	4	16.8
$[\text{Re}(\text{CO})_3\text{Cl}(\text{bipy})]$	NEt_4Cl	2	14.5
$[\text{Re}(\text{CO})_3\text{Cl}(\text{bipy})]$	NEt_4Cl	4	30.0
$[\text{Re}(\text{CO})_3\text{Br}(\text{bipy})]$	—	2	7.6
$[\text{Re}(\text{CO})_3\text{Br}(\text{bipy})]$	—	4	11.8
$[\text{Re}(\text{CO})_3\text{Br}(\text{bipy})]$	NEt_4Br	2	12.0
$[\text{Re}(\text{CO})_3\text{Br}(\text{bipy})]$	NEt_4Br	4	16.0


CO₂ and HCOO[−] by [Ru(bipy)₃]²⁺ in the presence of methylviologen (MV²⁺) in a TEOA–DMF mixture under irradiation of light (λ > 320 nm) [436]. Ziessel and co-workers have studied the photochemical reduction of CO₂ to formate by visible light irradiation (λ > 400 nm) of systems containing either the [Ru(bipy)₃]²⁺ complex which yields the active catalytic species by photolabilization of a bipy ligand, or a mixture of [Ru(L-L)₃]²⁺ (L-L ≡ 2,2′-bipyridyl or 1,10-phenanthroline) as photosensitizer and *cis*-[Ru(CO)X(bipy)₂]ⁿ⁺ (X ≡ Cl, H, n = 1 or X = CO, n = 2) or *cis*-[Ru(CO)₂Cl₂(bipy)] as homogeneous catalysts [437,438]. The solvent was either TEOA–DMF or TEOA–DMF–H₂O. Photocatalyst [Ru(CO)₂(bipy)₂]²⁺ and photosensitizer [Ru(bipy)₃]²⁺ (or [Ru(phen)₃]²⁺) reduce CO₂ to afford HCO₂[−] in the triethanolamine–DMF system under irradiation (λ > 320 nm) [439]. The complex *cis*-[Ru(CO)₂(bipy)₂]²⁺ is reduced by 2 mol [Ru(bipy)₃]⁺, a powerful monoelectronic reducing agent, to give a coordinatively unsaturated complex [Ru(CO)(bipy)₂] [439]. The generation of active species [Ru(CO)(bipy)₂] by double reduction [Ru(CO)Cl(bipy)₂]⁺ and [RuH(CO)(bipy)₂]⁺ has also been proposed by Lehn and Ziessel [438].





The CO₂ reduction to formate might proceed through protonation of bound CO₂:



The photochemical reduction of CO₂ to HCO₃[−] was observed using a cobalt(II)-carbonic hydrazase III [440]. Spreer and co-workers have reported photochemical reduction of CO₂ to CO and H₂ in water using [Ru(bipy)₃]²⁺ as a photosensitizer, ascorbate buffer as electron donor and the [Ni(cyclam)]²⁺ (cyclam ≡ 1,4,8,11-tetraazacyclotetra-decane) as catalyst. [Ni(cyclam)]²⁺ acts as a homogeneous catalyst for the photochemical reduction of CO₂ to CO, as well as for the reduction of H⁺ to H₂ [441]. The possible reaction mechanism is shown as follows:





Acknowledgments

This work was supported by the Department of Science and Technology, New Delhi. Professors K.M. Nicholas, A. Behr, S. Sakaki, J. Mascetti and H. Werner are thanked for articles supplied.

References

- [1] F.A. Cotton, Chemical Applications of Group Theory, Wiley Eastern, New Delhi, 1976, 2nd edn.
- [2] J.W. Rabalais, J.M. McDonald, V. Sherr and S.P.M. McGlynn, Chem. Rev., 71 (1971) 73.
- [3] K.S. Sidhu, I.G. Csizmadia, O.P. Strausz and H.E. Gunning, J. Am. Chem. Soc., 88 (1988) 2412.
- [4] P.C. Cross and L.O. Brockway, J. Chem. Phys., 3 (1935) 821.
- [5] D.W. Dakin, W.E. Good and D.K. Coles, Phys. Rev., 70 (1946) 560.
- [6] D.W. Dakin, W.E. Good and D.K. Coles, Phys. Rev., 71 (1947) 640.
- [7] R.W. Dornie, J. Am. Chem. Soc., 55 (1933) 4126.
- [8] L. Pauling, The Nature of the Chemical Bond, Cornell University Press, Ithaca, NY, 1969, p. 268.
- [9] K. Kuchitsu and K. Oyanagi, Faraday Discuss. Chem. Soc., (1977) 62.
- [10] I.L. Karle and J. Karle, J. Chem. Phys., 17 (1949) 1057.
- [11] J.M. McDonald, J.W. Rabalais and S.P.M. McGlynn, J. Chem. Phys., 52 (1970) 1332.
- [12] R.C. Brundle and D.W. Turner, Int. J. Mass Spectrom. Ion Phys., 2 (1969) 195.
- [13] D.W. Turner and D.P. May, J. Chem. Phys., 46 (1967) 1156.
- [14] M.I. Al-Joboury, D.P. May and D.W. Turner, J. Chem. Soc., (1965) 6350.
- [15] R.E. Honing, J. Chem. Phys., 16 (1948) 105.
- [16] J. Collin, Bull. Cl. Sci. Acad. R. Belg., 45 (1959) 105.
- [17] J.M. Sichel and M.A. Whitehead, Theor. Chim. Acta, 11 (1968) 239.
- [18] H.E. Hennecke and R.S. Drago, J. Am. Chem. Soc., 90 (1968) 5112.
- [19] R.N. Compton, P.W. Reinhardt and C.D. Cooper, J. Chem. Phys., 63 (1975) 3829.
- [20] K. Watanabe, J. Chem. Phys., 26 (1957) 542.
- [21] F.M. Field and J.L. Fracklin, Electron Impact Phenomena and the Properties of Gaseous Ions, Academic Press, New York, 1957.
- [22] V.M. Berdnikov, Russ. J. Phys. Chem., (1975) 11.
- [23] L.A. Pugh and K.N. Rao, Molecular Spectroscopy: Modern Research, Vol. II, Academic Press, New York, 1976.
- [24] W.B. Person, K.G. Brown, D. Steele and D. Peters, J. Phys. Chem., 85 (1981) 1998.
- [25] C.J. Allan, U. Gelius, D.A. Allison, G. Johansson, H. Siegbahn and K. Siegbahn, J. Electron Spectrosc. Relat. Phenom., 1 (1972) 131.
- [26] K.K. Pandey and H.L. Nigam, Rev. Inorg. Chem., 6 (1984) 69.
- [27] J.A. Ibers, Chem. Soc. Rev., 11 (1982) 57.
- [28] I.S. Butler and A.E. Fenster, J. Organomet. Chem., 66 (1974) 161.
- [29] P.V. Yaneff, Coord. Chem. Rev., 23 (1977) 183.
- [30] I.S. Butler, Acc. Chem. Res., 10 (1977) 359.
- [31] H. Werner, Coord. Chem. Rev., 43 (1982) 165.
- [32] C. Bianchini, C. Mealli, A. Meli and M. Sabat, in I. Bernal (ed.), Stereochemistry of Organometallic and Inorganic Compounds, Vol. 1, Elsevier, Amsterdam, 1986, p. 146.
- [33] M.E. Vol'pin and I.S. Kolomnikov, Pure Appl. Chem., 33 (1977) 567.

- [34] V.J. Choy and C.J. O'Connor, *Coord. Chem. Rev.*, 9 (1973) 145.
- [35] M.E. Vol'pin and I.S. Kolomnikov, in M. Tsutsui (ed.), *Organometallic Reactions*, Vol. 5, Wiley Interscience, New York, 1975, p. 313.
- [36] I.S. Kolomnikov and M.Kh. Grigoryan, *Russ. Chem. Rev.*, 47 (1978) 334.
- [37] R. Eisenberg and D.E. Hendrikson, *Adv. Catal.*, 28 (1979) 119.
- [38] A.L. Lapidus and Y.Y. Ping, *Russ. Chem. Rev.*, 50 (1981) 63.
- [39] R.P.A. Sneeden and C.N.R.S. Villeurbanne, in G. Wilkinson (ed.), *Comprehensive Organometallic Chemistry*, Pergamon, Oxford, 1982, p. 225.
- [40] R. Ziessel, *Nouv. J. Chim.*, 17 (1983) 613.
- [41] D.A. Palmer and R.V. Eldik, *Chem. Rev.*, 83 (1983) 651.
- [42] A. Behr, in W. Keim (ed.), *Catalysis in C₁ Chemistry*, Reidel, Dordrecht, 1983, p. 169.
- [43] D.J. Darensbourg and R.A. Kudarowski, *Adv. Organomet. Chem.*, 22 (1983) 129.
- [44] D.J. Darensbourg and C. Ovalles, *CHEMTECH.*, (1985) 636.
- [45] D. Walther, *Coord. Chem. Rev.*, 79 (1987) 135.
- [46] A. Behr, *Angew. Chem., Int. Ed. Engl.*, 27 (1988) 661.
- [47] P. Braunstein, D. Matt and D. Nobel, *Chem. Rev.*, 88 (1988) 747.
- [48] A.R. Cutler, P.K. Hanna and J.C. Vites, *Chem. Rev.*, 88 (1988) 1363.
- [49] E. Carmona, *J. Organomet. Chem.*, 358 (1988) 283.
- [50] A. Behr, in R. Ugo (ed.), *Aspects of Homogeneous Catalysis*, Vol. 6, Reidel, Dordrecht, 1988, p. 59.
- [51] J.-P. Collin and J.-P. Sauvage, *Coord. Chem. Rev.*, 93 (1989) 245.
- [52] I.S. Kolomnikov and T.V. Lysyak, *Russ. Chem. Rev.*, 59 (1990) 344.
- [53] S. Inoue and H. Koinuma, *Rev. Inorg. Chem.*, 6 (1984) 291.
- [54] K. Ogura and H. Uchida, *J. Chem. Soc., Dalton Trans.*, (1987) 1377.
- [55] M. Aresta, E. Quaranta and A. Cicarese, *J. Mol. Catal.*, 41 (1987) 355.
- [56] M. Aresta, C. Fragale, E. Quaranta and I. Tommasi, *J. Chem. Soc., Chem. Commun.*, (1992) 315.
- [57] E. Dunach and J. Perichon, *J. Organomet. Chem.*, 352 (1988) 239.
- [58] L. Garnier, Y. Rollin and J. Perichon, *J. Organomet. Chem.*, 367 (1989) 347.
- [59] L. Garnier, Y. Rollin and J. Perichon, *Nouv. J. Chim.*, 13 (1989) 53.
- [60] E. Dunach and J. Perichon, *Synlett.*, 1 (1990) 143.
- [61] S. Derien, E. Dunach and J. Perichon, *J. Organomet. Chem.*, 385 (1990) C45.
- [62] S. Derien, J.-C. Clinet, E. Dunach and J. Perichon, *J. Chem. Soc., Chem. Commun.*, (1991) 549.
- [63] S. Derien, J.-C. Clinet, E. Dunach and J. Perichon, *Tetrahedron*, 48 (1992) 5235.
- [64] T. Tsuda, S. Morikawa, R. Sumiya and T. Saegusa, *J. Org. Chem.*, 53 (1988) 3140.
- [65] T. Tsuda, K. Maruta and Y. Kitaie, *J. Am. Chem. Soc.*, 114 (1992) 1498.
- [66] T. Mitaudo, Y. Hori, Y. Yamakawa and Y. Watanabe, *Tetrahedron Lett.*, 28 (1987) 4417.
- [67] C. Amatore and A. Jutland, *J. Am. Chem. Soc.*, 113 (1991) 2819.
- [68] C. Amatore, A. Jutland, F. Khalil and M.L. Nelson, *J. Am. Chem. Soc.*, 114 (1992) 7076.
- [69] N. Komeda, H. Nagae, T. Matsui, G.-Y. Adachi and K. Tanaka, *J. Am. Chem. Soc.*, 114 (1992) 3652.
- [70] W.D. McGhee and D.P. Riley, *Organometallics*, 11 (1992) 900.
- [71] S.M. Pillai, R. Ohnishi and M. Ichikawa, *J. Chem. Soc., Chem. Commun.*, (1990) 246.
- [72] S. Schreiner, J.Y. Yu and L. Vaska, *J. Chem. Soc., Chem. Commun.*, (1988) 602.
- [73] S. Schreiner, J.Y. Yu and L. Vaska, *Inorg. Chim. Acta*, 147 (1988) 139.
- [74] E. Graf and W. Leitner, *J. Chem. Soc., Chem. Commun.*, (1992) 623.
- [75] J.W. Rathke, R.J. Klingler and T.R. Krause, *Organometallics*, 10 (1991) 1350.
- [76] J.W. Rathke, R.J. Klingler and T.R. Krause, *Organometallics*, 11 (1992) 585.
- [77] M. Koda, T. Tomohiro, A.L. Lee and H. Okuno, *J. Chem. Soc., Chem. Commun.*, (1989) 1479.
- [78] H. Hoberg and M. Minato, *J. Organomet. Chem.*, 406 (1991) C25.
- [79] A.K.F. Rahman, J.-C. Tsai and K.M. Nicholas, *J. Chem. Soc., Chem. Commun.*, (1992) 1334.
- [80] M. Rosi, A. Sgamellotti, F. Tarantelli and C. Floriani, *J. Organomet. Chem.*, 332 (1987) 153.
- [81] C.C. Hinkley, L.V. Dunkerton and J. Tyrell, Report DOE/FE/60339-T4, 1984; *Chem. Abstr.*, 102: 196803.
- [82] W. Poppitz and E. Uhlig, *Z. Anorg. Allg. Chem.*, 489 (1982) 67.
- [83] S. Gamarotta, M.L. Fiallo, C. Floriani, A.C. Villa and C. Guastini, *Inorg. Chem.*, 23 (1984) 3532.
- [84] C.C. Hinkley, L.V. Dunkerton and J. Tyrell, Report DOE/FE/60339-T20, 1985.

- [85] R.L. Cowan, D.B. Pourreau, A.L. Rheingold, S.J. Geib and W.C. Trogler, *Inorg. Chem.*, 26 (1987) 259.
- [86] D.L. Allen, M.L.H. Green and J.A. Bandy, *J. Chem. Soc., Dalton Trans.*, (1990) 541.
- [87] T.R. Gaffney and J.A. Ibers, *Inorg. Chem.*, 21 (1982) 2851.
- [88] K.K. Pandey and S.K. Tewari, *Polyhedron*, 9 (1989) 1149.
- [89] K.K. Pandey and S.K. Tewari, unpublished results, 1989.
- [90] S.K. Tewari and K.K. Pandey, *Syn. React. Inorg. Met.-Org. Chem.*, 19 (1989) 545.
- [91] T.R. Gaffney and J.A. Ibers, *Inorg. Chem.*, 21 (1982) 2854.
- [92] T.R. Gaffney and J.A. Ibers, *Inorg. Chem.*, 21 (1982) 2860.
- [93] M. Massoudipour, S.K. Tewari and K.K. Pandey, *Polyhedron*, 8 (1989) 1447.
- [94] H. Werner, M. Ebner and W. Bertlett, *Z. Naturforsch.*, 40b (1985) 1351.
- [95] C. Bianchini, A. Meli and G. Scapacci, *Cong. Nat. Chem. Inorg.*, 16 (1987) 51.
- [96] C. Bianchini, C. Mealli, A. Meli and G. Scapacci, *Organometallics*, 2 (1983) 141.
- [97] C. Bianchini, C.A. Ghilardi, A. Meli and A. Orlandini, *J. Organomet. Chem.*, 270 (1984) 251.
- [98] T.R. Gaffney and J.A. Ibers, *Inorg. Chem.*, 21 (1982) 2857.
- [99] J.M. Burke and J.P. Fackler, *Inorg. Chem.*, 11 (1972) 2744.
- [100] P.J. Hayward, D.M. Blake, G. Wilkinson and C.J. Nyman, *J. Am. Chem. Soc.*, 92 (1970) 5873.
- [101] H. Werner, W. Bertlett, B. Zimmer-Gasser and U. Schubert, *Chem. Ber.*, 115 (1982) 1004.
- [102] R. Alvarez, E. Carmona, E.G. Puebla, J.M. Marin, A. Monge and M.L. Poveda, *J. Chem. Soc., Chem. Commun.*, (1984) 1326.
- [103] R. Alvarez, E. Carmona, M.L. Poveda and R.S. Delgado, *J. Am. Chem. Soc.*, 106 (1984) 2731.
- [104] R. Alvarez, E. Carmona, J.M. Martin, M.L. Poveda, E.G. Puebla, A. Monge and M.L. Poveda, *J. Am. Chem. Soc.*, 108 (1986) 2286.
- [105] E. Carmona, A. Galindo, A. Monge, M.A. Munoz, M.L. Poveda and C. Ruiz, *Inorg. Chem.*, 29 (1990) 5074.
- [106] D.J. Darensbourg and A. Rokicki, *J. Am. Chem. Soc.*, 104 (1982) 349.
- [107] D.J. Darensbourg and A. Rokicki, *Organometallics*, 1 (1982) 1682.
- [108] C. Bianchini, C.A. Ghilardi, A. Meli, S. Midollini and A. Orlandini, *J. Organomet. Chem.*, 248 (1983) C13.
- [109] C. Bianchini, C.A. Ghilardi, A. Meli, S. Midollini and A. Orlandini, *J. Organomet. Chem.*, 255 (1983) C27.
- [110] C. Bianchini, C.A. Ghilardi, A. Meli, S. Midollini and A. Orlandini, *Inorg. Chem.*, 24 (1985) 924.
- [111] A. Mishra and U.C. Agarwala, *Inorg. Chim. Acta*, 145 (1988) 191.
- [112] K.K. Pandey, H. Garg and S.K. Tewari, *Polyhedron*, 11 (1992) 947.
- [113] G.J. Brehnan, R.A. Anderson and A. Zalkin, *Inorg. Chem.*, 25 (1986) 1761.
- [114] L. Ruiz-Ramirez, T.A. Stephenson and E.S. Switkes, *J. Chem. Soc., Dalton Trans.*, (1973) 1770.
- [115] F.G. Moers, R.W.M. ten Hoedt and J.P. Langhout, *Inorg. Chem.*, 12 (1973) 2196.
- [116] M.P. Yagupsky and G. Wilkinson, *J. Chem. Soc. A*, (1968) 2813.
- [117] M.C. Biard and G. Wilkinson, *J. Chem. Soc. A*, (1967) 865.
- [118] D. Brown and F. Hughes, *Inorg. Chim. Acta*, 1 (1967) 448.
- [119] S. Sakaki, K. Kitaura and K. Morokuma, *Inorg. Chem.*, 21 (1982) 760.
- [120] C. Mealli, R. Hoffmann and A. Stockis, *Inorg. Chem.*, 23 (1984) 56.
- [121] T. Ziegler, *Inorg. Chem.*, 25 (1986) 2721.
- [122] M. Rosi, A. Sgamellotti, F. Tarantelli and C. Floriani, *Inorg. Chem.*, 26 (1987) 3805.
- [123] H.G. Alt, K.-H. Schwind and M.D. Rausch, *J. Organomet. Chem.*, 321 (1987) C9.
- [124] E. Klei, J.H. Teuben, H.J.D. Meijer, E.J. Kwak and A.P. Bruins, *J. Organomet. Chem.*, 224 (1982) 327.
- [125] M.J. McGlinchey, J.L. Fletcher, B.G. Sayer, P. Bougeard, R. Faggiani, C.J.L. Lock, A.D. Bain, C. Rodger, E.P. Kundig, D. Astruc, J.R. Haman, P. Lemaux, S. Top and G. Jaouen, *J. Chem. Soc., Chem. Commun.*, (1983) 634.
- [126] G.R. Lee, J.M. Maher and N.J. Rooper, *J. Am. Chem. Soc.*, 109 (1987) 2956.
- [127] R.J. Angelici and J.W. Dunker, *Inorg. Chem.*, 24 (1985) 2215.
- [128] A. Gutierrez-Alonso and L.B. Reventos, *J. Coord. Chem.*, 22 (1991) 255.
- [129] M. Herberhold, A.F. Hill, N. McAuley and W.R. Roper, *J. Organomet. Chem.*, 310 (1986) 95.

- [130] J. Fortune and A.R. Manning, *Organometallics*, 2 (1983) 1719.
- [131] C. Bianchini, A. Meli and G. Scapacci, *Organometallics*, 2 (1983) 1834.
- [132] O. Kolb and H. Werner, *J. Organomet. Chem.*, 268 (1984) 49.
- [133] B. Klingert and H. Werner, *Chem. Ber.*, 116 (1983) 1450.
- [134] C. Bianchini, C. Mealli, A. Meli and M. Sabat, *J. Chem. Soc., Chem. Commun.*, (1984) 1647.
- [135] C. Bianchini, C. Mealli, A. Meli and M. Sabat, *Inorg. Chem.*, 23 (1984) 4125.
- [136] C. Bianchini, D. Masi, C. Mealli, A. Meli and M. Sabat, *Organometallics*, 4 (1985) 1014.
- [137] C. Bianchini, C. Mealli, A. Meli, M. Sabat and P. Zanello, *J. Am. Chem. Soc.*, 109 (1987) 185.
- [138] R. Uson, L.A. Ore, D. Carmona, M.J. Fernades and P. Lahuerta, *Rev. Acad. Cienc. Exactas Fis. Quim. Nat. Zaragoza*, 37 (1982) 61.
- [139] T. Hoffman, M. Harabit, P. Zdunneck and E. Werschuh, *Z. Chem.*, 28 (1988) 415.
- [140] W. Bertleff and H. Werner, *Z. Naturforsch.*, 37b (1982) 1294.
- [141] M. Ebner, H. Otto and H. Werner, *Angew. Chem., Int. Ed. Engl.*, 24 (1985) 518.
- [142] W.M. Hawling, A. Walker and M.A. Woitzik, *J. Chem. Soc., Chem. Commun.*, (1983) 11.
- [143] E. Ma, G. Semelhago, W. Walker, D.H. Farrar and R.R. Gukathasan, *J. Chem. Soc., Dalton Trans.*, (1985) 2595.
- [144] I.K. Gregor, *Inorg. Chim. Acta*, 176 (1990) 19.
- [145] L. Busetto, A. Palazzi and M. Monari, *J. Organomet. Chem.*, 228 (1982) C19.
- [146] H. Stolzenberg, W.P. Fehlhammer, *J. Organomet. Chem.*, 235 (1982) C7.
- [147] L. Busetto, A. Palazzi and M. Monari, *J. Chem. Soc., Dalton Trans.*, (1982) 1631.
- [148] L. Busetto, M. Monari, A. Palazzi, V. Albano and F. Demartin, *J. Chem. Soc., Dalton Trans.*, (1983) 1849.
- [149] H. Stolzenberg, W.P. Fehlhammer, M. Monari, V. Zanotti and L. Busetto, *J. Organomet. Chem.*, 272 (1984) 73.
- [150] M.E.G. Derry, B.E. Landrum, J.L. Shibley and A.R. Cutler, *J. Organomet. Chem.*, 378 (1989) 421.
- [151] C. Bianchini, S.A. Ghilardi, A. Meli, S. Midollini and A. Orlandini, *J. Chem. Soc., Chem. Commun.*, (1983) 753.
- [152] D.H. Farrar, R.R. Gukathasan and K. Won, *J. Organomet. Chem.*, 275 (1984) 263.
- [153] D.H. Farrar, R.R. Gukathasan and S.A. Morris, *Inorg. Chem.*, 23 (1984) 3258.
- [154] W. Deutsch, A. Gieren, C.R. Perz and K.E. Schwarhans, *J. Organomet. Chem.*, 355 (1988) 197.
- [155] M.L. Kullberg and C.P. Kubiak, *Inorg. Chem.*, 25 (1986) 26.
- [156] C.R. Langrick, P.G. Pringle and B.L. Shaw, *J. Chem. Soc., Dalton Trans.*, (1985) 1015.
- [157] P.G. Pringle and B.L. Shaw, *J. Chem. Soc., Dalton Trans.*, (1983) 889.
- [158] T.S. Cameron, P.A. Gardner and K.R. Grundy, *J. Organomet. Chem.*, 212 (1981) C19.
- [159] C. Bianchini, D. Masi, C. Mealli, A. Meli, M. Sabat and G. Scapacci, *J. Organomet. Chem.*, 273 (1984) 91.
- [160] D. Touchard, J.-L. Fillaut, P.H. Dixneuf and L. Toupet, *J. Organomet. Chem.*, 317 (1986) 291.
- [161] D. Touchard, C. Lelay, J.-L. Fillaut and P.H. Dixneuf, *J. Chem. Soc., Chem. Commun.*, (1986) 37.
- [162] W.A. Schenk, T. Schwietzke and H. Mueller, *J. Organomet. Chem.*, 232 (1982) C41.
- [163] P.J. Brothers and W.R. Roper, *J. Organomet. Chem.*, 258 (1983) 73.
- [164] P.W. Armit, T.A. Stephenson and E.S. Switkes, *Inorg. Synth.*, 21 (1982) 28.
- [165] P.V. Broadhurst, B.F.G. Johnson, J. Lewis and P.R. Raithby, *J. Chem. Soc., Chem. Commun.*, (1982) 140.
- [166] O.J. Scherer, R. Konard, E. Guggolz and M.L. Ziegler, *Chem. Ber.*, 116 (1983) 2676.
- [167] M. Massaudipour and K.K. Pandey, *Inorg. Chim. Acta*, 160 (1989) 115.
- [168] J.-S. Qi, P.W. Schrier, P.E. Fanwick and R.A. Walton, *J. Chem. Soc., Chem. Commun.*, (1991) 1737.
- [169] J.-S. Qi, P.W. Schrier, P.E. Fanwick and R.A. Walton, *Inorg. Chem.*, 31 (1992) 258.
- [170] E. Hoyer, *Comments Inorg. Chem.*, 2 (1983) 261.
- [171] J.J. Maj, A.D. Rae and L.F. Dahl, *J. Am. Chem. Soc.*, 104 (1982) 4278.
- [172] H.A. Harris, A.D. Rae and L.F. Dahl, 190th Natl. Meet. of the American Chemical Society, Chicago, IL, 1985.
- [173] H.A. Harris, A.D. Rae and L.F. Dahl, *J. Am. Chem. Soc.*, 109 (1987) 4739.
- [174] C. Bianchini, C. Mealli, A. Meli, M. Sabat and P. Zanello, *J. Am. Chem. Soc.*, 109 (1987) 185.
- [175] H. Werner, O. Kolb, R. Feser and U. Schubert, *J. Organomet. Chem.*, 191 (1980) 283.

- [176] E. Carmona, J. Campora, M.A. Munoz, M. Paneque and M.L. Poveda, *Pure Appl. Chem.*, 61 (1989) 1701.
- [177] M.G. Mason, P.N. Swebston and J.A. Ibers, *Inorg. Chem.*, 22 (1983) 411.
- [178] M. Cowie and S.K. Dwight, *J. Organomet. Chem.*, 214 (1981) 233.
- [179] C. Bianchini, C.A. Ghilardi, A. Meli, S. Midollini and A. Orlandini, *Inorg. Chem.*, 24 (1985) 932.
- [180] C. Bianchini, C.A. Ghilardi, A. Meli, S. Midollini and A. Orlandini, *J. Chem. Soc., Chem. Commun.*, (1983) 545.
- [181] C. Bianchini, A. Meli and A. Orlandini, *Inorg. Chem.*, 21 (1982) 4161.
- [182] C. Bianchini, A. Meli and A. Orlandini, *Inorg. Chem.*, 21 (1982) 4166.
- [183] S. Gopinathan, I.R. Unni, C. Gopinathan, V.G. Puranik, S.S. Tavele and T.N. Guru Row, *Polyhedron*, 6 (1987) 1859.
- [184] A. Campus, N. Marsich and G. Nardin, *J. Organomet. Chem.*, 188 (1980) 389.
- [185] A. Campus, N. Marsich and G. Pellizer, *J. Organomet. Chem.*, 259 (1980) 389.
- [186] A.M.M. Lanfredi, A. Tiripicchio, A. Campus and N. Marsich, *J. Chem. Soc., Chem. Commun.*, (1983) 1126.
- [187] A.M.M. Lanfredi, F. Ugozzoli, A. Campus and N. Marsich, *Inorg. Chim. Acta*, 99 (1985) 111.
- [188] A. Campus, N. Marsich, A.M.M. Lanfredi and F. Ugozzoli, *Inorg. Chim. Acta*, 175 (1990) 193.
- [189] H. Otto and H. Werner, *Chem. Ber.*, 120 (1987) 97.
- [190] M.R. Torres, A. Perales and J. Ros, *Organometallics*, 7 (1988) 1223.
- [191] M.I. Bruce, M.J. Liddell, M.R. Snow and E.R.J. Tiekink, *J. Organomet. Chem.*, 352 (1988) 199.
- [192] D.J. Darensbourg, H.P. Wiegrefe and J.H. Reibenspies, *Organometallics*, 10 (1991) 6.
- [193] M.A. Ansari, J. Chandrashekhara and S. Sarkar, *Inorg. Chem.*, 27 (1988) 763.
- [194] M.L. Ziegler, K. Blechschmitt, H. Bock, E. Guggolz and P.R. Korswagen, *Z. Naturforsch.*, 43b (1988) 590.
- [195] P.L. Bellon, S. Cenni, F. Demartin, M. Pizzotti and F. Porta, *J. Chem. Soc., Chem. Commun.*, (1982) 265.
- [196] S. Cenini, F. Porta, M. Pizzotti and C. Crotti, *J. Chem. Soc., Dalton Trans.*, (1985) 163.
- [197] T.R. Grafney and J.A. Ibers, *Inorg. Chem.*, 21 (1982) 2062.
- [198] L.I. Daniliana, A.V. Ireteki, Yu.N. Kukushkin, *Koord. Khim.*, 11 (1985) 1271.
- [199] H. Werner and W. Bertleff, *Chem. Ber.*, 113 (1980) 267.
- [200] E. Hey, M.F. Lappert, J.L. Atwood and S.G. Bott, *J. Chem. Soc., Chem. Commun.*, (1987) 421.
- [201] D. Coucouvanis and M.E. Draganjac, *J. Am. Chem. Soc.*, 104 (1982) 6820.
- [202] D. Coucouvanis, M.E. Draganjac, S.M. Koo, A. Toupadakis and A.I. Hadjikyriacou, *Inorg. Chem.*, 31 (1992) 1186.
- [203] K.N. Udpa and S. Barker, *Polyhedron*, 6 (1987) 627.
- [204] D. Jentsch, P.G. Jones, C. Thoenne and E. Schwarzmann, *J. Chem. Soc., Chem. Commun.*, (1989) 1495.
- [205] D. Touchard, J.-L. Fillout, P.H. Dixneuf, C. Mealli, M. Sabat and L. Toupet, *Organometallics*, 4 (1985) 1684.
- [206] H.G. Raubenheimer, G.J. Kruger and H.W. Viljoen, *J. Chem. Soc., Dalton Trans.*, (1985) 1963.
- [207] H.G. Raubenheimer, G.J. Kruger and A. Lombard, *J. Organomet. Chem.*, 240 (1982) C11.
- [208] H.G. Raubenheimer, S. Lotz, G.J. Kruger, A.A. Lombard and J.C. Viljoen, *J. Organomet. Chem.*, 336 (1987) 349.
- [209] G. Groetsch and W. Malisch, *J. Organomet. Chem.*, 262 (1984) C3.
- [210] M. Ebner and H. Werner, *Chem. Ber.*, 119 (1986) 482.
- [211] F. Ceconi, C.A. Ghilardi, S. Midollini, S. Monet and A. Orlandini, *J. Organomet. Chem.*, 323 (1987) C5.
- [212] A. Behr, W. Keim, W. Kipshagen, D. Vogt, E. Herdweck and W.A. Herrmann, *J. Organomet. Chem.*, 344 (1988) C15.
- [213] T. Kauffmann and J. Olbrich, *Tetrahedron Lett.*, 25 (1984) 1967.
- [214] S. Sakaki, K. Kitaura, K. Morokuma and K. Ohkubo, *Inorg. Chem.*, 22 (1983) 104.
- [215] O.-Z. Shi, T.G. Richmond, W.C. Troglor and F. Basolo, *Inorg. Chem.*, 23 (1984) 957.
- [216] S. Sakaki and A. Dedieu, *J. Organomet. Chem.*, 314 (1986) C63.

- [217] R.A. Wheeler, M.-H. Whangbo, T. Hughbanks, R. Hoffmann, J.K. Burdett and T.A. Albright, *J. Am. Chem. Soc.*, 108 (1986) 2222.
- [218] S. Sakaki, in A. Veillard (ed.), *Quantum Chemistry: The Challenge of Transition Metals and Coordination Chemistry*, NATO ASI Series, Reidel, Dordrecht, 1986, p. 319.
- [219] S. Sakaki and A. Dedieu, *Inorg. Chem.*, 26 (1987) 3278.
- [220] E.S. Marcos, R. Caballol, G. Trinquar and J.-C. Barthelat, *J. Chem. Soc., Dalton Trans.*, (1987) 1328.
- [221] R. Coballol, E.S. Marcos and J.-C. Barthelat, *J. Phys. Chem.*, 91 (1987) 1328.
- [222] V. Branchadell and A. Dedieu, *Inorg. Chem.*, 26 (1987) 3966.
- [223] S. Sakaki, T. Aizawa, N. Koga, K. Morokuma and K. Ohkubo, *Inorg. Chem.*, 28 (1989) 103.
- [224] S. Sakaki, *J. Am. Chem. Soc.*, 112 (1990) 7813.
- [225] S. Sakaki, N. Koga and K. Morokuma, *Inorg. Chem.*, 29 (1990) 3110.
- [226] S. Sakaki, *J. Am. Chem. Soc.*, 114 (1992) 2055.
- [227] S. Sakaki and K. Ohkubo, *Inorg. Chem.*, 27 (1988) 2020.
- [228] S. Sakaki and K. Ohkubo, *Inorg. Chem.*, 28 (1989) 2583.
- [229] C. Bo and A. Dedieu, *Inorg. Chem.*, 28 (1988) 304.
- [230] S. Sakaki and K. Ohkubo, *Organometallics*, 8 (1989) 2970.
- [231] J. Mascetti and M. Tranquilli, *J. Phys. Chem.*, 92 (1988) 2177.
- [232] C. Jegat, M. Fouassier, M. Tranquille and J. Mascetti, *Inorg. Chem.*, 30 (1991) 1529.
- [233] G.R. Lee and N.J. Cooper, *Organometallics*, 4 (1985) 794.
- [234] M.E. Giuseppetti and A.R. Cutler, *Organometallics*, 6 (1987) 970.
- [235] D.H. Gibson and T.-S. Ong, *J. Am. Chem. Soc.*, 109 (1987) 7191.
- [236] D.R. Senn, K. Emerson, R.D. Larsen and J.A. Gladysz, *Inorg. Chem.*, 26 (1987) 2737.
- [237] J.R. Sweet and W.A.G. Graham, *Organometallics*, 1 (1982) 982.
- [238] C.F. Barrientos-Penna, A.B. Gilchrist, A.H. Lkahn-Oliva, J.L.A. Hanlan and D. Sutton, *Organometallics*, 4 (1985) 478.
- [239] E. Fujita, D.J. Szalda, C. Creutz and N. Sutin, *J. Am. Chem. Soc.*, 110 (1988) 4870.
- [240] E. Fujita, C. Creutz, N. Sutin and D.J. Szalda, *J. Am. Chem. Soc.*, 113 (1991) 343.
- [241] C. Creutz, H.A. Schwartz, J.F. Wishart, E. Fujita and N. Sutin, *J. Am. Chem. Soc.*, 113 (1991) 3361.
- [242] S. Gambaratta, F. Arena, C. Floriani and P.F. Zanazzi, *J. Am. Chem. Soc.*, 104 (1982) 5082.
- [243] M.H. Schmidt, G.M. Miskelly and N.S. Lewis, *J. Am. Chem. Soc.*, 112 (1990) 3420.
- [244] D.A. Gangi and R.R. Durand, *J. Chem. Soc., Chem. Commun.*, (1986) 697.
- [245] T. Herskovitz, *Inorg. Synth.*, 21 (1982) 99.
- [246] J.C. Calabrese, T. Herskovitz and J.B. Kinney, *J. Am. Chem. Soc.*, 105 (1983) 5914.
- [247] T. Ergel and G. Ertl, *Adv. Catal.*, 28 (1979) 1.
- [248] H. Huber, D. McIntosh and G.A. Ozin, *Inorg. Chem.*, 16 (1977) 975.
- [249] A. Zecchina, G. Spoto, S. Coluccia and E. Guglielminotti, *J. Chem. Soc., Faraday Trans. I*, 80 (1984) 1975.
- [250] P.-F. Fu, M.A. Khan and K.M. Nicholas, *J. Am. Chem. Soc.*, 119 (1992) 6579.
- [251] G.S. Bristow, P.B. Hitchcock and M.F. Lappert, *J. Chem. Soc., Chem. Commun.*, (1981) 1145.
- [252] S. Gambarotta, C. Floriani, A.C. Villa and C. Guastini, *J. Am. Chem. Soc.*, 107 (1985) 2985.
- [253] K.A. Belmore, R.A. Vanderpool, J.-C. Tsai, M.A. Khan and K.M. Nicholas, *J. Am. Chem. Soc.*, 110 (1988) 2004.
- [254] R.S. Pilato, C.E. Housemekerides, P. Jernakoff, D. Rubin, G. Geoffroy and A.L. Rheingold, *Organometallics*, 9 (1990) 2333.
- [255] C.T. Tso and A.R. Cutler, *J. Am. Chem. Soc.*, 108 (1986) 6069.
- [256] V.C. Vites, B.D. Steffey, M.E. Giuseppetti-Dery and A.E. Cutler, *Organometallics*, 10 (1991) 2827.
- [257] A. Deohring, P.W. Jolly, C. Kruger and A.J. Romao, *Z. Naturforsch.*, 40b (1985) 484.
- [258] J.D. Audett, T.J. Collins, B.D. Santarsiero and G.H. Spies, *J. Am. Chem. Soc.*, 104 (1982) 7352.
- [259] M.A. Bennett, G.B. Robertson, A. Rokicki and W.A. Wickramasinghe, *J. Am. Chem. Soc.*, 110 (1988) 7098.
- [260] C.R. Eady, J.J. Guy, B.F.G. Johnson, J. Lewis, M.C. Malatesta and G.M. Sheldrick, *J. Chem. Soc., Chem. Commun.*, (1976) 602.
- [261] G.R. John, B.F.G. Johnson, J. Lewis and K.C. Wong, *J. Organomet. Chem.*, 169 (1979) C23.

- [262] B.K. Balbach, F. Helus, F. Oberdorfer and M.L. Ziegler, *Angew. Chem., Int. Ed. Engl.*, 20 (1981) 470.
- [263] B.K. Balbach, F. Helus, F. Oberdorfer and M.L. Ziegler, *Z. Naturforsch.*, 37b (1982) 1982.
- [264] W. Beck, K. Raab, V. Nagel and M. Steimann, *Angew. Chem., Int. Ed. Engl.*, 21 (1982) 526.
- [265] E.C. Lundquist, J.C. Huffman and K.G. Caulton, *J. Am. Chem. Soc.*, 108 (1986) 8309.
- [266] E.G. Lundquist, J.C. Huffman, K. Folting, B.E. Mann and K.G. Caulton, *Inorg. Chem.*, 29 (1990) 128.
- [267] W.B. Person, K.G. Brown, D. Steele and D. Peters, *J. Phys. Chem.*, 85 (1981) 1998.
- [268] C. Jegat, M. Fouassier and J. Mascetti, *Inorg. Chem.*, 30 (1991) 1521.
- [269] G.A. Luinstra and J.H. Teuben, *J. Chem. Soc., Chem. Commun.*, (1987) 849.
- [270] D. Lyons, G. Wilkinson, M. Thornton-Pelt and M.B. Hursthouse, *J. Chem. Soc., Dalton Trans.*, (1984) 695.
- [271] T. Ito and T. Matsubara, *J. Chem. Soc., Dalton Trans.*, (1988) 2241.
- [272] D.C. Brower, P.B. Winston, T.L. Tonker and J.L. Templeton, *Inorg. Chem.*, 25 (1986) 2883.
- [273] D. Lyons and G. Wilkinson, *J. Chem. Soc., Dalton Trans.*, (1985) 587.
- [274] F.J. Alonso, M.G. Sanz and V. Riera, *J. Organomet. Chem.*, 421 (1991) C12.
- [275] B.P. Sullivan and T.J. Meyer, *J. Chem. Soc., Chem. Commun.*, (1984) 1244.
- [276] D.J. Darensbourg, M.Y. Darensbourg, L.Y. Goh, M. Ludvig and P. Wiegrefe, *J. Am. Chem. Soc.*, 109 (1987) 7539.
- [277] A. Immirzi and A. Musco, *Inorg. Chim. Acta*, 22 (1977) L35.
- [278] G. LaMonica, M.A. Angaroni, F. Cariatì and S. Cenini, *Inorg. Chim. Acta*, 143 (1988) 239.
- [279] R.F. Johnston and J.C. Cooper, *Organometallics*, 6 (1987) 2448.
- [280] J. Nieman and J.H. Teuben, *J. Organomet. Chem.*, 287 (1985) 207.
- [281] D.J. Darensbourg and R. Kudarowski, *J. Am. Chem. Soc.*, 106 (1984) 3672.
- [282] D.J. Darensbourg, R. Kudarowski, C.G. Bauch, M. Pale, D. Simmons and J.N. White, *J. Am. Chem. Soc.*, 107 (1985) 7463.
- [283] D.J. Darensbourg, C.G. Bauch and A.I. Rheingold, *Inorg. Chem.*, 26 (1987) 977.
- [284] D.J. Darensbourg and G. Grotzsch, *J. Am. Chem. Soc.*, 107 (1985) 7473.
- [285] D.J. Darensbourg, C.G. Grotzsch and A.L. Rheingold, *Inorg. Chem.*, 26 (1987) 977.
- [286] D.J. Darensbourg, G. Grotzsch, P. Wiegrefe and A.R. Rheingold, *Inorg. Chem.*, 26 (1987) 3827.
- [287] M.M. Holl, G.L. Hillhouse, K. Folting and J.C. Huffman, *Organometallics*, 6 (1987) 1522.
- [288] P. Kundel and H. Berke, *J. Organomet. Chem.*, 339 (1988) 297.
- [289] D.J. Darensbourg, R. Kudarowski and T. Delord, *Organometallics*, 4 (1985) 1094.
- [290] M.H. Green, G. Parkin, D. O'Hare, L.L. Wong and A.E. Deroma, *J. Organomet. Chem.*, 317 (1986) 61.
- [291] J.F. Normant and G. Cahiez, in R. Scheffold (ed.), *Modern Synthetic Method*, Verlag Salle, Sauerlander, 1983, p. 173.
- [292] A. Behr, U. Kanne and G. Thelen, *J. Organomet. Chem.*, 269 (1984) C1.
- [293] I.M. Arafat, K. Shin and H.M. Goff, *J. Am. Chem. Soc.*, 110 (1988) 5228.
- [294] L. Dahlenburg and C. Pregnel, *J. Organomet. Chem.*, 308 (1986) 63.
- [295] A. Behr, W. Keim and G. Thelen, *J. Organomet. Chem.*, 249 (1983) C38.
- [296] R. Neidleim, A. Rufinska, H. Schwager and G. Wilke, *Angew. Chem., Int. Ed. Engl.*, 25 (1986) 641.
- [297] E. Carmona, P. Palma, M. Paneque, M.L. Poveda, E.G. Puebla and A. Monge, *J. Am. Chem. Soc.*, 108 (1986) 6424.
- [298] G.L. Monica, M.A. Angaroni and G.A. Ardizzoia, *J. Organomet. Chem.*, 348 (1988) 279.
- [299] A. Behr and G. Thelen, *C₁ Mol. Chem.*, 1 (1984) 137.
- [300] A. Behr, G. von Ilesmann, *J. Organomet. Chem.*, 276 (1984) C77.
- [301] T.W. Coffindaffer, I.P. Rothwell and J.C. Huffman, *Inorg. Chem.*, 23 (1984) 1433.
- [302] B. Dell'Amico, F. Calderazzo, U. Giurlani and G. Pelizzi, *Chem. Ber.*, 120 (1987) 955.
- [303] A. Belforte, D.B. Dell'Amico and F. Calderazzo, *Chem. Ber.*, 121 (1988) 1891.
- [304] M.B. Hursthouse, M.A. Malik, M. Motevalli and P. O'Brien, *J. Chem. Soc., Chem. Commun.*, (1991) 1690.
- [305] D.J. Darensbourg, K.M. Sanchez and A.L. Rheingold, *J. Am. Chem. Soc.*, 109 (1987) 290.
- [306] G.G. Sadler and T.P. Dasgupta, *Inorg. Chim. Acta*, 130 (1987) 185.

- [307] Y. Wakatsuki, M. Maniwa and H. Yamazaki, *Inorg. Chem.*, 29 (1990) 4204.
- [308] J.L. Newman and R.G. Bergman, *J. Am. Chem. Soc.*, 107 (1985) 5314.
- [309] S. Ganguly, J.T. Mague and D.M. Roundhill, *Inorg. Chem.*, 31 (1992) 3831.
- [310] R.A. Michelin, G. Strkul, N. Bresciani-Pahor, E. Zangrando and L. Randaccio, *Inorg. Chim. Acta*, 84 (1984) 2561.
- [311] M. Henary and J.I. Zink, *J. Am. Chem. Soc.*, 110 (1988) 5582.
- [312] G. Davies, M.A. El-Sayed and M. Henary, *Inorg. Chem.*, 26 (1987) 3266.
- [313] M. Kato and T. Ito, *Inorg. Chem.*, 24 (1985) 504.
- [314] M. Kato and T. Ito, *Inorg. Chem.*, 24 (1985) 509.
- [315] J. Arnold and T.D. Tilley, *J. Am. Chem. Soc.*, 109 (1987) 3318.
- [316] B.K. Campion, J. Falk and T.D. Tilley, *J. Am. Chem. Soc.*, 109 (1987) 2049.
- [317] D.J. Darensbourg, C.G. Bauch, J.H. Reibenspies and A.L. Rheingold, *Inorg. Chem.*, 27 (1988) 4203.
- [318] F.H. Elsner, T.D. Tilley, A.L. Rheingold and S.J. Geib, *J. Organomet. Chem.*, 358 (1988) 169.
- [319] B.K. Campion, R.H. Heyn and T.D. Tilley, *Inorg. Chem.*, 29 (1990) 4356.
- [320] R.T. Baker, 192nd Natl. Meet. of the American Chemical Society, Anaheim, CA, American Chemical Society, Washington, DC, 1986, p. 96.
- [321] W.E. Buhro, M.H. Chisholm, K. Folting and J.C. Huffman, *Inorg. Chem.*, 26 (1987) 3087.
- [322] W.E. Buhro, M.H. Chisholm, J.D. Martin, J.C. Huffman, K. Folting and W.E. Streib, *J. Am. Chem. Soc.*, 111 (1989) 8149.
- [323] S.A. Cohen and J.E. Bercaw, *Organometallics*, 4 (1985) 1006.
- [324] E. Klei, J.H. Teuben, H.J. DeLiefdemeijer, E.J. Kwak and A.P. Bruins, *J. Organomet. Chem.*, 224 (1982) 327.
- [325] B. Demerseman, R. Mahe and P.H. Dixneuf, *J. Chem. Soc., Chem. Commun.*, (1984) 1394.
- [326] H.G. Alt, G.S. Herrmann, M.D. Rausch and D.T. Mallin, *J. Organomet. Chem.*, 356 (1988) C53.
- [327] H. Yasuda and A. Nakamura, *Angew. Chem., Int. Ed. Engl.*, 26 (1987) 723.
- [328] M. Vivanco, J. Ruiz, C. Floriani, A.C. Villa and C. Guastini, *Organometallics*, 9 (1990) 2185.
- [329] J.A. Corella and N.J. Cooper, *J. Am. Chem. Soc.*, 112 (1990) 2832.
- [330] R. Alvarez, E. Carmona, D.J. Cole-Hamilton, A. Galindo, E. Gutierrez-Puebla, A. Monge, M.L. Poveda and C. Ruiz, *J. Am. Chem. Soc.*, 107 (1985) 5529.
- [331] M. Brookhart, S.K. Noh and F. Timmers, *J. Organomet. Chem.*, 6 (1987) 1829.
- [332] H. Hoberg, K. Jenni, K. Angermund and C. Kruger, *Angew. Chem., Int. Ed. Engl.*, 26 (1987) 193.
- [333] H. Hoberg, K. Jenni, C. Kruger and E. Raabe, *Angew. Chem., Int. Ed. Engl.*, 25 (1986) 810.
- [334] M. Farmer, M.A. Khan and K.M. Nicholas, *J. Chem. Soc., Chem. Commun.*, (1988) 1834.
- [335] A. Behr, E. Herdfweck, W.A. Herrmann, W. Keim and W. Kipshagen, *J. Chem. Soc., Chem. Commun.*, (1986) 1262.
- [336] A. Behr, E. Herdfweck, W.A. Herrmann, W. Keim and W. Kipshagen, *Organometallics*, 6 (1987) 2307.
- [337] G. Burkhardt and H. Hoberg, *Angew. Chem., Int. Ed. Engl.*, 21 (1982) 76.
- [338] H. Hoberg, D. Schaefer and G. Burkhardt, *J. Organomet. Chem.*, 228 (1982) C21.
- [339] H. Hoberg and D. Schaefer, *J. Organomet. Chem.*, 238 (1982) 383.
- [340] H. Hoberg and D. Schaefer, *J. Organomet. Chem.*, 236 (1982) C28.
- [341] H. Hoberg and D. Schaefer, *J. Organomet. Chem.*, 251 (1983) C51.
- [342] H. Hoberg, D. Schaefer, G. Burkhardt, C. Kruger and M.J. Romao, *J. Organomet. Chem.*, 266 (1984) 203.
- [343] H. Hoberg, Y. Peres and A. Milchereit, *J. Organomet. Chem.*, 307 (1986) C41.
- [344] H. Hoberg, Y. Peres and A. Milchereit, *J. Organomet. Chem.*, 307 (1987) C38.
- [345] H. Hoberg, Y. Peres, C. Kruger and Y.-H. Tsay, *Angew. Chem., Int. Ed. Engl.*, 26 (1987) 771.
- [346] D. Walther, E. Dinjus, J. Sieler, L. Andersen and O. Lindqvist, *J. Organomet. Chem.*, 276 (1984) 99.
- [347] D. Walther, E. Dinjus, J. Sieler, N.N. Thanh, W. Schade and I. Laben, *Z. Naturforsch.*, 38b (1983) 835.
- [348] H. Hoberg, and D. Schaefer, *J. Organomet. Chem.*, 255 (1983) C15.
- [349] H. Hoberg, D. Schaefer and B.W. Oster, *J. Organomet. Chem.*, 266 (1984) 313.
- [350] D. Walther, E. Dinjus, H. Goerls, J. Sieler, O. Lindqvist and L. Andersen, *J. Organomet. Chem.*, 286 (1985) 103.

- [351] W. Schroeder and K.R. Poerschke, *J. Organomet. Chem.*, 322 (1987) 385.
- [352] H. Hoberg and B. Apotecher, *J. Organomet. Chem.*, 270 (1984) C15.
- [353] A. Behr and U. Kanne, *J. Organomet. Chem.*, 317 (1986) C41.
- [354] H. Hoberg and B.W. Oster, *J. Organomet. Chem.*, 266 (1984) 321.
- [355] H.O. Froehlich and H. Schreer, *Z. Chem.*, 23 (1983) 348.
- [356] G.R. Lee, J.M. Mahar and N.J. Cooper, *J. Am. Chem. Soc.*, 109 (1987) 2956.
- [357] P. Fu, M.A. Khan and K.M. Nicholas, *Organometallics*, 11 (1992) 2607.
- [358] J.-C. Tsai, M.A. Khan and K.M. Nicholas, *Organometallics*, 8 (1989) 2967.
- [359] J.-C. Tsai, M.A. Khan and K.M. Nicholas, *Organometallics*, 10 (1991) 29.
- [360] R. Alvarez, J.L. Atwood, E. Carmona, P.J. Perez, M.L. Poveda and R.D. Rogers, *Inorg. Chem.*, 30 (1991) 1493.
- [361] J.C. Bryan, S.J. Geib, A.L. Rheingold and J.M. Mayer, *J. Am. Chem. Soc.*, 109 (1987) 2826.
- [362] T. Ito, S. Sugimoto, T. Ohki, T. Nakano and K. Osakada, *J. Organomet. Chem.*, 428 (1992) 69.
- [363] T.V. Sysyak, I.S. Kolomnikov and Yu. Ya. Kharitonov, *Koord. Khim.*, 9 (1983) 1000.
- [364] H.E. Ferkul, J.M. Berlie, D.J. Stanton, J.D. McCowan and M.C. Baird, *Can. J. Chem.*, 61 (1983) 1306.
- [365] W.R. Hasting, C.J. Cameron, M.J. Thomas and M.C. Baird, *Inorg. Chem.*, 27 (1988) 3024.
- [366] R.J. Klinger and J.W. Rathke, *J. Am. Chem. Soc.*, 106 (1984) 7650.
- [367] B.P. Sullivan and T.J. Meyer, *Organometallics*, 5 (1986) 1500.
- [368] C.T. Tso and A.R. Cutler, *Organometallics*, 4 (1985) 1442.
- [369] W.C. Kaska, S. Nemeh, A. Shirazi and S. Potuznik, *Organometallics*, 7 (1988) 13.
- [370] J. Wu, P.E. Fanwick and C.P. Kubiak, *Organometallics*, 6 (1987) 1805.
- [371] D.L. Delaet, R. Del Rosario, P.E. Fanwick and C.P. Kubiak, *J. Am. Chem. Soc.*, 109 (1987) 754.
- [372] M. Aresta, E. Quaranta and I. Tommasi, *J. Chem. Soc., Chem. Commun.*, (1988) 450.
- [373] T. Tanase, S. Nitta, S. Yoshikawa, K. Kobayashi, T. Sakurai and S. Yano, *Inorg. Chem.*, 31 (1992) 1058.
- [374] M.M.T. Khan, S.B. Halligudi and S. Shukla, *J. Mol. Catal.*, 5 (1989) 47.
- [375] J.-C. Tsai and K.M. Nicholas, *J. Am. Chem. Soc.*, 114 (1992) 5117.
- [376] D. Darensbourg and C. Ovalles, *J. Am. Chem. Soc.*, 106 (1984) 3750.
- [377] D. Darensbourg, C. Ovalles and M. Pale, *J. Am. Chem. Soc.*, 103 (1983) 5937.
- [378] D. Darensbourg and C. Ovalles, *J. Am. Chem. Soc.*, 109 (1987) 3330.
- [379] J.J. Anderson, D.J. Drury, J.E. Hamlin and A.G. Kent, WO 85-GB-443, 1985.
- [380] A.H.A. Tinnemans, T.P.M. Koster, D.H.M.W. Thewissen and A.M. Macker, *Recl. Trav. Chim. Pays-Bas*, 103 (1984) 288.
- [381] M. Beley, J.P. Collin, R. Ruppert and J.P. Sauvage, *J. Chem. Soc., Chem. Commun.*, (1984) 1315.
- [382] M. Beley, J.P. Collin, R. Ruppert and J.P. Sauvage, *J. Am. Chem. Soc.*, 108 (1986) 7461.
- [383] S. Kapusta and N. Hackerman, *J. Electrochem.*, 131 (1984) 1511.
- [384] C.M. Lieber and N.S. Lewis, *J. Am. Chem. Soc.*, 106 (1984) 5033.
- [385] P.A. Christensen, A. Hamnett and A.V.G. Muir, *J. Electroanal. Chem. Interfacial Electrochem.*, 24 (1988) 361.
- [386] J.Y. Becker, B. Vainas, R. Eger and L. Kaufman, *J. Chem. Soc., Chem. Commun.*, (1985) 1471.
- [387] X. Cao, Y. Mu, M. Wang and L. Luan, *Huaxue Xuebao*, 44 (1986) 220.
- [388] H. Tanabe and K. Ohno, *Electrochim. Acta*, 32 (1987) 1121.
- [389] K. Kusuda, R. Ishihara, H. Yamaguchi and I. Izumi, *Electrochim. Acta*, 31 (1986) 657.
- [390] N. Furuya and K. Matsui, *J. Electroanal. Chem.*, 271 (1989) 181.
- [391] N. Furuya and S. Koide, *Electrochim. Acta*, 36 (1991) 1309.
- [392] T. Atoguchi, A. Aramata, A. Kazusaka and M. Enyo, *J. Chem. Soc., Chem. Commun.*, (1991) 156.
- [393] M. Nakazawa, Y. Mizobe, Y. Matsumoto, Y. Uchida, M. Tesuka and M. Hidai, *Bull. Chem. Soc. Jpn.*, 59 (1986) 809.
- [394] K. Tanaka, R. Wakita and T. Tanaka, *Chem. Lett.*, (1987) 1951.
- [395] T. Tomohiro, K. Uoto and H. Okuno, *J. Chem. Soc., Chem. Commun.*, (1990) 194.
- [396] J. Hawecker, J.M. Lehn and R. Ziessel, *J. Chem. Soc., Chem. Commun.*, (1983) 536.
- [397] J. Hawecker, J.M. Lehn and R. Ziessel, *J. Chem. Soc., Chem. Commun.*, (1984) 328.
- [398] J. Hawecker, J.M. Lehn and R. Ziessel, *Helv. Chim. Acta*, 69 (1986) 1990.

- [399] B.P. Sullivan, C.M. Bolinger, D. Conard, W.J. Vining and T.J. Meyer, *J. Chem. Soc., Chem. Commun.*, (1985) 1414.
- [400] M.R.M. Bruce, E. Megehee, B.P. Sullivan, H. Thorp, T.R. O'Toole, A. Downart and T.J. Meyer, *Organometallics*, 7 (1988) 238.
- [401] H. Ishida, K. Tanaka and T. Tanaka, *Chem. Lett.*, (1985) 405.
- [402] H. Ishida, H. Tanaka, K. Tanaka and T. Tanaka, *J. Chem. Soc., Chem. Commun.*, (1987) 131.
- [403] H. Ishida, K. Tanaka and T. Tanaka, *Organometallics*, 6 (1987) 181.
- [404] C.M. Bollinger, B.P. Sullivan, D. Conard, J.A. Gilbert, N. Story and T.J. Meyer, *J. Chem. Soc., Chem. Commun.*, (1985) 796.
- [405] T.R. O'Toole, L.D. Margerum, T.D. Westmoreland, W.J. Vining, R.W. Murray and T.J. Meyer, *J. Chem. Soc., Chem. Commun.*, (1985) 1416.
- [406] C.R. Cabrera and H.D. Abruna, *J. Electroanal. Chem.*, 209 (1986) 101.
- [407] S. Cosnier, A. Derozier and J.C. Moutet, *J. Electroanal. Chem.*, 207 (1986) 315.
- [408] C.M. Bollinger, N. Story, B.P. Sullivan and T.J. Meyer, *Inorg. Chem.*, 27 (1988) 4582.
- [409] H. Ishida, K. Tanaka and T. Tanaka, *Chem. Lett.*, (1988) 339.
- [410] G. Grunwald and R.S. Drago, *J. Chem. Soc., Chem. Commun.*, (1988) 1206.
- [411] C. Arana, S. Yan, M. Keshavarz-K, K.T. Potts and H.D. Abruna, *Inorg. Chem.*, 31 (1992) 3680.
- [412] A.R. Guadalupe, D.A. Usifer, K.T. Potts, H.C. Hurrell, A.-L. Mogstad and H.D. Abruna, *J. Am. Chem. Soc.*, 110 (1988) 3462.
- [413] K.T. Potts, D.A. Usifer, A.R. Guadalupe and H.D. Abruna, *J. Am. Chem. Soc.*, 109 (1987) 3961.
- [414] H.D. Abruna, *Inorg. Chem.*, 28 (1989) 1080.
- [415] S. Daniele, P. Ugo, G. Bontempelli and M. Fiorani, *J. Electroanal. Chem.*, 219 (1987) 259.
- [416] H. Ishida, H. Tanaka, K. Tanaka and T. Tanaka, *Chem. Lett.*, (1987) 597.
- [417] S. Slater and J.H. Wagenknecht, *J. Am. Chem. Soc.*, 106 (1984) 5367.
- [418] D.L. Dubois and A. Miedaner, *J. Am. Chem. Soc.*, 109 (1987) 113.
- [419] A. Szymaszek and F.P. Pruchnik, *J. Organomet. Chem.*, 376 (1989) 133.
- [420] K.S. Ratliff, R.E. Lentz and C.P. Kubiak, *Organometallics*, 11 (1992) 1986.
- [421] D.L. Delaet, P.E. Fanwick and C.P. Kubiak, *J. Chem. Soc., Chem. Commun.*, (1987) 1412.
- [422] F.R. Lemke, D.L. Delaet, J. Gao and C.P. Kubiak, *J. Am. Chem. Soc.*, 110 (1988) 6904.
- [423] K. Ogura and H. Arima, *Chem. Lett.*, (1988) 311.
- [424] R.L. Cook, R.C. MacDuff and A. Sammells, *J. Electrochem. Soc.*, 135 (1988) 3070.
- [425] L. Willner and D. Mandler, *J. Am. Chem. Soc.*, 111 (1989) 1330.
- [426] S. Ikeda, Y. Saito, M. Yoshida, H. Noda, M. Maeda and K. Ito, *J. Electroanal. Chem.*, 260 (1989) 335.
- [427] M.W. Raphael and M.A. Malati, *J. Photochem. Photobiol. A*, 46 (1989) 367.
- [428] J.O'M. Bockris and J.C. Wass, *J. Electrochem. Soc.*, 136 (1989) 2521.
- [429] P. Fu, M.A. Khan and K.M. Nicholas, *Organometallics*, 10 (1991) 382.
- [430] K.A. Belmore, R.A. Vanderpool, J.-C. Tsai, M.A. Khan and K.M. Nicholas, *J. Am. Chem. Soc.*, 110 (1989) 2004.
- [431] W. Ziegler and K.M. Nicholas, *J. Organomet. Chem.*, 423 (1992) C35.
- [432] N.D. Silawe, A.S. Goldman, R. Ritter and D.R. Tyler, *Inorg. Chem.*, 28 (1989) 1231.
- [433] C. Kutal, M.A. Weber, G. Ferraudi and G. Geiger, *Organometallics*, 4 (1985) 2161.
- [434] H. Hukkanen and T.T. Pakkanen, *Inorg. Chim. Acta*, 114 (1986) L43.
- [435] C. Kutal, A.J. Corbin and G. Ferraudi, *Organometallics*, 6 (1987) 553.
- [436] N. Kitamura and S. Tazuke, *Chem. Lett.*, (1983) 1109.
- [437] J. Hawecker, J.M. Lehn and R. Ziessel, *J. Chem. Soc., Chem. Commun.*, (1985) 56.
- [438] J.M. Lehn and R. Ziessel, *J. Organomet. Chem.*, 382 (1990) 157.
- [439] H. Ishida, K. Tanaka and T. Tanaka, *Chem. Lett.*, (1987) 1035.
- [440] C.K. Tu and D.N. Silverman, *J. Am. Chem. Soc.*, 108 (1986) 6065.
- [441] J.L. Grant, K. Goswami, L.O. Spreer, J.W. Otvas and M. Calvin, *J. Chem. Soc., Dalton Trans.*, (1987) 2105.