

Wednesday, 22 March 2006

16:00–16:45

POSTER SESSION Nursing

168

Poster

Improving the quality and efficiency of breast cancer follow-up: a randomised controlled trial investigating nurse-led follow-up

M.L. Kimman¹, C. Dehing², P. Falger³, M. Hebly³, M.F. Von Meyenfeldt³, P. Hupperets³, B. Gijzen⁴, P. Lambin^{1,2,3}, C. Dirksen³, L.J. Boersma^{1,2,3}.
¹Maastricht University, GROW-MAASTRO, Maastricht, The Netherlands; ²MAASTRO Clinic, Radiotherapy, Heerlen, The Netherlands; ³University Hospital Maastricht, Maastricht, The Netherlands; ⁴Comprehensive Cancer Centre Zuid-Limburg, Maastricht, The Netherlands

Aim: To start a multicenter trial in order to investigate the cost-effectiveness of different, reduced follow-up (F-up) strategies in curatively treated breast cancer patients (BCPs).

Methods: In literature, randomised trials studying nurse-led (telephone) F-up in patients with other malignancies, had higher satisfactory ratings and lower costs compared to usual F-up, with similar survival rates. In addition, literature showed that group interventions, when mainly focussed on educational and informative aspects, can improve quality of life. Based on these data, we proposed to set-up a randomised multicenter trial with a 2x2 factorial design, investigating 4 different F-up strategies in the first 18 months after treatment: 1) standard F-up: 6 times in the first 18 months; 2) an outpatient F-up visit at 1 year, with 5 times nurse-led telephone F-up; 3) arm 1 combined with an educational group programme (EGP); 4) arm 2 combined with an EGP. The primary endpoint of this trial is cancer-specific QoL (CS QoL). Additionally, QoL, anxiety, feelings of control, patients' satisfaction and costs will be measured at randomisation, 3, 6, 12, and 18 months after inclusion. Based on the hypothesis that telephone F-up will not decrease CS QoL compared to usual F-up and that an EGP will show a clinically relevant increase in CS QoL, the sample size for the study was determined at 320 BCPs. Prior to start of the trial we developed the EGP and a training programme for nurses to perform telephone F-up.

Results: Fifteen nurses from 5 participating centers were successfully trained to perform telephone F-up by open discussion and a semi-structured questionnaire. An EGP was developed consisting of 2 group meetings for BCPs and their partners, given within 3 months after treatment. At this EGP, a nurse practitioner and health care psychologist discuss possible sequelae of the diagnosis and its treatment, using a standardized presentation, and information is given on where the patient can present problems if occurring. The EGP was tested in a group of 11 BCPs and partners, who were between 3 months and 5 years after treatment. Questionnaires filled out by the participants showed a 90% satisfactory rate.

In June 2005 the trial officially started; accrual is expected to be completed in 2 years. After 4 months, 32 out of approximately 100 eligible patients have signed informed consent of whom 4 have received their first telephone follow-up and 7 (with partners) joined the EGP. Reasons for refusal were 'wanting to see a doctor' (57%), 'no interest in the EGP' (22%) and 'too scared' or 'no interest' (21%). There have been no drop-outs.

Conclusion: We have successfully started a multicenter trial to study the cost-effectiveness of 4 different follow-up strategies. Apart from information on cost-effectiveness, this trial will also yield information on several psychosocial aspects.

169

Poster

An investigation of women's needs in and experiences of the provision, fitting and supply of external breast prostheses in Ireland

P. Gallagher¹, S. O'Carroll², A. Buckmaster¹, G. Kiernan¹, J. Geraghty³.
¹Dublin City University, School of Nursing, Dublin, Ireland; ²Action Breast Cancer, Irish Cancer Society, Dublin, Ireland; ³Tallaght Hospital, Department of Surgery, Dublin, Ireland

A good-quality prosthesis and prosthesis-fitting service is an essential part of the recovery process post-mastectomy. However, this is an area of care that has minimal information or research available to women. In Ireland, there has been no research in this area at all. In 2003, Healey argued that a rigorous, evidence based approach to the evaluation of external breast prostheses would enhance both the development of this service, as well as the adjustment, wellbeing and quality of life of breast cancer survivors. The overall aim of this research was to gain an insight into women's experiences of the provision, fitting, supply and use of breast prostheses in Ireland. The research had two stages, which involved women

requiring breast prostheses and the people who provide the service in order to affect the development of policy guidelines. The first stage used focus group methodology to investigate women's own personal and subjective experiences of the provision, fitting and supply of breast prostheses, and to gain a more complete understanding of the topic. Five focus groups were conducted with 47 women recruited through three national voluntary organisations and four Follow-up Breast Clinics throughout Ireland. The main findings emerging from the thematic analysis revealed the importance of the physical characteristics of the prosthesis; cost, affordability and entitlements; a lack of and perceived difficulty in getting information; the impact of the fitting experience, fitting environment and the qualities of the prosthesis fitter on a woman's experience in obtaining a first or replacement breast prosthesis; and the myriad of personal and social impacts of a breast prosthesis for the woman. Focus group data facilitated the development of a comprehensive questionnaire that was sent to 1240 women with a breast prosthesis. A questionnaire was also developed and sent to all breast care nurses (n=58), prosthesis fitters (n=20) and retail mastectomy bra fitting services (n=17) in Ireland. The findings from the surveys and the focus groups are facilitating the identification of factors that promote quality in the supply, fitting and aftercare of all breast prostheses and that impact on a woman's physical and psychosocial experience in obtaining and using a breast prosthesis. Overall, the paper will make recommendations for policy and practice in the fitting of breast prostheses, which, in turn, will ultimately benefit the women availing of these services.

170

Poster

Detecting risk factors by home-public health nurse in women with family history of breast cancer

A. Stanislawek¹, A. Bartoszek¹, H. Kachaniuk¹, M. Nowinska².

¹Skubiszewski Medical University of Lublin, Chair of Social Health Care, Lublin, Poland; ²Private Dermatology, Lublin, Poland

Women with family history of breast cancer require constant monitoring of an interdisciplinary team of therapists which also includes a home-public health nurse.

It is a home health nurse who detects factors increasing the risk of breast cancer in such women in 1st stage of nursing care. These factors can be divided as follows: factors which cannot be modified by a nurse's activities (these include facts which occurred earlier in a person's life: past diseases, education), factors which can be partially modified (e.g. family living conditions), factors that belong to the area of health behaviour which can undergo considerable direct modification (e.g. having prophylactic tests and examinations, breast-feeding, using hormonal contraception). Through prophylactic activities a nurse may eliminate or reduce such factors which are associated with health behaviour.

Material and Method: The aim of the study was to identify risk factors associated with hormonal activity and prophylaxis among women with family history of breast cancer. The study was performed by means of survey by questionnaire in the group of 70 women who had undergone tests for BRCA 1 gene mutations. Before genetic material collection each of the women gave her consent to the contact and co-operation with the authors of the study with the aim of receiving both medical and nursing care. In accordance with the aim of the study, the following factors were examined: age of menarche, pregnancy, miscarriage, birth of 1st child, breast-feeding period. In the area of prophylaxis the following factors were considered: the knowledge of the principles of breast self-examination (BSE) and BSE frequency, and the performance of such tests as: mammography, ultrasonography of the breast, ultrasonography of the ovaries, biopsy of the breast. The age of the women taken into the study varied from 23 to 84 years. University education was declared by 39% of the women, secondary education by 40%, and primary education by 21%. 45% of the women evaluated their financial and living conditions as good, 34% as average, and the remaining 21% as bad. The majority of the surveyed women lived in a town or city-76%.

Results: In order to examine risk factors associated with hormonal activity the surveyed women were divided into two groups: into women that had never been pregnant – 13 (19%), and who had been pregnant – 57 (81%). In the group of women who had been pregnant 65% were the women who had never had miscarriage, and out of the other 35%, 28% of the women had miscarried once and 7% – twice. Early age of menarche (before 12 years) was found in 21% of the women, however, the majority had their menarche after 12 years of age. Late primiparity after 35 years of age considerably increases breast cancer risk, but there were only 11% of such women in the studied group. Breast cancer risk reducing factor is the longest possible breast-feeding period whose total length after all births should exceed 18 months. Among the women who had been pregnant – 50 (88%) breast fed their babies. Among them, however, the majority were the women who performed breast-feeding for less than 18 months (62%). Prophylactic behaviour depends, among others, on women's knowledge. Most of the surveyed women 61(87%) declare the knowledge