

phos increase, hypokalemia, extremity pain, renal failure, urinary retention, pruritus, Gr 4 rash and purpura, and Gr 5 cerebrovascular accident at 160 mg/m<sup>2</sup>; Gr 3 melena, Gr 4 GI hemorrhage and thrombocytopenia at 200 mg/m<sup>2</sup>. The Gr 4 thrombocytopenia was a DLT. Rash developing in 3 patients after several cycles at 160 mg/m<sup>2</sup> has led to an ongoing evaluation of 140 mg/m<sup>2</sup>. Three patients have had stable disease for >3 months. Preliminary PK data document rapid conversion of GMX1777 to GMX1778 at all dose levels tested to date, and that exposure increases in a more than dose-proportional manner.

**Conclusions:** GMX1777 can be administered as a 24 hour infusion with a tolerable safety profile. The dose-limiting toxicities of thrombocytopenia and skin rash may be modified by the use of alternative schedules.

419

POSTER

**A phase I safety and pharmacokinetic (PK) study of 3 and 6 hours (h) intravenously administered belinostat (PXD101) plus carboplatin (C) and paclitaxel (P) in patients (pts) with advanced solid tumours**

M. Sorensen<sup>1</sup>, J. Tjørnelund<sup>2</sup>, P.B. Jensen<sup>2</sup>. <sup>1</sup>Rigshospitalet Copenhagen University Hospital, Dept of Oncology, Copenhagen, Denmark; <sup>2</sup>TopoTarget, NA, Copenhagen, Denmark

**Background:** Belinostat (Bel), an HDAC inhibitor of the hydroxamate class, has been well tolerated as a 30-min intravenous (iv) infusion in >100 pts treated at the MTD, 1000 mg/m<sup>2</sup>/day, days 1–5 q3 weeks. The main toxicity has been manageable nausea/vomiting. Prolonged exposure of tumor cell lines to Bel leads to improved cell kill in vitro, which was the rationale to prolong the daily exposure time in the clinic. The purpose of this study is to examine PK and safety of prolonged exposure of Bel.

**Methods:** The MTD for 30-min infusion of Bel (1000 mg/m<sup>2</sup>/day, days 1–5 q3 weeks) in combination with C (AUC 5) and P (175 mg/m<sup>2</sup> (delivered on day 3 has been determined (Sinha et al ASCO 2007). Based on a PK model derived from 30-min Bel infusion data, the present study included cohorts of 3–6 pts receiving C and P together with Bel administered as a 3 or 6 h iv infusion. Assessments included safety evaluations (NCI-CTCAE) and standard PK determinations. Acetylation status of histones H3 and H4 was assessed by Western blotting of extracted histones from peripheral blood mononuclear cells.

**Results:** The Bel PK model ( $V_1 = 0.015 \text{ ml/m}^2$ ;  $K_{21} = 2.082096 \text{ h}^{-1}$ ;  $K_{31} = 0.158794 \text{ h}^{-1}$ ;  $a = 5.6 \text{ h}^{-1}$ ;  $b = 1.21 \text{ h}^{-1}$ ;  $g = 0.156 \text{ h}^{-1}$ ) predicted that extension of the infusion time to 3 or 6 h would lead to biologic effective plasma concentrations (~1 mM) for ~5–7.5 h without accumulation of Bel. Seven pts were treated, 4 in the 3-h cohort (1 not evaluable due to early withdrawal) and 3 in the 6-h cohort. No major increase in clinical toxicity was seen as compared to pts treated with 30-min infusions, and no DLTs were observed. Initial PK data for the 3-h cohort confirm the simulation model with Bel concentrations remaining above biological active concentrations for 5.4 h. Further PK (including 6-h cohort) and clinical data will be presented at the meeting.

**Conclusion:** The combination of Bel with an infusion time of 3 or 6 h, with C and P was well tolerated, and the prolonged infusion of Bel resulted in an increased time above a biological effective plasma concentration as compared to 30-min infusion.

420

POSTER

**A phase I clinical trial of belinostat (PXD101) in combination with doxorubicin (BelDox) in advanced solid tumours, including soft tissue sarcomas (STS)**

A.T. Brunetto<sup>1</sup>, A. Krarup-Hansen<sup>2</sup>, O.S. Nielsen<sup>3</sup>, A. Norman<sup>4</sup>, A. Safwat<sup>3</sup>, J. Tjørnelund<sup>4</sup>, I. Judson<sup>1</sup>. <sup>1</sup>Royal Marsden Hospital, NHS Trust Cancer Research, London, United Kingdom; <sup>2</sup>Herlev Hospital, Department of Oncology, Herlev, Denmark; <sup>3</sup>Aarhus University Hospital, Department of Oncology, Aarhus, Denmark; <sup>4</sup>TopoTarget, Medical Department/ Drug Metabolism and Pharmacokinetics, Copenhagen, Denmark

**Background:** Belinostat (Bel) is a hydroxamate HDAC inhibitor with broad spectrum activity in vitro and in vivo and synergy in combination with doxorubicin (Dox). In Phase I studies in patients (pts) Bel is well-tolerated at a dose of 1000 mg/m<sup>2</sup> IV daily x 5, q 21 days. This study will determine the maximum tolerated dose (MTD), safety and pharmacokinetics (PK) of Bel plus doxorubicin (BelDox) in advanced solid tumours.

**Materials and Methods:** Sequential cohorts of 3–6 pts are planned to receive Bel at doses of 600, 800 and 1000 mg/m<sup>2</sup> daily x 5 plus Dox day 5, initially 50 mg/m<sup>2</sup> then 75 mg/m<sup>2</sup>, q21. PK studies, ECG monitoring for QTc prolongation, assessment of toxicity by CTCAEv3 and tumor imaging evaluated by RECIST was performed in all pts.

**Results:** 15 pts so far have been treated at 3 dose levels: 600/50, 600/75, 800/75 mg/m<sup>2</sup> of BelDox. 8 pts had received 0–1 previous treatment

regimens and 7 pts had received 2–4 regimens. Median age was 63 years (range 49–70) and the m:f ratio was 1.5: 1. The median number of cycles was 4.5 (range:1–7 cycles). Grade 3/4 adverse events (AE) included neutropenia in 5 pts, neutropenic fever in 2 pts, fatigue in 2 pts, and diarrhea, hand-foot syndrome, gram neg. bacteriemia and anemia in one pt each. The most common AE's were neutropenia, fatigue and nausea. Grade 4 neutropenia in cycle 1 was observed in 4 out of 11 pts in the 75 mg/m<sup>2</sup> Dox cohorts, and 3 of these pts had more than 2 prior regimens. There was one dose limiting toxicity (DLT) at the 600/75 cohort in a heavily pre-treated pt and 3 DLTs at 800/75. 3 out of 4 pts with DLT were heavily pretreated with other diagnoses rather than STS (i.e. pancreatic, renal and esophagus cancers). Currently, pts with no or limited previous treatment are being enrolled at 800/75. To date, the best clinical response observed has been partial response in cervical cancer. Mean values±SD for t<sub>1/2</sub>, AUClast and Cmax for Bel Day 4 were in cohort 600/50 (n=2): 1.2±0.1 hr, 11±2 hr µg/ml and 22±5 µg/ml; cohort 600/75 (n=6): 1.1±0.4 hr, 12±3 hr µg/ml, 20±8 µg/ml and cohort 800/75 (n=3) 1.4±2.6 hr, 12±4 hr µg/ml, 20±10 µg/ml, respectively. Bel PK results for Day 5 was similar to Day 4. **Conclusion:** Tolerance of BelDox was influenced by prior treatment. The regimen was best tolerated in patients with no or limited pre-treatment. Safety evaluation is ongoing in this population and a further escalation is planned.

421

POSTER

**Phase I dose escalation and pharmacokinetic study of oral enzastaurin in Japanese patients with advanced solid tumour**

T. Mukohara<sup>1</sup>, S. Nagai<sup>2</sup>, Y. Nambu<sup>3</sup>, K. Yoshizawa<sup>3</sup>, H. Minami<sup>1</sup>. <sup>1</sup>Kobe University Graduate School of Medicine, Medical Oncology, Kobe, Japan; <sup>2</sup>National Cancer Center Hospital East, Hematology/Oncology, Kashiwa, Japan; <sup>3</sup>Eli Lilly Japan K.K., TA Medical, Kobe, Japan

**Background:** Enzastaurin (ENZ) targets the protein kinase C and phosphoinositide 3-kinase/AKT pathways to inhibit tumor cell proliferation, induce tumor cell apoptosis, and suppress tumor-induced angiogenesis. A phase I study was conducted in Japan to evaluate the tolerability of ENZ and pharmacokinetic (PK) profile in patients (pts) with advanced solid tumor, and determine the recommended dose (RD) of ENZ for the phase II study.

**Material and Methods:** Eligible pts had incurable solid tumors, a PS of 0–2, and adequate organ function. Pts received ENZ once daily until disease progression (PD) or unacceptable toxicity occurred (1 cycle = 28 days). A six patient cohort for each dose level was used with a starting dose of 250 mg daily. The level was escalated in a stepwise manner based on the incidence of Dose Limiting Toxicities (DLTs). DLT was defined as any of the following: grade (G) 3, 4 or 5 non-hematologic toxicity, G4 neutropenia or thrombocytopenia, febrile neutropenia, G2 QTc prolongation lasting at least 1 week, or dose omission lasting at least 8 days in Cycle 1. Maximum tolerated dose (MTD) was defined as the lowest dose level where the incidence of DLTs was 33% or higher.

**Results:** 23 pts (7 pts: 250 mg, 6 pts: 375 mg, 6 pts: 500 mg, 4 pts: 750 mg) with median age of 55.0 years were enrolled and received ENZ. The major tumor types were non-small-cell lung cancer (n=5) and breast cancer (n=3).

Of the 23 pts, 4 pts discontinued the study for PD in Cycle 1 and 19 pts were evaluable for DLT (5: 250 mg, 6: 375 mg, 5: 500 mg, 3: 750 mg). No DLT was reported at doses of 500 mg or less, while 2 DLTs (G2 QTc prolongation) were observed at 750 mg. Based on these results, 750 mg ENZ was determined as the MTD and 500 mg as the RD for the phase II study.

The most common toxicities in the treated pts were chromaturia (n=15) and somnolence (n=9), which were all G1. The incidence of somnolence increased dose-dependently, however, no exacerbation was observed when the treatment was continued. The only G3 toxicity was lymphocytopenia (n=1). 3 pts (375 mg: 1, 750 mg: 2) required dose omission for G2 QTc prolongation. These pts showed signs of recovery from the event during dose omission while the events occurred again after receiving ENZ.

Multiple daily doses of 500 mg achieved the target plasma concentration (1400 nmol/L); the predicted average drug concentration of ENZ and its active metabolites at steady-state ranged from 1680 to 5040 nmol/L. Although none of the 23 pts achieved CR or PR in accordance with RECIST criteria, 9 pts (1: 250 mg, 4: 375 mg, 3: 500 mg, 1: 750 mg) achieved SD. A mesothelioma pt showed radiographical improvement.

**Conclusions:** Treatment with a daily dose of ENZ up to 500 mg was well tolerated in Japanese pts. Based on the incidence of DLTs (2 pts: G2 QTc prolongation) the RD of ENZ for the phase II study was determined as 500 mg/day for a 28-day-cycle. PK results support 500 mg as the RD to achieve the target plasma concentration of ENZ and its metabolites.