

As housekeeping we use 18S (Primer Design Co., Ltd.), such as was previously depicted.

**Results:** We have selected for this study three T-UCRs: uc.277, uc.301, and uc.440. We have found that the amount of these T-UCRs is generally low. T-UCR uc.440 was found over-expressed (nearly 16 times) in MDAMB231 breast cancer cell line versus healthy tissue ( $p < 0.01$ ). T-UCR uc.277 is also slightly over-expressed in MDAMB231 ( $p < 0.05$ ). In addition, T-UCR uc.440 was found slightly over-expressed in HT29 colon cancer cell line regarding to the others analyzed colon cancer cell lines ( $p < 0.05$ ), but there weren't found differences regarding to healthy tissue. It could not be detected significant expression of the T-UCR uc.301 in any tumour cell line. In melanoma cell lines could not be detected the expression of any assayed T-UCRs.

**Conclusion:** Our goal was to find a T-UCR with a constant and high expression in tumour cell lines regarding to healthy controls. We have found that T-UCR uc.440 is significantly upregulated in MDAMB231 breast cancer cell line regarding to healthy control, but also regarding to the others breast cancer cell lines analyzed. No relevant changes in the expression for the others T-UCRs tested were observed. For our knowledge, this is the first study where T-UCRs are assayed in colon, breast and melanoma tumour cell lines.

### P99

#### Rho C in melanoma: possible target for statin treatment?

L. Brochez<sup>1</sup>, T. Malfait<sup>1</sup>, B. Boone<sup>1</sup>, J. Lambert<sup>1</sup>, R. Kiss<sup>2</sup>, V. Mathieu<sup>2</sup>.  
<sup>1</sup>Dermatology Departement University Hospital Gent, Belgium; <sup>2</sup>Institute of Pharmacy ULB, Belgium

**Background:** There is large interest in statins as agents in cancer prevention and treatment. In more than 1300 melanoma cases statin use was associated with reduced Breslow thickness. Gene expression profiling experiments in melanoma cell lines revealed RhoC to be upregulated in cell lines with high metastatic capacity. We demonstrated that RhoC immunohistochemical expression in primary cutaneous melanoma was strongly associated with thicker and ulcerated tumors. Specific inhibition of Rho C in the A375M melanoma cell line reversed migration and invasion. Statins can block activation of RhoC by blocking geranylgeranyl pyrophosphate and farnesyl pyrophosphate which are important for isoprenylation of Rho proteins. Atorvastatin reverted the metastatic phenotype of several human melanoma cell lines and inhibited in vivo metastasis in SCID mice injected with A375M melanoma cells.

We explored if the statin simvastatin can influence the growth rate of melanoma cells of the BLM melanoma cell line.

**Materials and Methods:** The effect of different concentrations simvastatin (50, 500 and 5000 nM) on the global growth rate of BLM melanoma cell lines was investigated and compared to untreated BLM cells (control). By using 12 semi-automatic phase-contrast microscopes the relative increase of BLM cells was measured at 24, 48 and 72 hours.

**Results:** The simvastatin 5000 nM concentration significantly reduced global growth rate of the BLM cells after 48 and 72 hours ( $p < 0.001$ ) with stabilization of the cell population. There were no significant differences in global growth rate of the 50 nM and 500 nM simvastatin groups compared to control.

**Conclusion:** These observations support a growth inhibitory effect of high concentrations of simvastatin on the BLM melanoma cell line. Further research will focus on the determination of growth inhibitory concentrations of different statins on different melanoma cell lines.

### P10

#### Detection of circulating tumor cells (CTCs) in Stage cT3-4 or N+ rectal cancer patients (pts) undergoing combined neoadjuvant therapy plus curative surgery. Preliminary data

G. Contino<sup>1</sup>, E. Bertani<sup>1</sup>, G.M. Zampino<sup>2</sup>, L. Zorzino<sup>3</sup>, E. Magni<sup>2</sup>, B. Andreoni<sup>1</sup>, M.T. Sandri<sup>3</sup>, A. Chiappa<sup>1</sup>. <sup>1</sup>Div. of General and Laparoscopic Surgery, European Institute of Oncology, Milan, Italy; <sup>2</sup>Medical Oncology, European Institute of Oncology, Milan, Italy; <sup>3</sup>Laboratory Medicine, European Institute of Oncology, Milan, Italy

**Background:** CTCs detected at baseline and at disease-evaluation time-point during treatment seem to be an independent prognostic factor in metastatic colorectal cancer (Cohen SJ JCO, 2008). CTCs' role as predictive marker in early stages after radical surgery is under investigation, while no data are available in locally advanced rectal cancer suitable for neoadjuvant chemoradiotherapy. Aim of the study is to investigate the role of CTCs in local advanced rectal cancer pts undergoing neo-adjuvant chemo-radiotherapy (CT-RT).

**Materials and Methods:** In a prospective single institution study, cT3-4 or N+ rectal cancer pts staged by transrectal ultrasound and/or pelvic MRI and chest-abdomen CT scan, are submitted to capecitabine

(825 mg/mq, orally, twice daily continuous) with concomitant radiotherapy (50.4 Gy/fractions to the primary tumor and perirectal nodes), followed by two cycles of capecitabine (1250 mg/mq, orally, tid 14/21 days). Primary endpoint is evaluation of CTCs at baseline (t0), after neoadjuvant therapy, before surgery (t1), after surgery (t2), and at 6-month follow-up (t3) and its correlation with survival parameters. CTCs are enumerated with immunomagnetic separation in 7.5 ml peripheral blood at over-mentioned time-points (CellSearch System, Veridex Inc).

**Results:** Twenty-six pts (16M; 10F; median age: 63±13 yrs; range: 44–83 yrs) underwent t0 sampling, 8 pts completed CT/RT and therefore underwent t1 and t2 sampling. At baseline (t0) three pts presented 1 CTC (12%), one 2 CTCs (3.5%), one 27 CTCs (3.5%) while in twenty-one (81%) no CTCs were detected. At t1 and t2 none of the eight pts analyzed showed CTCs.

**Conclusion:** CTCs ≥1 are present in 15% of our patients, but the sample is too small for statistical analysis. The study is still ongoing; more data will allow to assess prognostic and predictive significance of CTCs during treatment in this setting.

### P30

#### Assessment of oxidative stress in tumor cells and normal mucosa cells from head and neck squamous cell carcinoma patients

D. Dequanter, K. Zouaoui, V. Nuyens, A. Rousseau, D. Brohée, M. Vanhaeverbeek, Ph. Lothaire. CHU Charleroi – Hôpital André Vésale, Belgium

**Background:** One of the cancers particularly linked to oxidative damage and oxidative stress is the head and neck squamous cell carcinoma (HNSCC). Tobacco and alcohol are well defined etiological factors.

Concurrent radiation and chemotherapy, a recognized alternative treatment to surgery for patients with advanced HNSCC, can induce a systemic oxidative stress. Oxidative damage is the main mechanism mediating the clinical effect of radiotherapy, and an increased resistance to oxidative stress by malignant cells is associated with treatment failure. Response to (chemo) radiation treatment varies from patient to patient.

The purpose of this study was to compare the tissue levels of glutathione in HNSCC tumoral tissue (Tum) and adjacent normal mucosa (Muc) biopsies as a potential factor of variability in (chemo) radiosensitivity.

**Materials and Methods:** 27 newly diagnosed HNSCC patients were prospectively studied. All were current smokers. 27 tumoral biopsies and an equal number of biopsies from normal mucosa were analysed. The oxidised/reduced glutathione ratio was measured with the capillary electrophoresis Ceofix GSH/GSSG kit (Analisis, Namur, Belgium). Two hundred µl of whole blood, normal and tumoral tissues were immediately grinded with 600 µl of 5% metaphosphoric acid. After centrifugation (within 3 hours), 100 µl of the supernatant was mixed with 400 µl of the kit diluent containing naphthalene sulfonic acid as an internal standard. Analysis was done on a P/ACE 5000 series with a 37 cm and 75 µm i.d. capillary maintained at 25°C. The separation was realized at 8 kV with a pH 8.2 borate buffer containing SDS. The glutathione peaks were detected at 200 nm and integrated as under-the-curve areas (AUC). The results are expressed as the ratio of oxidised GSSG AUC to the reduced GSH AUC. Clinico-pathological parameters were also considered as potential factors of variability in oxidative stress status of HNSCC tumoral tissue.

**Results:** The GSSG to GSH ratio was higher in the tumoral tissue than in the adjacent normal tissue in 12/27 of the cases.

**Conclusion:** In 44% of our cases, HNSCC tumoral tissue from untreated patients had a GSSG/GSH ratio different than that found in the normal adjacent cancer free mucosa. This difference was not related with clinico-pathological parameters. Heterogeneity in HNSC cancer is emphasized. The ability of the GSSG/GSH ratio to predict differential chemoradiosensitivity will be evaluated by long-term survival data.

### P34

#### Reduced p63 expression in myoepithelium correlating with increased invasiveness in epithelium

Y.-H. Hsiao<sup>1</sup>, C. Deng<sup>2</sup>, Y.A. Su<sup>3</sup>, H.-D. Tsai<sup>4</sup>, X. Wang<sup>5</sup>, J.T. Mason<sup>6</sup>, M.-C. Chou<sup>1</sup>, Y.-g. Man<sup>7</sup>. <sup>1</sup>Institute of Medicine, Chung Shan Medical University, Taiwan; <sup>2</sup>Mammalian Genetics Section, GDBB, NIDDK, National Institutes of Health, USA; <sup>3</sup>George Washington University School of Medicine and Health Sciences, USA; <sup>4</sup>Department of Obstetrics and Gynecology, Changhua Christian Hospital, Taiwan; <sup>5</sup>The George Washington University School of Medicine and Health Sciences, USA; <sup>6</sup>Department of Biophysics, Armed Forces Institute of Pathology, USA; <sup>7</sup>Armed Forces Institute of Pathology and American Registry of Pathology, USA

**Background:** Our previous studies revealed that pregnancy associated breast cancer (PABC) had significantly reduced nuclear p63 expression