

patient groups had poorer functioning and more depression ( $P < 0.001$ ). Among the patient groups, the LD group reported lower scores for body image ( $P = 0.007$ ) and future perspective ( $P = 0.023$ ) than the BCS group. In the LD group, patients who received neoadjuvant chemotherapy reported lower scores for future perspective and higher scores for depression than those who did not receive neoadjuvant chemotherapy ( $P < 0.001$ ).

**Conclusion:** The BCS and LD groups did not differ in oncologic outcome, and the QoL of patients in the LD group was not always good. Mastectomy with immediate reconstruction should be considered carefully and tailored to the patient's needs and characteristics.

## 273

**Skin reducing mastectomy**

Poster

N. Bruno<sup>1</sup>, A.E. Pennati<sup>1</sup>, J. Ottolenghi<sup>1</sup>, A. Spano<sup>1</sup>, V. Visintini<sup>1</sup>, G. Catanuto<sup>1</sup>, M.B. Nava<sup>1</sup>. <sup>1</sup>Fondazione IRCCS Istituto Nazionale Tumori, U.O. Chirurgia Plastica e Ricostruttiva, Milan, Italy

**Background:** we propose an alternative technique to reconstruct large and medium sized ptotic breasts in a single stage procedure using anatomical permanent implants and a dermal-muscle pocket.

**Material and Methods:** we enrolled 60 patients fulfilling the oncological criteria for skin-sparing mastectomy and presenting with large or medium size breasts whose areola-to-inframmary fold distance was more than 8 cm and notch-to-areola distance was more than 25 cm. All operations were performed as a single-stage procedure. After a careful preoperative assessment, conventional incisions for breast reduction (wise pattern) were used with the preservation of an inferior dermal flap. Subsequently this was sutured to the inferomedial fibers of the pectoralis major muscle to create a dermo-muscular pocket. An anatomical prosthesis was then allocated into the pouch, that was closed laterally with the serratus anterior fascia. Skin flaps were finally closed down to the inframmary fold.

**Results:** a total of 71 procedures were performed on 60 patients. The medium size of the anatomical implants employed was 444.3 cc. A contra-lateral procedure was carried out on 37 patients. In 83.7% of them a reduction of the contralateral breast was performed. At a median follow-up of 35 months, the implant removal rate was 16.9%. The overall complication rate was 22.5% (infection-skin necrosis). When the procedure was accomplished in smokers a higher, although not statistically significant, complication rate was demonstrated (27.2% smokers vs. 22.7% non smokers).

**Conclusion:** The results of this study confirm our previous reports about implant based breast reconstruction. All breast, irrespectively of mammary shape and size, can be reconstructed with medium size implants and, if required, contralateral adjustments. A one stage procedure could also allow immediate radiotherapy on permanent prosthesis if post-operatively required. The overall complication rate is in keeping with data on inverted T mastectomies reported in literature. Improvement in this field could be obtained with more restricted inclusion criteria. It is advisable to avoid this procedure in smokers or women with skin of poor quality. Alternatively a temporary expander in the first stage of a two stage procedure may avoid tension and sufferings of the dermal flap at the T junction.

## 274

**A prospective observational study on the oncological outcome and patient satisfaction with skin sparing mastectomy and immediate breast reconstruction**

Poster

S. Al-Reefy<sup>1</sup>, N. Patani<sup>2</sup>, A. Anderson<sup>3</sup>, G. Burgoyne<sup>3</sup>, C. Chao<sup>3</sup>, H. Osman<sup>4</sup>, K. Mokbel<sup>5</sup>. <sup>1</sup>London Breast Institute, Oncoplastic Breast Surgery, London, United Kingdom; <sup>2</sup>St. Georges University, General Surgery, London, United Kingdom; <sup>3</sup>London Breast Institute, Breast Surgery, London, United Kingdom; <sup>4</sup>St Georges University, General Surgery, London, United Kingdom; <sup>5</sup>London Breast Institute & St Georges University, Breast Surgery, London, United Kingdom

**Introduction and Background:** The management of early breast cancer (BC) with skin-sparing mastectomy (SSM) and immediate breast reconstruction (IBR) is not based on level-1 evidence. In this study, the oncological outcome, post-operative morbidity and patients' satisfaction with SSM and IBR using the latissimus dorsi (LD) myocutaneous flap and/or breast prosthesis is evaluated.

**Methods:** 137 SSMs with IBR (10 bilateral) were undertaken in 127 consecutive women, using the LD flap plus implant ( $n = 86$ ) or implant alone ( $n = 51$ ), for early BC ( $n = 130$ ) or prophylaxis ( $n = 7$ ). Nipple reconstruction was performed in 69 patients, using the trefoil local flap technique ( $n = 61$ ), nipple sharing ( $n = 6$ ), skin graft ( $n = 1$ ) and Monocryl mesh ( $n = 1$ ). Thirty patients underwent contra-lateral procedures to enhance symmetry, including 19 augmentations and 11 mastopexy/reduction mammoplasties. A linear visual analogue scale was used to assess patient satisfaction with surgical outcome, ranging from 0 (not satisfied) to 10 (most satisfied).

**Results:** After a median follow-up of 36 months (range = 6–101 months) there were no local recurrences. Overall breast cancer specific survival was 99.2%, 8 patients developed distant disease and 1 died of metastatic BC. There were no cases of partial or total LD flap loss. Morbidities included infection, requiring implant removal in 2 patients and 1 patient developed marginal ischaemia of the skin envelope. Chemotherapy was delayed in 1 patient due to infection. Significant capsule formation, requiring capsulotomy, was observed in 87% of patients who had either post-mastectomy radiotherapy (PMR) or prior radiotherapy (RT) compared with 13% for those who had not received RT. The outcome questionnaire was completed by 82 (64.6%) of 127 patients with a median satisfaction score of 9 (range = 5–10).

**Conclusion:** SSM with IBR is associated with low morbidity, high levels of patient satisfaction and is oncologically safe for T(is), T1 and T2 tumours without extensive skin involvement.

## 275

**Extended indications for skin conservation of the nipple areola complex during subcutaneous mastectomy**

Poster

M. Niemeier<sup>1</sup>, R. Backe<sup>2</sup>, S. Paepke<sup>1</sup>, D. Paepke<sup>2</sup>, B. Niemeier<sup>2</sup>, M. Kiechle<sup>2</sup>. <sup>1</sup>Klinikum rechts der Isar, Gynaekology/Senology, Munich, Germany; <sup>2</sup>Klinikum rechts der Isar, Gynaekology, Munich, Germany

**Background:** Conventional breast cancer surgery in case of a non breast conserving procedure still includes the routine excision of the nipple areolar complex (NAC) optional followed by nipple reconstruction. Numerous authors consider multiple predictive factors to decide whether the Nipple Areola Complex (NAC) can be conserved or not during mastectomy. These are: tumour-nipple distance, tumour size, axillary lymph-node status and lymph vascular invasion. Thus only a very limited percentage of patients can keep their NAC.

If the breast gland tissue and all galactophoric ducts can be separated completely from the NAC during subcutaneous mastectomy (SCM), conservation of the NAC is feasible even in large, central and retroareola tumours.

**Patients and Methods:** (www.clinicaltrials.com ID: NCT00641628) From July 2003 to May 2006, we performed 109 SCMs on 96 patients. 94 of these breasts showed indications for modified radical mastectomy (MRM) or ablatio simplex. 78 due to invasive carcinoma (MRM indication) and 16 of them had extensive ductal carcinoma in situ (DCIS), (indication for simple ablation). At least 33 of the breasts had malignancy underneath the skin within the areola margin (central located tumours). After dissection of the complete mammary gland tissue, the skin envelope with the areola is folded inside out an all galactophoric ducts and any subjacent tissue on the areola base are precisely dissected under the surgeons' visual control. Of this skin-closest retroareola tissue, frozen sections and HE-histopathologic examinations are requested to decide whether the NAC can be preserved or not.

**Results:** We found the Need to dissect the NAC in 13 of 109 breasts (12%), altering the procedure to a skin sparing mastectomy (SSM). Necrosis of the NAC which had to be subjected to surgical intervention occurred in only one of the conserved 96. After a follow-up of 20–54, median 34 months no recurrence within the NAC was observed. One local recurrence on the thoracic wall and one axillary recurrence were detected. 2 of 96 patients developed distant metastases. One death was recorded. Occasionally, partial necrosis of the nipple occurred, leaving a depigmentation of the skin but maintaining a good or excellent cosmetic result in most cases. Necrosis of the NAC which had to be subjected to surgical intervention occurred in only one patient.

**Conclusion:** SCM, including NAC conservation, may be performed according to MRM indications if an intraoperatively frozen section (and the corresponding HE-histopathology) of the NAC closest tissue is free of tumour. The remaining contraindications for SCM are: extensive tumour involvement of the skin, inflammatory breast cancer and clinical suspicious nipple.

## 276

**Criteria of oncoplastic approach of local advanced breast cancer after neoadjuvant chemotherapy**

Poster

A.G. Zucca Matthes<sup>1</sup>, R.A.C. Vieira<sup>1</sup>, G. Uemura<sup>1</sup>, R.L. Haikel<sup>1</sup>, M.A.A.K. Folgeira<sup>1</sup>, C.M.S. Fregnani<sup>1</sup>. <sup>1</sup>Hospital do Cancer de Barretos, Senology and Breast Reconstructive Surgery, Barretos, Brazil

**Introduction and Background:** It is considered that the neoadjuvant chemotherapy (NC) offers several advantages in the treatment of local advanced breast carcinoma (LABC), especially as the possibility of conservative surgical treatment. Yet the literature is scarce and controversial and this study seeks to clarify criteria that indicate an oncoplastic treatment after neoadjuvant therapy.