

clinical signs of recurrence or metastasis, knowing that more sensitive means of early diagnosis are available.

Methods: We retrospectively selected women on the database of the PET center of the Bordet Institute. Only were selectionated patients with proven breast cancer who had undergone prior curative intent treatment; an asymptomatic elevation of the circulating tumor marker (CA15-3 and/or CEA); no known recurrent disease; and a follow-up for at least 12 months. PET(CT) images were assessed by 2 experienced nuclear medicine specialists. Both were blinded to the results of the other observer and to the medical records of the patients. A pre-established patient-based classification was used. Finally, both investigators in consensus assessed the results.

Results: Twenty-four asymptomatic women were included. Previously performed diagnostic imaging revealed no evidence of tumour at all. Mean age was 61.1 years. Median CEA level was 10.0 ng/ml (reference >2.5 ng/mL) and median CA 15-3 level was 100.4 U/ml (reference >30.0 U/mL). Four patients presented elevated levels in both tumour markers. Ten patients presented only CEA elevated levels. The follow-up of 12 months after the date of the PET demonstrated a tumoral recurrence in thirteen patients (54.2%). PET(CT) was true positive in ten cases, false positive in one patient and false negative in three cases. The patient-based sensitivity and specificity of PET(CT) was 77% and 91% respectively. The negative predictive value was 77%; the positive predictive value was 91% and accuracy of 83%.

Conclusion: Our data demonstrates the usefulness of PET/CT in the follow-up of asymptomatic women suffering from breast cancer who have elevated levels of tumour markers, by providing an accurate method for detecting metastases and/or an alternative diagnostic for example a second primary. Nowadays, combined PET/CT seems to be the best imaging modality (all in one) detecting in a reliable way disease recurrence in the follow-up of women breast cancer.

593

Poster

Accuracy of breast cancer detection with full-field digital mammography (FFDM) and integral computer-aided detection (CAD) correlated with breast density as assessed by a new automated volumetric breast density measurement system

K. Pinker¹, S. Milner², D. Stephen³, K. Mokbel², N.M. Perry². ¹Medical University Vienna, Department of Radiology, Vienna, Austria; ²The Princess Grace Hospital, The London Breast Institute, London, United Kingdom; ³Cancer Research Centre of Epidemiology Mathematics and Statistics, Wolfson Institute of Preventive Medicine Barts and the London School of Medicine and Dentistry, London, United Kingdom

Background: To retrospectively assess the diagnostic performance of computer-aided detection (CAD) for full-field digital mammography (FFDM) in terms of sensitivity and specificity correlated with breast density as assessed by an automated breast density measurement system in histopathologically proven breast cancers and age-matched healthy controls.

Materials and Methods: 200 consecutive histopathologically proven breast cancers imaged with FFDM (Senographe DS or GE Essential, GE Healthcare) and 200 age-matched healthy controls identified from June 2005 through February 2009 were evaluated retrospectively using CAD (Hologic R2, version 8.3.17). Each case included a craniocaudal and mediolateral oblique view. Each cancer case was matched to one control case by date of birth, age at examination and laterality of mammogram used for density determination. In all cancer cases the malignant lesion was identifiable in at least one projection on radiologist review. A CAD mark was scored true-positive, if it correctly indicated a malignant lesion. All other CAD marks were considered false. CAD sensitivity and specificity were calculated and correlated with mammographic breast density (percentage of fibroglandular tissue) as assessed by an integral automated volumetric breast density measurement system (Hologic, Quantra).

Results: CAD correctly identified 157 of the 200 cancers, a sensitivity of 79%. Sensitivity was suggestively but non-significantly lower with increased density ($p=0.09$). In those cancer cases with density at or below the median of 20%, sensitivity was 82%, compared to 75% in those with density above the median. The presence of one or more false CAD prompts was suggestively but not significantly more likely in controls than cases (87% vs 80%, $p=0.06$). The number of false prompts was significantly higher in controls (average 3.6 vs 2.6, $p<0.001$). False prompts were significantly less likely with higher density ($p=0.008$). False prompts were present in 86% of cases and controls with density at or below the median, and in 81% of those with density above the median.

Conclusion: Increased breast density is significantly associated with higher specificity of CAD, and there is suggestive evidence that it is also associated with lower sensitivity. Radiologists should be aware that CAD is likely to be less sensitive in mammography screening of younger women with denser breast tissue.

594

Poster

Quantitative multivoxel magnetic resonance spectroscopy in 13 breast lesions, a pilot study

M.D. Dorrius¹, R.M. Pijnappel², M.C. Jansen-van der Weide¹, P. Kappert¹, M. Oudkerk¹, P.E. Sijens¹. ¹University Medical Center Groningen, Radiology, Groningen, The Netherlands; ²Martini Hospital, radiology, Groningen, The Netherlands

Background: The purpose of this study is to determine whether a new quantitative multivoxel Magnetic Resonance Spectroscopy (MRS) can differentiate benign from malignant breast lesions by measuring the highest choline concentration.

Materials and Methods: Twelve patients with 13 breast lesions (7 mammographic BIRADS-3 lesions and 6 mammographic BIRADS-4 lesions) underwent an MRI and MRS at 1.5 Tesla using a Magnetom Avanto system (Siemens, Erlangen). The multivoxel MRS technique used was 2D-chemical shift imaging (CSI) with point resolved spectroscopy (PRESS), first without suppression of the water and fat signals (repetition time (TR) 1500ms, echo time (TE) 30ms) to serve as a reference measurement, and subsequently with suppression of the water and fat signals (TR 1500ms, TE 135ms) to be able to detect choline. The choline concentrations were measured in 36 voxels of $0.5 \times 0.5 \times 1 \text{ cm}^3$.

Results: The highest choline concentrations for the seven mammographic BIRADS-3 lesions were found in 2 to 4 voxels: 0.05mM, 0.13mM, 0.13mM, 0.10mM, 0.15mM, 0.15mM and 0.23mM, respectively. The work-up of the last one (0.23mM) showed invasive ductal carcinoma, the other mammographic BIRADS-3 lesions showed no malignancy. The highest choline concentrations for six mammographic BIRADS-4 lesions were found in 4 to 6 voxels: 0.22mM, 0.41mM, 0.46mM, 0.25mM, 0.68mM and 0.10mM, respectively. The last lesion with 0.10mM highest choline concentration was not malignant. Pathology confirmed malignancy in all other mammographic BIRADS-4 lesions: 4 lesions were invasive ductal carcinomas and 1 lesion was a medullary carcinoma.

Conclusion: The new quantitative multivoxel MRS method can differentiate benign from malignant lesions by indicating that a breast lesion with a highest choline concentration exceeding 0.20mM is malignant.

595

Poster

Comparison of synchrotron images of Paget's disease of the breast with their pathologic findings

S. Park¹, J. Bong¹, J. Hong¹, H. Wang¹, H. Kim², J. Kim³, H. Oh⁴, S. Jheon⁵, J. Huang⁶. ¹Daegu Catholic University Hospital, Department of General Surgery, Daegu, Korea; ²Daegu Catholic University Hospital, Department of Anatomy, Daegu, Korea; ³Daegu Catholic University Hospital, Department of Radiology and Biomedical Engineering, Daegu, Korea; ⁴Daegu Catholic University Hospital, Department of Pathology, Daegu, Korea; ⁵Seoul National University, Department of Thoracic and Cardiovascular Surgery, Seoul, Korea; ⁶Pohang University of Science and Technology, Pohang Accelerator Laboratory, Pohang, Korea

Background: Synchrotron radiation x-ray imaging has revealed its possibilities to evaluate various breast diseases non-invasively. Using a phase contrast technique, we received monochromated synchrotron images of Paget's disease of the breast tissue section. To figure out the relation with their optical microscopic features, we compared the synchrotron images of the Paget's disease with their histopathologic findings of the same stained section.

Material and Methods: An x-ray microscope was installed on 1B2 beamline of Pohang Light Source, a third generation synchrotron radiation facility with operating energy of 2.5 GeV in Pohang, Korea. The x-ray energy was set at 11.1 keV, and the x-ray beam was monochromatized by a W/B4C monochromator. Zernike phase-shifter was adapted for phase contrast x-ray microscopy. Formalin-fixed 5µm-thick breast tissue sample was attached onto the Kapton film, and positioned 25 m away from the beam source. The synchrotron image of the sample was converted into a visual image on the CsI(Tl) scintillation crystal, and this visual image was captured by a full frame CCD camera. After scanning, we put together these images one another to show the large area of the tissue section. For the comparative analysis with their synchrotron image, synchrotron-scanned breast tissue sections were stained, and the histopathologic findings of the samples were captured by the image analyzer. To identify the association of synchrotron image with its histopathologic findings, we compared them with each other. The magnifying power of this microscope was 100x.

Results: The monochromated x-ray microscopic images of Paget's disease of the breast tissue sections were obtained with a good contrast and high visibility by phase contrast technique. These images showed the large lacuna-shaped Paget cells within the epidermis of nipple. The Paget cells revealed large electron-dense nuclei and electron-lucent abundant