

Results: Of the 18 community based programmes carried out (2006–09) 886 completed the questionnaire and its factor analysis revealed: limited knowledge/lack of education (n = 250), Limited funds (n = 200), Difficulty in accessing facilities (n = 135), family priorities (n = 122), unhelpful attitude of health workers (n = 103), religious/spiritual delays (n = 76). Out of 886 females who completed the questionnaire, 421 attended the breast screening service giving an uptake of 50% compared with an uptake of 35% in previous screening rounds (2003–06). Sample age was 20–55 yrs and 78% were married. Clinical breast examination was carried out in all while mammogram was done in (n = 324). Cases detected were ductal carcinoma in situ (n = 2) and cancer breast (n = 2). Pathologically both were infiltrating ductal carcinoma. Tumor staging: Stage II: n = 1, Stage IIIB: n = 1.

Conclusion: Factor analysis emphasizes regular screening awareness programmes, providing education, allocation of funds, accessible health care and trained health workers as major interventions to increase uptake of breast cancer screening, and to diagnosis this disease early.

624

Poster

Physical examination is a valuable tool in the follow up of young women with a history of early breast cancer

G. de Bock¹, W.L. Lu¹, M. Schaapveld², P.C. Baas³, L.M. Hansen⁴, H.C. van der Mijle⁵, T. Wiggers⁶, L. Jansen⁶. ¹Groningen University Hospital, Department of Epidemiology, Groningen, The Netherlands; ²Comprehensive Cancer Centre, Northeast-Netherlands, Groningen, The Netherlands; ³Martiniziekenhuis, Department of Surgery, Groningen, The Netherlands; ⁴Diaconessenhuis, Department of Surgery, Meppel, The Netherlands; ⁵Nije Smellinghe, Maatschap Heelkunde Zuid Friesland, Drachten, The Netherlands; ⁶Groningen University Hospital, Department of Surgery, Groningen, The Netherlands

Background: Regular physical examination is recommended in follow up guidelines for breast cancer patients. The objective of this study is to assess the contribution of physical examination in addition to mammography in the early diagnosis of breast cancer recurrences.

Methods: The medical follow-up documents of 669 patients were reviewed. 127 contra-lateral breast cancers and 65 loco-regional recurrences in 169 patients were included. The contribution of physical examination over mammography was evaluated with the proportions of loco-regional recurrences or contra-lateral breast cancers detected by physical examination alone and were assessed stratified for type of recurrences and surgical modalities. The potential impact of patients' age and time from first tumour on the contribution of physical examination was evaluated with Chi-square tests.

Results: Seven (5.5%) out of 127 contra-lateral breast cancer recurrences and 13 (20.0%) out of 65 loco-regional recurrences were detected by physical examination alone. The contribution of physical examination in detecting loco-regional recurrences was not statistically different between patients after mastectomy and patients after breast conserving treatment (25.9% vs. 15.8%; Chi-square=1.014, P=0.314). There was a trend that the contribution of physical examination is higher in women under 60 years of age than in patients over 60 years of age (14.8% vs. 6.7%; Chi-square=3.304, P=0.069). There is no significant difference in the contribution of physical examination during the first 5 year and after the first 5 years since diagnosis of the primary tumour (8.0% vs. 13.3%; Chi-square=1.430, P=0.232).

Conclusions: Some breast cancer recurrences would have been detected later without physical examination. Physical examination has a higher contribution in younger patients (<60).

625

Poster

The effectiveness of breast cancer screening with MRI and mammography in women with a BRCA1/2 mutation

G. de Bock¹, F.S. Hassankhan¹, J.C. Oosterwijk², M.J. Mourits³, L. Jansen⁴. ¹Groningen University Hospital, Department of Epidemiology, Groningen, The Netherlands; ²Groningen University Hospital, Department Clinical Genetics, Groningen, The Netherlands; ³Groningen University Hospital, Department of Gynecologic Oncology, Groningen, The Netherlands; ⁴Groningen University Hospital, Department of Surgical Oncology, Groningen, The Netherlands

Background: The objective of this study is to evaluate the effectiveness of screening with mammography and MRI in detecting breast cancer in BRCA1 or BRCA2 mutation carriers.

Methods: Women who were screened by a surgeon of the Family Cancer Clinic at least once in 2004–2006 were included. Breast cancer screening consisted of clinical breast examination twice a year and annual alternating MRI or mammography, where BIRADS ≥3 was considered as positive. Sensitivity, specificity, positive and negative predicting values (PPV and

NPV) as well as the number needed to screen (NNS) to detect 1 early stage breast cancer, were calculated.

Results: During the screening period 305 mammographies and 256 MRIs were performed in 173 consecutive BRCA1/2 carriers. A total of 13 invasive ductal carcinomas were found of which 3 prevalent, 5 interval and 5 screen-detected carcinomas. The screen-detected and prevalent carcinomas were all diagnosed in stage I/II. Of the 5 interval carcinomas 1 was in stage III. The sensitivities of mammography and MRI were 67% and 71%, respectively. The PPV of mammography and MRI was 60% and 12%, respectively. The NPV was 99% for both tests. The NNS to detect one breast cancer for mammography as well as for MRI were about 50.

Conclusion: MRI has a higher sensitivity than mammography. However, as there are still carcinomas detected with mammography only, mammography is still warranted in breast cancer screening. Given the early stages of detected breast cancers, the current screening policy of BRCA 1/2 mutation carriers seems effective.

626

Poster

The usefulness of telemammography using soft-copy computed radiography (CR) in screening program for Japanese women

T. Funakoshi¹, T. Sugimoto¹, Y. Nakauchi², F. Suehiro², M. Takechi³, Y. Okamoto³, W. Hamada³, Y. Okada³, K. Hanazaki¹. ¹Kochi Medical School, Surgery 1, Kochi, Japan; ²Kochi Kenshin Clinic, Screening, Kochi, Japan; ³Kochi Kenshin Clinic, Radiology, Kochi, Japan

Background: Digital mammography machines are widely used across Japan, and the majority of them are computed radiography (CR) systems. Therefore, telemammography using soft-copy CR can compensate for the uneven national distribution of mammographers and radiologists in Japan. However, the evaluation of soft-copy interpretation of CR is still controversial because it is usually diagnosed using hard-copy. The purpose of this study is to elucidate the usefulness of telemammography using soft-copy CR in breast cancer screening.

Screenings and Methods: The mammograms of 18,549 screenees had been taken at Kochi Kenshin Clinic using Phase Contrast Mammography (PCM) systems (Konica Minolta Health Care Co. Ltd.) between July 2005 and September 2008. Digital data of them were made by Regius 190 (Konica Minolta), and then processed and compressed to the transferable sized soft-copy by Vox-base II (J-Mac System Inc.). Thereafter, the soft-copy CR were transferred to Kochi Medical School via optic fiber (provided by NTT West co.) and interpreted on 5 mega pixel monitor by the mammographers who were licensed by the central committee on quality control of mammographic screening.

Results: The recall rate was 6.1%, the cancer detection rate 0.33% and the positive predictive value 5.4%. These results were not inferior to those of Kochi prefectural screening program using analogue mammograms of 26,747 screenees (8.7%, 0.34% and 3.9%, respectively).

Conclusion: The long-term results of our telemammographic screening using soft-copy CR had not been inferior to those of analogue mammography. Telemammography using soft-copy CR might be one of promising strategies to overcome the uneven distribution of human resources that participate mammographic screening in Japan. Recently, we structured a new telemammography network between our institute and five screening facilities including Kochi Kenshin Clinic.

627

Poster

Ethnicity is a high risk for breast cancer: should we target screening of high-risk groups earlier?

M. Uzzaman¹, M.S. Nair¹, T. Wahab¹, O. Fafemi¹, A. Atthow¹. ¹North Middlesex University Hospital, General Surgery, London, United Kingdom

Background: Breast cancer screening in the United Kingdom is currently indicated in the 50–70 year age group. Previous studies have suggested that patients from minor ethnic groups, especially Afro-Caribbean's, have a worse prognosis after being diagnosed with breast cancer compared to the Caucasian population. We explored whether there are any ethnic differences in the incidence of breast cancer rates among women attending a busy breast clinic at a London University Hospital. We were especially interested in whether there are any benefits of screening the 41–50 year age group in the various ethnic groups.

Materials and Methods: This was a retrospective analysis identifying patients who attended the breast clinic with newly-diagnosed breast cancer. The data was retrieved from the Breast Cancer Registered Database. Our centre has observed one of the highest ethnic population attendance amongst the London Hospitals. The period of study was from March 2002 to March 2009. Patients with previous diagnosis of breast cancer, male patients and those patients not presenting at the Breast clinic were excluded from the study. From the data, the age specific breast cancer distributions (age 21–100 years) of various ethnicities were compared and statistically analysed. Odds Ratio, chi-squared, 95% confidence interval