

35.

HALF-BODY IRRADIATION (HBI) IN THE TREATMENT OF METASTATIC EWING'S SARCOMA RESISTENT TO CHEMOTHERAPY. M.Gasparini, F.Lombardi, A.Lattuada, F.Fossati-Bellani, C.Gianni. Istituto Nazionale per lo Studio e la Cura dei Tumori, 20133 Milan, Italy.

Upper and lower HBI in two subsequent sessions were delivered to 21 consecutive patients with disseminated Ewing's sarcoma relapsing after or while on radiotherapy and multidrug chemotherapy. A 6 MeV linear accelerator was utilized to deliver a midplane dose of 6 Gy in one single fraction. The interval between the two sessions varied according to the degree of myelosuppression, ranging from 4 to 8 weeks. The median age of patients was 16 years (range 8 to 47 years). Sites of metastatic deposits before HBI were as follows: multiple skeletal lesions (8 patients), multiple pulmonary metastases (7), one single bone lesion (4), disseminated bone and lung metastases (2). Thirteen of 21 patients (62%) received both sessions of radiotherapy. Only one session of HBI, either upper or lower, was given to 6 and 2 patients, respectively. Seventeen sessions were performed to treat painful metastases. A complete pain control was obtained in 15 cases (88%). This lasted for 2 to more than 17 months. Twenty-five of the 34 total sessions of HBI were employed to treat overt metastases. The overall response rate was 43%. At the time of the present analysis, 7 of 21 patients (33%) are alive 7 to 36 months from their first HBI session. Three patients are free from progressive disease. Besides a transient pneumonitis, which cleared after steroid therapy, no major or fatal complications were recorded. HBI resulted to be mostly effective in patients relapsing while off chemotherapy with metastases confined to the lungs or to one single bone segment. This is an ongoing clinical trial.

36.

TREATMENT BY SEQUENTIAL RADIOCHEMOTHERAPY OF SOFT TISSUE SARCOMA. RADIOBIOLOGICAL BASES AND PRELIMINARY CLINICAL RESULTS. C. Dionet, F. Deméocq. Centre Jean Perrin, Place Henri Dunant, 63011 Clermont-Ferrand Cédex, France

Based on cellular kinetics data obtained by monolayer cell culture, we have shown, on L1210 ascitic tumour in the mouse, the potentialisation of X-rays and of the chemotherapeutic association of 5-Fluorouracil (5-Fu) and cis-platinum (cis-DDP). We have developed a sequential treatment that can be applied in human therapy, and which has given us encouraging results in soft tissue sarcoma. The aim of the protocol is to treat, with a minimum of side effects, lesions that have resisted normal therapy. It is applied to patients who have never been treated with large doses of cis-platinum. The treatment takes place over 7 days and consists of: D₁ to D₇: 5-Fu at 370 mg/m²; D₁ and D₂ then from D₂ to D₇: cis-platinum 15 mg/m²; D₂ and D₄: X-rays, total dose 8 to 10 Grays depending on the extent and position of the lesions. 5-Fu is used as inhibitor of X-ray potential lethal damages (PLD) repair, and as potentialiser of cis-platinum (own work). Cis-DDP administered over several days is as effective as a single, large dose (own work). The choice of D₂ and D₄ for administration of X-rays allows maximum benefit of potentialisation of X-rays by cis-DDP and of that of cis-DDP by X-rays. This treatment is repeated after a rest period of 3 weeks. Disadvantages are cumulative dose of cis-DDP and long-term thrombopenia. Advantages are the practicability, which is the same as for sequential chemotherapy, and the tolerance, which is usually good both from the haematological and clinical points of view. The results are still in their preliminary stages, but we have always obtained a good response on soft tissue sarcoma: on pulmonary metastasis of synovial sarcoma, volume reduction was at least 80% after 3 cycles; an inoperable malignant abdominal schwannoma became, after 4 cycles, an entirely necrotic, operable mass; a pelvic leiomyosarcoma disappeared after 5 cycles. We are currently treating other cases, but it would be premature to discuss them at the moment. The encouraging results incite us to continue this therapy, and to extend it to other lesions.

37.

CHEMORADIO THERAPEUTIC CONSERVATIVE MANAGEMENT IN 23 PATIENTS WITH LOCALLY EXTENDED BILATERAL RETINOBLASTOMA. J.M. Zucker, N. Lemerrier, P. Schlienger, E. Margulis, C. Hays. Institut Curie.

To reduce the number of enucleated eyes and to improve the useful vision in locally advanced disease, we developed a conservative approach in 23 patients (pts), 9 boys and 14 girls aged of 3 to 36 mths (median = 8 mths) with bilateral non metastatic retinoblastoma. 36/46 eyes were Reese stage V and in 10/23 pts unilateral enucleation had been mandatory at first. All pts received two courses of VAC (vincristine 1.5 mg/m² day I, actinomycin D 10 gammas/kg day I to 5, cyclophosphamide 200 mg/m² day I to 5) at a three wks interval, followed by irradiation with a 22 MeV electron beam delivering 45 grays in 5 wks; after one mth, 6 monthly courses of VAC were resumed. **Results:** overall follow up is 12 to 54 mths (median = 24 mths). 2/23 pts died from neuromeningeal involvement at 18 and 22 mths. 21/23 pts are alive and well 2 to 48 mths after cessation of therapy (median = 14 mths). 5 of them are blind. 12 keep one eye, 4 keep both eyes. There was, on the basis of fundoscopy, a good partial response to initial chemotherapy in 9 patients, a moderate

decrease of the tumor in 8, no response without progression in 6.

. Later on 15 secondary enucleations -12/15 in the first 10 months- were done in 14/23 pts due to tumor progression in 11 cases. 7/14 pts kept one useful eye which was initially staged II to V in 4 cases.

. A useful vision was kept in 12/13 eyes (II/I2 stage V) of 9/23 pts (4/9 keeping both eyes).

Conclusions: 1°- These preliminary results are encouraging both in terms of visual capacity and cosmetics. 2° A careful fundoscopic supervision must be extended to detect late local relapses.

38.

MANAGEMENT OF RETINOBLASTOMA (R.B.) BY PRECISION MEGAVOLTAGE IRRADIATION. J. Schipper, K.E.W.P. Tan. Dept. of Radiotherapy Utrecht, The Netherlands.

The conservative management of R.B. in The Netherlands is centralized in Utrecht. The principal concept in the treatment of R.B. in this centre is radiation therapy (R.Th.) followed by lightcoagulation (L.C.) and/or cryotherapy (C.T.) if there is some doubt as to whether the tumour is still active. R.Th. is administered by means of a highly accurate megavoltage X-ray beam technique previously described (1). The dose of radiation is standardized at 45 Gy (4500 rad) given in 15 fractions of 3 Gy each, 3 fractions per week. Between 1971 and 1980, 31 children with R.B. have been irradiated to at least one eye. Of the 58 affected eyes, 16 were primarily enucleated, one was lightcoagulated only and 41 were irradiated. Of the 41 irradiated eyes, 27 were additionally treated by L.C. or C.T. and 7 were ultimately enucleated. The percentages of cure of the irradiated eyes with a minimum follow-up of two years were 100%(8/8), 100%(9/9), 75%(6/8), 85%(11/13), and 0%(0/3) in stages 1 to V, respectively. Twelve eyes developed a clinically detectable radiation cataract (R.C.); in 5 of these the lens was aspirated. R.C. developed exclusively in those lenses of which a major part was included in the treatment field. In a preliminary study (1) the likelihood and extent of cataract formation was found to be directly related with the dose of radiation to the germinative zone of the lens epithelium. Irradiation of a posterior portion of 1 mm of the human lens with a sharply edged irradiation beam will not produce a radiation cataract.

References: (1) Schipper, J.: Retinoblastoma. A medical and experimental study. Thesis, State University of Utrecht, 1980.

39.

INTESTINAL MICROBIAL FLORA SUPPRESSION IN PREVENTION OF INFECTIONS IN PATIENTS UNDERGOING CYTOTOXIC OR RADIATION THERAPY. F. Waldvogel, Department of Medicine, Geneva, Switzerland.

Besides tumor growth, or therapy induced destruction of the mucosal barrier, the absolute granulocyte count is one of the best predictors of bacterial or fungal infection in patients with malignancies. 50% of these infections are caused by flora, acquired subsequent to initial hospitalization. Major pathogens include *S. aureus*, *E. coli*, *Klebsiella species*, *Pseudomonas aeruginosa*, *Candida species* and *Aspergillus species*. Viruses contribute to the morbidity of these patients, but rarely to their mortality except for cytomegalovirus (bone marrow transplants). Herpes zoster/varicella virus (lymphomas), and the newly recognized gastrointestinal viruses.

Gastrointestinal bacterial and fungal suppression by non absorbable antibiotics has therefore been used extensively as a major means to prevent infections in neutropenic patients. Full suppression with large spectrum antibiotics has given contradictory results. Selective suppression with Co-trimoxazole, or partial suppression with a combination of Co-trimoxazole and an antifungal agents represents promising direction for the future. These results have to be compared to other possible means of preventing infections in granulocytopenic patients, such as reversed precautions, strict hygiene, vaccinations and antisera, and new compounds decreasing the colonization with pathogenic bacteria and fungi.

40.

INFECTIONS AND OTHER TOXIC DEATHS IN THE SECOND NATIONAL WILMS' TUMOR STUDY. B.Jones, N.Breslow, J.Takahashi, for the National Wilms' Tumor Committee, West Virginia