

exposed on the surfaces of syngeneic murine tumour cells of known malignancy and with well defined metastasis characteristics.

We used 5 lectins conjugated to either 125 IUDR or fluorescein isothiocyanate: Con A, WGA, PNA, SBA and UEA. The binding pattern was characteristically changed by treatment with the proteolytic enzyme (pronase) and with neuraminidase. The data were used as basis for attempts to separate cells in subfractions on Pharmacia Sepharose 6 MB columns. PNA (peanut agglutinin) was found to be the only suitable ligand in terms of cell yield and specificity.

SERUM TOCOPHEROL AND INCIDENCE OF CANCER IN FINLAND

M.Hakama(1,3), A.Aromaa(2), P.Knekt(2), J.Maatala(2), R.K.Aaran(3), T.Nikkari(3), T.Hakulinen(1), R.Peto(1), E.Saxen(1) and L.Teppo(1)

(1)Finnish Cancer Registry Helsinki, (2)The Social Insurance Institution Helsinki and Turku, (3)Departments of Biomedical Sciences and Public Health University of Tampere, Finland

The 35,000 persons belonging to the Finnish Social Insurance Institution Mobile Clinic Survey were linked to the Finnish Cancer Registry. During a follow-up of 6 to 10 years 766 cancer patients were identified. Stored serum samples were available for the patients and 1:2 matched controls. The alpha-tocopherol levels were higher for controls. The association was strongest for smoking unrelated cancers. It persisted after exclusion of cases diagnosed shortly (<2 years) after the serum sample was drawn as well as after adjusting for the confounding effects of smoking, cholesterol and socioeconomic status.

17 BETA-ESTRADIOL MEDIATES GLYCOSYLATION OF HUMAN BREAST CARCINOMA GROWTH FACTOR RECEPTOR (BCGF-R)

Anwar A.Hakim

Evanston Hospital, Evanston, and Loyola University Medical Center, Maywood Ill, U.S.A.

Malignant cells produce growth factors to which they respond because of specific receptors at their surface, thus conferring a growth advantage to these cells (Hakim, *Expt. Cell Biol.* 54: 193-211, 1986). Estradiol mediated the release of a serine protease (Hakim, *Cancer Biochem. Biophys.*, 4: 173-185, 1980) from breast carcinoma, and of an immunosuppressive agent from

malignant melanoma (Hakim, *Annales d'Immunol.*, 131C: 155-170, 1980). The present investigation reveals that estradiol modulates glycosylation of the growth factor receptor on human breast carcinoma cell membrane. Short term cultures were developed from biopsies of human breast carcinomas confirmed as ductal (BDC), lobular (BLC) and colloidal (BOC). These cell lines were grown in estrogen-free (EFM) medium and in EFM supplemented with 10^{-7} M estradiol and/or tunicamycin (Tn). The cells were examined for mitogenicity, clonogenicity, 125 I-EGF binding ability and 3 H-galactosamine uptake capacity. The cells were also grown in athymic mice. Extraction of the BCGF-R followed by alkaline borohydride treatment and chromatography on Sepharose-4B showed that BCGF-R from *in vitro* cultured cells in presence of estradiol, or grown in nude mice treated with estradiol contained significantly higher fucosyl- and sialyl BCGF-R.

TATI, CEA AND CA 125 IN HUMAN OVARIAN CYSTS

H.Halila(1), C.Haglund(2), S.Nordling(3) and U.-H.Sterman(1)

(1)I-II Departments of Obstetrics and Gynaecology; (2)IV Department of Surgery; and (3)Department of Pathology, University of Helsinki, Finland

Tumour-associated trypsin inhibitor (TATI) is a 6kD peptide isolated from the urine of an ovarian cancer patient. Highly increased excretion of this inhibitor has been found in the urine and serum of cancer patients. We have now examined the concentration of TATI in human ovarian cysts. Very high cyst fluid concentrations of TATI were found in all mucinous cystadenomas, both benign and malignant. The mean concentration was about 100-fold compared to serum levels. This strongly suggests that this tumour-associated peptide was actually produced by the tumour. In serous cyst fluids low levels of TATI were found, similar to concentrations in serum. By immunohistochemistry TATI was demonstrated in most benign mucinous cystadenomas and in some semi-malignant and malignant tumours. Positive staining was predominantly seen in the apical parts of the cells. All serous tumours were negative in TATI staining. Very high cyst fluid levels of CEA and CA 125 were also detected, CEA exclusively in the mucinous type, CA 125 in both mucinous and serous cysts.

TUMOUR PROMOTING ACTIVITY OF TGF β

E.Hamel, F.Katch, G.Muller(1), W.Birchmeier(1) and H.Yamasaki