

202

POSTER

### Assessment of pulmonary reaction after carbon ion irradiation in the patients with stage I non-small cell lung cancer

H. Nishimura<sup>1,2</sup>, T. Miyamoto<sup>1</sup>, N. Yamamoto<sup>1</sup>, M. Koto<sup>1</sup>, K. Sugimura<sup>2</sup>, H. Tujii<sup>1</sup>. <sup>1</sup>National Institute of Radiological Sciences, Research Center of Charged Particle Therapy, Chiba, Japan; <sup>2</sup>Kobe University, Department of Radiology, Kobe, Japan

Experiences about phase I/II clinical trial of carbon ion irradiation against stage I non-small cell lung cancer (NSCLC) was prospectively evaluated with especial respect to pulmonary reaction found on computed tomography (CT) in terms of treatment dose, volume and dose-volume histogram (DVH). 46 patients (35 male, 11 female) aged ranging from 49 to 82, with medically inoperable stage I NSCLC received carbon ion radiation from 1994 to 1998. The follow-up duration was 6 to 54 months (median:23). Since one patient had two different primaries, 47 NSCLCs were evaluated. 27 tumors were smaller than 3cm, while 20 tumors were ranging from 3 to 7cm. In the phase I/II study, the dose escalation in 18 fractions in 6 weeks was performed from 59.4 photon gray-equivalent dose (GyE) to 95.4GyE by the step of 10% increase. Clinical target volume (CTV) was determined to cover gross tumor volume with more larger than 5mm margin (median; 46.8ml). Prophylactic irradiation to the hilus and mediastinum was not applied. Chest CT was serially performed before and 1, 3, 6, 9, 12 and every 6 months after the end of treatment. CT findings classified into two discrete categories: "Pleural" (pleural thickening and/or effusion) and "Pulmonary" reaction. The pulmonary reactions were classified according to the modified scoring system proposed by Bush et al. (grade0: no remarkable change, grade1: slight increase in density with no consolidation, grade2: consolidation involves less than 50% of the irradiated portion, grade3: consolidation involves more than 50% of the irradiated portion, grade4: changes spreading beyond the boundaries of radiation field) (AJR, 1999;172:753-9). DVH was calculated for every patient and the volume irradiated more than 20GyE (V20) and 40GyE (V40) was used for analysis. Overall and cause-specific 5-year survivals were 48% and 66%, respectively (median follow up: 43 months) (Kaplan-Meier). Pleural reaction was observed in 23 patients (49%) at 1 to 12 months after the initiation of the treatment (median: 4 months). Such pleural reaction was observed in the patients with large CTV ( $r=0.552$ ,  $p=0.00018$ ). Pulmonary reaction was observed in 45 (98%) patients (grade0: 2, grade1: 14, grade2: 19, grade3: 12, grade4: 0). No correlation was found between the grade of pulmonary reaction and CTVs or irradiated doses, but the grade of pulmonary reaction showed a statistically significant correlation with V20 ( $r=0.4$ ,  $p=0.007$ ) and V40 ( $r=0.43$ ,  $p=0.0037$ ).

203

POSTER

### Impact of neoadjuvant chemotherapy on RT treatment planning for stage III NSCLC patients

J.A. Soriano, G.T. Henning, J.A. Hayman, M.K. Martel. University of Michigan Health System, Radiation Oncology, Ann Arbor, USA

**Purpose:** Patients with inoperable NSCLC who respond to neoadjuvant chemotherapy have a reduction in the tumor size as compared to the pre-chemotherapy volume, changing the targeting of the subsequent RT treatment plan. This may lead to a decrease in the amount of irradiated normal lung and surrounding normal tissues and a corresponding decrease in the risk of developing normal tissue toxicity. The aim of this study was to compare RT plans designed to target the pre- and post-chemotherapy tumor volumes and evaluate the clinical benefit with dose-volume parameters.

**Methods:** Ten patients with Stage III NSCLC underwent CT scan of the chest prior to and after the administration of a two month course of neoadjuvant chemotherapy consisting of cisplatin and Navelbine. These patients received an average of 73Gy (60-84) on a dose escalation trial. The gross tumor volume (GTV) (including the involved lymph nodes) decreased in size after chemotherapy in all patients an average (range) of 77cc (11-210). For each patient, the original treatment plan for the post-chemotherapy planning target volume (PTV) was compared to a plan with identical beam arrangement but with altered field shapes for the pre-chemotherapy PTV. Treatment planning goals included minimum target volume dose of 95% of the prescription dose, minimizing the amount of normal lung irradiated, limiting cord dose to <50Gy, esophagus volume (Veff) to 33% < 80Gy, and heart Veff to 100% and 33% < 40 and 65Gy respectively. For each patient, the changes in the lung dose-volume parameters such as mean lung dose, lung volume receiving >20 Gy (V20) and normal tissue complication probability (NTCP) were compared. NTCP was calculated with the Lyman, Kutcher, Burman method with Kwa's parameter fit (Kwa IJROBP 42, 1998)

**Results:** Targeting the post-chemotherapy PTV resulted in the following average decreases (range): mean lung dose by 4Gy (0-10), NTCP by 8% (0.3-17), V20 by 6% (0.2-13), maximum cord dose by 4Gy (0-15), esophagus Veff by 13% (0-74), whole heart Veff by 22% (0-65) and partial heart Veff by 5.5% (0-16).

**Conclusion:** Radiation treatment plans for patients who respond to neoadjuvant chemotherapy result in a clinically significant lower probability (8% on average) of developing radiation pneumonitis. Radiation doses to other surrounding normal tissues can also be decreased, specifically the spinal cord, esophagus and heart, thereby decreasing the risk of late complications involving these structures.

204

POSTER

### A phase I study of irinotecan and carboplatin with concurrent thoracic radiotherapy for unresectable stage III non-small cell lung cancer

T. Kurata<sup>1</sup>, M. Yamada<sup>2</sup>, S. Kudoh<sup>2</sup>, N. Yamamoto<sup>1</sup>, K. Nakagawa<sup>1</sup>, K. Takeda<sup>3</sup>, S. Negoro<sup>3</sup>, M. Fukuoka<sup>1</sup>. <sup>1</sup>Kinki University School of Medicine, Fourth Department of Internal Medicine, Osakasayama, Japan; <sup>2</sup>Osaka City University Medical School, First Department of Internal Medicine, Osaka, Japan; <sup>3</sup>Osaka City General Hospital, 3Department of Pulmonary Medicine, Osaka, Japan

**Purpose:** A phase I clinical trial was performed to evaluate the efficacy and toxicity of weekly irinotecan with daily carboplatin and concurrent thoracic radiotherapy in patients with locally advanced non-small-cell lung cancer (NSCLC).

**Patients and Methods:** Thirty-one previously untreated patients with unresectable stage III NSCLC were enrolled in a phase I study. Patients received weekly irinotecan plus carboplatin (20 mg/m<sup>2</sup> daily for 5 days a week) for 4 weeks and thoracic radiotherapy (60 Gy in 30 fractions). The irinotecan dose was escalated from 30 mg/m<sup>2</sup> in increments of 10 mg/m<sup>2</sup>.

**Results:** Four irinotecan dose levels were given and 30 patients were assessable. Their median age was 62 years (range: 52-72 years), 28 had a performance status of 0-1 and 2 had a performance status of 2, and 12 had stage IIIA disease and 18 had IIIB disease. There were 19 squamous cell carcinomas, 10 adenocarcinomas, and 1 large cell carcinoma. The dose-limiting toxicities were pneumonitis and thrombocytopenia. The maximum tolerated dose of irinotecan was 60 mg/m<sup>2</sup>, with 3 patients developing grade 4 pulmonary toxicity (1 died of pneumonitis). Three patients achieved complete remission and 15 had partial remission, for an objective response rate of 60.0%. The median survival time was 14.9 months, and the 1- and 2-year survival rates were 51.6% and 34.2%, respectively.

**Conclusion:** The major toxicity of this regimen was pneumonitis. This therapy may be active against unresectable NSCLC and a phase II study is warranted.

205

POSTER

### Phase II study of low-dose weekly paclitaxel as second-line treatment for advanced non-small cell lung cancer

O. Juan, V. Alberola, A. Rocher, R. Casany, V. Carañana, J.M. Campos, A. Albert, V. Alberola Jr. Hospital Arnau de Vilanova, Oncology, Valencia, Spain

**Purpose:** To investigate the activity and toxicity of low-dose weekly paclitaxel (WP) as single-agent in patients with advanced non-small cell lung cancer (NSCLC) who experienced recurrence or failure to previous chemotherapy regimen.

**Patients and methods:** From August-98 to August-00, twenty-nine patients with metastatic or unresectable NSCLC were treated with 80 mg/m<sup>2</sup> WP as 1-hour infusion, without rest period, until disease progression, unacceptable drug toxicity or after achievement of best response. The median age was 66 (range 42 to 81); 19 patients had ECOG-PS 1 and 10 PS 2. The predominant histology was squamous (21 patients); twenty-one patients had stage IV disease, 7 patients had stage IIIB and 1 had stage IIIA. Sixteen of the 29 patients had initially responded to platinum-based therapy, 6 patients had achieved stable disease (SD) and 7 had disease progression (DP) prior to the present chemotherapy regimen. The median time between the last dose of the previous chemotherapy and the start of the weekly paclitaxel was 25 weeks (range = 2 to 106 weeks).

**Results:** Two (6.9%) patients achieved a complete response (CR) and 8 (27.6%) partial response (PR) for an overall response rate of 34.5% (10 of 29 patients). There were also 12 (41.4%) cases of SD and 5 (17.2%) of DP. Median survival (intent to treat) was 51 weeks (95% CI: 29-73). Treatment was well tolerated. There were no episodes of grade 4