

available at www.sciencedirect.comjournal homepage: www.ejconline.com

Membrane type matrix metalloproteinases (MMPs) show differential expression in non-small cell lung cancer (NSCLC) compared to normal lung: Correlation of MMP-14 mRNA expression and proteolytic activity

J.M. Atkinson^a, C.J. Pennington^b, S.W. Martin^a, V.A. Anikin^c, A.J. Mearns^c, P.M. Loadman^a, D.R. Edwards^b, J.H. Gill^{a,*}

^aInstitute of Cancer Therapeutics, University of Bradford, Tumbling Hill Street, Bradford, West Yorkshire, BD7 1DP, UK

^bSchool of Biological Sciences, University of East Anglia, Norwich, Norfolk, NR4 7TJ, UK

^cDepartment of Cardiothoracic Surgery, Bradford Royal Infirmary, Bradford, West Yorkshire, BD9 6RJ, UK

ARTICLE INFO

Article history:

Received 25 September 2006

Received in revised form 10 May 2007

Accepted 14 May 2007

Available online 27 June 2007

Keywords:

Matrix metalloproteinases

Membrane-type metalloproteinases

Lung cancer

Real-time PCR

Carcinoma

Non-small cell lung

Cancer therapeutics

ABSTRACT

Improved understanding of the involvement of matrix metalloproteinases (MMPs), including membrane-type MMPs (MT-MMPs), in human tumours has potential diagnostic, prognostic and therapeutic implications. We assessed the relationship between MT-MMP expression and clinicopathological parameters in human non-small cell lung cancer (NSCLC) and histologically normal lung tissue by quantitative Real Time PCR (qRT-PCR). All MT-MMPs (MMPs 14–17, 24 and 25) were detected by qRT-PCR with significantly higher MMP-14, -15 and -17 expression observed in tumour relative to normal lung specimens. MMP-16 was undetectable in normal lung but expressed in 8% tumours. MMP-15 demonstrated significant overexpression in adenocarcinomas relative to squamous cell carcinomas and normal lung tissue. MMP-14 mRNA expression strongly correlated to MMP-14 proteolytic activity in preclinical tumour models, indicating that qRT-PCR may predict MMP-14 activity levels in NSCLC. These data suggest that MMP-14, -15 and -17 may be good markers of disease, or therapeutic targets for treatment of human NSCLC.

© 2007 Elsevier Ltd. All rights reserved.

1. Introduction

Non-small cell lung cancer (NSCLC) is the major cancer type in developed countries. Early stage NSCLC, without distant metastases or heavy regional nodal involvement, is often amenable to surgical resection and adjuvant chemotherapy. However, response to chemotherapy in NSCLC and prediction of tumour pathogenesis is often poor. As a consequence, better understanding of the molecular pathology of NSCLC, determination of novel tumour markers and identification

of targets for therapeutic exploitation are central to improving diagnosis and treatment.

Matrix metalloproteinases (MMPs) are a family of at least 24 zinc-dependent endopeptidases, with the collective ability to degrade the basement membrane and all protein components of the extracellular matrix (ECM).^{1–3} However, MMP activity is not restricted to ECM degradation and is essential for several processes involved in tumourigenesis, including neovascularisation, induction of the epithelial-mesenchymal transition and regulation of growth factor and chemokine

* Corresponding author.

E-mail address: j.gill1@bradford.ac.uk (J.H. Gill).

0959-8049/\$ - see front matter © 2007 Elsevier Ltd. All rights reserved.

doi:10.1016/j.ejca.2007.05.009

activity to name but a few.^{1–8} The human MMP family is divided into eight distinct structural groups: five of which are secreted and three of which are membrane-type MMPs (MT-MMPs)². MT-MMPs are membrane-tethered proteolytic enzymes and constitute the largest family of MMPs identified so far. The type I transmembrane structural class of MT-MMPs contains MMP-14 (MT1-MMP), MMP-15 (MT2-MMP), MMP-16 (MT3-MMP) and MMP-24 (MT5-MMP). The glycosyl phosphatidylinositol (GPI)-anchored structural group encompasses MMP-17 (MT4-MMP) and MMP-25 (MT6-MMP). Very little is known regarding the third MT-MMP group, the type II transmembrane MMPs, which contains only MMP-23 (cysteine-array MMP), a member structurally different from all other MMPs.⁹ Overexpression of MT-MMPs has been observed in many forms of cancer.¹⁰ The functions of MT-MMPs are known to include roles in activation of other MMP family members, pericellular proteolysis, modulation of cellular signalling, cellular migration, regulation of cellular proliferation and apoptosis, the angiogenic response and regulation of tumour invasion metastasis.^{4,10–13} Given the vast range of cellular processes influenced by the MT-MMPs, evaluation of their expression and activity in human tumours may be important in terms of diagnosis, prognosis and treatment.

A number of recent studies have analysed genetic expression of MMP-14 in several human tumour types, including tumours of the breast, colon, head and neck, oral cavity, prostate and importantly for this study, pulmonary tumours including bronchopulmonary and NSCLC.^{14–20} The majority of these studies were semiquantitative and did not evaluate expression of the other MT-MMP members.

Data regarding the expression of the MT-MMP subfamily in NSCLC are limited; therefore, the aim of the present study was to: (a) assess the mRNA expression of the MT-MMPs by qRT-PCR in human preclinical tumour models and paired clinical human NSCLC tumour and histologically normal lung tissue; (b) to evaluate the correlation between MMP-14 mRNA expression and proteolytic activity.

2. Materials and methods

2.1. Human tissue samples

A total of 39 freshly resected and snap frozen specimens of NSCLC were used for this study. In addition, the study also included paired histologically normal human lung tissue specimens for 15 of the tumours, excised distant to the tumour mass. Patient ages ranged from 47 to 85 years (median 68 years). Informed consent was obtained from all patients prior to specimen collection and all patient details were anonymised to ensure confidentiality. All experiments were performed after first obtaining consent from the local research and ethics committee according to Medical Research Council Regulations.

The presence of tumour (or lack in terms of normal tissue specimens) and histopathological classification was confirmed by pathologists at Bradford Royal Infirmary, UK. The panel of tumours comprised 28 adenocarcinomas and 11 squamous cell carcinomas (SCCs) and a range of tumour stages (4 pT1; 30 pT2; 4 pT3 and one pTx).

2.2. Cell culture

Cell lines were grown in monolayer culture in complete RPMI 1640 containing 10% v/v FBS, 1% v/v L-glutamine and 1% v/v sodium pyruvate (Sigma, Dorset, UK). All cultured cells were incubated at 37 °C in a humidified atmosphere of 5% CO₂. Cells were maintained in continuous exponential growth by passage every 5–7 days. Cells were harvested when subconfluent by trypsinisation (Trypsin-EDTA; Sigma) and stored as pellets at –20 °C. Cell lines used in these studies include HT1080 (fibrosarcoma); MCF-7 and MDA-MB-231 (breast carcinoma); HT29, BE, HCT116, COLO205, DLD1 (colorectal carcinoma); A549, H661, H460 (NSCLC); PC3 and LNCAP (prostate carcinoma). All cell lines were obtained from ECACC (Sailsbury, Wiltshire, UK) or ATCC (Manassas, VA, USA).

2.3. Human tumour xenografts

Human tumour cell lines were xenografted in mice under a project license issued by the UK Home Office, following UKCCCR guidelines. Female mice (nu/nu from an inbred colony, B and K Universal, Hull, UK) 6–8 weeks old were implanted subcutaneously with 2–3 mm³ fragments of H460, A549 (NSCLC); MCF7 (breast carcinoma); BE, COLO205, DLD1, HCT116, HT29 (colorectal carcinoma); PC3 (prostate adenocarcinoma) or HT1080 (fibrosarcoma) tumours. The resultant tumours were removed, snap-frozen in liquid nitrogen and stored at –70 °C.

2.4. Determination of MMP-14 mRNA expression by semi-quantitative RT-PCR analysis

TRI REAGENT™ (Sigma) was used to isolate RNA from cell pellets and tissue samples. Reverse transcription was performed using Advantage™ RT-for-PCR kit (BD Bioscience, Oxford, UK) according to the manufacturer's instructions. PCR was performed using the Titanium™ Taq PCR kit (BD Bioscience) according to the manufacturer's instructions and the primers described below (Invitrogen, Paisley, UK). Primers were based on previous studies,^{21,22} and were MMP-14 (sense) 5'-CGC TAC GCC ATC CAG GGT CTC AAA-3'; MMP-14 (antisense) 5'-CGG TCA TCA TCG GGC AGC ACA AAA-3'; GAPDH (sense) 5'-CCA CCC ATG GCA AAT TCC ATG GCA-3'; GAPDH (antisense) 5'-TCT AGA CGG CAG GTC AGG TCC ACC-3'. PCR amplification of MMP-14 involved initial denaturation of 94 °C for 4 min followed by 30 cycles of 94 °C for 30 s, 62 °C for 30 s and 72 °C for 30 s. PCR for GAPDH involved denaturation at 94 °C for 4 min followed by 25 cycles of 94 °C for 30 s, 60 °C for 30 s and 72 °C for 90 s. All cycles were terminated with a 10 min incubation at 72 °C to ensure complete elongation prior to incubation and storage at 4 °C. RT-PCR products were separated through a 1% w/v agarose gel containing 0.01% v/v ethidium bromide and visualised under ultraviolet (UV) illumination.

2.5. Quantification of MT-MMP expression by real-time RT-PCR (qRT-PCR)

For qRT-PCR, RNA extraction was carried out as for RT-PCR. For cDNA synthesis, a total of 1 µg of total RNA was reverse transcribed using random hexamers (Amersham,

Buckinghamshire, UK) and Superscript II reverse transcriptase (Invitrogen), according to the supplier's instructions. For PCRs, specific primers and fluorogenic probes for all MT-MMPs were designed using Primer Express 1.0 software (Applied Biosystems, Warrington, UK) and synthesised by Applied Biosystems. Where possible, to prevent the amplification of genomic DNA and to ensure that the PCR signal was generated from cDNA, primers were generated with sequences within different exons, close to intron-exon boundaries. BLASTN searches²³ were conducted on all primer/probe nucleotide sequences to ensure gene specificity. The 18S ribosomal RNA gene was used as an endogenous control to normalise for differences in the amount of total RNA in each sample, using previously validated procedures²⁴; 18S rRNA primers and probes were purchased from Applied Biosystems. PCRs were performed as described previously.²⁴ To determine relative RNA levels within the samples, standard curves for the PCR were prepared by using cDNA from one sample and making twofold serial dilutions covering the range equivalent to 20–0.625 ng RNA (for 18S analyses, the range was 4–0.125 ng).

2.6. Determination of MMP-14 protein levels by ELISA assay

MMP-14 protein activity levels were measured using the Biotrak Activity Assay System (Amersham Biosciences), carried out according to the manufacturer's instructions. Briefly, xenograft tissues were weighed, incubated for 15 min at 4 °C in membrane extraction buffer (50 mM Tris-HCl, pH 7.6, 1.5 mM NaCl, 0.5 mM CaCl₂, 1 µM ZnCl₂, 0.01% v/v Brij-35 and 0.25% v/v Triton X-100), homogenised on ice and incubated for a further 15 min at 4 °C. The resultant supernatants and MMP-14 standards (ranging from 0.125 ng/ml to 32 ng/ml) were incubated overnight at 4 °C on an anti-MMP-14 antibody coated microplate. Following incubation, all wells were washed (phosphate buffered saline containing 0.05% v/v Tween-20), the detection reagent added and the plate agitated and incubated at 37 °C for 6 h. The absorbance of each sample was read at 405 nm and the concentration of MMP-14 in the samples calculated from the standard curve.

2.7. Data analysis

For comparison of MT-MMP expression between NSCLC specimens and histologically normal tissue, the data did not satisfy normality or equal variance, so non-parametric statistical tests were used. The Mann-Whitney U-test was carried out to compare malignant and non-malignant samples. For all tests, a P-value of less than 0.01 was considered statistically significant.

3. Results

3.1. Expression of MT-MMPs in human tumour cell lines and xenografts

The expression of the MT-MMPs was assessed in a panel of human cell lines and human tumour xenografts representing several different human tumour types by qRT-PCR. All MT-MMPs

were detected successfully (Fig. 1) and were in agreement with expression observed by semi-quantitative RT-PCR (data not shown). MMP-14 and MMP-15 were expressed at high levels in all the tumour types represented in the screen (Fig. 1). MMP-16, -17, -24 and -25 demonstrated differential expression throughout the preclinical tumour panel, suggesting potential associations between these MMPs and tumour origin or tumour type (Fig. 1). In contrast to the other MT-MMPs, the type II transmembrane MMP, MMP-23, was expressed at relatively low levels in both lung cell lines and xenografts (Fig. 1) and was not therefore analysed in subsequent studies.

3.2. MT-MMP expression in NSCLC specimens and histologically normal lung tissue

The high level and range of MT-MMP expression observed in preclinical models of lung cancer (Fig. 1) led us to evaluate the expression of MMP-14, -15, -16, -17, -24 and -25 MT(1–6)-MMP in clinical specimens of NSCLC and associated histologically normal lung tissue by qRT-PCR. MMP-14 was expressed in 100% of NSCLC tumour specimens analysed in our study at levels significantly higher than that observed in histologically normal lung tissue (Fig. 2; $P < 0.01$). Of the other type I transmembrane MMPs, MMP-15 demonstrated significantly higher expression in NSCLC compared to histologically normal lung tissue (Fig. 2; $P < 0.01$), MMP-16 was absent or expressed at extremely low levels in all clinical specimens evaluated (Fig. 2) and MMP-24 demonstrated no significant difference in expression between NSCLC and histologically normal lung tissue (Fig. 2; $P = 0.5$).

The GPI-anchored MT-MMPs, MMP-17 and -25 were also expressed in both NSCLC and the associated histologically normal lung tissue (Fig. 2), although a significant difference between tumour and normal was detected for MMP-17 but not MMP-25 ($P < 0.01$ and $P = 0.9$, respectively).

3.3. Relationship between MT-MMP expression and tumour pathology

In contrast to the other MT-MMPs evaluated, expression of MMP-15 demonstrated differential expression between NSCLC tumour subtypes (Fig. 3). In addition to being significantly higher than that observed in histologically normal lung tissue ($P < 0.01$), the levels of MMP-15 in adenocarcinoma ($n = 28$) were also significantly higher than that detected in squamous cell carcinoma (SCC) ($n = 11$; $P < 0.01$) (Fig. 3). In contrast, although MMP-14 also demonstrated significantly higher expression in NSCLC compared to histologically normal lung, no significant difference was observed between tumour types (Fig. 3).

Although expressions of MMP-14, -16, -17, -24 and -25 were detected in human NSCLC, no significant correlation was observed between the expression of the individual MMPs and either tumour stage (pT status), tumour grade or presence of lymph node invasion (pN status) (data not shown).

3.4. Correlation between MMP-14 activity and MMP-14 qRT-PCR expression

Levels of proteolytically active MMP-14 were determined in human tumour cell lines and xenografts using the fluorogenic

Origin	Cell line	MMP						
		14	15	16	17	23	24	25
Fibrosarcoma	HT1080							
Breast	MCF7							
	MDA-MB-231							
Colon	HT29							
	BE							
	HCT116							
	COLO205							
Lung	DLD1							
	A549							
	H661							
	H460							
Prostate	PC3							
	LNCAP							

Origin	Xenograft	14	15	16	17	23	24	25
Fibrosarcoma	HT1080							
Breast	MCF7							
	BE							
Colon	HCT116							
	COLO205							
	DLD1							
	HT29							
Lung	A549							
	H460							
Prostate	PC3							

Expression key:

 Negative
 Low
 Moderate
 High
 Very high

Fig. 1 – Matrix metalloproteinases expression in human tumour cell lines and xenografts as measured by quantitative RT-PCR (qRT-PCR). Values of gene output are probe, and therefore gene specific, thus precluding comparison of expression between genes.

MT1-MMP (MMP-14) Biotrak assay. MMP-14 activity levels detected using this assay ranged from 1.1 to 0.02 ng MMP-14 per mg tissue, with the highest levels detected in HT1080 (1.12 ± 0.10 ng MMP-14 per mg tissue) and lowest in H460 (0.02 ± 0.01 ng MMP-14 per mg tissue) (Fig. 4). Comparison of MMP-14 activity levels with MMP-14 mRNA expression, as measured by qRT-PCR, demonstrated an excellent correlation between the levels of active MMP-14 and MMP-14 mRNA expression in both cell lines (data not shown) and xenografts (Fig. 4; $R^2 = 0.9$).

4. Discussion

MMPs are central regulators in the tumourigenic process and the major proteases involved in ECM breakdown and remodelling.¹ The original hypothesis that MMPs were involved purely with ECM degradation and consequently tumour invasion and metastasis was demonstrated to be too simplistic and the MMPs are now known to be a major force in the phenotypic evolution of cancer.²⁵ Growth-factor receptors, cell adhesion molecules, chemokines, cytokines, apoptotic ligands and angiogenic factors are just some examples of the diversity of substrates targeted by the MMPs.³

Identification of proteolytic enzyme expression and localisation in human cancers is critical in understanding the characteristics of individual cancer types, referred to as the cancer degradome.²⁶ Identification of specific MMPs important in the various stages of cancer progression will lead to improved understanding of different cancers and may repre-

sent new markers of disease or may lead to the identification and validation of new therapeutic targets.

The purpose of this study was to comprehensively analyse expression of the membrane-type MMP (MT-MMP) subfamily in human NSCLC and to evaluate the relationship between genetic expression of MMP-14 and the presence of active MMP-14. We have shown differential expression of these MMPs in NSCLC and demonstrated significantly higher levels of several family members in tumours compared to histologically normal lung tissue (Fig. 2).

MMP-14 was expressed in 100% of NSCLC tumour specimens analysed in our study (Fig. 2). The high levels of MMP-14 expression quantified in NSCLC in this study reinforce previous studies, implicating increased MMP-14 expression in lung cancer by semi-quantitative *in situ* hybridisation and cDNA array analysis.^{15,27} Although in our study MMP-14 was detected at low levels in histologically normal lung tissue, expression was significantly higher in NSCLC specimens ($P < 0.01$; Fig. 2). The elevated levels of MMP-14 we observed in NSCLC demonstrated no relationship to tumour stage, tumour grade, presence of tumour lymph node invasion or differential between histological tumour type (data not shown).

Data focussing on the expression of MMP-15 (MT2-MMP) in NSCLC are sparse. As for MMP-14, MMP-15 displayed a statistically significant difference ($P < 0.01$) in expression between tumour tissue and histologically normal lung (Fig. 2). This is supported by a previous study in which MMP-15 levels were suggested to be higher in adenocarcinoma cells than either type II alveolar cells or bronchiolar epithelial cells.²⁸ Further




Fig. 2 – Differential expression of the MT-MMP subfamily in human NSCLC and histologically normal lung tissue by qRT-PCR. Values are after normalisation to 18S-rRNA and are therefore gene specific. Expression data are plotted as mean expression \pm standard error (SE).

to demonstrating increased MMP-15 expression in NSCLC relative to histologically normal lung tissue, we have also shown significantly higher expression in lung adenocarcinoma relative to lung SCC ($P < 0.01$; Fig. 3), a previously unreported observation. No such differential was observed for MMP-14 (Fig. 3). This strongly suggests that MMP-15 may be a viable marker for molecular diagnostics of NSCLC and as such warrants further investigation. Similarly, the suggestion that MMP-15 has an anti-apoptotic mechanism and may connect tumour metastasis and apoptotic resistance¹¹ is also of relevance to our study and worthy of further investigation due to the high levels of MMP-15 we observed in NSCLC specimens.

There is very little data concerning the expression of MMP-16, MMP-17, MMP-24 or MMP-25 (MT(3–6)-MMP) in human tumours and no specific evidence for their involvement in NSCLC. MMP-16 was absent or expressed at extremely low levels in all clinical samples evaluated (Fig. 2). It is interesting to note the expression profile of MMP-16 in preclinical human tumour models, the differential in expression observed between tumour types and between cells grown *in vitro* or as *in vivo* tumour xenografts (Fig. 1). These observations could be the consequence of the degree of xenograft vascularisation or invasiveness, as supported by the recent associations between MMP-16 and both tissue vascularisation²⁹ and β -catenin mediated tumour invasion.³⁰ Identification of any such




Fig. 3 – Elevated expression of MMP-15 by qRT-PCR in lung adenocarcinomas compared to SCC ($P < 0.01$) and histologically normal lung ($P < 0.01$).

relationship and their clinical significance have yet to be undertaken. In the case of the lung cell lines, MMP-16 was detected in the H460 cells grown *in vitro* but not *in vivo* (Fig. 1), which is interesting in light of the fact that MMP-16 was absent in the majority of NSCLC samples.

Increased expression of MMP-17 was observed in NSCLC and did demonstrate a statistically significant difference from histologically normal lung (Fig. 2). Due to the low levels of MMP-17 detected in the majority of samples, caution should be adopted in the significance of this difference and the relationship to protein levels. MMP-24 and MMP-25 displayed differential expression between tumour and normal tissue although these failed to reach statistical significance ($P = 0.5$ and $P = 0.9$, respectively). Whilst we would predict that the expression of the MT-MMPs might be related to the stage and development of cancers due to their known involvement in metastasis, cell migration and activation of other MMP family members, no correlation was observed between tumour stage, grade or lymph node invasion and the expression of individual MT-MMPs (data not shown). These findings are perhaps surprising given the well documented role of the MMPs, including MT-MMPs, in various stages of cancer development and tumour spread¹³.




Fig. 4 – MMP-14 activity levels determined by ELISA (a) correlate strongly to MMP-14 expression by qRT-PCR (b), as measured in human tumour xenografts. A close correlation ($R^2 = 0.9$) is observed between activity and expression (c).

The excellent correlation ($R^2 = 0.9$) noted in this study between qRT-PCR and activity data for MMP-14 indicates that qRT-PCR is a proportional measurement of the level of MMP-14 protein present in a sample (Fig. 4). This correlation thereby adds value to the qRT-PCR clinical sample data for the MT-MMPs. Due to the nature of this assay, it does not take into account any potential intrinsic regulation of MT-MMP activity, such as by TIMPs, and may therefore signify the levels of MMP-14 protein and total potential MMP-14 activity rather than actual activity. Since both the qRT-PCR and

activity assay capture a single snapshot in the tumour, it would be interesting to assess the potential changes in expression and activity during tumourigenesis and tumour progression. In support of this, it would be desirable to evaluate the expression and sub-tumoural distribution of specific MT-MMPs using *in situ* hybridisation or immunohistochemistry, should specific and appropriate antibodies become available. In addition, it would also be interesting to differentiate the levels of actual MMP-14 activity from total potential MMP-14 activity in this context, to further assess MMP-14 as a disease marker or target for therapeutic exploitation in NSCLC.

Currently, there are no assays available to study the activity of MT(2–6)MMP in samples and we have not found any commercially available antibodies suitable for the development of such assays (data not shown). It will be important to investigate the relationship between qRT-PCR data and protein expression for all MT-MMPs to ascertain if a similar relationship is observed as is seen for MMP-14.

This study is the first analysis of all 6 MT-MMPs in NSCLC clinical specimens. Our data identify a number of MT-MMPs as being differentially expressed in NSCLC relative to histologically normal lung tissue, suggesting roles in tumourigenesis and/or as markers of disease. Using an MMP-14 activity assay, we have demonstrated the value of qRT-PCR as a predictor of protein expression. In summary, this study indicates the potential use of MMP-14 and MMP-15 as disease markers and therapeutic targets.

Conflict of interest statement

None of the authors disclose any financial or personal relationships with other people or organisations that could inappropriately influence the present work.

Acknowledgements

This work was supported by Cancer Research UK (Programme Grant A5953), The Big C Appeal and the Cancer Degradome Integrated Project Grant LSHC-CT-2003-503297 from the European Union Framework Programme 6.

REFERENCES

- Deryugina EI, Quigley JP. Matrix metalloproteinases and tumor metastasis. *Cancer Metast Rev* 2006;**25**:9–34.
- Egeblad M, Werb Z. New functions for the matrix metalloproteinases in cancer progression. *Nat Rev Cancer* 2002;**2**:161–74.
- Overall CM, Kleinfeld O. Tumour microenvironment – opinion: validating matrix metalloproteinases as drug targets and anti-targets for cancer therapy. *Nat Rev Cancer* 2006;**6**:227–39.
- Genis L, Galvez BG, Gonzalo P, Arroyo AG. MT1-MMP: Universal or particular player in angiogenesis? *Cancer Metast Rev* 2006;**25**:77–86.
- Handsley MM, Edwards DR. Metalloproteinases and their inhibitors in tumor angiogenesis. *Int J Cancer* 2005;**115**:849–60.
- McQuibban GA, Gong JH, Tam EM, McCulloch CA, Clark-Lewis CM, Overall CM. Inflammation dampened by gelatinase A cleavage of monocyte chemoattractant protein-3. *Science* 2000;**289**:1202–6.
- Pirila E, Sharabi A, Salo T, et al. Matrix metalloproteinases process the laminin-5 gamma 2-chain and regulate epithelial cell migration. *Biochem Biophys Res Commun* 2003;**303**:1012–7.
- Radisky DC, Bissell MJ. Matrix metalloproteinase-induced genomic instability. *Curr Opin Genet Dev* 2006;**16**:45–50.
- Velasco G, Pendas AM, Fueyo A, Knauper V, Murphy G, Lopez-Otin C. Cloning and characterization of human MMP-23, a new matrix metalloproteinase predominantly expressed in reproductive tissues and lacking conserved domains in other family members. *J Biol Chem* 1999;**274**:4570–6.
- Yana I, Seiki M. MT-MMPs play pivotal roles in cancer dissemination. *Clin Exp Metast* 2002;**19**:209–15.
- Abraham R, Schafer J, Rothe M, Bange J, Knyazev P, Ullrich A. Identification of MMP-15 as an anti-apoptotic factor in cancer cells. *J Biol Chem* 2005;**280**:34123–32.
- Plaisier M, Kapiteijn K, Koolwijk P, et al. Involvement of membrane-type matrix metalloproteinases (MT-MMPs) in capillary tube formation by human endometrial microvascular endothelial cells: role of MT3-MMP. *J Clin Endocrinol Metab* 2004;**89**:5828–36.
- Sounni NE, Noel A. Membrane type-matrix metalloproteinases and tumor progression. *Biochimie* 2005;**87**:329–42.
- Nawrocki B, Polette M, Marchand V, et al. Expression of matrix metalloproteinases and their inhibitors in human bronchopulmonary carcinomas: quantitative and morphological analyses. *Int J Cancer* 1997;**72**:556–64.
- Yamamura T, Nakanishi K, Hiroi S, et al. Expression of membrane-type-1-matrix metalloproteinase and metalloproteinase-2 in nonsmall cell lung carcinomas. *Lung Cancer* 2002;**35**:249–55.
- Myoung H, Kim MJ, Hong SD, Lee JI, Lim CY, Hong SP. Expression of membrane type I-matrix metalloproteinase in oral squamous cell carcinoma. *Cancer Lett* 2002;**185**:201–9.
- Udayakumar TS, Chen ML, Bair EL, et al. Membrane type-1-matrix metalloproteinase expressed by prostate carcinoma cells cleaves human laminin-5 beta3 chain and induces cell migration. *Cancer Res* 2003;**63**:2292–9.
- Nabeshima K, Inoue T, Shimao Y, et al. Front-cell-specific expression of membrane-type 1 matrix metalloproteinase and gelatinase A during cohort migration of colon carcinoma cells induced by hepatocyte growth factor/scatter factor. *Cancer Res* 2000;**60**:3364–9.
- Munaut C, Noel A, Hougrand O, Foidart JM, Boniver J, Deprez M. Vascular endothelial growth factor expression correlates with matrix metalloproteinases MT1-MMP, MMP-2 and MMP-9 in human glioblastomas. *Int J Cancer* 2003;**106**:848–55.
- Okada A, Bellocq JP, Rouyer N, et al. Membrane-type matrix metalloproteinase (MT-MMP) gene is expressed in stromal cells of human colon, breast, and head and neck carcinomas. *Proc Natl Acad Sci USA* 1995;**92**:2730–4.
- Daja MM, Niu X, Zhao Z, Brown JM, Russell PJ. Characterization of expression of matrix metalloproteinases and tissue inhibitors of metalloproteinases in prostate cancer cell lines. *Prostate Cancer Prostat Dis* 2003;**6**:15–26.
- Jung M, Romer A, Keyszer G, et al. mRNA expression of the five membrane-type matrix metalloproteinases MT1-MT5 in human prostatic cell lines and their down-regulation in human malignant prostatic tissue. *Prostate* 2003;**55**:89–98.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol* 1990;**215**:403–10.

24. Nuttall RK, Pennington CJ, Taplin J, et al. Elevated membrane-type matrix metalloproteinases in gliomas revealed by profiling proteases and inhibitors in human cancer cells. *Mol Cancer Res* 2003;1:333–45.
25. Overall CM, Dean RA. Degradomics: systems biology of the protease web. Pleiotropic roles of MMPs in cancer. *Cancer Metast Rev* 2006;25:69–75.
26. Lopez-Otin C, Overall CM. Protease degradomics: a new challenge for proteomics. *Nat Rev Mol Cell Biol* 2002;3:509–19.
27. Kettunen E, Anttila S, Seppanen JK, et al. Differentially expressed genes in nonsmall cell lung cancer: expression profiling of cancer-related genes in squamous cell lung cancer. *Cancer Genet Cytogenet* 2004;149:98–106.
28. Kobayashi K, Nishioka M, Kohno T, et al. Identification of genes whose expression is upregulated in lung adenocarcinoma cells in comparison with type II alveolar cells and bronchiolar epithelial cells in vivo. *Oncogene* 2004;23:3089–96.
29. Plaisier M, Koolwijk P, Hanemaaijer R, et al. Membrane-type matrix metalloproteinases and vascularization in human endometrium during the menstrual cycle. *Mol Hum Reprod* 2006;12:11–8.
30. Lowy AM, Clements WM, Bishop J, et al. beta-Catenin/Wnt signaling regulates expression of the membrane type 3 matrix metalloproteinase in gastric cancer. *Cancer Res* 2006;66:4734–41.