

DOI: 10.1002/ejic.201101120

Synthesis and Structural Features of New Ruthenium(II) Complexes Containing the Scorpionate Ligands Tris(pyrazol-1-yl)methanesulfonate (Tpms) and Tris(pyrazol-1-yl)methane (Tpm)

Sara Miguel,^[a] Josefina Diez,^[a] M^a Pilar Gamasa,^[a] and M^a Elena Lastra^{*[a]}

Keywords: Ruthenium / Scorpionate ligands / Phosphane ligands / N,O ligands

In order to provide a correct state of the art on PTA metal complexes, the following references must be included in the original paper:^[1]

A. D. Phillips, L. Gonsalvi, A. Romerosa, F. Vizza, M. Peruzzini, *Coord. Chem. Rev.* **2004**, 248, 955–993.

J. Bravo, S. Bolaño, L. Gonsalvi, M. Peruzzini *Coord. Chem. Rev.* **2010**, 254, 555–607.

The Authors

[1] S. Miguel, J. Diez, M. P. Gamasa, M. E. Lastra, *Eur. J. Inorg. Chem.* **2011**, 4745–4755.

Received: October 14, 2011

Published Online: October 31, 2011

[a] Departamento de Química Orgánica e Inorgánica, Instituto de Química Organometálica “Enrique Moles” (Unidad Asociada al C.S.I.C.), Universidad de Oviedo, 33006 Oviedo, Principado de Asturias, Spain
Fax: +34-985103446
E-mail: elb@uniovi.es