

## Adenosine receptors: synthesis, structure–activity relationships and biological activity of new 6-amino purine derivatives

Giovannella Strappaghetta<sup>a\*</sup>, Stefano Corsano<sup>a</sup>, Roberta Barbaro<sup>a</sup>, Antonio Lucacchini<sup>b</sup>,  
Gino Giannaccini<sup>b</sup>, Laura Betti<sup>b</sup>

<sup>a</sup>*Istituto di Chimica e Tecnologia del Farmaco, Università di Perugia, via del Liceo 1, 06123 Perugia, Italy*

<sup>b</sup>*Dipartimento di Psichiatria, Neurobiologia, Farmacologia e Biotecnologie, Università di Pisa, via Bonanno 6, 56126 Pisa, Italy*

(Received 13 June 1997; accepted 21 November 1997)

**Abstract** – The synthesis and evaluation of the biological activity of a series of pyridazin-3(2H)-one derivatives is reported. The compounds were tested in radioligand binding assays for affinity at A<sub>1</sub> and A<sub>2A</sub> adenosine receptors in bovine brain cortical membranes, and bovine brain striatal membranes, respectively. None of the compounds shows any affinity towards A<sub>2A</sub> receptor, while compounds in which the 6-chloro-pyridazin-3(2H)-one or 6-phenyl-pyridazin-3(2H)-one group is linked through a chain of two carbon atoms in the 6 position of the adenosine, show a good affinity towards A<sub>1</sub> adenosine receptor, particularly compound **8** in which a phenyl-pyridazinone group is present shows highest affinity with K<sub>i</sub> values 6.6 nM. © Elsevier, Paris

**A<sub>1</sub>-adenosine affinity / pyridazinones / anticonvulsant activity**

### 1. Introduction

Adenosine is a neuromodulator which acting through specific cell-surface receptors, regulates several physiological responses, including vasodilatation [1], antilipolytic effect, inhibition of platelet aggregation [2], inhibition of lymphocyte functions, depression of central nervous system activity [3–5]. Moreover the adenosine has inhibitory effects on neurotransmission and on spontaneous activity of central neurons [6].

Potent and selective agonists have been developed for these receptors, of which at least three subclasses are known (A<sub>1</sub>, A<sub>2</sub> and A<sub>3</sub>) [7]. These receptor types have been cloned and characterized as belonging to the superfamily of receptors with seven transmembrane helices that couple to G proteins [8]. A<sub>1</sub> receptors may couple to a variety of second messenger systems, including inhibition of adenylate cyclase, inhibition or stimulation of phosphoinositol turnover, and activation of ion channels [9]. A<sub>2</sub> receptors stimulate adenylate cyclase and can be further subdivided into high-affinity A<sub>2A</sub> and low-affinity A<sub>2B</sub> subtypes. It

is well known that N<sup>6</sup>-substituted adenosine derivatives such as (R)N<sup>6</sup>-[2-(1-phenyl)-propyl]adenosine **1** [10] have generally proved to be A<sub>1</sub>-receptor selective.

On the other hand, 3(2H)-pyridazinone derivatives show many pharmacological activities: reduction of blood pressure like compound **1a** [11], anticonvulsant activity like compound **1b** [12] (see *figure 1*). Moreover there are pyridazinone derivatives that have shown an interesting affinity towards A<sub>1</sub>-receptor [13].

We believed that combining purine riboside with the pyridazinone ring through a chain of two or three carbon atoms or through a piperazine ring it would be possible to obtain new compounds with interesting affinity towards A<sub>1</sub>-receptors, utilizable as cerebro-protective agents. In order to confirm the importance of the riboside in these 6-amino purine derivatives, the synthesis of a second series of compounds without the riboside fragment has been realized.

### 2. Chemistry

The compounds here reported were prepared as shown in *figures 2* and *3*. The reaction of the 6-chloropurine riboside **2** or of the 6-chloropurine **2a** with the amino derivatives in absolute ethanol in triethylamine, gave compounds **3–12** and **13–17** respectively.

\*Correspondence and reprints




Figure 1.

### 3. Biochemistry

Affinity of the new adenosine derivatives **3–17** towards adenosine receptors was evaluated using radioligand binding technique. Competition assays for A<sub>1</sub> and A<sub>2A</sub> adenosine receptors were determined in bovine brain cortical membranes, and bovine striatal membranes, respectively.

N<sup>6</sup>-(cyclohexyl)-adenosine ([<sup>3</sup>H]CHA), was used as A<sub>1</sub> ligand, and 2-[[p-(2-carboxyethyl)-phenylethyl]-amino]-5'-(N-ethylcarbamoyl)adenosine ([<sup>3</sup>H]CGS 21680) as A<sub>2A</sub> ligand. The biological results expressed in *K<sub>i</sub>* are reported in *table 1*. Thermodynamic analysis of the temperature-dependent affinities of agonist and antagonist have revealed fundamental differences in the interaction of agonist and antagonist with receptors. The temperature dependence of agonist and antagonist interactions varies at several neurotransmitters. At dopamine and  $\beta$ -adrenergic receptors agonists become more potent at lower temperatures [14, 15].

In initial studies of adenosine receptors binding was found that [<sup>3</sup>H]CHA binding was enhanced at higher temperatures, whereas [<sup>3</sup>H]DPX binding was favoured by lower temperature [16]. We have used two-temperature measurements of *in vitro* inhibitory binding constant as a simple method of discriminating between *in vivo* agonistic and antagonistic behaviour of A<sub>1</sub> adenosine receptor ligand [17], agonists and antagonists being characterized by *K<sub>i</sub>* (0 °C)/*K<sub>i</sub>* (25 °C) ratios respectively greater or less than unity. Our finding suggest that at adenosine receptors agonists are more potent at higher temperatures and antagonists at lower temperatures. To explore this phenomenon in greater detail, we evaluated the potencies of a series of adenosine agonists and antagonists in inhibi-

ting [<sup>3</sup>H]CHA binding at 0 °C and 25 °C (*table 1*). All tested compounds showed an agonist behaviour on A<sub>1</sub> receptors. The Hill coefficients, *nH* are reported for the temperature of 25 °C and are not significantly different from unity for either class of compounds suggesting the presence, at least in our experimental conditions, of a single class of binding sites.

### 4. Results and discussion

Compounds **7, 8, 9, 10** in which the adenosine is linked in 6-position with the pyridazinone fragment through a chain of two carbon atoms, show high affinity towards A<sub>1</sub> adenosine receptor, particularly compound **8** in which a phenyl-pyridazinone group is present shows highest affinity with a *K<sub>i</sub>* values 6.6 nM. Similar result has been obtained with the replacement of phenyl group with a chlorine atom (compound **7**), while replacement of phenyl group with a hydrogen atom (compound **9**) or with a imidazole group (compound **10**) gives a reduction of the affinity. These results show that there is a decrease of the affinity with the lengthening of the chain from two to three carbon atoms (compounds **11** and **12**), the affinity lowers also when the pyridazinone ring was substituted with a phthalazinone ring (compound **6**). Moreover in compounds **3, 4, 5** in which the ethylenic or propylenic chain is substituted by a rigid structure such as a piperazine ring, the affinity towards A<sub>1</sub> receptor disappears. Compounds **13–17**, adenine derivatives, do not show affinity towards A<sub>1</sub> receptor. Finally for all compounds the affinity at A<sub>2A</sub> receptor was also determined and none of the compounds showed any affinity.





Figure 2.



**Figure 3.**

In conclusion, the introduction of a pyridazinone system in 6-position of the purine riboside through an ethylenic chain, maintains a good affinity towards  $A_1$  receptor in comparison with CHA used as reference drug, moreover the

importance of the riboside in these structures for the affinity towards  $A_1$ -receptor has been confirmed. On the basis of these results compounds 7 and 8 may be useful as cerebroprotective agents.

**Table I.** Affinity towards A<sub>1</sub> and A<sub>2A</sub> adenosine receptor of new 6-amino purine derivatives.

Compound	$K_i$ [25 °C](A <sub>1</sub> ) <sup>a</sup> (nM ± SEM)	nH <sup>b</sup>	$K_i$ [0 °C]/ $K_i$ [25 °C] <sup>c</sup>	$K_i$ [25 °C](A <sub>2</sub> ) <sup>d</sup> (nM ± SEM)	nH <sup>b</sup>
3	> 10000	–	–	> 10000	–
4	> 10000	–	–	> 10000	–
5	> 10000	–	–	> 10000	–
6	86 ± 7.3	0.98	7.3	> 10000	–
7	8.50 ± 0.7	1.01	7.3	> 10000	–
8	6.6 ± 0.5	0.95	4.8	> 10000	–
9	49.6 ± 3.5	0.96	5.4	> 10000	–
10	35.6 ± 2.7	0.99	4.1	> 10000	–
11	82.6 ± 8.1	0.97	22	> 10000	–
12	37.4 ± 2.5	1.05	17	> 10000	–
13	> 10000	–	–	> 10000	–
14	> 10000	–	–	> 10000	–
15	> 10000	–	–	> 10000	–
16	> 10000	–	–	> 10000	–
17	> 10000	–	–	> 10000	–
<i>Agonists</i>					
CHA	1.3 ± 0.2	1.03	4.5	750 ± 65	0.97
R-PIA	1.2 ± 0.1	0.95	3.5	500 ± 40	1.01
CGS 21680	2600 ± 220	0.98	–	10 ± 0.9	0.96
NECA	6 ± 0.5	0.96	2.5	15 ± 1.3	0.99
<i>Antagonists</i>					
CPX	0.9 ± 0.08	0.96	0.3	470.2 ± 40.1	0.98
8-PT	86.0 ± 8.5	0.97	0.4	850 ± 82.8	0.96

<sup>a</sup>A<sub>1</sub> binding was measured as inhibition of [<sup>3</sup>H]-CHA binding as described in the Experimental protocols. The  $K_i$  values are means ± SEM of four separate assays, each performed in triplicate; <sup>b</sup>nH = Hill coefficient; <sup>c</sup> $K_i$  ratio =  $K_i$ [0 °C]/[25 °C]; <sup>d</sup>A<sub>2A</sub> binding was measured as inhibition of [<sup>3</sup>H]-CGS 21680 binding as described in the Experimental protocols. The  $K_i$  values are means ± SEM of four separate assays, each performed in triplicate.

## 5. Experimental protocols

### 5.1. Biological methods

#### 5.1.1. A<sub>1</sub> receptor binding

Bovine cerebral cortex was homogenized in 10 volumes ice-cold 0.25 M sucrose containing 5 mM EDTA and protease inhibitors (0.1 mM of phenylmethylsulfonylfluoride (PMSF), 200 µg/mL bacitracine, and 160 µg/mL benzamidine) in an ultra-turrax homogenizer. The homogenate was centrifuged at 1000 g for 10 min at 4 °C and the supernatant again centrifuged at 46000 g for 20 min at 4 °C. The resulting pellet was suspended in 10 volumes of ice-cold 50 mM Tris-HCl buffer at pH 7.7 containing 1 mM EDTA, 5 mM MgCl<sub>2</sub> and protease inhibitors (buffer T<sub>1</sub>). It was then homogenized and centrifuged at 46000 g for 20 min at 4 °C.

The pellet was dispersed in 5 volumes of fresh T<sub>1</sub> buffer and incubated with adenosine deaminase (2UI/mL) at 37 °C for 30 min, then recentrifuged at 46000 g for 20 min at 4 °C. The resulting pellet was frozen at –80 °C until the time of assay. The pellet was suspended in 20 volumes of ice-cold T<sub>1</sub> buffer and A<sub>1</sub> binding assay was performed in triplicate by incubating at 25 °C for 120 min or 0 °C for 150 min in 0.5 mL T<sub>1</sub> buffer containing aliquots of the membrane fraction (0.2–0.3 mg protein) and 1.3 nM[<sup>3</sup>H]CHA in the absence or presence of unlabelled 15 µM R-PIA.

The binding reaction was terminated by filtering through Whatman GF/C glass fiber filters under suction and washing twice with 5 mL ice-cold Tris-buffer. The filters were placed in scintillation vials and 4 mL Gold MN Cocktail-Packard solvent scintillation fluid was added. The radioactivity was counted with an Packard 1600 TR scintillation counter. Specific binding was obtained by subtracting non-specific binding from total binding and was approximated to 85–90% of the total binding.

#### 5.1.2. A<sub>2A</sub> receptor binding

Striatum was dissected from bovine brain and the tissue was homogenized in 20 volumes of ice-cold 50 mM Tris-HCl buffer at pH 7.4 containing 10 mM MgCl<sub>2</sub>, 1 mM EDTA, and protease inhibitors as reported above (buffer T<sub>2</sub>). The homogenate was centrifuged at 46000 g for 10 min at 4 °C. The pellet was then suspended in 20 volumes of Tris-HCl buffer (T<sub>2</sub>) containing adenosine deaminase (2UI/mL) and incubated for 30 min at 37 °C. The resulting pellet was diluted in 20 volumes of 50 mM Tris-HCl buffer at pH 7.5 containing 10 mM MgCl<sub>2</sub> and used in the binding assay.

Binding assay was performed in triplicate, by incubating aliquots of the membrane fraction (0.2–0.3 mg protein) in Tris-HCl at pH 7.5, with approximately 5 nM [<sup>3</sup>H]CGS 21680 in a final volume of 0.5 mL. Incubation was carried out at 25 °C for 90 min. Non-specific binding was defined in the presence of 50 µM NECA. The binding reaction was concluded by filtrac-

tion through Whatman GF/C glass fiber filters under reduced pressure. Filters were washed four times with 5 mL aliquots of ice-cold buffer and placed in scintillation vials. Specific binding was obtained by subtracting non-specific binding from total binding and approximated to 85–90% of the total binding. The receptor-bound radioactivity was measured as described above.

Compounds were dissolved in ethanol or DMSO (buffer/concentration of 2%) and added to the assay mixture. Blank experiment were carried out to determine the effect of the solvent on binding.

Protein estimation was based on a reported method [18], after solubilization with 0.75 N sodium hydroxide, using bovine serum albumin as standard.

The concentration of tested compound that produce 50% inhibition of specific [ $^3\text{H}$ ]CHA or [ $^3\text{H}$ ]CGS 21680 binding ( $\text{IC}_{50}$ ) was determined by log-probit analysis with seven concentrations of the displacer, each performed in triplicate. Inhibition constants ( $K_i$ ) were calculated according to the equation of Cheng and Prousoff [19];  $K_i = \text{IC}_{50}/([L]/K_d)$ , where  $[L]$  is the ligand concentration and  $K_d$  its dissociation constant.  $K_d$  of [ $^3\text{H}$ ]CHA binding to cortex membranes was 1.2 nM and  $K_d$  of [ $^3\text{H}$ ]CGS 21680 binding to striatal membranes was 10 nM.

## 5.2. Chemistry

Melting points were determined using a Kofler hot-stage apparatus and are uncorrected. The NMR spectra were recorded with a Bruker AC 200 MHz instrument in the solvent indicated below. The chemical shift values (ppm) are relative to tetramethylsilane as internal standard. Elemental analyses are within  $\pm 0.4\%$  of theoretical values. Precoated Kiesegel 60 F 254 plates (Merck) were used for TLC.

### 5.2.1. 6-Substituted amino-9H-purines – General method for the preparation of compounds 3–17 [20]

**6-[4-(pyridazin-3(2H)-one-6-yl)-piperazin-1-yl]-9- $\beta$ -D-ribofuranosyl-9H-purine 3:** A mixture of 6-chloropurine riboside 1.50 g (5.2 mmol), 1.20 g (6.5 mmol) of 6-(piperazin-1-yl)-pyridazin-3(2H)-one **18** [21], and a 3-fold excess of triethylamine in 50 mL of absolute ethanol was heated a reflux for 20 h. (until TLC showed disappearance of starting material). After cooling the mixture was filtered and the collected solid was crystallized from ethanol. Yield: 55%; m.p.: 274–278 °C;  $^1\text{H-NMR}$  ( $\text{DMSO-}d_6$ )  $\delta$ : 3.4–3.8 (m, 6H,  $\text{CH}_2$ -5', 4H-pip.), 3.9 (m, 1H, H-4'), 4.2 (d, 1H, H-3'), 4.3 (m, 4H, H-pip.), 4.6 (m, 1H, H-2'), 5.2 (d, 1H, OH-3'), 5.4 (m, 1H, OH-5'), 5.5 (d, 1H, OH-2'), 6.0 (d, 1H, H-1'), 6.8 (d, 1H, H-4 pyrid.), 7.5 (d, 1H, H-5 pyrid.), 8.3 (s, 1H, H-2 purin), 8.4 (s, 1H, H-8 purin.), 12.2 (s, 1H, H-2 pyrid.).

**6-[4-(pyridazin-3(2H)-one-5-yl)-piperazin-1-yl]-9- $\beta$ -D-ribofuranosyl-9H-purine 4:** This compound was prepared by reaction of 6-chloropurine riboside **2** with 5-(piperazin-1-yl)-pyridazin-3(2H)-one **19** [22]. Yield: 50%; m.p.: 264–267 °C (methanol- $\text{H}_2\text{O}$ );  $^1\text{H-NMR}$  ( $\text{DMSO-}d_6$ )  $\delta$ : 3.4–3.8 (m, 6H,  $\text{CH}_2$ -5', 4H-pip.), 3.9 (m, 1H, H-4'), 4.2 (m, 1H, H-3'), 4.3 (m, 4H, H-pip.), 4.6 (m, 1H, H-2'), 5.2 (d, 1H, OH-3'), 5.3 (d, 1H, OH-5'), 5.5 (d, 1H, OH-2'), 5.8 (s, 1H, H-4 pyrid.), 6.0 (d, 1H, H-1'), 8.0 (s, 1H, H-6 pyrid.), 8.3 (s, 1H, H-2 purin), 8.4 (s, 1H, H-8 purin.), 12.2 (s, 1H, H-2 pyrid.).

**6-[4-(4-chloro-pyridazin-3(2H)-one-5-yl)piperazin-1-yl]-9- $\beta$ -D-ribofuranosyl-9H-purine 5:** This compound was prepared by reaction of 6-chloropurine riboside **2** with 4-chloro-5-(piperazin-1-yl)-pyridazin-3(2H)-one **19a** [22]. Yield: 65%; m.p.:

258–261 °C (ethanol- $\text{H}_2\text{O}$ );  $^1\text{H-NMR}$  ( $\text{DMSO-}d_6$ )  $\delta$ : 3.4–3.8 (m, 6H,  $\text{CH}_2$ -5', 4H-pip.), 3.9 (m, 1H, H-4'), 4.2 (m, 1H, H-3'), 4.3 (m, 4H, H-pip.), 4.6 (m, 1H, H-2'), 5.2 (d, 1H, OH-3'), 5.3 (d, 1H, OH-5'), 5.5 (d, 1H, OH-2'), 5.6 (d, 1H, H-1'), 8.0 (s, 1H, H-6 pyrid.), 8.3 (s, 1H, H-2 purin), 8.4 (s, 1H, H-8 purin.), 12.9 (s, 1H, H-2 pyrid.).

**6-[2-(phthalazin-1(2H)-one-2-yl)-aminoethyl]-9- $\beta$ -D-ribofuranosyl-9H-purine 6:** This compound was prepared by reaction of 6-chloropurine riboside **2** with 2-(2-aminoethyl)-phthalazin-1(2H)-one **21**. Yield: 50%; m.p.: 140–143 °C (ethanol);  $^1\text{H-NMR}$  ( $\text{DMSO-}d_6$ )  $\delta$ : 3.5 (m, 2H,  $\text{CH}_2$ -5'), 3.8–5.0 (m, 3H,  $\text{CH}_2$ , H-4'), 4.2 (m, 1H, H-3'), 4.4 (t, 2H,  $-\text{CH}_2$ ), 4.6 (m, 1H, H-2'), 5.2 (d, 1H, OH-3'), 5.3–5.5 (m, 2H, OH-5', 2'), 6.0 (d, 1H, H-1'), 7.8–7.9 (m, 4H, H-arom.), 8.0 (m, 2H, NH, 1H-arom.), 8.3 (m, 2H, H-2,8 purin).

**6-[2-(6-chloro-pyridazin-3(2H)-one-2-yl)-aminoethyl]-9- $\beta$ -D-ribofuranosyl-9H-purine 7:** This compound was prepared by reaction of 6-chloropurine riboside **2** with 2-(2-aminoethyl)-6-chloro-pyridazin-3(2H)-one **20**. Yield: 40%; m.p.: 230–233 °C (ethanol- $\text{H}_2\text{O}$ );  $^1\text{H-NMR}$  ( $\text{DMSO-}d_6$ )  $\delta$ : 3.5 (m, 2H,  $\text{CH}_2$ -5'), 3.8–4.0 (m, 3H,  $\text{CH}_2$ , H-4'), 4.2 (m, 1H, H-3'), 4.3 (m, 2H,  $\text{CH}_2$ ), 4.6 (m, 1H, H-2'), 5.2 (d, 1H, OH-3'), 5.3–5.5 (m, 2H, OH-5', 2'), 6.0 (d, 1H, H-1'), 7.0 (d, 1H, H-4 pyrid.), 7.5 (d, 1H, H-5 pyrid.), 8.0 (s, 1H, NH), 8.3 (s, 1H, H-2 purin), 8.4 (s, 1H, H-8 purin.).

**6-[2-(6-phenyl-pyridazin-3(2H)-one-2-yl)-aminoethyl]-9- $\beta$ -D-ribofuranosyl-9H-purine 8:** This compound was prepared by reaction of 6-chloropurine riboside **2** with 2-(2-aminoethyl)-6-phenyl-pyridazin-3(2H)-one **20a** [23]. Yield: 45%; m.p.: 206–209 °C (ethanol);  $^1\text{H-NMR}$  ( $\text{DMSO-}d_6$ )  $\delta$ : 3.6 (m, 2H,  $\text{CH}_2$ -5'), 3.9 (m, 3H,  $\text{CH}_2$ , H-4'), 4.2 (m, 1H, H-3'), 4.4 (t, 2H,  $-\text{CH}_2$ ), 4.6 (m, 1H, H-2'), 5.2 (d, 1H, OH-3'), 5.3–5.5 (m, 2H, OH-5', 2'), 6.0 (d, 1H, H-1'), 7.0 (d, 1H, H-4 pyrid.), 7.4–7.5 (m, 3H, H-arom.), 7.6 (m, 2H, H-arom.), 8.0 (m, 2H, NH, H-5 pyrid.), 8.2 (s, 1H, H-2 purin), 8.4 (s, 1H, H-8 purin.).

**6-[2-(pyridazin-3(2H)-one-2-yl)-aminoethyl]-9- $\beta$ -D-ribofuranosyl-9H-purine 9:** Prepared by reaction of 6-chloropurine riboside **2** with 2-(2-aminoethyl)-pyridazin-3(2H)-one **20b**. Yield: 60%; m.p.: 195–198 °C (ethanol);  $^1\text{H-NMR}$  ( $\text{DMSO-}d_6$ )  $\delta$ : 3.6 (m, 2H,  $\text{CH}_2$ -5'), 3.9 (m, 3H,  $\text{CH}_2$ , H-4'), 4.0 (m, 1H, H-3'), 4.3 (t, 2H,  $-\text{CH}_2$ ), 4.6 (m, 1H, H-2'), 5.2 (d, 1H, OH-3'), 5.3–5.5 (m, 2H, OH-5', 2'), 6.0 (d, 1H, H-1'), 6.9 (d, 1H, H-4 pyrid.), 7.4 (dd., 1H, H-5 pyrid), 7.8 (d, 1H, H-6 pyrid.), 8.0 (s, 1H, H-2 purin.), 8.2 (s, 1H, H-8 purin.).

**6-[2-(imidazol-1-yl)-pyridazin-3(2H)-one-2-yl)-aminoethyl]-9- $\beta$ -D-ribofuranosyl-9H-purine 10:** Prepared by reaction of 6-chloropurine riboside **2** with 2-(2-aminoethyl)-6-(imidazol-1-yl)-pyridazin-3(2H)-one **20c**. Yield: 30%; m.p.: 95–100 °C (ethanol);  $^1\text{H-NMR}$  ( $\text{DMSO-}d_6$ )  $\delta$ : 3.6 (m, 2H,  $\text{CH}_2$ -5'), 3.8–4.0 (m, 3H,  $\text{CH}_2$ , H-4'), 4.2 (m, 1H, H-3'), 4.3 (m, 2H,  $\text{CH}_2$ ), 4.6 (m, 1H, H-2'), 5.2 (d, 1H, OH-3'), 5.3 (d, 1H, OH-5'), 5.5 (d, 1H, OH-2'), 6.0 (d, 1H, H-1'), 7.0 (s, 1H, H-imid.), 7.2 (d, 1H, H-4 pyrid.), 7.5 (s, 1H, H-imid.), 7.9 (d, 1H, H-5 pyrid.), 8.0 (m, 2H, NH, H-imid.), 8.2 (s, 1H, H-2 purin.), 8.4 (s, 1H, H-8 purin.).

**6-[3-(6-phenyl-pyridazin-3(2H)-one-2-yl)-aminopropyl]-9- $\beta$ -D-ribofuranosyl-9H-purine 11:** Prepared by reaction of 6-chloropurine riboside **2** with 2-(3-aminopropyl)-6-phenyl-pyridazin-3(2H)-one **20d** [23]. Yield: 50%; m.p.: 116–118 °C (ethanol);  $^1\text{H-NMR}$  ( $\text{DMSO-}d_6$ )  $\delta$ : 2.2 (m, 2H,  $\text{CH}_2$ ), 3.5–3.8

(m, 4H, CH<sub>2</sub>-5', CH<sub>2</sub>), 4.0 (d, 1H, H-4'), 4.1–4.3 (m, 3H, CH<sub>2</sub>, H-3'), 4.6 (m, 1H, H-2'), 5.2 (d, 1H, OH-3'), 5.3–5.5 (m, 2H, OH-5', 2'), 6.0 (d, 1H, H-1'), 7.0 (d, 1H, H-4 pyrid.), 7.4–7.6 (m, 3H, H-arom.), 7.8–7.9 (m, 3H, 2H-arom., NH.), 8.0 (s, 1H, H-5 pyrid.), 8.2 (s, 1H, H-2 purin), 8.4 (s, 1H, H-8 purin).

**6-[3-(6-chloro-pyridazin-3(2H)-one-2-yl)-aminopropyl]-9-β-D-ribofuranosyl-9H-purine 12:** This compound was prepared by reaction of 6-chloropurine riboside **2** with 2-(3-aminopropyl)-6-chloro-pyridazin-3(2H)-one **20e**. Yield: 40%; m.p.: 112–115 °C (ethanol); <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>) δ: 2.0 (m, 2H, CH<sub>2</sub>), 3.4–3.7 (m, 4H, CH<sub>2</sub>, CH<sub>2</sub>-5'), 4.0 (d, 1H, H-4'), 4.1 (t, 2H, CH<sub>2</sub>), 4.2 (m, 1H, H-3'), 4.6 (m, 1H, H-2'), 5.2 (d, 1H, OH-3'), 5.3–5.4 (d, 2H, OH-5', 2'), 6.0 (d, 1H, H-1'), 7.0 (d, 1H, H-4 pyrid.), 7.5 (d, 1H, H-5 pyrid.), 7.8 (s, 1H, NH), 8.2 (s, 1H, H-2 purin), 8.4 (s, 1H, H-8 purin.).

**6-[4-(pyridazin-3(2H)-one-5-yl)]-piperazin-1-yl]-9H-purine 13:** This compound was prepared by reaction of 6-chloropurine **2a** with 5-piperazin-1-yl-pyridazin-3(2H)-one **19** [22]. Yield: 50%; m.p. > 300 °C (DMF); <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>) δ: 4.0 (m, 4H, H-pip.), 4.6 (m, 4H, H-pip.), 6.6 (s, 1H, H-4 pyrid.), 8.4 (s, 1H, H-6 pyrid.), 8.8 (s, 2H, H-2, 8 purin), 12.2 (s, 1H, H-2 pyrid.), 13.0 (s, 1H, H-9 purin.).

**6-[4-(4-chloro-pyridazin-3(2H)-one-5-yl)-piperazin-1-yl]-9H-purine 14:** This compound was prepared by reaction of 6-chloropurine **2a** with 4-chloro-5-piperazin-1-yl-pyridazin-3(2H)-one **19a** [22]. Yield: 50%; m.p. > 300 °C (DMF); <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>) δ: 3.5 (m, 4H, H-pip.), 4.3 (m, 4H, H-pip.), 7.9 (s, 1H, H-6 pyrid.), 8.2 (s, 1H, H-2 purin), 8.3 (s, 1H, H-8 purin.), 12.5 (s, 1H, H-2 pyrid.), 13.0 (s, 1H, H-9 purin.).

**6-[4-(pyridazin-3(2H)-one-6-yl)-piperazin-1-yl]-9H-purine 15:** This compound was prepared by reaction of 6-chloropurine **2a** with 6-piperazin-1-yl-pyridazin-3(2H)-one **18** [21]. Yield: 30%; m.p. > 300 °C (ethanol); <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>) δ: 3.5 (m, 4H, H-pip.), 4.2 (m, 4H, H-pip.), 6.8 (d, 1H, H-4 pyrid.), 7.6 (d, 1H, H-5 pyrid.), 8.2 (s, 1H, H-2 purin), 8.3 (s, 1H, H-8 purin.), 12.2 (s, 1H, H-2 pyrid.), 13.0 (s, 1H, H-9 purin.).

**6-[2-(phthalazin-1(2H)-one-2-yl)-aminoethyl]-9H-purine 16:** Prepared by reaction of 6-chloropurine **2a** with 2-(2-aminoethyl)-phthalazin-1(2H)-one **21**. Yield: 60%; m.p.: 268–272 °C (ethanol); <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>) δ: 3.8 (m, 2H, CH<sub>2</sub>), 4.4 (t, 2H, CH<sub>2</sub>), 7.7–8.0 (m, 4H, H-arom.), 8.0–8.2 (m, 2H, NH, H-pyrid.), 8.2–8.4 (m, 2H, H-8,2 purin), 13.0 (s, 1H, H-9 purin.).

**6-[2-(6-chloro-pyridazin-3(2H)-one-2-yl)-aminoethyl]-9H-purine 17:** Prepared by reaction of 6-chloropurine **2a** with 2-(2-aminoethyl)-6-chloro-pyridazin-3(2H)-one **20**. Yield: 50%; m.p.: 270–273 °C (ethanol); <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>) δ: 3.8 (m, 2H, CH<sub>2</sub>), 4.3 (t, 2H, CH<sub>2</sub>), 7.0 (d, 1H, H-4 pyrid.), 7.5 (d, 1H, H-5 pyrid.), 7.8 (s, 1H, NH), 8.1–8.2 (m, 2H, H-8,2 purin.), 13.0 (s, 1H, H-9 purin.).

**2-(2-aminoethyl)-6-chloro-pyridazin-3(2H)-one 20:** 2.6 g (0.02 mol) of 6-chloro-pyridazin-3(2H)-one was added to an ethanolic solution (50 mL) of sodium 0.46 g (0.02 g atom). On the other hand, a solution of 2-aminoethyl chloride hydrochloride 2.3 g (0.02 mol) in absolute ethanol (40 mL) was added to an ethanolic solution (50 mL) of sodium (0.46 g, 0.02 g atom).

The two solutions were mixed at room temperature and stirring for 1 h, under the mixture was refluxed for 18 h, and the mixture was filtered. The organic layer was evaporated, and the residue was diluted with water acidified with HCl 1 N and extracted by CH<sub>2</sub>Cl<sub>2</sub>.

The water solution was alkalized with NaOH 1 N, and extracted with CH<sub>2</sub>Cl<sub>2</sub>, the organic phase dried over dry sodium sulphate, and removed under reduced pressure. The crude product was used without further purification. Yield: 30%, m.p.: 83–85 °C; <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.6 (s, 2H, NH<sub>2</sub>), 3.1 (t, 2H, CH<sub>2</sub>), 4.15 (t, 2H, CH<sub>2</sub>), 6.9 (d, 1H, H-4 pyrid.), 7.2 (d, 1H, H-5 pyrid.).

**2-(2-aminoethyl)-pyridazin-3(2H)-one 20b:** This compound was prepared by alkylation of pyridazin-3(2H)-one with N-(2-bromoethyl)-phthalimide in the presence of K<sub>2</sub>CO<sub>3</sub>, followed by hydrolysis with hydrazine hydrate, using the method of Yamada [24]. The residue was used without further purification, m.p.: 232–235 °C. <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.4 (s, 2H, NH<sub>2</sub>), 3.15 (t, 2H, CH<sub>2</sub>), 4.2 (t, 2H, CH<sub>2</sub>), 6.9 (d, 1H, H-4 pyrid.), 7.2 (m, 1H, H-5 pyrid.), 7.8 (d, 1H, H-6 pyrid.).

**2-(2-aminoethyl)-6-(imidazol-1-yl)-pyridazin-3(2H)-one 20c:** This compound was prepared using the method described for compound **20b**. The residue was used without further purification, m.p.: 180–185 °C. <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.3 (s, 2H, NH<sub>2</sub>), 3.1 (t, 2H, CH<sub>2</sub>), 4.15 (t, 2H, CH<sub>2</sub>), 7.05 (d, 1H, H-pyrid.), 7.1 (s, 1H, H-imid.), 7.3–7.5 (m, 2H, H-5 pyrid., H-imid.), 8.0 (s, 1H, H-imid.).

**2-(3-aminopropyl)-6-chloro-pyridazin-3(2H)-one 20e:** This compound was prepared from 6-chloro-pyridazin-3(2H)-one with 1,3-dibromopropane in benzene, potassium hydroxide and tetrabutyl ammonium bromide (TBAB). The corresponding 3-bromopropyl derivate was treated with an ethanolic solution of 25% ammonium hydroxide, using the method of Yamada [23]. <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 2.0 (m, 4H, CH<sub>2</sub>, NH<sub>2</sub>), 2.7 (t, 2H, CH<sub>2</sub>), 4.2 (t, 2H, CH<sub>2</sub>), 6.9 (d, 1H, H-4 pyrid.), 7.2 (m, 1H, H-5 pyrid.).

**2-(2-aminoethyl)-phthalazin-1(2H)-one 21:** A solution of 0.660 g (4.5 mmol) of 1(2H)-phthalazinone in 15 mL of dry ethanolic and 0.240 g (6 mmol) of NaOH was stirred under reflux for 30 min.

After a solution of 0.922 g (4.5 mmol) of 2-aminoethyl-bromide hydrobromide in 15 mL of dry ethanol and 0.2 g (5 mmol) of NaOH was added. The mixture was refluxed for 24 h, filtered hot, and then the organic layer was evaporated in vacuo. The residue was diluted with water acidified with HCl 1 N and extracted by CH<sub>2</sub>Cl<sub>2</sub>. The water solution was alkalized with NaOH 1 N, and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic phase dry on sodium sulphate was evaporated under reduced pressure, the residue was used without further purification. m.p.: 84–88 °C, <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.5 (s, 2H, NH<sub>2</sub>), 3.2 (t, 2H, CH<sub>2</sub>), 4.3 (t, 2H, CH<sub>2</sub>), 7.6–7.8 (m, 3H, H-arom.), 8.1 (s, 1H, H-4-pyrid.), 8.4–8.5 (m, 1H, H-arom.).

## Acknowledgement

This work was supported by MURST (Ministry of University and Scientific Research). Thanks are due to Dr. C. Santi for NMR analysis and the aid in the structure elucidation.

## References

- [1] Conti A., Monopoli A., Gamba M. et al. *Naunyn Schmiedeberg's Arch. Pharmacol.* 348 (1993) 108–112.
- [2] Lohse M.J., Klotz K.N., Schwabe U., Cristalli G., Vittori S., Grifantini M., *Naunyn Schmiedeberg's Arch. Pharmacol.* 337 (1988) 687–689.
- [3] Radulovacky M., Virus R.M., Djuricic Nedelson M., Green R.D., *J. Pharmacol. Exp. Ther.* 228 (1984) 268–274.
- [4] Daly J.W., *J. Med. Chem.* 25 (1982) 187–207.
- [5] Church M.K., Bernyon R.C., Hughes P.I. et al., in: Stone T.W. (Ed.), *Purines: Pharmacology and Physiological Role*, VCH, Weinheim, 1985, pp. 175–184.
- [6] von Lubitz D.K.J.E., Lin R.C.-S., Melman N., Ji X.-d., Jacobson K.A., *Eur. J. Pharmacol.* 256 (1994) 161–167.
- [7] Fredholm B., Abbracchio M.P., Burnstock G., Daly J.W., Harden T.K., Jacobson K.A., Leff P., Williams M., *Pharmacol. Rev.* 46(2) (1994) 143–213.
- [8] Maenhaut C., van Sande S.J., Libert F., Abramowicz M., Parmentier M., Vanderhaegen J.J., Dumont J.E., Vassart G., Schiffmann S., *Biochem. Biophys. Res. Commun.* 173 (1990) 1169–1178.
- [9] Stiles G.L., *J. Biol. Chem.* 267 (1992) 6451–6454.
- [10] Kusachi S., Thompson R.D., Olsson R.A., *J. Pharmacol. Exp. Ther.* 227 (1983) 316–322.
- [11] Narimatsu A., Katida J., Satoh N., Suzuki R., Kobayashi M., Okushima H., *Jpn. J. Pharmacol.* 40 (1986) 234–240; Narimatsu A., Katiba J. Satoh N., Suzuki R., Kobayashi M., *Fed. Proc.* 45 (1986) 810–815.
- [12] Coudert P., Rubat C., Couquelet J. et al., *Eur. J. Med. Chem.* 24 (1989) 551–555.
- [13] Corsano S., Strappaghetti G., Scapicchi R., Lucacchini A., Senatore G., *Arch. Pharm.* 328 (1995) 654–658.
- [14] Niehoff D.L., Palacios J.M., Kuhar M.J., *Neurosci. Abstr.* 6 (1980) 252–256.
- [15] Weiland G.A., Minneman K.P., Molinoff P.B., *Nature* 281 (1979) 114–117.
- [16] Bruns R.F., Daly J.W., Snyder S.H., *Proc. Natl. Sci. USA* 77 (1980) 5547–5551.
- [17] Borea P.A., Varani K., Malaguti V., Gilli G., *J. Pharm. Pharmacol.* 43 (1991) 866–868.
- [18] Lowry O.H., Rosenbrough N.J., Farr A.L., Randall R.J., *J. Biol. Chem.* 193 (1951) 265–275.
- [19] Cheng Y.C., Prusoff W.H., *Biochem. Pharmacol.* 22 (1973) 3099–3108.
- [20] Kusachi S., Thompson R.D., Bugni W.J., Yamada N., Olsson R.A., *J. Med. Chem.* 28 (1985) 1636–1643.
- [21] Corsano S., Strappaghetti G., Codagnone A., Scapicchi R., Marucci G., *Eur. J. Med. Chem.* 27 (1992) 545–549.
- [22] Corsano S., Scapicchi R., Strappaghetti G., Marucci G., Paparelli F., *Eur. J. Med. Chem.* 28 (1993) 647–651.
- [23] Yamada T., Tsukamoto Y., Shimamura H., Banno S., Sapo M., *Eur. J. Med. Chem.* 18 (1983) 209–214.
- [24] Yamada T., Nobuhara Y., Shimamura H., Tsukamoto Y., Yosmimara K. et al., *J. Med. Chem.* 26 (1983) 373–381.