


### Metal–Organic Framework

J.-N. Rebilly, J. Bacsá, M. J. Rosseinsky\*

#### 1D Tubular and 2D Metal–Organic Frameworks Based on a Flexible Amino Acid Derived Organic Spacer

**Flexible, yet rigid!** A ligand derived from glutamic acid generates chiral MOF architectures ranging from 1D to 2D that are organized in the crystal through a complex hierarchy of coordination and hydrogen bonds that include encapsulation of solvent molecules. The flexibility of the ligand backbone gives access to several topologies and dimensionalities depending on the type of metal center used in the synthesis.



*Chem. Asian J.*  
DOI: 10.1002/asia.200900078




### Inhibitors


V. Ahmed, Y. Liu, S. D. Taylor\*

#### Multiple Pathways for the Irreversible Inhibition of Steroid Sulfatase with Quinone Methide-Generating Suicide Inhibitors

**Unexpected inhibition:** 2- and 4-mono- and difluoromethyl estrone sulfate derivatives are suicide inhibitors of steroid sulfatase (STS). Kinetic studies suggest that inhibition by the monofluoro derivatives is a result of a quinone methide intermediate that reacts with active-site nucleophiles, whereas the main inhibition pathway of the 4-difluoromethyl derivative is a result of decomposition of the initial quinone methide to an aldehyde that acts as potent, almost irreversible inhibitor.



*ChemBioChem*  
DOI: 10.1002/cbic.200900143




### Molecular Dynamics


E. C. Beret, R. R. Pappalardo, D. Marx, E. Sánchez Marcos\*

#### Characterizing Pt-Derived Anticancer Drugs from First Principles: The Case of Oxaliplatin in Aqueous Solution

**Anionic hydration:** Ab initio molecular dynamics simulations help to obtain a microscopic description of the behavior of oxaliplatin anticancer drug in bulk water (see picture). Its hydration structure is closely related to that previously found for  $\text{Pt}^{\text{II}}$  aqua ion and derives from the presence of a square-planar motif around the metal center.



*ChemPhysChem*  
DOI: 10.1002/cphc.200900027




### Silver Nanoparticles


K. K. Y. Wong,\* S. O. F. Cheung, L. Huang, J. Niu, C. Tao, C.-M. Ho, C.-M. Che, P. K. H. Tam

#### Further Evidence of the Anti-inflammatory Effects of Silver Nanoparticles

**The production of pure silver** in nanoparticle size has opened new dimensions in the clinical use of this precious metal. We and others have demonstrated previously that silver nanoparticles (nAg) possess efficient antimicrobial activity. Herein we show they may also have significant anti-inflammatory effects in a postoperative peritoneal adhesion model. This finding provides further insight into the biological actions of nAg as well as a potentially novel therapy for peritoneal adhesions in clinical surgery.



*ChemMedChem*  
DOI: 10.1002/cmdc.200900049



*Eur. J. Inorg. Chem.*  
DOI: 10.1002/ejic.200900123

### Copper(I) Complexes


L. Shi, B. Li\*

#### A Series of Cu<sup>I</sup> Complexes Containing 1,10-Phenanthroline Derivative Ligands: Synthesis, Characterization, Photophysical, and Oxygen-Sensing Properties

A series of copper(I) complexes was synthesized and characterized. Their photophysical and oxygen-sensing properties were studied systematically.



Single-Crystalline Fe<sub>3</sub>O<sub>4</sub>? No!





*Angew. Chem. Int. Ed.*  
DOI: 10.1002/anie.200900083

### Nanoparticle Crystallinity

H. Kim, M. Lee, Y. Kim, J. Huh, H. Kim, M. Kim, T. Kim, V. N. Phan, Y.-B. Lee, G.-R. Yi, S. Haam, K. Lee\*

#### Quantitative Assessment of Nanoparticle Single Crystallinity: Palladium-Catalyzed Splitting of Polycrystalline Metal Oxide Nanoparticles

**Crystal gazing:** A simple Pd-catalyzed site-specific nanoetching method was developed to visualize the polycrystalline nature of Fe<sub>3</sub>O<sub>4</sub> (see picture), Fe<sub>2</sub>O<sub>3</sub>, MnFe<sub>2</sub>O<sub>4</sub>, CoFe<sub>2</sub>O<sub>4</sub>, and MnO nanoparticle systems. The technique relies on the very fast etching speed of the grain interface within bi- or polycrystalline nanocrystals.



*Chem. Eur. J.*  
DOI: 10.1002/chem.200900310

### Natural Products

F. Yoshimura, M. Sasaki, I. Hattori, K. Komatsu, M. Sakai, K. Tanino, M. Miyashita\*

#### Synthetic Studies of the Zoanthamine Alkaloids: The Total Syntheses of Norzoanthamine and Zoanthamine

**Highly effective:** We report herein the first and highly efficient total syntheses of norzoanthamine and zoanthamine in full detail, which involves stereoselective synthesis of the requisite triene for an intramolecular Diels–Alder reaction via three-component coupling reactions, the intramolecular Diels–Alder reaction, and subsequent crucial bis-aminoacetalization as the key steps.




*ChemSusChem*  
DOI: 10.1002/cssc.200900047

### Biofuels

F. Zaccheria, S. Brini, R. Psaro, N. Scotti, N. Ravasio\*

#### Esterification of Acidic Oils over a Versatile Amorphous Solid Catalyst

**An amorphous SiO<sub>2</sub>–ZrO<sub>2</sub> catalyst** shows high activity in the esterification of free fatty acids contained in vegetable oils while at the same time promoting the transesterification of triglycerides. The catalyst is hence a good candidate for a low-waste deacidification pretreatment or for a one-pot biodiesel production process starting from oils with a high acid content.



On these pages, we feature a selection of the excellent work that has recently been published in our sister journals. If you are reading these pages on a com-

puter, click on any of the items to read the full article. Otherwise please see the DOIs for easy online access through Wiley InterScience.