

Approach to Trisubstituted 3-Aminopyrrole Derivatives by Yb(OTf)₃-Catalyzed [4+1] Annulation of 2-Azadiene with Me₃SiCN

Toshiaki Sasada,^[a] Takuya Sawada,^[a] Reiko Ikeda,^[a] Norio Sakai,^[a] and Takeo Konakahara*^[a]

Keywords: Multicomponent reactions / Nitrogen heterocycles / Annulation / Ytterbium


In the present study, we found a novel approach to the simple and practical synthesis of 3-aminopyrrole derivatives through the four-component coupling reaction of a functionalized silane, a nitrile, an aldehyde, and trimethylsilyl cyanide

by Yb(OTf)₃-catalyzed annulation of a 2-azabutadiene with trimethylsilyl cyanide. In this paper, we also disclose a single-step transformation of the 3-aminopyrrole framework into the pyrrolo[3,4-*b*]pyridin-4-one skeleton.

Introduction


3-Aminopyrrole derivatives are found in many natural products and in a variety of biologically active substances,^[1] such as distamycin A^[2] and netropsin,^[3] and they are useful building blocks in synthetic chemistry.^[4] To date, a number of approaches to the construction of the pyrrole skeleton have been reported by many organic and pharmaceutical chemists,^[5–9] and general synthetic methods for the synthesis of 3-aminopyrrole derivatives are shown in Scheme 1: (i) functional group transformation of pyrrole derivatives such as 3-hydroxypyrroles,^[10] 3-nitropyrroles,^[11] 3-arylazopyrroles,^[12] and pyrrol-3-carboxylic acids^[13] (path a); (ii) 2,3-bond-forming^[14] or 3,4-bond-forming^[15] cyclization (paths b and c); and (iii) 1,2- and 2,3-bond-forming,^[16] 2,3- and 4,5-bond-forming,^[17] 1,5- and 2,3-bond-forming,^[18] and 1,5- and 3,4-bond-forming^[19] annulation (paths d–g). To the best of our knowledge, there are few examples of the preparation of a 3-aminopyrrole derivative by 2,3- and 3,4-bond-forming annulation (path h).^[20] Moreover, the previous methods require two equivalents of TiCl₄ as an additive,^[20a] unavailable reagents such as a tungsten carbene complex^[20b] and a multifunctionalized triazine,^[20c] a multistep synthesis, and low temperature,^[20a] which lead to low selectivity and a decline in product yields.

Thus, on the basis of our previous work,^[21] we envisioned a facile and efficient approach for the preparation of 3-aminopyrrole derivatives, in which the intermolecular coupling reaction of a functionalized silane, a nitrile, and an aldehyde would initially form a 2-azabutadiene derivative followed by the Lewis acid catalyzed annulation of the



Scheme 1. Approach to the synthesis of 3-aminopyrrole derivative.

formed 2-azabutadiene with trimethylsilyl cyanide to produce the desired 3-aminopyrrole derivative (Scheme 2). Herein, we report the efficient and facile synthesis of 3-aminopyrrole derivatives by a four-component coupling reaction from a functionalized silane, a nitrile, an aldehyde, and trimethylsilyl cyanide. We also disclose a novel approach to the single-step synthesis of pyrrolo[3,4-*b*]pyridin-4-one de-



Scheme 2. This work.


[a] Department of Pure and Applied Chemistry, Faculty of Science and Technology, Tokyo University of Science (RIKADAI), Noda, Chiba 278-8510, Japan
Fax: +81-4-7123-9326
E-mail: konaka@rs.noda.tus.ac.jp

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/ejoc.201000241>.

rivatives starting from the 3-aminopyrrole containing an isoxazole group by the reductive ring-opening reaction of the isoxazole ring.

Results and Discussion

On the basis of our previous work, we first examined the one-pot, three-component coupling reaction of a functionalized silane, an aromatic nitrile, and an aromatic aldehyde, leading to the preparation of 2-azabutadiene derivatives (Scheme 3).^[22,23] For example, when the reaction of 3-methyl-5-(trimethylsilyl)methylisoxazole (**1a**) with benzonitrile (**2a**) was carried out in THF at -70°C for 1 h in the presence of *n*BuLi, followed by the addition of benzaldehyde (**3a**), desired 2-azadiene **4a** was obtained in 53% yield. The structure of the 2-azadiene framework was unambiguously determined by spectroscopy and by X-ray structure analysis of isolated compound **4d**.




Scheme 3. One-pot synthesis of 2-azadiene **4a**.

The annulation of prepared 2-azabutadiene **4a** with trimethylsilyl cyanide (Me_3SiCN) to afford 3-aminopyrrole derivative **6a** was examined, and the results are summarized in Table 1. When the model reaction was performed without a Lewis acid catalyst, desired pyrrole derivative **6a** was not obtained (Table 1, Entry 1). Addition of a typical Lewis acid as an additive, such as InCl_3 , ZnCl_2 , $\text{Cu}(\text{OTf})_2$, and AlCl_3 , promoted the annulation to give desired product **6a**, but the yields were rather low (Table 1, Entries 2–5). The structure of the pyrrole skeleton was unambiguously determined by X-ray diffraction of crystallized compound **6g**. To enhance the yield, when $\text{Yb}(\text{OTf})_3$ was employed, the product yield was drastically increased (Table 1, Entries 7–9). Because both Et_2O and toluene gave a slightly decent yield, THF was the most suitable reaction solvent (Table 1, Entries 10 and 11). Consequently, the reaction described in Entry 7 proved to be the optimal conditions for annulation, as shown in Table 1.

To extend the scope of the $\text{Yb}(\text{OTf})_3$ -catalyzed annulation, we next examined the reaction of various 2-azabutadienes with trimethylsilyl cyanide, leading to 3-aminopyrrole derivatives under our optimal conditions. These results are summarized in Table 2. When R^3 was a 4-chlorophenyl group, desired pyrrole derivative **6b** was produced in excellent yield (Table 2, Entry 2). By contrast, when R^3 was a 4-methoxyphenyl group, the desired reaction proceeded, but the yield of product **6c** was moderate (Table 2, Entry 3). Similarly, the use of 2-azabutadienes **4e–i**, containing a 4-methoxyphenyl group or a 4-chlorophenyl group as R^2 ,

Table 1. Examination of synthesis of 3-aminopyrrole **6a**.



Entry	Lewis acid (equiv.)	Solvent	Time [h]	Yield [%] ^[a]
1	none	THF	24	NR
2	InCl_3 (0.2)	THF	24	30
3	ZnCl_2 (0.2)	THF	24	31
4	$\text{Cu}(\text{OTf})_2$ (0.2)	THF	24	trace
5	AlCl_3 (0.2)	THF	24	20
6	$\text{Hf}(\text{OTf})_4$ (0.2)	THF	24	53
7	$\text{Yb}(\text{OTf})_3$ (0.2)	THF	24	(75)
8	$\text{Yb}(\text{OTf})_3$ (0.1)	THF	48	76
9	$\text{Yb}(\text{OTf})_3$ (0.4)	THF	12	72
10	$\text{Yb}(\text{OTf})_3$ (0.2)	Et_2O	24	68
11	$\text{Yb}(\text{OTf})_3$ (0.2)	PhMe	24	65

[a] NMR yield (isolated yield); NR = no reaction.

produced the corresponding pyrrole derivatives **6e–i** in good yields (Table 2, Entries 5–9). On the other hand, 2-azabutadiene **4j**, with a pyridin-3-yl group as R^3 , gave desired product **6j**, but the yield was low (Table 2, Entry 10). Surprisingly, when 2-azabutadiene **4k** having a pyridin-4-yl group instead of a 3-methylisoxazol-5-yl group as R^1 was employed, the reaction was complete in the presence of only 0.05 equiv. of $\text{Yb}(\text{OTf})_3$ within only 2 h to give pyrrole **6k** in 83% yield (Table 2, Entry 11).

Table 2. Single-step synthesis of 3-aminopyrrole **6**.


Reaction scheme showing the conversion of 2-azadiene **4** to 3-aminopyrrole **6** using $\text{Yb}(\text{OTf})_3$ (0.2 equiv.) and Me_3SiCN (1.2 equiv.) in THF at r.t. for 24 h.

Entry	Azadiene 4			Pyrrole 6	Yield [%] ^[a]	
	R^1	R^2	R^3			
1		Ph	Ph	4a	6a	75
2		Ph	4-ClC ₆ H ₄	4b	6b	87
3		Ph	4-MeOC ₆ H ₄	4c	6c	51
4		4-ClC ₆ H ₄	Ph	4d	6d	77
5		4-ClC ₆ H ₄	4-ClC ₆ H ₄	4e	6e	73
6		4-ClC ₆ H ₄	4-MeOC ₆ H ₄	4f	6f	80
7		4-MeOC ₆ H ₄	Ph	4g	6g	75
8		4-MeOC ₆ H ₄	4-ClC ₆ H ₄	4h	6h	83
9		4-MeOC ₆ H ₄	4-MeOC ₆ H ₄	4i	6i	62
10		Ph	3-Py	4j	6j	30
11 ^[b]		4-MeOC ₆ H ₄	Ph	4k	6k	83

[a] Isolated yield. [b] Reaction time = 2 h; $\text{Yb}(\text{OTf})_3$ = 0.05 equiv.


To apply the four-component coupling reaction using a functionalized silane, a nitrile, an aldehyde, and trimethylsilyl cyanide, we next attempted the one-pot synthesis of 3-aminopyrrole derivative **6** (Scheme 4). When the reaction

was carried out under the above optimal conditions, the desired reaction successfully proceeded to give pyrrole derivatives **6a** and **6b** in 53 and 41 % yield, respectively.




Scheme 4. One-pot synthesis of 3-aminopyrrole **6**.

Finally, to illustrate the utility of an isoxazole group, we examined the skeletal transformation of prepared 3-aminopyrroles **6** into pyrrolo[3,4-*b*]pyridin-4-one derivatives **8** (Scheme 5). When product **6a** was treated with 0.5 equiv. of Mo(CO)₆,^[21f] rather than expected pyrrole derivative **7a**, pyrrolopyridine derivative **8a** was obtained in 88 % yield through the reductive ring opening of the isoxazole followed by intramolecular cyclization.



Scheme 5. Synthesis of pyrrolo[3,4-*b*]pyridin-4-ones **8**.

A plausible mechanism for the Yb(OTf)₃-catalyzed annulation is shown in Scheme 6. First, coordination of Yb(OTf)₃ to the nitrogen atom of the 2-azadiene facilitated the attack of the azadiene by a cyanide ion, which led to the formation of intermediate **9**. Practically, when the reaction



Scheme 6. A plausible mechanism.

of azadiene **4a** with Me₃SiCN was treated with a catalytic amount of Yb(OTf)₃, intermediate **9a** was isolated in 70 % yield.^[24] Yb(OTf)₃-catalyzed intramolecular cyclization of intermediate **9** then occurred to produce corresponding intermediate **10** with a shift of the silyl group.^[25] Finally, aromatization of intermediate **10** worked as the driving force to produce the desired 3-aminopyrrole derivative after conventional aqueous workup.

Conclusions

Thus far, we have outlined the process for a novel, simple, and practical synthesis of 3-aminopyrrole derivatives through the Yb(OTf)₃-catalyzed annulation of 2-azabutadiene with trimethylsilyl cyanide, and the four-component coupling reaction of a functionalized silane, a nitrile, an aldehyde, and trimethylsilyl cyanide. Also detailed is the transformation of 3-aminopyrroles with an isoxazolyl group into pyrrolo[3,4-*b*]pyridin-4-one derivatives through a reductive ring-opening reaction and subsequent intramolecular cyclization.

Experimental Section

General Methods: Column chromatography was performed by using Silica gel 60 and aluminum oxide 90 active neutral. THF, Et₂O, and toluene were distilled from sodium–benzophenone and dried with 5 Å MS or 4 Å MS. Prior to use, 3,5-dimethylisoxazole, trimethylsilyl chloride, benzonitrile, and benzaldehyde were distilled. Other materials were commercially available and were used without further purification. Using a previously reported procedure,^[26] 4-(trimethylsilyl)methylpyridine (**1b**) was prepared. All reactions were carried out under a nitrogen atmosphere, unless otherwise noted. ¹H NMR spectra were measured at 500 or 300 MHz by using tetramethylsilane as an internal standard. ¹³C NMR spectra were measured at 125 or 75 MHz by using the center peak of chloroform (δ = 77.0 ppm) as an internal standard. High-resolution mass spectra were measured by using NBA (3-nitrobenzyl alcohol) as a matrix.

1b:^[26] Colorless liquid. B.p. 76 °C/5 Torr (ref.^[27] 73–74 °C/4.5 Torr). ¹H NMR (300 MHz, CDCl₃, 25 °C): δ = −0.05 (s, 9 H, −SiMe₃), 2.03 (s, 2 H, Ar-CH₂-SiMe₃), 6.84 (d, ³J_{H,H} = 4.5 Hz, 2 H, Ar-*H*), 8.32 (d, ³J_{H,H} = 4.5 Hz, 2 H, Ar-*H*) ppm. ¹³C NMR (75 MHz, CDCl₃, 25 °C): δ = −2.1, 27.3, 123.4, 149.2, 150.1 ppm. MS (FA): *m/z* (%) = 166 (100) [M + H].

Procedure for the Synthesis of 1a: To a THF solution (120 mL) of diisopropylamine (55.6 g, 0.550 mol) was added *n*BuLi (1.5 M in hexane, 0.550 mol) at −70 °C, and the mixture was stirred at the same temperature. After 30 min, 3,5-dimethylisoxazole (48.6 g, 0.500 mol) was added dropwise, and the mixture was stirred for 1 h at −70 °C. Trimethylsilyl chloride (54.3 g, 0.500 mol) was gradually added to the solution at −70 °C, and the mixture was stirred for 1 h at the same temperature and then for 1 h at room temperature. To quench the reaction, a saturated aqueous solution of NH₄Cl was added to the mixture. The mixture was extracted several times with Et₂O, and the combined organic extracts were dried with Na₂SO₄, filtered, and then concentrated under reduced pressure. The residue was purified by distillation under reduced pressure to give 3-methyl-5-(trimethylsilyl)methylisoxazole (**1a**; 68.6 g, 81 %) as

a colorless liquid. B.p. 109–110 °C/32 Torr (ref.^[28] 43 °C/0.6 Torr). ¹H NMR (500 MHz, CDCl₃, 25 °C): δ = 0.03 (s, 9 H, -SiMe₃), 2.12 (s, 2 H, Ar-CH₂-SiMe₃), 2.18 (s, 3 H, Ar-Me), 5.57 (s, 1 H, Ar-H) ppm. ¹³C NMR (125 MHz, CDCl₃, 25 °C): δ = -1.8, 11.3, 17.6, 99.9, 159.7, 171.9 ppm. MS (FA): m/z (%) = 170 (100) [M + H]⁺.

General Procedure for the Synthesis of 2-Azadienes 4a–j: To a THF solution (15 mL) of **1a** (847 mg, 5.00 mmol) was added *n*BuLi (1.5 M in hexane, 5.50 mmol) at -70 °C, and the reaction mixture was stirred at the same temperature. After 1 h, benzonitrile (**2a**; 516 mg, 5.00 mmol) was added to the solution by syringe, and the solution was further stirred for 1 h at -70 °C, then for 1 h at room temperature. Benzaldehyde (**3a**; 531 mg, 5.00 mmol) was added to the solution at -70 °C, and the mixture was stirred for 1 h at the same temperature, then for 1 h at room temperature. To quench the reaction, water (30 mL) was added to the mixture at 0 °C. The mixture was extracted several times with AcOEt, and the combined organic extracts were dried with Na₂SO₄, filtered, and then concentrated under reduced pressure. The residue was purified by neutral Al₂O₃ chromatography (hexane/AcOEt) to give **4a** (764 mg, 53%) as a yellow solid (CHCl₃/hexane).

4a: M.p. 87.2–87.4 °C. ¹H NMR (500 MHz, CDCl₃, 25 °C): δ = 2.14 (s, 3 H, Ar-Me), 6.04 (s, 1 H), 6.35 (s, 1 H), 7.26–7.31 (m, 3 H, Ar-H), 7.40–7.48 (m, 5 H, Ar-H), 7.84 (d, ³*J*_{H,H} = 6.5 Hz, 2 H, Ar-H), 8.18 (s, 1 H, -N=CH-Ar) ppm. ¹³C NMR (125 MHz, CDCl₃, 25 °C): δ = 11.5, 101.0, 103.6, 127.0, 128.7, 129.0, 129.1, 132.2, 135.4, 137.1, 153.7, 159.8, 163.9, 167.6 ppm. MS (FA): m/z (%) = 288 (100) [M]⁺. C₁₉H₁₆N₂O (288.34): calcd. C 79.14, H 5.59, N 9.72; found C 79.06, H 5.50, N 9.73.

4b: Yield: 662 mg, 41%, yellow solid (CHCl₃/hexane). M.p. 120.1–121.0 °C. ¹H NMR (500 MHz, CDCl₃, 25 °C): δ = 2.25 (s, 3 H, Ar-Me), 6.09 (s, 1 H), 6.44 (s, 1 H), 7.38–7.42 (m, 3 H, Ar-H), 7.41–7.51 (m, 4 H, Ar-H), 7.87 (d, ³*J*_{H,H} = 8.0 Hz, 2 H, Ar-H), 8.25 (s, 1 H, -N=CH-Ar) ppm. ¹³C NMR (125 MHz, CDCl₃, 25 °C): δ = 11.5, 101.4, 103.7, 127.0, 128.7, 129.2, 129.4, 130.1, 133.9, 137.0, 138.4, 153.4, 159.8, 164.5, 167.4 ppm. MS (FA): m/z (%) = 323 (100) [M + H]⁺. C₁₉H₁₅ClN₂O (322.79): calcd. C 70.70, H 4.68, N 8.68; found C 70.83, H 4.62, N 8.62.

4c: Yield: 1.10 g, 69%, yellow solid (CHCl₃/hexane). M.p. 119.5–120.7 °C. ¹H NMR (300 MHz, CDCl₃, 25 °C): δ = 2.24 (s, 3 H, Ar-Me), 3.90 (s, 3 H, Ar-OMe), 6.13 (s, 1 H), 6.43 (s, 1 H), 7.03 (d, ³*J*_{H,H} = 9.0 Hz, 2 H, Ar-H), 7.37–7.39 (m, 3 H, Ar-H), 7.49–7.53 (m, 2 H, Ar-H), 7.88 (d, ³*J*_{H,H} = 9.0 Hz, 2 H, Ar-H), 8.19 (s, 1 H, -N=CH-Ar) ppm. ¹³C NMR (125 MHz, CDCl₃, 25 °C): δ = 11.5, 55.5, 100.9, 103.4, 114.5, 127.1, 128.4, 128.7, 129.0, 130.8, 137.4, 154.0, 159.8, 163.0, 163.0, 167.8 ppm. MS (FA): m/z (%) = 319 (100) [M + H]⁺. C₂₀H₁₈N₂O₂ (318.37): calcd. C 75.45, H 5.70, N 8.80; found C 75.45, H 5.71, N 8.80.

4d: Yield: 775 mg, 48%, yellow solid (CHCl₃/hexane). M.p. 92.8–92.9 °C. ¹H NMR (500 MHz, CDCl₃, 25 °C): δ = 2.23 (s, 3 H, Ar-Me), 6.11 (s, 1 H), 6.42 (s, 1 H), 7.35 (d, ³*J*_{H,H} = 8.5 Hz, 2 H, Ar-H), 7.44 (d, ³*J*_{H,H} = 8.5 Hz, 2 H, Ar-H), 7.52–7.59 (m, 3 H, Ar-H), 7.93 (d, ³*J*_{H,H} = 7.5 Hz, 2 H, Ar-H), 8.27 (s, 1 H, -N=CH-Ar) ppm. ¹³C NMR (75 MHz, CDCl₃, 25 °C): δ = 11.4, 100.9, 103.7, 128.1, 128.9, 129.0, 132.4, 135.1, 135.2, 135.5, 152.5, 159.8, 164.2, 167.2 ppm. MS (FA): m/z (%) = 323 (100) [M + H]⁺. C₁₉H₁₅ClN₂O (322.79): calcd. C 70.70, H 4.68, N 8.68; found C 70.84, H 4.71, N 8.69.

4e: Yield: 679 mg, 38%, yellow solid (CHCl₃/hexane). M.p. 174.1–175.1 °C. ¹H NMR (500 MHz, CDCl₃, 25 °C): δ = 2.24 (s, 3 H, Ar-Me), 6.07 (s, 1 H), 6.40 (s, 1 H), 7.36 (d, ³*J*_{H,H} = 8.5 Hz, 2 H, Ar-H), 7.43 (d, ³*J*_{H,H} = 8.5 Hz, 2 H, Ar-H), 7.50 (d, ³*J*_{H,H} = 8.5 Hz,

2 H, Ar-H), 7.86 (d, ³*J*_{H,H} = 8.5 Hz, 2 H, Ar-H), 8.23 (s, 1 H, -N=CH-Ar) ppm. ¹³C NMR (125 MHz, CDCl₃, 25 °C): δ = 11.4, 101.3, 103.8, 128.1, 128.9, 129.4, 130.2, 133.7, 135.2, 135.4, 138.6, 152.3, 159.8, 162.8, 167.1 ppm. MS (FA): m/z (%) = 358 (100) [M + H]⁺. C₁₉H₁₄Cl₂N₂O (357.23): calcd. C 63.88, H 3.95, N 7.84; found C 63.95, H 3.97, N 7.80.

4f: Yield: 759 mg, 43%, yellow solid (CHCl₃/hexane). M.p. 158.7–160.1 °C. ¹H NMR (300 MHz, CDCl₃, 25 °C): δ = 2.24 (s, 3 H, Ar-Me), 3.90 (s, 3 H, Ar-OMe), 6.11 (s, 1 H), 6.40 (s, 1 H), 7.03 (d, ³*J*_{H,H} = 8.7 Hz, 2 H, Ar-H), 7.35 (d, ³*J*_{H,H} = 8.7 Hz, 2 H, Ar-H), 7.45 (d, ³*J*_{H,H} = 8.7 Hz, 2 H, Ar-H), 7.88 (d, ³*J*_{H,H} = 8.7 Hz, 2 H, Ar-H), 8.17 (s, 1 H, -N=CH-Ar) ppm. ¹³C NMR (75 MHz, CDCl₃, 25 °C): δ = 11.5, 55.5, 100.8, 103.5, 114.5, 128.2, 128.8, 130.9, 135.0, 135.8, 152.8, 159.8, 163.1, 163.3, 167.5 ppm. MS (FA): m/z (%) = 353 (100) [M + H]⁺. C₂₀H₁₇ClN₂O₂ (352.81): calcd. C 68.09, H 4.86, N 7.94; found C 67.76, H 4.93, N 7.86.

4g: Yield: 796 mg, 50%, yellow solid (CHCl₃/hexane). M.p. 123.8–124.9 °C. ¹H NMR (500 MHz, CDCl₃, 25 °C): δ = 2.22 (s, 3 H, Ar-Me), 3.84 (s, 3 H, Ar-OMe), 6.05 (s, 1 H), 6.38 (s, 1 H), 6.91 (d, ³*J*_{H,H} = 8.5 Hz, 2 H, Ar-H), 7.46 (d, ³*J*_{H,H} = 8.5 Hz, 2 H, Ar-H), 7.51–7.57 (m, 3 H, Ar-H), 7.94 (d, ³*J*_{H,H} = 7.0 Hz, 2 H, Ar-H), 8.28 (s, 1 H, -N=CH-Ar) ppm. ¹³C NMR (75 MHz, CDCl₃, 25 °C): δ = 11.5, 55.3, 99.0, 103.0, 114.1, 128.3, 129.0, 129.0, 129.5, 132.2, 135.5, 153.6, 159.8, 160.4, 163.9, 167.9 ppm. MS (FA): m/z (%) = 319 (100) [M + H]⁺. HRMS (FAB): calcd. for C₂₀H₁₉N₂O₂ [M + H]⁺ 319.1447; found 319.1467.

4h: Yield: 529 mg, 30%, yellow solid (CHCl₃/hexane). M.p. 178.7–179.3 °C. ¹H NMR (500 MHz, CDCl₃, 25 °C): δ = 2.23 (s, 3 H, Ar-Me), 3.84 (s, 3 H, Ar-OMe), 6.01 (s, 1 H), 6.36 (s, 1 H), 6.92 (d, ³*J*_{H,H} = 8.5 Hz, 2 H, Ar-H), 7.43 (d, ³*J*_{H,H} = 8.5 Hz, 2 H, Ar-H), 7.50 (d, ³*J*_{H,H} = 8.5 Hz, 2 H, Ar-H), 7.87 (d, ³*J*_{H,H} = 8.5 Hz, 2 H, Ar-H), 8.25 (s, 1 H, -N=CH-Ar) ppm. ¹³C NMR (125 MHz, CDCl₃, 25 °C): δ = 11.5, 55.4, 99.4, 103.1, 114.1, 128.3, 129.3, 129.4, 130.1, 133.9, 138.4, 153.3, 159.8, 160.5, 162.4, 167.7 ppm. MS (FA): m/z (%) = 353 (100) [M + H]⁺. C₂₀H₁₇ClN₂O₂ (352.81): calcd. C 68.09, H 4.86, N 7.94; found C 67.99, H 5.18, N 7.94.

4i: Yield: 610 mg, 35%, yellow oil. ¹H NMR (500 MHz, CDCl₃, 25 °C): δ = 2.22 (s, 3 H, Ar-Me), 3.82 (s, 3 H, Ar-OMe), 3.89 (s, 3 H, Ar-OMe), 6.05 (s, 1 H), 6.35 (s, 1 H), 6.90 (d, ³*J*_{H,H} = 9.0 Hz, 2 H, Ar-H), 7.03 (d, ³*J*_{H,H} = 9.0 Hz, 2 H, Ar-H), 7.45 (d, ³*J*_{H,H} = 9.0 Hz, 2 H, Ar-H), 7.88 (d, ³*J*_{H,H} = 9.0 Hz, 2 H, Ar-H), 8.18 (s, 1 H, -N=CH-Ar) ppm. ¹³C NMR (125 MHz, CDCl₃, 25 °C): δ = 11.4, 55.3, 55.4, 98.8, 102.7, 114.0, 114.4, 128.2, 128.4, 129.7, 130.7, 153.9, 159.7, 160.3, 162.9, 168.0 ppm. MS (FA): m/z (%) = 349 (100) [M + H]⁺. HRMS (FAB): calcd. for C₂₁H₂₀N₂O₃ [M]⁺ 348.1474; found 348.1501.

4j: Yield: 477 mg, 33%, yellow oil. ¹H NMR (500 MHz, CDCl₃, 25 °C): δ = 2.25 (s, 3 H, Ar-Me), 6.15 (s, 1 H), 6.44 (s, 1 H), 7.38–7.42 (m, 3 H, Ar-H), 7.46–7.49 (m, 3 H, Ar-H), 8.31 (td, ³*J*_{H,H} = 8.0 Hz, ⁴*J*_{H,H} = 2.0 Hz, 1 H, Ar-H), 8.34 (s, 1 H, -N=CH-Ar), 8.76 (dd, ³*J*_{H,H} = 4.5 Hz, ⁴*J*_{H,H} = 2.0 Hz, 1 H, Ar-H), 9.05 (d, ⁴*J*_{H,H} = 2.0 Hz, 1 H, Ar-H) ppm. ¹³C NMR (125 MHz, CDCl₃, 25 °C): δ = 11.3, 102.1, 104.0, 123.9, 126.9, 128.6, 129.1, 130.9, 135.1, 136.6, 150.8, 152.6, 153.0, 159.6, 161.1, 167.0 ppm. MS (FA): m/z (%) = 290 (100) [M + H]⁺. HRMS (FAB): calcd. for C₁₈H₁₆N₃O [M + H]⁺ 290.1293; found 290.1289.

Procedure for the Synthesis of 2-Azadiene 4k: To a THF solution (15 mL) of 4-(trimethylsilyl)methylpyridine (**1b**, 457 mg, 5.00 mmol) was added MeLi (1 M in Et₂O, 5.50 mmol) at room temperature, and the reaction mixture was heated at 60 °C. After 1 h, anisonitrile (666 mg, 5.00 mmol) was added to the solution by

syringe at room temperature, and the solution was stirred at the same temperature. After 1 h, benzaldehyde (531 mg, 5.00 mmol) was added to the mixture, and the reaction mixture was stirred for 1 h at room temperature. To quench the reaction, water (30 mL) was added to the mixture at 0 °C. The mixture was extracted several times with AcOEt, and the combined organic extracts were dried with Na₂SO₄, filtered, and then concentrated under reduced pressure. The residue was purified by neutral Al₂O₃ chromatography (hexane/AcOEt) to give **4k** (629 mg, 40%) as a yellow solid (CHCl₃/hexane). M.p. 191.2–192.4 °C. ¹H NMR (500 MHz, CDCl₃, 25 °C): δ = 3.84 (s, 3 H, Ar-OMe), 6.24 (s, 1 H, Ar-CH=CAr-), 6.92 (d, ³J_{H,H} = 9.0 Hz, 2 H, Ar-H), 7.42–7.45 (m, 4 H, Ar-H), 7.49–7.54 (m, 3 H, Ar-H), 7.89 (d, ³J_{H,H} = 6.0 Hz, 2 H, Ar-H), 8.26 (s, 1 H, -N=CH-Ar), 8.47 (d, ³J_{H,H} = 6.0 Hz, 2 H, Ar-H) ppm. ¹³C NMR (125 MHz, CDCl₃, 25 °C): δ = 55.3, 112.6, 114.0, 123.9, 128.6, 128.9, 129.0, 130.7, 132.0, 135.7, 144.2, 149.5, 153.9, 160.1, 163.3 ppm. MS (FA): *m/z* (%) = 315 (100) [M + H]⁺. C₂₁H₁₈N₂O (314.38): calcd. C 80.23, H 5.77, N 8.91; found C 80.04, H 6.15, N 8.75.

General Procedure for the Synthesis of 3-Aminopyrrole 6: To a THF (1 mL) solution of 2-azadiene **4** (0.500 mmol) and trimethylsilyl cyanide (**5**, 59.5 mg, 0.600 mmol) was added Yb(OTf)₃ (62.0 mg, 0.100 mmol) at room temperature, and the reaction mixture was stirred for 24 h at the same temperature. To quench the reaction, a saturated aqueous solution of NaHCO₃ (5 mL) was added to the mixture. The mixture was extracted several times with CHCl₃, and the combined organic extracts were dried with anhydrous Na₂CO₃, filtered, and then concentrated under reduced pressure. The crude product was purified by SiO₂ column chromatography (hexane/AcOEt) to afford **6** in the yields shown in Table 2.

6a: Yield: 118 mg, 75%, colorless solid (CHCl₃/hexane). M.p. 157.9–158.0 °C. ¹H NMR (500 MHz, CDCl₃, 25 °C): δ = 2.20 (s, 3 H, Ar-Me), 3.83 (br., 2 H, Ar-NH₂), 5.79 (s, 1 H, Ar-H), 7.20 (t, ³J_{H,H} = 7.5 Hz, 1 H, Ar-H), 7.33 (t, ³J_{H,H} = 7.5 Hz, 1 H, Ar-H), 7.36–7.45 (m, 6 H, Ar-H), 7.50 (t, ³J_{H,H} = 7.5 Hz, 2 H, Ar-H), 8.24 (br., 1 H, -NH-) ppm. ¹³C NMR (125 MHz, CDCl₃, 25 °C): δ = 11.3, 100.0, 101.8, 116.1, 124.8, 125.7, 127.9, 128.0, 128.7, 129.1, 129.2, 130.5, 132.1, 132.2, 159.6, 166.7 ppm. MS (FA): *m/z* (%) = 315 (100) [M]⁺. HRMS (FAB): calcd. for C₂₀H₁₇N₃O [M]⁺ 315.1372; found 315.1380.

6b: Yield: 152 mg, 87%, pale-yellow solid (CHCl₃/hexane). M.p. 165.7–165.8 °C. ¹H NMR (500 MHz, CDCl₃, 25 °C): δ = 2.21 (s, 3 H, Ar-Me), 3.86 (br., 2 H, Ar-NH₂), 5.79 (s, 1 H, Ar-H), 7.33–7.45 (m, 9 H, Ar-H), 8.16 (br., 1 H, -NH-) ppm. ¹³C NMR (125 MHz, CDCl₃, 25 °C): δ = 11.3, 100.1, 102.1, 115.2, 125.8, 127.9, 128.2, 128.8, 129.3, 129.5, 130.7, 130.9, 131.1, 131.9, 159.7, 166.5 ppm. MS (FA): *m/z* (%) = 349 (100) [M]⁺. C₂₀H₁₆ClN₃O (349.81): calcd. C 68.67, H 4.61, N 12.01; found C 68.51, H 4.61, N 12.01.

6c: Yield: 88.1 mg, 51%, green solid (CHCl₃/hexane). M.p. 148.0–148.7 °C. ¹H NMR (500 MHz, CDCl₃, 25 °C): δ = 2.19 (s, 3 H, Ar-Me), 3.75 (br., 2 H, Ar-NH₂), 3.80 (s, 3 H, Ar-OMe), 5.79 (s, 1 H, Ar-H), 6.94 (d, ³J_{H,H} = 8.5 Hz, 2 H, Ar-H), 7.30 (t, ³J_{H,H} = 7.5 Hz, 1 H, Ar-H), 7.36 (t, ³J_{H,H} = 7.5 Hz, 2 H, Ar-H), 7.40–7.43 (m, 4 H, Ar-H), 8.25 (br., 1 H, -NH-) ppm. ¹³C NMR (125 MHz, CDCl₃, 25 °C): δ = 11.3, 55.3, 100.0, 101.8, 114.6, 116.3, 124.9, 126.4, 127.8, 127.8, 128.0, 128.7, 129.7, 132.2, 157.8, 159.6, 166.8 ppm. MS (FA): *m/z* (%) = 345 (100) [M]⁺. C₂₁H₁₉N₃O₂ (345.39): calcd. C 73.03, H 5.54, N 12.17; found C 73.03, H 5.76, N 12.08.

6d: Yield: 135 mg, 77%, pale-yellow solid (CHCl₃/hexane). M.p. 234.3–234.4 °C. ¹H NMR (500 MHz, CDCl₃, 25 °C): δ = 2.27 (s, 3 H, Ar-Me), 3.87 (br., 2 H, Ar-NH₂), 5.85 (s, 1 H, Ar-H), 7.24 (t, ³J_{H,H} = 8.0 Hz, 1 H, Ar-H), 7.36–7.40 (m, 4 H, Ar-H), 7.44 (t,

³J_{H,H} = 7.5 Hz, 2 H, Ar-H), 7.52 (d, ³J_{H,H} = 7.5 Hz, 2 H, Ar-H), 7.97 (br., 1 H, -NH-) ppm. ¹³C NMR (125 MHz, [D₆]DMSO, 25 °C): δ = 11.0, 101.2, 101.3, 116.5, 124.8, 125.0, 128.2, 128.6, 128.6, 129.5, 129.8, 130.8, 131.7, 132.4, 159.4, 165.4 ppm. MS (FA): *m/z* (%) = 349 (100) [M]⁺. C₂₀H₁₆ClN₃O (349.81): calcd. C 68.67, H 4.61, N 12.01; found C 68.53, H 4.38, N 11.70.

6e: Yield: 140 mg, 73%, pale-yellow solid (CHCl₃/hexane). M.p. 233.8–234.0 °C. ¹H NMR (500 MHz, CDCl₃, 25 °C): δ = 2.26 (s, 3 H, Ar-Me), 3.85 (br., 2 H, Ar-NH₂), 5.83 (s, 1 H, Ar-H), 7.37–7.41 (m, 6 H, Ar-H), 7.45 (d, ³J_{H,H} = 8.5 Hz, 2 H, Ar-H), 7.98 (br., 1 H, -NH-) ppm. ¹³C NMR (125 MHz, CDCl₃, 25 °C): δ = 11.4, 100.4, 102.4, 115.5, 125.9, 129.1, 129.1, 129.5, 129.5, 129.7, 130.3, 130.6, 131.4, 134.2, 159.8, 166.1 ppm. MS (FA): *m/z* (%) = 383 (100) [M]⁺. C₂₀H₁₅Cl₂N₃O (384.26): calcd. C 62.51, H 3.93, N 10.94; found C 62.31, H 3.78, N 10.81.

6f: Yield: 152 mg, 80%, pale-yellow solid (CHCl₃/hexane). M.p. 199.7–200.5 °C. ¹H NMR (500 MHz, [D₆]DMSO, 25 °C): δ = 2.22 (s, 3 H, Ar-Me), 3.78 (s, 3 H, Ar-OMe), 4.01 (br., 2 H, Ar-NH₂), 6.22 (s, 1 H, Ar-H), 6.99 (d, ³J_{H,H} = 9.0 Hz, 2 H, Ar-H), 7.39 (d, ³J_{H,H} = 9.0 Hz, 2 H, Ar-H), 7.44 (d, ³J_{H,H} = 9.0 Hz, 2 H, Ar-H), 7.59 (d, ³J_{H,H} = 9.0 Hz, 2 H, Ar-H), 11.09 (br., 1 H, -NH-) ppm. ¹³C NMR (125 MHz, CDCl₃, 25 °C): δ = 11.0, 55.1, 101.2, 101.2, 114.1, 116.8, 125.1, 126.6, 127.7, 128.2, 128.6, 129.3, 131.0, 131.5, 157.0, 159.4, 165.6 ppm. MS (FA): *m/z* (%) = 380 (100) [M + H]⁺. C₂₁H₁₈ClN₃O₂ (379.84): calcd. C 66.40, H 4.78, N 11.06; found C 66.02, H 4.54, N 10.77.

6g: Yield: 130 mg, 75%, pale-yellow solid (CHCl₃/hexane). M.p. 195.3–195.6 °C. ¹H NMR (300 MHz, CDCl₃, 25 °C): δ = 2.23 (s, 3 H, Ar-Me), 3.84 (s, 3 H, Ar-OMe), 3.91 (br., 2 H, Ar-NH₂), 5.77 (s, 1 H, Ar-H), 6.94 (d, ³J_{H,H} = 7.5 Hz, 2 H, Ar-H), 7.23 (t, ³J_{H,H} = 7.5 Hz, 1 H, Ar-H), 7.36–7.44 (m, 4 H, Ar-H), 7.50 (d, ³J_{H,H} = 7.5 Hz, 2 H, Ar-H), 7.99 (br., 1 H, -NH-) ppm. ¹³C NMR (75 MHz, CDCl₃, 25 °C): δ = 12.4, 55.3, 99.6, 101.7, 114.2, 115.5, 124.5, 124.6, 125.6, 129.0, 129.2, 129.4, 130.6, 132.4, 159.6, 166.9 ppm. MS (FA): *m/z* (%) = 346 (100) [M + H]⁺. HRMS (FAB): calcd. for C₂₁H₂₀N₃O₂ [M + H]⁺ 346.1556; found 346.1585.

6h: Yield: 158 mg, 83%, pale-yellow solid (CHCl₃/hexane). M.p. 212.5–212.9 °C. ¹H NMR (500 MHz, [D₆]DMSO, 25 °C): δ = 2.20 (s, 3 H, Ar-Me), 3.78 (s, 3 H, Ar-OMe), 4.20 (br., 2 H, Ar-NH₂), 6.08 (s, 1 H, Ar-H), 6.97 (d, ³J_{H,H} = 9.0 Hz, 2 H, Ar-H), 7.34 (d, ³J_{H,H} = 9.0 Hz, 2 H, Ar-H), 7.41 (d, ³J_{H,H} = 9.0 Hz, 2 H, Ar-H), 7.68 (d, ³J_{H,H} = 9.0 Hz, 2 H, Ar-H), 11.03 (br., 1 H, -NH-) ppm. ¹³C NMR (125 MHz, [D₆]DMSO, 25 °C): δ = 11.0, 55.1, 100.6, 100.7, 113.7, 114.3, 124.3, 126.0, 128.4, 128.4, 129.4, 130.1, 130.8, 131.6, 158.8, 159.2, 165.9 ppm. MS (FA): *m/z* (%) = 379 (100) [M]⁺. C₂₁H₁₈ClN₃O₂ (379.84): calcd. C 66.40, H 4.78, N 11.06; found C 66.13, H 4.57, N 10.79.

6i: Yield: 116 mg, 62%, pale-yellow solid (CHCl₃/hexane). M.p. 158.7–160.0 °C. ¹H NMR (500 MHz, [D₆]DMSO, 25 °C): δ = 2.20 (s, 3 H, Ar-Me), 3.77 (s, 3 H, Ar-OMe), 3.78 (s, 3 H, Ar-OMe), 3.99 (br., 2 H, Ar-NH₂), 6.08 (s, 1 H, Ar-H), 6.96 (d, ³J_{H,H} = 9.0 Hz, 2 H, Ar-H), 6.98 (d, ³J_{H,H} = 9.0 Hz, 2 H, Ar-H), 7.33 (d, ³J_{H,H} = 9.0 Hz, 2 H, Ar-H), 7.58 (d, ³J_{H,H} = 9.0 Hz, 2 H, Ar-H), 10.91 (br., 1 H, -NH-) ppm. ¹³C NMR (125 MHz, [D₆]DMSO, 25 °C): δ = 11.0, 55.1, 55.1, 100.3, 100.5, 113.7, 114.0, 115.6, 124.7, 125.5, 126.3, 128.2, 129.3, 129.3, 156.7, 158.6, 159.1, 166.3 ppm. MS (FA): *m/z* (%) = 375 (100) [M]⁺. C₂₂H₂₁N₃O₃ (375.42): calcd. C 70.38, H 5.64, N 11.19; found C 70.01, H 5.52, N 10.88.

6j: Yield: 47.5 mg, 30%, pale-yellow solid (CHCl₃/hexane). M.p. 242.4–142.5 °C. ¹H NMR (500 MHz, [D₆]DMSO, 25 °C): δ = 2.21 (s, 3 H, Ar-Me), 4.33 (br., 2 H, Ar-NH₂), 6.16 (s, 1 H, Ar-H), 7.32–

7.43 (m, 5 H, Ar-*H*), 8.03 (d, $^3J_{\text{H,H}} = 8.0$ Hz, 1 H, Ar-*H*), 8.32 (d, $^3J_{\text{H,H}} = 5.0$ Hz, 1 H, Ar-*H*), 8.92 (s, 1 H, Ar-*H*), 11.24 (br 1 H, -*NH*-) ppm. ^{13}C NMR (125 MHz, $[\text{D}_6]\text{DMSO}$, 25 °C): $\delta = 11.0$, 101.1, 101.2, 113.1, 123.5, 127.5, 128.0, 128.3, 128.7, 131.0, 131.2, 131.4, 131.8, 145.2, 145.9, 159.3, 165.5 ppm. MS (FA): m/z (%) = 317 (100) $[\text{M}]^+$. $\text{C}_{19}\text{H}_{16}\text{N}_4\text{O}$ (316.36): calcd. C 72.13, H 5.10, N 17.71; found C 71.91, H 4.90, N 17.46.

6k: Yield: 142 mg, 83%, yellow solid ($\text{CHCl}_3/\text{hexane}$). M.p. 219.4–220.0 °C. ^1H NMR (500 MHz, CDCl_3 , 25 °C): $\delta = 3.35$ (br., 2 H, Ar-*NH*₂), 3.80 (s, 3 H, Ar-*OMe*), 6.83 (d, $^3J_{\text{H,H}} = 8.0$ Hz, 2 H, Ar-*H*), 7.20–7.22 (m, 3 H, Ar-*H*), 7.26–7.28 (m, 2 H, Ar-*H*), 7.44 (t, $^3J_{\text{H,H}} = 8.0$ Hz, 2 H, Ar-*H*), 7.55 (d, $^3J_{\text{H,H}} = 8.0$ Hz, 2 H, Ar-*H*), 8.12 (br., 1 H, -*NH*-), 8.50 (d, $^3J_{\text{H,H}} = 5.0$ Hz, 2 H, Ar-*H*) ppm. ^{13}C NMR (125 MHz, $[\text{D}_6]\text{DMSO}$, 25 °C): $\delta = 55.1$, 111.0, 113.9, 116.0, 124.3, 124.4, 124.7, 124.8, 128.5, 128.9, 129.0, 129.1, 133.1, 143.3, 149.6, 158.2 ppm. MS (FA): m/z (%) = 342 (100) $[\text{M} + \text{H}]^+$. HRMS (FAB): calcd. for $\text{C}_{22}\text{H}_{20}\text{N}_3\text{O}$ $[\text{M} + \text{H}]^+$ 342.1606; found 342.1607.

General Procedure for the One-pot Synthesis of 3-Aminopyrrole 6:

To a THF solution (5 mL) of **1a** (339 mg, 2.00 mmol) was added *n*BuLi (1.5 M in hexane, 2.20 mmol) at –70 °C, and the reaction mixture was stirred at the same temperature. After 1 h, benzonitrile (**2a**, 206 mg, 2.00 mmol) was added to the solution by syringe, and the solution was further stirred for 1 h at –70 °C, then for 1 h at room temperature. Aldehyde **3** (2.00 mmol) was added to the solution at –70 °C, and the mixture was stirred for 1 h at the same temperature, then for 1 h at room temperature. Trimethylsilyl cyanide (**5**, 59.5 mg, 0.600 mmol) and $\text{Yb}(\text{OTf})_3$ (62.0 mg, 0.100 mmol) were added to the mixture at room temperature, and the reaction mixture was stirred for 24 h at the same temperature. To quench the reaction, a saturated aqueous solution of NaHCO_3 (10 mL) was added to the mixture. The mixture was extracted several times with CHCl_3 , and the combined organic extracts were dried with anhydrous Na_2CO_3 , filtered, and then concentrated under reduced pressure. The crude product was purified by SiO_2 column chromatography (hexane/AcOEt) to afford **6a** and **6b** in 53 and 41 % yield, respectively.

General Procedure for the Synthesis of Pyrrolo[3,4-*b*]pyridin-4-ones 8:

To an MeCN (9 mL) and H_2O (1 mL) solution of pyrrole **5** (0.500 mmol) was added $\text{Mo}(\text{CO})_6$ (66.0 mg, 0.250 mmol). The mixture was stirred at 100 °C. After 2.5 h, the volatile compounds were removed by evaporation *in vacuo*. The residue was purified by SiO_2 column chromatography (AcOEt) to give **8**.

8a: Yield: 132 mg, 88%, pale-yellow solid ($\text{CHCl}_3/\text{hexane}$). M.p. 188.6–189.7 °C. ^1H NMR (300 MHz, $[\text{D}_6]\text{DMSO}$, 25 °C): $\delta = 2.28$ (s, 3 H Ar-*Me*), 5.49 (s, 1 H, Ar-*H*), 7.24–7.49 (m, 6 H, Ar-*H*), 7.69 (d, $^3J_{\text{H,H}} = 7.5$ Hz, 2 H, Ar-*H*), 8.04 (d, $^3J_{\text{H,H}} = 7.5$ Hz, 2 H, Ar-*H*), 10.46 (br., 1 H, -*NH*-), 12.04 (br., 1 H, -*NH*-) ppm. ^{13}C NMR (75 MHz, $[\text{D}_6]\text{DMSO}$, 25 °C): $\delta = 19.4$, 106.4, 113.5, 114.1, 126.1, 126.6, 126.8, 127.3, 127.5, 127.8, 128.6, 129.0, 131.0, 131.7, 149.0, 176.5 ppm. MS (FA): m/z (%) = 301 (100) $[\text{M} + \text{H}]^+$. HRMS (FAB): calcd. for $\text{C}_{20}\text{H}_{17}\text{N}_2\text{O}$ $[\text{M} + \text{H}]^+$ 301.1341; found 301.1347.

8b: Yield: 151 mg, 90%, pale-yellow solid ($\text{CHCl}_3/\text{hexane}$). M.p. 292 °C (decomp.). ^1H NMR (500 MHz, $[\text{D}_6]\text{DMSO}$, 25 °C): $\delta = 2.27$ (s, 3 H, Ar-*Me*), 5.48 (s, 1 H, Ar-*H*), 7.27 (t, $^3J_{\text{H,H}} = 8.0$ Hz, 1 H, Ar-*H*), 7.38 (t, $^3J_{\text{H,H}} = 8.0$ Hz, 2 H, Ar-*H*), 7.51 (d, $^3J_{\text{H,H}} = 8.0$ Hz, 2 H, Ar-*H*), 7.69 (d, $^3J_{\text{H,H}} = 8.0$ Hz, 2 H, Ar-*H*), 8.01 (d, $^3J_{\text{H,H}} = 8.0$ Hz, 2 H, Ar-*H*), 10.45 (br., 1 H, -*NH*-), 12.07 (br., 1 H, -*NH*-) ppm. ^{13}C NMR (125 MHz, $[\text{D}_6]\text{DMSO}$, 25 °C): $\delta = 19.4$, 106.6, 112.8, 113.4, 126.8, 127.4, 127.8, 128.1, 128.4, 128.5, 129.0, 129.8, 130.3, 131.6, 148.9, 176.3 ppm. MS (FA): m/z (%) = 335

(100) $[\text{M} + \text{H}]^+$. HRMS (FAB): calcd. for $\text{C}_{20}\text{H}_{16}\text{ClN}_2\text{O}$ $[\text{M} + \text{H}]^+$ 335.0951; found 335.0956.

8c: Yield: 164 mg, 99%, pale-yellow solid ($\text{CHCl}_3/\text{hexane}$). M.p. 256.6–257.0 °C. ^1H NMR (500 MHz, $[\text{D}_6]\text{DMSO}$, 25 °C): $\delta = 2.26$ (s, 3 H, Ar-*Me*), 3.79 (s, 3 H, Ar-*OMe*), 5.45 (s, 1 H, Ar-*H*), 6.94 (d, $^3J_{\text{H,H}} = 8.5$ Hz, 2 H, Ar-*H*), 7.26 (t, $^3J_{\text{H,H}} = 7.5$ Hz, 1 H, Ar-*H*), 7.46 (t, $^3J_{\text{H,H}} = 7.5$ Hz, 2 H, Ar-*H*), 7.67 (d, $^3J_{\text{H,H}} = 7.5$ Hz, 2 H, Ar-*H*), 7.99 (d, $^3J_{\text{H,H}} = 8.5$ Hz, 2 H, Ar-*H*), 10.37 (br., 1 H, -*NH*-), 11.88 (br., 1 H, -*NH*-) ppm. ^{13}C NMR (125 MHz, $[\text{D}_6]\text{DMSO}$, 25 °C): $\delta = 19.4$, 55.1, 106.3, 112.9, 113.1, 124.4, 125.7, 126.6, 127.5, 127.6, 128.6, 130.3, 131.1, 148.8, 158.3, 176.6 ppm. MS (FA): m/z (%) = 331 (100) $[\text{M} + \text{H}]^+$. HRMS (FAB): calcd. for $\text{C}_{21}\text{H}_{19}\text{N}_2\text{O}_2$ $[\text{M} + \text{H}]^+$ 331.1447; found 331.1449.

Supporting Information (see footnote on the first page of this article): Detailed experimental procedures and characterization data for novel compounds, ORTEP diagram of **4d** and **6g**, X-ray data for **4d** and **6g**, and copies of the ^1H and ^{13}C NMR spectra of the novel products.


Acknowledgments

This work was partially supported by a grant from the Japan Private School Promotion Foundation (2008 and 2009) and a Grant-in-Aid for Scientific Research from Ministry of Education, Culture, Sports, Science and Technology (MEXT) (16550148, 2004–2005; 21590025, 2009–2011) and a fund for the High-Tech Research Center Project for Private Universities, which is a matching fund subsidy from MEXT (2000–2004 and 2005–2007).


- [1] For selected recent reviews and papers, see: a) P. G. Baraldi, D. Preti, F. Fruttarolo, M. A. Tabrizi, R. Romagnoli, *Bioorg. Med. Chem.* **2007**, *15*, 17–35; b) T. Bando, H. Sugiyama, *Acc. Chem. Res.* **2006**, *39*, 935–944; c) P. G. Baraldi, A. N. Zaid, D. Preti, F. Fruttarolo, M. A. Tabrizi, A. Iaconinoto, M. G. Pavani, M. D. Carrion, C. L. Cara, R. Romagnoli, *ARKIVOC* **2006**, 20–34; d) C. Dose, M. E. Farkas, D. M. Chenoweth, P. B. Dervan, *J. Am. Chem. Soc.* **2008**, *130*, 6859–6866; e) K. Schneider, S. Keller, F. E. Wolter, L. Roeglin, W. Beil, O. Seitz, G. Nicholson, C. Bruntner, J. Riedlinger, H.-P. Fiedler, R. D. Suessmuth, *Angew. Chem. Int. Ed.* **2008**, *47*, 3258–3261; f) K.-i. Wakabayashi, H. Miyachi, Y. Hashimoto, A. Tanatani, *Bioorg. Med. Chem.* **2005**, *13*, 2837–2846; g) W. Hu, R. W. Buerli, J. A. Kaizerman, K. W. Johnson, M. I. Gross, M. Iwamoto, P. Jones, D. Lofland, S. Difuntorum, H. Chen, B. Bozdogan, P. C. Appelbaum, H. E. Moser, *J. Med. Chem.* **2004**, *47*, 4352–4355; h) I. Beria, P. G. Baraldi, P. Cozzi, M. Caldarelli, C. Geroni, S. Marchini, N. Mongelli, R. Romagnoli, *J. Med. Chem.* **2004**, *47*, 2611–2623; i) T. Bando, A. Narita, I. Saito, H. Sugiyama, *J. Am. Chem. Soc.* **2003**, *125*, 3471–3485; j) J. A. Kaizerman, M. I. Gross, Y. Ge, S. White, W. Hu, J.-X. Duan, E. E. Baird, K. W. Johnson, R. D. Tanaka, H. E. Moser, R. W. Buerli, *J. Med. Chem.* **2003**, *46*, 3914–3929; k) M. T. Cocco, C. Congiu, V. Onnis, *Bioorg. Med. Chem.* **2003**, *11*, 495–503; l) P. G. Baraldi, I. Beria, P. Cozzi, N. Bianchi, R. Gambari, R. Romagnoli, *Bioorg. Med. Chem.* **2003**, *11*, 965–975.
- [2] F. Arcamone, S. Penco, P. Orezzi, V. Nicoletta, A. Pirelli, *Nature* **1964**, *203*, 1064–1065.
- [3] A. C. Finlay, F. A. Hochstein, B. A. Sobin, F. X. Murphy, *J. Am. Chem. Soc.* **1951**, *73*, 341–343.
- [4] For selected recent papers, see: a) D. M. Chenoweth, D. A. Harki, P. B. Dervan, *J. Am. Chem. Soc.* **2009**, *131*, 7175–7181; b) F. Brucoli, P. W. Howard, D. E. Thurston, *J. Comb. Chem.* **2009**, *11*, 576–586; c) A. Gangjee, W. Li, J. Yang, R. L. Kisliuk, *J. Med. Chem.* **2008**, *51*, 68–76; d) G. Fridkin, W. D. Lubell,

- Org. Lett.* **2008**, *10*, 849–852; e) P. Deaudelin, W. D. Lubell, *Org. Lett.* **2008**, *10*, 2841–2844; f) H. Bekolo, G. Kirsch, *Can. J. Chem.* **2007**, *85*, 1–6; g) C. Rochais, V. Lisowski, P. Dallemagne, S. Rault, *Bioorg. Med. Chem.* **2006**, *14*, 8162–8175; h) T. L. Smalley, A. J. Peat, J. A. Boucheron, S. Dickerson, D. Garrido, F. Preugschat, S. L. Schweiker, S. A. Thomson, T. Y. Wang, *Bioorg. Med. Chem. Lett.* **2006**, *16*, 2091–2094; i) G. Fridkin, W. D. Lubell, *J. Comb. Chem.* **2005**, *7*, 977–986; j) F. J. R. Rombouts, G. Fridkin, W. D. Lubell, *J. Comb. Chem.* **2005**, *7*, 589–598; k) E. Patane, V. Pittala, F. Guerrero, L. Salerno, G. Romeo, M. A. Siracusa, F. Russo, F. Manetti, M. Botta, I. Mereghetti, A. Cagnotto, T. Mennini, *J. Med. Chem.* **2005**, *48*, 2420–2431; l) B. K. S. Yeung, D. L. Boger, *J. Org. Chem.* **2003**, *68*, 5249–5253.
- [5] For selected recent books and reviews, see: a) J. J. Li (Ed.), *Name Reactions in Heterocyclic Chemistry*, Wiley, Hoboken, NJ, **2005**, pp. 69–97; b) D. S. Black, *Sci. Synth.* **2002**, *9*, 441–552; c) H. Fan, J. Peng, M. T. Hamann, J.-F. Hu, *Chem. Rev.* **2008**, *108*, 264–287; d) H. Kagoshima, K. Fuchibe, T. Akiyama, *Chem. Rev.* **2007**, *7*, 104–114; e) R. A. Stockman, *Ann. Rep. Prog. Chem. Sect. B: Org. Chem.* **2007**, *103*, 107–124; f) J. S. Fisk, R. A. Mosey, J. J. Tepe, *Chem. Soc. Rev.* **2007**, *36*, 1432–1440; g) F. Bellina, R. Rossi, *Tetrahedron* **2006**, *62*, 7213–7256; h) S. Manginckx, N. Giubellina, N. De Kimpe, *Chem. Rev.* **2004**, *104*, 2353–2399; i) B. Stanovnik, J. Svete, *Chem. Rev.* **2004**, *104*, 2433–2480; j) A. Fürstner, *Angew. Chem. Int. Ed.* **2003**, *42*, 3582–3603.
- [6] For recent selected reviews on the multicomponent synthesis of pyrroles, see: a) D. M. D'Souza, T. J. J. Müller, *Chem. Soc. Rev.* **2007**, *36*, 1095–1108; b) G. Balme, *Angew. Chem. Int. Ed.* **2004**, *43*, 6238–6241.
- [7] For recent selected papers on the multicomponent synthesis of pyrroles, see: a) Y. Lu, B. A. Arndtsen, *Org. Lett.* **2009**, *11*, 1369–1372; b) B. Tan, Z. Shi, P. J. Chua, Y. Li, G. Zhong, *Angew. Chem. Int. Ed.* **2009**, *48*, 758–761; c) Y. Lu, B. A. Arndtsen, *Org. Lett.* **2009**, *11*, 1369–1372; d) P. Fontaine, G. Masson, J. Zhu, *Org. Lett.* **2009**, *11*, 1555–1558; e) A. Alizadeh, M. Babaki, N. Zohreh, *Tetrahedron* **2009**, *65*, 1704–1707; f) I. Yavari, E. Kowsari, *Synlett* **2008**, 897–899; g) J. S. Yadav, B. V. S. Reddy, M. Srinivas, C. Divyavani, S. J. Basha, A. V. S. Sarma, *J. Org. Chem.* **2008**, *73*, 3252–3254; h) B. Khalili, P. Jajarmi, B. Eftekhari-Sis, M. M. Hashemi, *J. Org. Chem.* **2008**, *73*, 2090–2095; i) G. Dou, C. Shi, D. Shi, *J. Comb. Chem.* **2008**, *10*, 810–813; j) M. D'Hooghe, C. Buyck, J. Contreras, N. De Kimpe, *Org. Biomol. Chem.* **2008**, *6*, 3667–3669; k) S. Dey, C. Pal, D. Nandi, V. S. Giri, M. Zaidlewicz, M. Krzeminski, L. Smentek, B. A. Hess Jr., J. Gawronski, M. Kwit, N. J. Babu, A. Nangia, P. Jaisankar, *Org. Lett.* **2008**, *10*, 1373–1376; l) J. H. Chang, H. Shin, *Org. Process Res. Dev.* **2008**, *12*, 291–293; m) A. Alizadeh, A. Rezvanian, H. R. Bijanzadeh, *Synthesis* **2008**, 725–728; n) D. J. St. Cyr, N. Martin, B. A. Arndtsen, *Org. Lett.* **2007**, *9*, 449–452; o) D. J. St. Cyr, B. A. Arndtsen, *J. Am. Chem. Soc.* **2007**, *129*, 12366–12367; p) B. C. Milgram, K. Eskildsen, S. M. Richter, W. R. Scheidt, K. A. Scheidt, *J. Org. Chem.* **2007**, *72*, 3941–3944; q) C. V. Galliford, K. A. Scheidt, *J. Org. Chem.* **2007**, *72*, 1811–1813; r) V. Cadierno, J. Gimeno, N. Nebra, *Chem. Eur. J.* **2007**, *13*, 9973–9981; s) I. Bergner, T. Opatz, *J. Org. Chem.* **2007**, *72*, 7083–7090; t) M. Shimizu, A. Takahashi, S. Kawai, *Org. Lett.* **2006**, *8*, 3585–3587; u) A. E. Mattson, A. R. Bharadwaj, A. M. Zuhl, K. A. Scheidt, *J. Org. Chem.* **2006**, *71*, 5715–5724; v) I. Bianchi, R. Forlani, G. Minetto, I. Peretto, N. Regalia, M. Taddei, L. F. Raveglia, *J. Comb. Chem.* **2006**, *8*, 491–499; w) D. Suzuki, Y. Nobe, Y. Watai, R. Tanaka, Y. Takayama, F. Sato, H. Urabe, *J. Am. Chem. Soc.* **2005**, *127*, 7474–7479; x) G. Minetto, L. F. Raveglia, M. Taddei, *Org. Lett.* **2004**, *6*, 389–392; y) K. A. Hansford, V. Zanzarova, A. Doerr, W. D. Lubell, *J. Comb. Chem.* **2004**, *6*, 893–898; z) R. Dhawan, B. A. Arndtsen, *J. Am. Chem. Soc.* **2004**, *126*, 468–469; aa) A. R. Bharadwaj, K. A. Scheidt, *Org. Lett.* **2004**, *6*, 2465–2468; Y. Nishibayashi, M. Yoshikawa, Y. Inada, M. D. Milton, M. Hidai, S. Uemura, *Angew. Chem. Int. Ed.* **2003**, *42*, 2681–2684; ac) R. U. Braun, K. Zeitler, T. J. J. Müller, *Org. Lett.* **2001**, *3*, 3297–3300.
- [8] For recent selected reviews on the metal-mediated synthesis of pyrroles, see: a) N. T. Patil, Y. Yamamoto, *Chem. Rev.* **2008**, *108*, 3395–3442; b) S. F. Kirsch, *Synthesis* **2008**, 3183–3204; c) N. T. Patil, Y. Yamamoto, *ARKIVOC* **2007**, 121–141; d) G. Zeni, R. C. Larock, *Chem. Rev.* **2006**, *106*, 4644–4680; e) A. L. Odom, *Dalton Trans.* **2005**, 225–233; f) G. Zeni, R. C. Larock, *Chem. Rev.* **2004**, *104*, 2285–2309; g) M. Mori, *J. Organomet. Chem.* **2004**, 689, 4210–4227; h) K. Narasaka, *Pure Appl. Chem.* **2003**, *75*, 19–28; i) M. Kitamura, K. Narasaka, *Chem. Rec.* **2002**, *2*, 268–277.
- [9] For recent selected papers on the metal-mediated synthesis of pyrroles, see: a) A. V. Lygin, O. V. Larionov, V. S. Korotkov, A. de Meijere, *Chem. Eur. J.* **2009**, *15*, 227–236; b) S. Kiren, X. Hong, C. A. Leverett, A. Padwa, *Org. Lett.* **2009**, *11*, 1233–1235; c) Y. Kim, J. Kim, S. B. Park, *Org. Lett.* **2009**, *11*, 17–20; d) P. W. Davies, N. Martin, *Org. Lett.* **2009**, *11*, 2293–2296; e) Y.-J. Bian, X.-Y. Liu, K.-G. Ji, X.-Z. Shu, L.-N. Guo, Y.-M. Liang, *Tetrahedron* **2009**, *65*, 1424–1429; f) L. Ackermann, R. Sandmann, L. T. Kaspar, *Org. Lett.* **2009**, *11*, 2031–2034; g) R. Surmont, G. Verniest, F. Colpaert, G. Macdonald, J. W. Thuring, F. Deroose, N. De Kimpe, *J. Org. Chem.* **2009**, *74*, 1377–1380; h) A. S. Dudnik, A. W. Sromek, M. Rubina, J. T. Kim, A. V. Kel'in, V. Gevorgyan, *J. Am. Chem. Soc.* **2008**, *130*, 1440–1452; i) Y. Lu, B. A. Arndtsen, *Angew. Chem. Int. Ed.* **2008**, *47*, 5430–5433; j) J.-L. Zhu, Y.-H. Chan, *Synlett* **2008**, 1250–1254; k) Z. Zhang, J. Zhang, J. Tan, Z. Wang, *J. Org. Chem.* **2008**, *73*, 5180–5182; l) G. E. Veitch, K. L. Bridgwood, K. Rands-Trevor, S. V. Ley, *Synlett* **2008**, 2597–2600; m) M.-J. R. P. Queiroz, A. Begouin, G. Pereira, P. M. T. Ferreira, *Tetrahedron* **2008**, *64*, 10714–10720; n) V. Lavallo, G. D. Frey, B. Donnadieu, M. Soleilhavoup, G. Bertrand, *Angew. Chem. Int. Ed.* **2008**, *47*, 5224–5228; o) S. Chiba, Y.-F. Wang, G. Lapointe, K. Narasaka, *Org. Lett.* **2008**, *10*, 313–316; p) S. Cacchi, G. Fabrizi, E. Filisti, *Org. Lett.* **2008**, *10*, 2629–2632; q) M. R. Rivero, S. L. Buchwald, *Org. Lett.* **2007**, *9*, 973–976; r) X. Yuan, X. Xu, X. Zhou, J. Yuan, L. Mai, Y. Li, *J. Org. Chem.* **2007**, *72*, 1510–1513; s) R. Martin, C. H. Larsen, A. Cuenca, S. L. Buchwald, *Org. Lett.* **2007**, *9*, 3379–3382; t) F. M. Istrate, F. Gagosz, *Org. Lett.* **2007**, *9*, 3181–3184; u) H. Dong, M. Shen, J. E. Redford, B. J. Stokes, A. L. Pumphrey, T. G. Driver, *Org. Lett.* **2007**, *9*, 5191–5194; v) X. Wan, D. Xing, Z. Fang, B. Li, F. Zhao, K. Zhang, L. Yang, Z. Shi, *J. Am. Chem. Soc.* **2006**, *128*, 12046–12047; w) R. Martin, M. R. Rivero, S. L. Buchwald, *Angew. Chem. Int. Ed.* **2006**, *45*, 7079–7082; x) L. Lu, G. Chen, S. Ma, *Org. Lett.* **2006**, *8*, 835–838; y) T. Ishikawa, T. Aikawa, S. Watanabe, S. Saito, *Org. Lett.* **2006**, *8*, 3881–3884; z) K. Hiroya, S. Matsumoto, M. Ashikawa, K. Ogawara, T. Sakamoto, *Org. Lett.* **2006**, *8*, 5349–5352; aa) B. Gabriele, G. Salerno, A. Fazio, L. Veltri, *Adv. Synth. Catal.* **2006**, *348*, 2212–2222; K. Fuchibe, D. Ono, T. Akiyama, *Chem. Commun.* **2006**, 2271–2273; ac) C. Dong, G. Deng, J. Wang, *J. Org. Chem.* **2006**, *71*, 5560–5564; ad) M. L. Crawley, I. Goljer, D. J. Jenkins, J. F. Mehlmann, L. Nogle, R. Dooley, P. E. Mahaney, *Org. Lett.* **2006**, *8*, 5837–5840; ae) J. T. Binder, S. F. Kirsch, *Org. Lett.* **2006**, *8*, 2151–2153; af) O. V. Larionov, A. de Meijere, *Angew. Chem. Int. Ed.* **2005**, *44*, 5664–5667; ag) S. Kamijo, C. Kanazawa, Y. Yamamoto, *J. Am. Chem. Soc.* **2005**, *127*, 9260–9266; ah) D. J. Gorin, N. R. Davis, F. D. Toste, *J. Am. Chem. Soc.* **2005**, *127*, 11260–11261; ai) A. I. Siriwardana, K. K. A. D. S. Kathirarachchi, I. Nakamura, I. D. Gridnev, Y. Yamamoto, *J. Am. Chem. Soc.* **2004**, *126*, 13898–13899; aj) B. Ramanathan, A. J. Keith, D. Armstrong, A. L. Odom, *Org. Lett.* **2004**, *6*, 2957–2960; ak) Y. Zhang, J. W. Herndon, *Org. Lett.* **2003**, *5*, 2043–2045; al) M. Yu, B. L. Pagenkopf, *Org. Lett.* **2003**, *5*, 5099–5101; am) Y. Wang, S. Zhu, *Org. Lett.* **2003**, *5*, 745–748; an) Y. Nishibayashi, M. Yoshikawa, Y. Inada, M. D. Milton, M. Hidai, S. Uemura, *Angew. Chem. Int. Ed.* **2003**, *42*, 2681–2684;

- ao) J. T. Kim, A. V. Kel'in, V. Gevorgyan, *Angew. Chem. Int. Ed.* **2003**, *42*, 98–101; ap) H. Takaya, S. Kojima, S.-I. Murahashi, *Org. Lett.* **2001**, *3*, 421–424; aq) A. V. Kel'in, A. W. Sromek, V. Gevorgyan, *J. Am. Chem. Soc.* **2001**, *123*, 2074–2075.
- [10] For selected papers, see: a) Y. Brouillette, F. J. R. Rombouts, W. D. Lubell, *J. Comb. Chem.* **2006**, *8*, 117–126; b) F.-A. Marcotte, W. D. Lubell, *Org. Lett.* **2002**, *4*, 2601–2603; c) M. Glaser, M. Gruner, H. Viola, *J. Prakt. Chem.* **1994**, *336*, 363–365.
- [11] For selected papers, see: a) D. Breen, A. R. Kennedy, C. J. Suckling, *Org. Biomol. Chem.* **2009**, *7*, 178–186; b) L. Fu, G. W. Gribble, *Synthesis* **2008**, 788–794; c) L. Fu, G. W. Gribble, *Tetrahedron Lett.* **2008**, *49*, 3545–3548; d) M. Valik, J. Malina, L. Palivec, J. Foltynova, M. Tkadlecova, M. Urbanova, V. Brabec, V. Kral, *Tetrahedron* **2006**, *62*, 8591–8600; e) A. M. Almerico, G. Cirrincione, E. Aiello, G. Dattolo, *J. Heterocycl. Chem.* **1989**, *26*, 1631–1633; f) G. Cirrincione, G. Dattolo, A. M. Almerico, E. Aiello, R. A. Jones, W. Hinz, *Tetrahedron* **1987**, *43*, 5225–5228; g) M. Nagarajan, H. Shechter, *J. Org. Chem.* **1984**, *49*, 62–74; h) E. Aiello, G. Dattolo, G. Cirrincione, *J. Chem. Soc. Perkin Trans. 1* **1981**, 1–3; i) L. Grehn, *Chemica Scripta* **1979**, *13*, 78–95.
- [12] For selected papers, see: a) H. Onda, H. Toi, Y. Aoyama, H. Ogoshi, *Tetrahedron Lett.* **1985**, *26*, 4221–4224; b) G. Dattolo, G. Cirrincione, A. M. Almerico, G. Presti, E. Aiello, *Heterocycles* **1984**, *22*, 2269–2276.
- [13] For selected papers, see: F. Fabis, P. Dallemagne, S. Rault, M. Robba, *Org. Prep. Proced. Int.* **1995**, *27*, 236–239.
- [14] For selected papers, see: a) G. Tsolomiti, K. Tsolomiti, A. Tsolomitis, *Heterocycl. Commun.* **2007**, *13*, 143–146; b) C. Rochais, V. Lisowski, P. Dallemagne, S. Rault, *Tetrahedron* **2004**, *60*, 2267–2270; c) L. Selic, B. Stanovnik, *Helv. Chim. Acta* **1998**, *81*, 1634–1639; d) M. Rehwal, H. Schafer, K. Gewald, *Monatsh. Chem.* **1997**, *128*, 933–943.
- [15] For selected papers, see: a) E. Toja, J. Kettenring, B. Goldstein, G. Tarzia, *J. Heterocycl. Chem.* **1986**, *23*, 1561–1564; b) S. M. Bayomi, K. E. Price, J. W. Sowell, *J. Heterocycl. Chem.* **1985**, *22*, 729–734; c) T. Murata, T. Sugawara, K. Ukawa, *Chem. Pharm. Bull.* **1978**, *26*, 3080–3100; d) G. Tarzia, G. Panzone, *Ann. Chim.* **1974**, *64*, 807–824.
- [16] For selected papers, see: a) E. Barnea, S. Majumder, R. J. Staples, A. L. Odom, *Organometallics* **2009**, *28*, 3876–3881; b) A. M. Salaheldin, Z. *Naturforsch., Teil B: Chem. Sci.* **2008**, *63*, 564–570; c) A. W. Erian, S. M. Sherif, N. R. Mohamed, *Synth. Commun.* **2003**, *33*, 1563–1573; d) A. W. Erian, S. M. Sherif, N. R. Mohamed, *ARKIVOC* **2001**, 85–94; e) S. A. Ryndina, A. V. Kadushkin, N. P. Solov'eva, V. G. Granik, *Chem. Heterocycl. Compd.* **2000**, *36*, 1409–1420; f) M. T. Cocco, C. Congiu, V. Onnis, *J. Heterocycl. Chem.* **1995**, *32*, 1679–1682; g) K. Gewald, H. Schaefer, P. Bellmann, U. Hain, *J. Prakt. Chem.* **1992**, *334*, 491–496.
- [17] For selected papers, see: G. Tsolomiti, K. Tsolomiti, A. Tsolomitis, *Heterocycl. Commun.* **2007**, *13*, 235–238.
- [18] For selected papers, see: a) T. Semeraro, C. Mugnaini, F. Corelli, *Tetrahedron Lett.* **2008**, *49*, 5965–5967; b) T. Semeraro, C. Mugnaini, F. Manetti, S. Pasquini, F. Corelli, *Tetrahedron* **2008**, *64*, 11249–11255; c) G. L. Sommen, A. Comel, G. Kirsch, *Synth. Commun.* **2005**, *35*, 693–699; d) N. Chen, Y. Lu, K. Gadamasetti, C. R. Hurt, M. H. Norman, C. Fotsch, *J. Org. Chem.* **2000**, *65*, 2603–2605; e) R. H. Furneaux, P. C. Tyler, *J. Org. Chem.* **1999**, *64*, 8411–8412; f) B. Erdmann, A. Knoll, J. Liebscher, *J. Prakt. Chem.* **1988**, *330*, 1015–1021.
- [19] For selected papers, see: a) Y. Cheng, Z.-M. Kang, Y.-G. Ma, J.-H. Peng, M.-F. Liu, *Tetrahedron* **2008**, *64*, 7362–7368; b) A. El-Sharief, A. Al-Amri, S. Al-Raqa, *J. Sulfur Chem.* **2006**, *27*, 245–263; c) O. A. Attanasi, G. Favi, P. Filippone, A. Golobic, B. Stanovnik, J. Svete, *J. Org. Chem.* **2005**, *70*, 4307–4313; d) Y. A. Ammar, A. M. S. El-Sharief, M. M. Ali, Y. A. Mohamed, S. I. Mohamed, *Phosphorus Sulfur Silicon Relat. Elem.* **2000**, *166*, 173–185; e) G. Dattolo, G. Cirrincione, A. M. Almerico, E. Aiello, *J. Heterocycl. Chem.* **1989**, *26*, 1747–1749.
- [20] a) E. Bandini, G. Martelli, G. Spunta, A. Bongini, M. Panunzio, G. Piersanti, *Tetrahedron: Asymmetry* **1999**, *10*, 1445–1449; b) R. Aumann, E. Kuckert, C. Krueger, K. Angermund, *Angew. Chem.* **1987**, *99*, 587–588; c) G. Seitz, S. Dietrich, R. Dhar, W. Massa, G. Baum, *Arch. Pharm.* **1986**, *319*, 798–807.
- [21] a) T. Sasada, M. Moriuchi, N. Sakai, T. Konakahara, *Eur. J. Org. Chem.* **2009**, 5738–5743; b) T. Sasada, F. Kobayashi, N. Sakai, T. Konakahara, *Org. Lett.* **2009**, *11*, 2161–2164; c) T. Sasada, N. Sakai, T. Konakahara, *J. Org. Chem.* **2008**, *73*, 6905–6908; d) H. Suzuki, N. Sakai, R. Iwahara, T. Fujiwaka, M. Satoh, A. Kakehi, T. Konakahara, *J. Org. Chem.* **2007**, *72*, 5878–5881; e) N. Sakai, D. Aoki, T. Hamajima, T. Konakahara, *Tetrahedron Lett.* **2006**, *47*, 1261–1265; f) N. Sakai, Y. Aoki, T. Sasada, T. Konakahara, *Org. Lett.* **2005**, *7*, 4705–4708.
- [22] See details in the Supporting Information.
- [23] For selected papers on the synthesis of 2-azadienes, see: a) J. Capra, T. Le Gall, *Synlett* **2010**, 441–444; b) N. De Kimpe, M. Nagy, M. Boeykens, D. Van der Schueren, *J. Org. Chem.* **1992**, *57*, 5761–5764; c) N. De Kimpe, Z. Yao, M. Boeykens, M. Nagy, *Tetrahedron Lett.* **1990**, *31*, 2771–2774; d) N. De Kimpe, E. Stanoeva, R. Verhe, N. Schamp, *Synthesis* **1988**, 587–592.
- [24] When the reaction of 2-azadiene **4a** with trimethylsilyl cyanide (**5**) was carried out in THF at room temperature for 2 h in the presence of Yb(OTf)₃ (0.2 equiv.), intermediate **9a** and 3-aminopyrrole **6a** were obtained in 70 and 25% yield, respectively. Also, see details in the Supporting Information.



- [25] When the intramolecular cyclization of intermediate **9a** was carried out in THF at room temperature for 24 h in the presence of Yb(OTf)₃ (0.2 equiv.), 3-aminopyrrole **6a** was obtained in 80% yield. Also, see details in the Supporting Information.



- [26] A. Happer, J. Ng, B. Pool, J. White, *J. Organomet. Chem.* **2002**, *659*, 10–14.
- [27] C. Eaborn, R. A. Shaw, *J. Chem. Soc.* **1955**, 3306–3308.
- [28] R. Nesi, A. Ricci, M. Taddei, P. Tedeschi, *J. Organomet. Chem.* **1980**, *195*, 275–283.

Received: February 22, 2010
Published Online: June 9, 2010