

Short communication

Morphine causes a delayed increase in glutamate receptor functioning in the nucleus accumbens core

Edwin H. Jacobs^{a,1}, George Wardeh^a, August B. Smit^b, Anton N.M. Schoffelmeer^{a,*}

^aResearch Institute Neurosciences Vrije Universiteit, Drug Abuse Program, Department of Medical Pharmacology, Free University Medical Center, VUmc, Van der Boechorststraat 7, 1081 BT Amsterdam, The Netherlands

^bDepartment of Molecular and Cellular Neurobiology, Faculty of Earth and Life Sciences, Vrije Universiteit, De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands

Received 1 November 2004; received in revised form 8 February 2005; accepted 10 February 2005

Abstract

Enhanced excitatory neurotransmission in the mesocorticolimbic system may contribute to the persistence of addiction behaviour. Here, we demonstrated that glutamate-, *N*-methyl-D-aspartate (NMDA)- and α -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid (AMPA)-induced [³H]- γ -aminobutyric acid (GABA) release from superfused rat nucleus accumbens core slices is profoundly enhanced 3 weeks, but not 3 days, after a single s.c. morphine injection. This delayed increase in glutamate receptor functioning is associated with enhanced gene transcript levels of ionotropic NMDA and AMPA/kainate receptor subunits. These data reveal that morphine may progressively enhance glutamate neurotransmission within the nucleus accumbens core subsequent to drug exposure.

© 2005 Elsevier B.V. All rights reserved.

Keywords: Ionotropic glutamate receptor; Gene expression; GABA release; Nucleus accumbens core

1. Introduction

Exposure to psychostimulants and opiates causes time-dependent neuroadaptive changes within the mesocorticolimbic system that may underlie certain behavioural characteristics of drug addiction such as sensitization and stimulus-induced drug-seeking (Vanderschuren and Kalivas, 2000; Nestler, 2001; Robinson and Berridge, 2003). The nucleus accumbens, a point of convergence integrating dopamine and glutamate neurotransmission, is of particular importance in this respect. Dopamine and glutamate afferents coordinate the activity of

nucleus accumbens neurones that, for the major part, consist of γ -aminobutyric acid (GABA) producing medium-sized spiny neurones (Pierce and Kalivas, 1997).

Whereas most neurochemical studies thus far addressed the role of dopamine in addiction behaviour, recent evidence indicates that long-lasting expression of major features of addiction behaviour, such as sensitization and propensity to relapse, may be due to changes in pre- and postsynaptic glutamate neurotransmission particularly within the core region of the nucleus accumbens. This long-term neuroplasticity may strengthen the excitatory control of glutamate afferents onto nucleus accumbens neurones (Tzschentke and Schmidt, 2003; Kalivas, 2004). Therefore, we examined the occurrence of time-dependent alterations in the functioning of ionotropic glutamate receptors in nucleus accumbens core slices following a single exposure of rats to morphine, i.e., representing the context-independent effect of the opiate.

* Corresponding author. Tel.: +31 20 4448104; fax: +31 20 4448100.

E-mail address: anm.schoffelmeer@vumc.nl (A.N.M. Schoffelmeer).

¹ Present address: Erasmus Medical Center, Department of Cell Biology and Genetics, P.O. Box 1738, 3000 DR Rotterdam, The Netherlands.

2. Materials and methods

2.1. Drug administration

In their home cage, male Wistar rats (180–200 g) received a single s.c. injection with saline (1 ml/kg) or morphine-HCl (30 mg/kg, dissolved in saline), inducing long-lasting behavioural and neurochemical sensitization (Vanderschuren et al., 2001). Three days or 3 weeks after pretreatment, the animals were decapitated and the nucleus accumbens core was dissected from a coronal slice (bregma +1.2 to +2.2 mm), used for neurochemical experiments or stored at -80°C for real-time quantitative polymerase chain reaction (PCR) analysis.

2.2. Neurotransmitter release

Neurotransmitter release was measured as previously described (Schoffelemeier et al., 2000). In short, nucleus accumbens core slices ($0.3 \times 0.3 \times 1$ mm) were washed twice with 5 ml Krebs–Ringer's bicarbonate medium (containing 121 mM NaCl, 1.87 mM KCl, 1.17 mM KH_2PO_4 , 25 mM NaHCO_3 , 0.1 mM CaCl_2 , 10 mM D-(+)-glucose and 10 μM aminooxyacetic acid, pH 7.4), incubated for 15 min in this medium containing 0.1 μM [^3H]-GABA under an atmosphere of 95% O_2 /5% CO_2 at 37°C . After labelling, the slices were washed and transferred to each of 24 chambers of a superfusion apparatus (approximately 3 mg tissue in 0.2 ml volume per chamber) and superfused (0.25 ml/min) with medium gassed with 95% O_2 /5% CO_2 at 37°C . The superfusate was collected as 10-min samples after 40 min of superfusion ($t=40$ min). [^3H]-GABA release was induced during superfusion by exposing the slices to medium containing a maximally effective concentration of glutamate (100 μM), *N*-methyl-D-aspartate (NMDA, 100 μM), α -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid (AMPA, 30 μM) or to KCl (10 mM) (NaCl was reduced accordingly to maintain isomolarity) for 10 min at $t=50$ min. Radioactivity remaining at the end of the experiment was extracted from the tissue with 0.1 N HCl. Radioactivity in superfusion fractions and tissue extracts was determined by liquid scintillation counting. The efflux of radioactivity during each collection period was expressed as a percentage of the amount of radioactivity in the slices at the beginning of the respective collection period. Evoked neurotransmitter release was calculated by subtracting the spontaneous efflux of radioactivity from the total overflow of radioactivity during stimulation and the following 10 min. Because the neurotransmitter release was returned to basal levels during the next 10-min period, a linear decline from the 10-min interval before that to 20–30 min after the start of stimulation was assumed for calculation of the spontaneous efflux of radioactivity. The evoked release was expressed as percentage of the

content of radioactivity of the slices at the start of the stimulation period. Statistical significance of differences in evoked neurotransmitter release between experimental groups was evaluated using Student–Newman–Keuls tests.

2.3. Real-time quantitative PCR

Total RNA of each treatment group was pooled, divided into six equal quantities, treated with DNase I (20 U/ μg RNA), random primed with 50 pmol hexanucleotides and reverse transcribed using 200 U MMLV H^- reverse transcriptase. Quantitative PCR was performed in triplicate on each plate and repeated three to six times in separate experiments on an ABI PRISM 7700 Sequence detector as described previously (Jacobs et al., 2002). PCR conditions and SYBR green reagents were used in a reaction of 20 μl using transcript-specific primers (0.3 μM) and 0.7 μl cDNA according to guidelines of the manufacturer with the exception of 45 cycles and elongation at 59°C . Transcript-specific primers were generated with Primer Express software based on GenBank sequence information, verified by NCBI BLAST search and custom synthesized. Primers corresponded to nucleotides (nt) 4091–4114 and 4173–4195 of NMDA NR1 (GenBank accession number X63255.1), nt 6050–6066 and 6117–6135 of NR2A (AF001423), nt 3531–3553 and 3586–3606 of NR2B (M91562.1), nt 3667–3687 and 3723–3745 of NR2C (M91563.1), nt 3211–3231 and 3259–3281 of NR2D (L31612.1), nt 3809–3828 and 3857–3879 of NR3 (AF073379.1), nt 2866–2886 and 2919–2941 of GluR1 (X17184.1), nt 2995–3015 and 3057–3077 of GluR2 (M85035.1), nt 2877–2897 and 2948–2969 of GluR3 (M85036.2), nt 2831–2850 and 2918–2937 of GluR4 (M85037.1), nt 2933–2955 and 2977–2993 of GluR5 (M83560.1), nt 2938–2958 and 3014–3033 of GluR6 (Z11548.1), nt 2710–2730 and 2799–2817 of GluR7 (M83552.1), nt 3001–3024 and 3081–3101 of kainate (KA) KA1 (U08257.1), nt 3392–3408 and 3450–3470 of KA2 (Z11581.1) and nt 267–286 and 322–343 of hypoxanthine phosphoribosyltransferase (HPRT, M63983.1) rat mRNA sequences. Primer sets were tested on nucleus accumbens cDNA by PCR and gel electrophoresis in addition to melting curve analysis using Dissociation Curves 1.0 software for the absence of primer–dimer artefacts and multiple products. Amplification efficiency was determined by quantitative PCR using repetitive dilution series of nucleus accumbens cDNA; an efficiency of ~ 2 was accepted. The cycle of threshold (C_t) was determined as the value 3 cycles above noise. For each primer set, a no template control was performed. C_t values with a difference smaller than 6 cycles were discarded. C_t values were used to calculate the relative level of expression normalized to the transcript for HPRT (Spijker et al., 2004). Student–Newman–Keuls analysis was performed on relative




Fig. 1. Effect of a single morphine injection on glutamate-, NMDA-, AMPA- and potassium-induced $[^3\text{H}]$ -GABA release from rat nucleus accumbens core slices 3 or 21 days after opiate exposure. Spontaneous tritium efflux amounted to about 3.5% of total tissue content of both saline and morphine pretreated rats. Glutamate (100 μM)-, NMDA (100 μM)-, AMPA (30 μM)- and K^+ (10 mM)-induced $[^3\text{H}]$ -GABA release in excess of spontaneous efflux amounted to about 5%, 3.5%, 2% and 2% of total tissue tritium 3 as well as 21 days after saline pretreatment (control release). Evoked neurotransmitter release is expressed as percentage of control (dashed line) and represents mean \pm S.E.M. of three to four experiments. Per experiment four to six observations were made. Significantly different from control: $*P < 0.01$.

values of each treatment group in the nucleus accumbens core for each time point.

3. Results

Rats received a single injection of morphine (30 mg/kg). Superfusion experiments of rat nucleus accumbens core slices showed that after 3 days there was no significant difference in $[^3\text{H}]$ -GABA release induced by maximally effective concentrations of glutamate (100 μM), AMPA (30 μM), NMDA (100 μM) or K^+ (10 mM) between morphine- and saline-treated rats (Fig. 1). However, after 3 weeks glutamate-, AMPA- and NMDA-induced $[^3\text{H}]$ -GABA release in the nucleus accumbens core was increased by, respectively, 42%, 27% and 47% in rats treated with morphine compared to those treated with saline. The K^+ -

induced release of $[^3\text{H}]$ -GABA was unaltered by prior morphine administration.

Three days after a single injection, mRNA levels of 8 out of 15 measured NMDA, AMPA and kainate receptor subunits were significantly reduced in the nucleus accumbens core of morphine- compared to saline-treated rats (Fig. 2). In contrast, mRNA levels of all 15 receptor subunits were significantly enhanced in the nucleus accumbens core 3 weeks after morphine exposure.

4. Discussion

The long-term expression of certain characteristic features of addiction in rats following exposure to cocaine, such as behavioural sensitization and cue- or cocaine-induced relapse to drug-seeking behaviour, may be mediated by an increase in the release of glutamate in the nucleus accumbens core, involving activation of glutamate efferents from the amygdala and prefrontal cortex (Tzschentke and Schmidt, 2003; Kalivas, 2004). A number of cocaine-induced alterations in postsynaptic glutamate transmission in the nucleus accumbens core has also been reported considering the gene expression of glutamate receptor subunits. However, the direction of the changes in glutamate receptors following psychostimulant administration reported thus far is rather inconsistent (Lu et al., 2002; Churchill et al., 1999; Ghasemzadeh et al., 1999).

Here, we examined whether time-dependent alterations at the level of postsynaptic glutamate receptors is apparent in slices of the nucleus accumbens core following exposure of rats to morphine. To that end, rats received a single injection of morphine (30 mg/kg), previously shown to cause long-lasting sensitization (Vanderschuren




Fig. 2. Relative gene expression levels of transcripts encoding ionotropic glutamate receptor subunits in the nucleus accumbens core after a single exposure of rats to morphine. Shown are expression levels relative to saline controls as determined by quantitative PCR, 3 or 21 days after morphine exposure. Values were normalized to the HPRT transcript. Student–Newman–Keuls analysis was performed on relative values of each drug treatment group (4–6 rats per group). Expression levels are presented as fold change. All values are means \pm S.E.M. Significantly different from control: $*P < 0.05$.

et al., 2001). Our *ex vivo* experiments revealed that the stimulatory effect of (maximally effective concentrations of) glutamate (100 μ M), AMPA (30 μ M) and NMDA (100 μ M) on [3 H]-GABA release was profoundly enhanced in superfused nucleus accumbens core slices from morphine pretreated rats as compared to that in slices from saline pretreated animals, 3 weeks (but not 3 days) after morphine exposure. The observation that this long-term increase in AMPA/kainate receptor efficacy was almost twice that of NMDA receptor efficacy suggests that morphine may cause adaptive changes at the level of ionotropic glutamate receptors rather than at the level of the neurotransmitter release mechanism. Accordingly, the K^+ -induced release of [3 H]-GABA remained unaffected by prior morphine administration. In this respect, we also examined the gene expression of the distinct subunits of these ionotropic receptors in the nucleus accumbens core. Interestingly, the expression of gene transcripts encoding these receptor subunits was strongly increased long after a single morphine exposure. In contrast, transcript levels were generally decreased after 3 days of opiate absence. Since the maximal effect of glutamate receptor activation was unaltered at this time point, it is conceivable that enough unaltered glutamate receptors remained for normal functioning (receptor reserve). Moreover, the observation that the degree of enhanced gene expression varied between the distinct transcripts studied suggests the occurrence of adaptive alterations in the subunit composition of postsynaptic glutamate receptors. Clearly, however, further work is needed to address these issues at the level of glutamate receptors.

An intriguing aspect of the drug-induced neuroplasticity observed here is its delayed occurrence subsequent to drug exposure. Such time dependence has also been observed pre- and postsynaptically regarding the effects of morphine on striatal dopamine release and dopamine D1 receptor functioning (Tjon et al., 1994). As dopamine D1 receptor activation has been shown to increase the surface expression of glutamate receptors in nucleus accumbens neurones (Chao et al., 2002), the delayed morphine-induced increase in glutamate receptor functioning observed here may well be related to enhanced dopamine neurotransmission. These delayed pharmacological effects of morphine may play a crucial role in the persistence of opiate addiction.

References

- Chao, S.Z., Ariano, M.A., Peterson, D.A., Wolf, M.E., 2002. D1 dopamine receptor stimulation increases GluR1 surface expression in nucleus accumbens neurons. *J. Neurochem.* 83, 704–712.
- Churchill, L., Swanson, C.J., Urbina, M., Kalivas, P.W., 1999. Repeated cocaine alters glutamate receptor subunit levels in the nucleus accumbens and ventral tegmental area of rats that develop behavioral sensitization. *J. Neurochem.* 72, 2397–2403.
- Ghasemzadeh, M.B., Nelson, L.C., Lu, X.Y., Kalivas, P.W., 1999. Neuroadaptations in ionotropic and metabotropic glutamate receptor mRNA produced by cocaine treatment. *J. Neurochem.* 72, 157–165.
- Jacobs, E.H., Spijker, S., Verhoog, C.W., Kamprath, K., de Vries, T.J., Smit, A.B., Schoffelmeer, A.N.M., 2002. Active heroin administration induces specific genomic responses in the nucleus accumbens shell. *FASEB J.* 16, 1961–1963.
- Kalivas, P.W., 2004. Glutamate systems in cocaine addiction. *Curr. Opin. Pharmacol.* 4, 23–29.
- Lu, W., Monteggia, L.M., Wolf, M.E., 2002. Repeated administration of amphetamine or cocaine does not alter AMPA receptor subunit expression in the rat midbrain. *Neuropsychopharmacology* 26, 1–13.
- Nestler, E.J., 2001. Molecular basis of long-term plasticity underlying addiction. *Nat. Rev. Neurosci.* 2, 119–128.
- Pierce, R.C., Kalivas, P.W., 1997. A circuitry model of the expression of behavioral sensitization to amphetamine-like psychostimulants. *Brain Res. Rev.* 25, 192–216.
- Robinson, T.E., Berridge, K.C., 2003. Addiction. *Annu. Rev. Psychol.* 54, 25–53.
- Schoffelmeer, A.N.M., Vanderschuren, L.J.M.J., De Vries, T.J., Hogenboom, F., Wardeh, G., Mulder, A.H., 2000. Synergistically interacting dopamine D1 and NMDA receptors mediate nonvesicular transporter-dependent GABA release from rat striatal medium spiny neurons. *J. Neurosci.* 20, 3496–3503.
- Spijker, S., Houtzager, S.W., De Gunst, M.C., De Boer, W.P., Schoffelmeer, A.N.M., Smit, A.B., 2004. Morphine exposure and abstinence define specific stages of gene expression in rat nucleus accumbens. *FASEB J.* 18, 848–850.
- Tjon, G.H.K., De Vries, T.J., Ronken, E., Hogenboom, F., Wardeh, G., Mulder, A.H., Schoffelmeer, A.N.M., 1994. Intermittent and chronic morphine administration causes differential long-lasting changes in dopaminergic neurotransmission in rat striatum without changing its delta- and kappa-opioid receptor regulation. *Eur. J. Pharmacol.* 252, 205–212.
- Tzschentke, T.M., Schmidt, W.J., 2003. Glutamatergic mechanisms in addiction. *Mol. Psychiatry* 8, 373–382.
- Vanderschuren, L.J.M.J., Kalivas, P.W., 2000. Alterations in dopaminergic and glutamatergic transmission in the induction and expression of behavioral sensitization: a critical review of preclinical studies. *Psychopharmacology* 151, 99–120.
- Vanderschuren, L.J.M.J., De Vries, T.J., Wardeh, G., Hogenboom, F.A.C.M., Schoffelmeer, A.N.M., 2001. A single exposure to morphine induces long-lasting behavioural and neurochemical sensitization in rats. *Eur. J. Neurosci.* 14, 1533–1538.