

A cDNA coding for human sex hormone binding globulin

Homology to vitamin K-dependent protein S

Sten Gershagen, Per Fernlund and Åke Lundwall

Department of Clinical Chemistry, University of Lund, Malmö General Hospital, S-214 01 Malmö, Sweden

Received 16 March 1987

Affinity purified antibodies to human sex hormone binding globulin (SHBG) were used in screening a human liver cDNA library, constructed in the expression vector λ gt11. One clone, identified as producing recombinant SHBG, carried a cDNA insert of 1.1 kb. The nucleotide sequence of the insert had an open reading frame coding for 356 amino acid residues. The coding sequence was followed by a short 3'-region of 19 non-translated nucleotides and a poly(A) tail. Confirmation that the cDNA clone represented human SHBG was obtained by the finding of a complete agreement in amino acid sequence with several peptide fragments generated from purified SHBG by proteolytic cleavage. The primary structure of SHBG shows a considerable homology to that of protein S, a vitamin K-dependent protein with functions in the coagulation system.

Sex hormone binding globulin; Sex hormone; Carrier protein; Plasma protein; Blood coagulation

1. INTRODUCTION

Sex hormone binding globulin (SHBG) is an important carrier of sex steroids in plasma [1]. It is thought to be synthesized in the liver [2] and the concentration of SHBG in plasma is dependent on several hormones, being increased by estrogens and thyroid hormones and decreased by androgens. Human SHBG is a dimeric protein with a molecular mass of approx. 100 kDa. It is composed of two identical or very similar, non-covalently associated protomers. Only one steroid molecule is bound per dimeric protein, but little else is known about the mechanisms for the steroid binding [3-5].

The protomers of SHBG vary in size, two main size classes having usually been observed, one heavier and one lighter occurring at an approximate ratio of 10:1 [6,7]. However, it has recently

been demonstrated that this only represents the prevailing pattern, occurring in 80% of the population. In the remaining 20% there are three different sizes of the SHBG protomer [8].

Previous reports on the carbohydrate structure [4,9,10] and partial amino acid sequences of human SHBG [3,7] have recently been complemented with the complete amino acid sequence of the human SHBG protomer [4]. None of these reports, however, explains the observed variation in size of the SHBG protomer.

To gain further knowledge of the structure of SHBG and its possible variation, and to permit studies of its gene structure and of its biosynthesis in various organs, we have isolated a cDNA clone coding for SHBG from a human liver cDNA library.

2. MATERIALS AND METHODS

2.1. Reagents

Chemicals used for amino acid analysis were obtained from Beckman Instruments, and for protein

Correspondence address: S. Gershagen, Dept Clinical Chemistry, Malmö General Hospital, S-21401 Malmö, Sweden

sequence determination from Applied Biosystems. Solvents for HPLC were from Rathburn. TPCK-trypsin was obtained from Sigma and *Staphylococcus aureus* protease V8 from Miles. A λ gt11 screening kit was purchased from Promega Biotechniques. The Klenow fragment of DNA polymerase 1 was obtained from Boehringer Mannheim. The restriction enzymes *EcoRI* and *StuI* were from Amersham, who also supplied α -[35 S]thio-dATP (22 TBq/mmol). Deoxynucleotides, dideoxynucleotides, T₄ DNA polymerase and the replicative form of the phage M13 mp18 were purchased from Pharmacia. An antiserum to human SHBG has been described [11]. It was affinity purified on a column of agarose coupled with SHBG, purified as described [12].

2.2. Isolation of cDNA clones and DNA sequencing

A λ gt11 cDNA library prepared from human liver poly(A)⁺ RNA was kindly provided by Dr Savio L. Woo. The library was screened according to the procedures described by Young and Davis [13,14], using affinity purified rabbit anti-human SHBG as first antibody and an alkaline phosphatase labelled goat anti-rabbit immunoglobulin for detection. Recombinant phages were purified from plate lysates by centrifugation in a CsCl gradient, and the DNA was extracted by standard procedures [15]. Sequencing of DNA was carried out on material subcloned in M13 mp18 according to Sanger et al. [16] as modified by Biggin et al. [17]. Nucleotide sequences were aligned and analyzed with computer programs kindly provided by R. Staden, Medical Research Council Unit, Cambridge [18,19] and the Protein Identification Resource [20].

2.3. Isolation and sequencing of peptide fragments

Human SHBG (1.4 mg), purified as described [12], was dialyzed at 4°C against 6 M guanidine HCl and 5 mM EDTA in 0.5 M Tris-HCl buffer, pH 8.0. The dialyzed sample was flushed with N₂ and reduced by adding 3.1 μ mol of dithiothreitol followed by incubation at 37°C for 4 h. Alkylation was done with 7.8 μ mol of solid, recrystallized, iodoacetic acid and the mixture was left for 2 h in the dark at room temperature with gentle agita-

tion. The sample was dialyzed against 0.1 M ammonium bicarbonate buffer, pH 7.5, followed by evaporation. The residue was dissolved in 200 μ l of the ammonium bicarbonate buffer and treated with 28 μ g of trypsin or *S. aureus* protease V8 for 2 h at 37°C. The sample was evaporated to dryness, which was repeated twice after the addition of redistilled water to remove salt thoroughly.

High-performance liquid chromatography (HPLC) was performed at room temperature on a μ Bondapak column (0.39 \times 30 cm) eluted at a flow rate of 0.8 ml/min with a linear gradient from solvent A:solvent B (95:5, v/v) to solvent A:solvent B (30:70, v/v) developed over 1 h. Solvent A was 0.08% trifluoroacetic acid in water and solvent B acetonitrile. Fractions corresponding to peptide peaks were pooled and lyophilized. One half of the material from each isolated peptide was subjected to amino acid analysis and the other to sequence determination. Amino acid sequence and composition of peptides were determined on an Applied Biosystems gas-phase sequencer and a Beckman 6300 amino acid analyzer, respectively, as described [21].




Fig.1. Restriction map and sequencing strategy (a) and nucleotide sequence with predicted amino acid sequence (b) of human SHBG cDNA. (a) Restriction endonuclease cleavage sites are shown above the linear map; arrows, DNA fragments sequenced. (b) Nucleotide sequence of the cDNA presented in the 5' to 3' orientation. The longest translational open reading frame is shown above the nucleotide sequence. Numbers to the right of the sequences indicate amino acid residue (upper) and nucleotide (lower). Number 18 is assigned to the first amino acid residue for reasons given in the text. Amino acid sequences also determined in peptides (generated from purified SHBG by proteolytic cleavage) are underlined.

(b)

(b)	Val	His	Ser	Ala	Ala	Gln	Thr	Thr	Leu	Ile	Ala	Val	Met	Thr	Phe	Asp	<u>Leu</u>	<u>Thr</u>	<u>Lys</u>	36
CT	GTC	CAC	TCG	GCG	GCT	CAG	ACA	ACT	CTT	ATC	GCT	GTC	ATG	ACC	TTT	GAC	CTC	ACC	AAG	59
<u>Ile</u>	<u>Thr</u>	<u>Lys</u>	<u>Thr</u>	<u>Ser</u>	Ser	Ser	Phe	Glu	<u>Val</u>	<u>Arg</u>	<u>Thr</u>	<u>Trp</u>	<u>Asp</u>	<u>Pro</u>	<u>Glu</u>	<u>Gly</u>	<u>Val</u>	<u>Ile</u>	<u>Phe</u>	56
ATC	ACA	AAA	ACC	TCC	TCC	TCC	TTT	GAG	GTT	CGA	ACC	TGG	GAC	CCA	GAG	GGA	GTG	ATT	TTT	119
<u>Tyr</u>	<u>Gly</u>	<u>Asp</u>	<u>Thr</u>	<u>Asn</u>	<u>Pro</u>	Lys	<u>Asp</u>	<u>Asp</u>	<u>Trp</u>	<u>Phe</u>	<u>Met</u>	Leu	Gly	Leu	Arg	<u>Asp</u>	<u>Gly</u>	<u>Arg</u>	<u>Pro</u>	76
TAT	GGG	GAT	ACC	AAC	CCT	AAG	GAT	GAC	TGG	TTT	ATG	CTG	GGA	CTT	CGA	GAC	GGC	AGG	CCT	179
<u>Glu</u>	<u>Ile</u>	<u>Gln</u>	<u>Leu</u>	<u>His</u>	<u>Asn</u>	His	Trp	Ala	Gln	Leu	Thr	Val	Gly	Ala	Gly	Pro	Arg	<u>Leu</u>	<u>Asp</u>	96
GAG	ATC	CAA	CTG	CAC	AAT	CAC	TGG	GCC	CAG	CTT	ACG	GTG	GGT	GCT	GGA	CCA	CGG	CTG	GAT	239
<u>Asp</u>	<u>Gly</u>	<u>Arg</u>	<u>Trp</u>	<u>His</u>	<u>Gln</u>	<u>Val</u>	<u>Glu</u>	<u>Val</u>	Lys	<u>Met</u>	<u>Glu</u>	<u>Gly</u>	<u>Asp</u>	<u>Ser</u>	<u>Val</u>	<u>Leu</u>	<u>Leu</u>	<u>Glu</u>	<u>Val</u>	116
GAT	GGG	AGA	TGG	CAC	CAG	GTG	GAA	GTC	AAG	ATG	GAG	GGG	GAC	TCT	GTG	CTG	CTG	GAG	GTG	299
<u>Asp</u>	<u>Gly</u>	<u>Glu</u>	<u>Glu</u>	<u>Val</u>	Leu	Arg	Leu	Arg	Gln	Val	Ser	Gly	Pro	Leu	Thr	Ser	Lys	Arg	His	136
GAT	GGG	GAG	GAG	GTG	CTG	CGC	CTG	AGA	CAG	GTC	TCT	GGG	CCC	CTG	ACC	AGC	AAA	CGC	CAT	359
Pro	Ile	Met	Arg	<u>Ile</u>	<u>Ala</u>	<u>Leu</u>	<u>Gly</u>	<u>Gly</u>	<u>Leu</u>	<u>Leu</u>	<u>Phe</u>	<u>Pro</u>	<u>Ala</u>	<u>Ser</u>	<u>Asn</u>	<u>Leu</u>	<u>Arg</u>	<u>Leu</u>	<u>Pro</u>	156
CCC	ATC	ATG	AGG	ATT	GCG	CTT	GGG	GGG	CTG	CTC	TTC	CCC	GCT	TCC	AAC	CTT	CGG	TTG	CCG	419
<u>Leu</u>	<u>Val</u>	<u>Pro</u>	<u>Ala</u>	<u>Leu</u>	<u>Asp</u>	<u>Gly</u>	<u>Cys</u>	<u>Leu</u>	Arg	Arg	Asp	Ser	Trp	Leu	Asp	Lys	Gln	Ala	Glu	176
CTG	GTT	CCT	GCC	CTG	GAT	GGC	TGC	CTG	CGC	CGG	GAT	TCC	TGG	CTG	GAC	AAA	CAG	GCC	GAG	479
Ile	Ser	Ala	Ser	Ala	Pro	Thr	Ser	Leu	Arg	<u>Ser</u>	<u>Cys</u>	<u>Asp</u>	<u>Val</u>	<u>Glu</u>	<u>Ser</u>	<u>Asn</u>	<u>Pro</u>	<u>Gly</u>	<u>Ile</u>	196
ATC	TCA	GCA	TCT	GCC	CCC	ACT	AGC	CTC	AGA	AGC	TGT	GAT	GTA	GAA	TCA	AAT	CCC	GGG	ATA	539
<u>Phe</u>	<u>Leu</u>	<u>Pro</u>	<u>Pro</u>	<u>Gly</u>	<u>Thr</u>	<u>Gln</u>	<u>Ala</u>	<u>Glu</u>	<u>Phe</u>	<u>Asn</u>	<u>Leu</u>	<u>Arg</u>	<u>Asp</u>	<u>Ile</u>	<u>Pro</u>	<u>Gln</u>	<u>Pro</u>	<u>His</u>	<u>Ala</u>	216
TTT	CTC	CCT	CCA	GGG	ACT	CAG	GCA	GAA	TTC	AAT	CTC	CGA	GAC	ATT	CCC	CAG	CCT	CAT	GCA	599
<u>Glu</u>	<u>Pro</u>	<u>Trp</u>	<u>Ala</u>	<u>Phe</u>	<u>Ser</u>	<u>Leu</u>	<u>Asp</u>	<u>Leu</u>	Gly	Leu	Lys	Gln	Ala	Ala	Gly	Ser	Gly	His	Leu	236
GAG	CCC	TGG	GCC	TTC	TCT	TTG	GAC	CTG	GGA	CTC	AAG	CAG	GCA	GCA	GGC	TCA	GGC	CAC	CTC	659
Leu	Ala	Leu	Gly	Thr	Pro	Glu	Asn	Pro	Ser	Trp	Leu	Ser	Leu	His	Leu	Gln	Asp	Gln	Lys	256
CTT	GCT	CTT	GGG	ACA	CCA	GAG	AAC	CCA	TCT	TGG	CTC	AGT	CTC	CAC	CTC	CAA	GAT	CAA	AAG	719
Val	Val	Leu	Ser	Ser	Gly	Ser	Gly	Pro	Gly	Leu	Asp	Leu	Pro	Leu	Val	Leu	Gly	Leu	Pro	276
GTG	GTG	TTG	TCT	TCT	GGG	TCG	GGG	CCA	GGG	CTG	GAT	CTG	CCC	CTG	GTC	TTG	GGA	CTC	CCT	779
Leu	Gln	Leu	Lys	Leu	Ser	Met	Ser	Arg	Val	Val	Leu	Ser	Gln	Gly	Ser	Lys	Met	Lys	<u>Ala</u>	296
CTT	CAG	CTG	AAG	CTG	AGT	ATG	TCC	AGG	GTG	GTC	TTG	AGC	CAA	GGG	TCG	AAG	ATG	AAG	GCC	839
<u>Leu</u>	<u>Ala</u>	<u>Leu</u>	<u>Pro</u>	<u>Pro</u>	<u>Leu</u>	<u>Gly</u>	<u>Leu</u>	<u>Ala</u>	<u>Pro</u>	<u>Leu</u>	<u>Leu</u>	<u>Asn</u>	<u>Leu</u>	<u>Trp</u>	<u>Ala</u>	<u>Lys</u>	<u>Pro</u>	<u>Gln</u>	<u>Gly</u>	316
CTT	GCC	CTG	CCT	CCC	TTA	GGC	CTG	GCT	CCC	CTC	CTT	AAC	CTC	TGG	GCC	AAG	CCT	CAA	GGG	899
Arg	Leu	Phe	Leu	Gly	Ala	Leu	Pro	Gly	Glu	Asp	Ser	Ser	Thr	Ser	Phe	Cys	Leu	Asn	Gly	336
CGT	CTC	TTC	CTG	GGG	GCT	TTA	CCA	GGA	GAA	GAC	TCT	TCC	ACC	TCT	TTT	TGC	CTG	AAT	GGC	959
Leu	Trp	Ala	Gln	Gly	Gln	Arg	Leu	Asp	Val	Asp	Gln	Ala	Leu	Asn	Arg	<u>Ser</u>	<u>His</u>	<u>Glu</u>	<u>Ile</u>	356
CTT	TGG	GCA	CAA	GGT	CAG	AGG	CTG	GAT	GTG	GAC	CAG	GCC	CTG	AAC	AGA	AGC	CAT	GAG	ATC	1019
<u>Trp</u>	<u>Thr</u>	<u>His</u>	<u>Ser</u>	<u>Cys</u>	<u>Pro</u>	<u>Gln</u>	<u>Ser</u>	<u>Pro</u>	<u>Gly</u>	<u>Asn</u>	<u>Gly</u>	<u>Thr</u>	<u>Asp</u>	<u>Ala</u>	Ser	His				
TGG	ACT	CAC	AGC	TGC	CCC	CAG	AGC	CCA	GGC	AAT	GGC	ACT	GAC	GCT	TCC	CAT	TAA	AGCTCCA		1080

CCTAAGAACCCCAAAAAAAAAAAAAAAAAA

3. RESULTS

3.1. Isolation and sequencing of a human SHBG cDNA clone

Approximately 1.2×10^6 recombinants, in a λ gt11 human liver cDNA library were screened. One clone, designated λ SHBG-1, was isolated, the DNA of which was digested with *Eco*RI and then analyzed by agarose gel electrophoresis. In addition to the vector bands, two bands were seen, one representing a fragment of approximately 560 nucleotides, the other of 450. The *Eco*RI fragments were isolated and subcloned in bacteriophage M13 mp18. The ends of the subcloned DNA were sequenced, yielding approximately 90% of the sequence of the whole cDNA insert. From analysis of the partial sequence, two suitably located *Stu*I cleavage sites were found. By combined *Eco*RI and *Stu*I digestion, fragments were generated that yielded enough data to complete the sequence of the λ SHBG-1 cDNA insert. Fig. 1a gives an outline of the sequencing strategy. The nucleotide and derived amino acid sequences for the cloned SHBG cDNA are given in fig. 1b. The length is 1110 base pairs, including a short poly(A) tail of 18 nucleotides. The coding region, encoding 356 amino acids, extends from the beginning of the insert to a stop codon, TAA, at position 1071–1073. The stop codon is also part of the polyadenylation signal, ATTAAA. A stretch of 19 non-translated nucleotides separates the coding region from the poly(A) tail.

3.2. Comparison of the deduced amino acid sequence with peptide fragments of SHBG

A trypsin digest of reduced and alkylated human SHBG was separated by HPLC. Nine pure peptides and two fractions, each a mixture of two peptides, were obtained. Similarly, six pure peptides were isolated from a digest of reduced and alkylated SHBG obtained with the *S. aureus* protease V8. For both digests, the yield of each peptide was about 10%, which probably should not be regarded as low when considering the small scale on which the isolations had to be carried out.

The isolated peptides were sequenced and all of them could be identified as parts of the amino acid sequence deduced from the nucleotide sequence of the cDNA. The parts of the deduced sequence corresponding to the peptides are underlined in fig. 1b.

The peptide sequence data accounted for altogether 49% of the entire deduced amino acid sequence (fig. 1b) and there were no peptide sequences that could not be found in the deduced sequence.

The SHBG cDNA contains one internal *Eco*RI restriction site (fig. 1b), and this site is not overlapped by any of the DNA sequences (fig. 1a). It is, however, overlapped by one long tryptic peptide (fig. 1b) thus excluding the presence of more than one *Eco*RI cleavage site.

Comparison of the deduced amino acid sequence with the recently published protein sequence of the SHBG protomer [4] shows a complete agreement from residue 27 in the published sequence to its end. The amino acid residues in fig. 1b have therefore been numbered to coincide with the published sequence, the only discrepancy being in the amino-terminal part; the first nine amino acids, which should correspond to amino acid residues 18–26 in the published sequence, are completely different.

3.3. Homology to vitamin K-dependent protein S

An alignment of human SHBG with bovine and human protein S [22,23] is shown in fig. 2. The alignment shown was obtained by first aligning human SHBG separately with human and bovine protein S by means of a computer program [20]. All three sequences were then aligned with each other by making a few arbitrary adjustments which increased residue correspondence. The computer alignments with human and bovine protein S gave the maximal scores 252 and 264, respectively, when 40 was used as a penalty for a break. The mean scores obtained with 100 random sequences, generated from an amino acid composition identical to that of human SHBG, were 11.4 and 11.7 for alignments with human and bovine protein S, respectively, with standard deviations 6.2 and 4.7.

4. DISCUSSION

Except for the nine amino acid residues coded for by the 5'-end, the amino acid sequence deduced from the cloned cDNA is in complete agreement with the recently published sequence of human SHBG determined by protein chemical methods [4]. Although the first 29 nucleotides of the cDNA (coding for nine amino acids) might




Fig.2. Amino acid sequence comparison between human SHBG (lower) and the carboxy-terminal portions of human (middle) and bovine (upper) protein S. Amino acid residues matched in SHBG and in at least one of the protein S species are boxed. Spaces represent gaps inserted to maximize the number of matches in the comparison.

represent a cloning artefact, another explanation that cannot be ruled out is that the variant sequence represents an SHBG subunit partly different from that sequenced by Walsh et al. [4].

It is a well established fact that SHBG, which is a dimeric protein, contains subunits differing slightly in structure [3,6,7] and it has recently been shown that the SHBG subunits are not uniform in

all humans, 20% of the population having an extra SHBG subunit which is heavier than the usual two subunits [8]. The structure differences between the SHBG subunits may be due to different degrees of glycosylations or other posttranslational modifications, but a difference in amino acid sequence is also a possibility. Such a variation in primary structure might result from an alternative splicing, for instance.

If SHBG were purified from pooled plasma containing a minor amount of a subunit with a sequence slightly different from that of the predominant form of the protein, sequence analysis would probably not detect the variant sequence. In contrast, cloning of a cDNA would give one unique species, one that could represent a variant polypeptide. The resolution of the question of whether variation in the primary structure of the SHBG subunit exists must await the isolation of additional SHBG cDNA clones, studies of the SHBG gene and/or the sequencing of individual subunits of SHBG.

When the amino acid sequence of human SHBG was reported SHBG was found not to be homologous to any other protein, thus implying it to be the single member of a new protein family [4]. Shortly afterwards, however, androgen binding protein (ABP) from rat testis was shown to have a high degree of homology to human SHBG [24]. Our comparison of the sequence of human SHBG with the sequences of bovine and human protein S, two proteins recently cloned and sequenced [22,23,25], reveals a homology of human SHBG to protein S, not unexpected in the light of the observation [24] that rat ABP seems to be related to the carboxy-terminal region of protein S.

Protein S is an important cofactor in the anticoagulation system [26] and has connections with the complement system [27]. Structurally, protein S has been subdivided into four different regions, the Gla region, the thrombin-sensitive region, the growth factor region, and the large carboxy-terminal region [22,23]. The homology between human SHBG and protein S is confined to the carboxy-terminal region of protein S, and SHBG is almost comparable in size to this region. It is noteworthy that the four half-cystine residues in SHBG seem to correspond to four of the five half-cystine residues present in the carboxy-terminal region of protein S. These four residues in protein

S form disulphide loops within the carboxy-terminal region. By analogy, we suggest that the corresponding half-cystines are disulphide bonded in SHBG, i.e. Cys₁₆₄ to Cys₁₈₈ and Cys₃₃₃ to Cys₃₆₁.

The lowest degree of homology between SHBG and protein S is seen in the parts corresponding to residues 240 to 290 in SHBG (fig.2). This is noteworthy since this part of the SHBG molecule has been suggested to be responsible for the steroid binding property of SHBG [4]. It contains unusual structural features not recognizable in protein S, such as a long stretch of alternating leucine residues (residues 267–281) and a sequence repeat (residues 277–292 being a repeat of residues 248–263). It is too early to speculate whether this reflects a more pronounced difference in function between these parts of the molecules since the steroid binding site of SHBG is not yet accurately mapped and the function of the carboxy-terminal region of protein S is largely unknown. Nonetheless, the finding of amino acid sequence homology between human SHBG and protein S from both the human and the cow throws new light on these proteins and may help in the elucidation of their function.

ACKNOWLEDGEMENTS

We thank Dr Savio L. Woo for his gift of a human liver λ gt11 cDNA library and Professor P.H. Petra for letting us read his pre-publication manuscript on the SHBG sequence. The skillful technical assistance of Ingrid Dahlkvist is gratefully acknowledged. This work was supported by grants from the Swedish Medical Research Council (project no.5913). S.G. received grants from the Fertility Research Institute and Sterling Winthrop Scandinavia.

REFERENCES

- [1] Anderson, D.C. (1974) Clin. Endocrinol. 3, 69–79.
- [2] Kahn, M.S., Knowles, B.B., Aden, D.P. and Rosner, W. (1981) J. Clin. Endocrinol. Metab. 53, 448–449.
- [3] Petra, P.H., Stanczyk, F.Z., Seneor, D.F., Namkung, P.C., Novy, M.J., Ross, J.B.A., Turner, E. and Brown, J.A. (1983) J. Steroid Biochem. 19, 699–706.

- [4] Walsh, K.H., Titani, K., Takio, K., Kumar, S., Hayes, R. and Petra, P.H. (1986) *Biochemistry* 25, 7584–7590.
- [5] Örstad, A., Lulka, M.F., Eide, B., Petra, P.H. and Ross, J.B.A. (1986) *Biochemistry* 25, 2686–2692.
- [6] Cheng, C.Y., Musto, N.A., Gunsalus, G.L. and Bardin, C.W. (1983) *J. Steroid Biochem.* 19, 1379–1389.
- [7] Hammond, G.L., Robinson, P.A., Sugino, H., Ward, D.N. and Finne, J. (1986) *J. Steroid Biochem.* 24, 815–824.
- [8] Gershagen, S., Henningsson, K. and Fernlund, P. (1987) *J. Biol. Chem.*, in press.
- [9] Avvakumov, G.V., Matveentseva, J.V., Akhrem, L.V., Shelchyonok, O.A. and Akhrem, A.A. (1983) *Biochim. Biophys. Acta* 760, 104–110.
- [10] Cheng, C.Y., Musto, N.A., Gunsalus, G.L., Frick, J. and Bardin, C.W. (1985) *J. Biol. Chem.* 260, 5631–5640.
- [11] Fernlund, P., Gershagen, S., Kjellberg, M. and Rannevik, G. (1984) *Scand. J. Clin. Lab. Invest.* 44, 25–31.
- [12] Fernlund, P. and Laurell, C.-B. (1981) *J. Steroid Biochem.* 14, 545–552.
- [13] Young, R.A. and Davis, R.W. (1983) *Proc. Natl. Acad. Sci. USA* 80, 1194–1198.
- [14] Young, R.A. and Davis, R.W. (1983) *Science* 222, 778–782.
- [15] Maniatis, T., Fritsch, E.F. and Sambrook, J. (1982) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- [16] Sanger, F., Nicklen, S. and Coulson, A.R. (1977) *Proc. Natl. Acad. Sci. USA* 74, 5463–5467.
- [17] Biggin, M.D., Gibson, T.T. and Hong, G.F. (1983) *Proc. Natl. Acad. Sci. USA* 80, 3963–3965.
- [18] Staden, R. (1982) *Nucleic Acids Res.* 10, 2951–2961.
- [19] Staden, R. (1982) *Nucleic Acids Res.* 10, 4731–4751.
- [20] Orcutt, B.C., Dayhoff, M.O., George, D.G. and Barker, W.C. (1984) *Protein Identification Resource*, National Biomedical Research Foundation, Georgetown University Medical Center, Washington, DC 20007, USA.
- [21] Dahlbäck, B., Lundwall, Å. and Stenflo, J. (1986) *J. Biol. Chem.* 261, 5111–5115.
- [22] Dahlbäck, B., Lundwall, Å. and Stenflo, J. (1986) *Proc. Natl. Acad. Sci. USA* 83, 4199–4203.
- [23] Lundwall, Å., Dackowski, W., Cohen, E., Shaffer, M., Mahr, A., Dahlbäck, B., Stenflo, J. and Wydro, R. (1986) *Proc. Natl. Acad. Sci. USA* 83, 6716–6720.
- [24] Joseph, D.R., Hall, S.H. and French, F.S. (1987) *Proc. Natl. Acad. Sci. USA* 84, 339–343.
- [25] Hoskins, J., Norman, K.D., Beckmann, R.J. and Long, G.L. (1987) *Proc. Natl. Acad. Sci. USA* 84, 349–353.
- [26] Walker, F.J. (1981) *J. Biol. Chem.* 256, 11128–11131.
- [27] Dahlbäck, B. (1984) *Semin. Thromb. Hemostasis* 10, 139–148.