

Cyclic AMP-dependent phosphorylation and regulation of the cardiac dihydropyridine-sensitive Ca channel

Akira Yoshida^a, Masami Takahashi^a, Seiichiro Nishimura^b, Hiroshi Takeshima^b and Shinichiro Kokubun^c

^aMitsubishi Kasei Institute of Life Sciences, Machida, Tokyo 194, Japan, ^bDepartment of Medical Chemistry, Kyoto University, Faculty of Medicine, Kyoto 606, Japan and ^cDepartment of Physiology, Nihon University School of Medicine, Tokyo 173, Japan

Received 5 June 1992

A polyclonal antibody, CR2, prepared using the C-terminal peptide of the $\alpha 1$ subunit of the rabbit cardiac DHP-sensitive Ca channel, specifically immunoprecipitated the [³H]PN200-110-labeled Ca channel solubilized from cardiac microsomes. The antibody recognized 250 and 200-kDa cardiac microsomal proteins as determined by immunoblotting, and cAMP-dependent protein kinase phosphorylated the 250-kDa, but not the 200-kDa protein *in vitro*. CHO cells, transfected with the cardiac $\alpha 1$ subunit cDNA carried by an expression vector, synthesized a 250-kDa protein which was recognized by CR2. Adding db-cAMP or forskolin to the transformed CHO cells induced phosphorylation of the 250-kDa protein and stimulated the DHP-sensitive Ba current under patch-clamp conditions. These results suggested that the cardiac DHP-sensitive Ca channel was regulated by cAMP-dependent phosphorylation of the $\alpha 1$ subunit.

Cardiac Ca channel; Dihydropyridine; cAMP-dependent phosphorylation; Ca channel modulation

1. INTRODUCTION

Stimulation of Ca²⁺ influx through the DHP-sensitive Ca channel into cardiac muscle is essential for the isotropic effect caused by β -adrenergic drugs [1]. The activation of the β -adrenergic receptor induces the elevation of intracellular cAMP followed by the activation of PKA [2]. There is much evidence indicating that the PKA-mediated phosphorylation participates in the stimulation of the Ca current [3,4], however, phosphoproteins involved in this regulation remain to be identified.

The DHP-sensitive Ca channel has been purified from skeletal muscle and is composed of $\alpha 1$, $\alpha 2$, β , γ and δ subunits [5]. All subunits have been investigated by cDNA cloning and their primary amino-acid sequences have been determined [6–9]. The $\alpha 1$ and β subunits of the purified skeletal muscle Ca channel are phosphorylated by PKA accompanied by stimulation of the Ca channel function [5,10–12]. The $\alpha 1$ subunit was further shown to be phosphorylated in Ser-687 by PKA [13]. The cardiac DHP-sensitive Ca channel has been partially purified and contains $\alpha 1$ and $\alpha 2$ subunits, however the precise subunit composition has not been estab-

lished [14–16]. Several reports have shown that the $\alpha 1$ subunit in partially purified cardiac Ca channels is not phosphorylated by PKA [14–16] and its corresponding Ser-687 region of the skeletal muscle $\alpha 1$ subunit is not a phosphate acceptor [17]. From the amino acid sequence, there are several possible sites of PKA mediated phosphorylation in the C-terminal region of the cardiac $\alpha 1$ subunit [17]. The molecular mass of the $\alpha 1$ subunit of partially purified cardiac Ca channel is about 200 kDa [14–16] and is much smaller than that deduced from the amino acid sequence (M_r 242,771), leading to the notion that the C-terminal region containing potential phosphorylation sites is cleaved from the $\alpha 1$ subunit during partial purification. We investigated the phosphorylation of the $\alpha 1$ subunit in rabbit heart and CHO cells expressing the $\alpha 1$ subunit and the effect of db-cAMP and forskolin on the Ca current in CHO cells.

2. EXPERIMENTAL

2.1. Preparation of the polyclonal antibody, CR2

The 2.0-kb *Bgl*II (4892)/*Bam*HI (vector) fragment from pCARD3 [17] was cloned into the *Bam*HI site of pAR3038 [18] to yield pCAS9, in which the cDNA insert was positioned downstream of and in the same orientation as the ϕ 10 promoter for T7 RNA polymerase. The generated expression plasmid pCAS9 contains the 549 amino acid-coding region of the C terminus in the $\alpha 1$ subunit of the rabbit cardiac Ca channel, fused in-frame, with the translation initiation codon provided on the pAR3038. *E. coli* BL21 transformed with pCAS9 was cultured and the fusion protein obtained as an insoluble pellet [19]. The fusion protein (0.22 mg per rabbit) was further purified by SDS-PAGE. The fusion protein, which located on the gel by copper chloride staining [20], was fragmented and injected into multiple sites on

Correspondence address: M. Takahashi, Mitsubishi Kasei Institute of Life Sciences, Machida, Tokyo 194, Japan. Fax: (81) (427) 29-1252.

Abbreviations: db-cAMP, dibutyryl cAMP; CHO, Chinese hamster ovary; DHP, dihydropyridine; I_{Ba} , Ba current; PKA, cAMP-dependent protein kinase; PN200-110, isopropyl 4-(2,1,3-benzoxadiazol-4-yl)-1,4-dihydro-2,6-dimethyl-5-(methoxycarbonyl)pyridine-3-carboxylate.

the back of rabbit with Freund's complete adjuvant. Immunization without adjuvant was repeated every 3–4 weeks and blood was taken 7 to 10 days after each booster.

2.2. Isolation of CHO cells expressing the $\alpha 1$ subunit of rabbit cardiac DHP-sensitive Ca channel

The 7.0-kb *HindIII* fragment containing the entire protein-coding sequence of the cardiac $\alpha 1$ subunit of DHP-sensitive Ca channel from pCARD1 [17] was cloned into the *HindIII* site of pKNH [21] to yield pCCAR. CHO cells were transfected with *PvuII*-cleaved pCCAR. Clones CCAR3217 and CCAR2823 were isolated by screening G418-resistant clones by RNA blotting using the cDNA of cardiac $\alpha 1$ subunit as the probe. Cells were maintained in MEM α medium (Gibco, no. 410-1900) supplemented with 10% calf serum.

2.3. Membrane preparation

Cardiac membranes were prepared from rabbit ventricles [22]. CHO and CCAR cells were homogenized in NEH solution (125 mM NaCl, 1 mM EDTA, and 25 mM HEPES-Tris, pH 7.4) by sonication. The homogenates were centrifuged at $540,000 \times g$ for 10 min at 4°C and the membrane pellet was washed once with 0.5 M KCl/NEH, resuspended in NEH and stored at -80°C. Protease inhibitors (1 mM phenylmethanesulfonyl fluoride, 1 μ M pepstatin A, 1 mM 1,10-phenanthroline, 1 μ g/ml antipain and 1 μ g/ml leupeptin) were added throughout the preparation.

2.4. Immunoprecipitation and immunoblotting of the Ca channel proteins with CR2

The [3 H]PN200-110-labeled rabbit cardiac Ca channels were precipitated as previously described [23]. Immunoblotting was performed as described in [24] except using a semi-dry transblotting apparatus. Transblot SD (Bio-Rad) and a blotting solution containing 20% methanol, 48 mM Tris, 39 mM glycine, and 1.3 mM SDS at pH 9.2. The ECL Western blotting detection reagent (Amersham) was used for detection.

2.5. Phosphorylation of the $\alpha 1$ subunit in vitro and in vivo

Membranes were solubilized with 1% digitonin in NEH and the Ca channel proteins were immunoprecipitated with CR2 or control rabbit IgG and phosphorylated with a catalytic subunit of PKA (35.7 U) as described in [14]. The phosphorylated proteins were analyzed by SDS-PAGE and a Fuji Bioimage-analyzer BAS 2000 (Fuji Photo Film Co.) using an imaging plate [25].

Phosphorylation of the $\alpha 1$ subunit in vivo was measured by backphosphorylation [26]. CCAR cells (2×10^4 per 35 mm plastic culture dish, Corning) were incubated in a low- K^+ solution (140 mM NaCl, 4.7 mM KCl, 1.2 mM KH_2PO_4 , 2.5 mM $CaCl_2$, 1.2 mM $MgSO_4$, 11 mM glucose, and 15 mM HEPES-Tris, pH 7.4) in the presence or absence of db-cAMP (5 mM) or forskolin (100 μ M) for various periods at 37°C. The cells were solubilized with 3% CHAPS in NEHDPF (75 mM NaCl, 50 mM Na-phosphate, 2.5 mM EDTA, 20 mM NaF, 50 mM HEPES-Tris, and the protease inhibitors, pH 7.4) and incubated with either CR2 or control rabbit IgG for 2 h at 4°C. The antigen-antibody complex was absorbed onto protein A-Sepharose, washed three times by centrifugation with the same solution and three times with phosphorylation buffer (0.1% CHAPS, 6 mM EGTA, 6 mM $MgCl_2$, and 50 mM HEPES-Tris, pH 7.4), then incubated in phosphorylation buffer containing 2 μ M [γ - 32 P]ATP (5 μ Ci) and the catalytic subunit of PKA (16 U) for 10 min at 30°C. The reaction was stopped by adding cold 0.3% CHAPS in NEHDPF and the phosphorylated proteins were analyzed as described above.

2.6. Electrophoresis

Proteins were separated by SDS-PAGE as described previously [23] using a linear 4–12% acrylamide gradient gel (Tefco).

2.7. Electrophysiological measurements

The membrane current was measured by means of the patch-clamp method [27] using a pipette solution containing 65 mM CsCl, 60 mM

CsOH, 55 mM aspartic acid, 5 mM $MgCl_2$, 5 mM EGTA, 5 mM ATP, 5 mM phosphocreatine and 10 mM HEPES at pH 7.4 with CsOH. Cells were seeded in the chamber mounted on the inverted microscope, and perfused with Ca^{2+} -free Tyrode's solution (136.9 mM NaCl, 5.4 mM KCl, 0.5 mM $MgCl_2$, 0.33 mM NaH_2PO_4 , 5 mM glucose and 5 mM HEPES at pH 7.4 with NaOH). When the whole cell configuration was achieved, the solution was replaced with BA^{2+} -recording solution (136.9 mM tetrabutyl ammonium chloride, 5.4 mM CsCl, 0.5 mM $MgCl_2$, 0.33 mM NaH_2PO_4 , 10 mM BaCl₂ and 5 mM HEPES at pH 7.4 with tetrabutyl ammonium hydroxide) to obtain a high enough amplitude of I_{Ca} through the DHP-sensitive Ca channels. We used an experimental set-up connected on line to a computer (Atari MEGA ST-4) to supply command potentials and to acquire data immediately. A patch clamp amplifier, EPC-7 (List electronics, DA-Eberstadt, Germany) was used to record currents. Currents were low pass filtered (1 kHz) by an eight-pole Bessel filter, viewed on a CRT computer terminal, and recorded on a hard disk for further analysis.

2.8. Materials

The sources of drugs and chemicals were as follows: the catalytic subunit of type I cAMP-dependent protein kinase, Sigma; [3 H]PN200-110 and [γ - 32 P]ATP, New England Nuclear; peroxidase-conjugated goat anti-rabbit IgG, Zymed Laboratories; protein A-Sepharose CL-4B, Pharmacia, LKB Biotechnology; db-cAMP and 8-bromo-cAMP, Yamasa; digitonin, Wako Pure Chemical Industries; CHAPS, Dojin; ECL Western blotting detection reagent, Amersham; and all other chemicals, Wako, Sigma, or Bio-Rad Laboratories.

3. RESULTS AND DISCUSSION

A polyclonal antibody, CR2, was generated by immunizing a rabbit with the C-terminal region of the $\alpha 1$ subunit of the rabbit DHP-sensitive cardiac Ca channel expressed in *E. coli*. As shown in Fig. 1, CR2 im-




Fig. 1. Immunoprecipitation of [3 H]PN200-110-labeled cardiac calcium channels by CR2. The [3 H]PN200-110-labeled calcium channels solubilized and partially purified from rabbit heart were incubated with the indicated amounts of CR2 (●) or control rabbit IgG (○), then the antigen-antibody complexes were precipitated by absorption onto protein A-Sepharose. The radioactivity recovered in the precipitate was expressed as a percentage of the total counts added.




Fig. 2. Immunoblotting of $\alpha 1$ subunit of rabbit heart and CCAR membranes with CR2. Membrane proteins were separated by SDS-PAGE, transblotted to nitrocellulose and immunostained with control IgG (A, lane 1 and B, lanes 1-3) or CR2 (A, lane 2 and B, lanes 4-6) at a concentration of 10 $\mu\text{g}/\text{ml}$. (A) Rabbit heart (4 μg per lane). (B) Lanes 1 and 2, CHO cells (20 μg per lane); lanes 2 and 5, CCAR2823 (10 μg per lane); lanes 3 and 6, CCAR3217 (4 μg per lane). The migration positions of molecular mass standards (kDa , $\times 10^{-3}$) are indicated on the right.

munoprecipitated more than 60% of [^3H]PN200-110-labeled Ca channel solubilized from rabbit cardiac membranes.

A polypeptide of 200 kDa was identified as the major antigen in the rabbit cardiac membranes by immunoblotting with CR2 (Fig. 2A). In addition to the 200 kDa band, a faint band at 250 kDa was detected by blotting. Since the molecular mass of the $\alpha 1$ subunit of the rabbit cardiac Ca channel deduced from the amino acid sequence is 242,771 [17], these results suggest that a large part of the $\alpha 1$ subunit in the cardiac membrane preparation is partially proteolysed. In order to confirm this assumption, we prepared two stable transformed CHO cells (CCAR2823 and CCAR3217) using an expression plasmid which carries the entire protein-coding sequence of the rabbit cardiac $\alpha 1$ subunit. As shown in Fig. 2B, a polypeptide of 250-kDa was identified by immunoblotting in the transformants, but not in untransformed CHO cells. There was a marked difference in the amount of expression between these two transformants and much more 250-kDa protein was expressed in CCAR3217 than in CCAR2823 cells.

To examine the phosphorylation, the $\alpha 1$ subunit was solubilized from the membrane fractions of rabbit heart and the two transformants, immunoprecipitated with CR2 and incubated with a catalytic subunit of PKA and [$\gamma\text{-}^{32}\text{P}$]ATP. As shown in Fig. 3, only the 250-kDa form of the $\alpha 1$ subunit from rabbit cardiac membranes was phosphorylated by PKA and no incorporation of P_i was detected in the 200-kDa component. These results are in good agreement with those previously obtained by us




Fig. 3. Phosphorylation of the cardiac $\alpha 1$ subunit from the rabbit heart and CCAR cell by PKA in vitro. The $\alpha 1$ subunits were solubilized from CHO cells (lanes 1 and 2), CCAR3217 cells (lanes 3 and 4) and rabbit hearts (lanes 5 and 6), and immunoprecipitated with either control IgG (lanes 1, 3 and 5) or CR2 (lanes 2, 4 and 6). The immunoprecipitates were incubated with [$\gamma\text{-}^{32}\text{P}$]ATP and a catalytic subunit of PKA for 10 min at 30°C. The phosphorylated proteins were analyzed by SDS-PAGE then image analysed. The migration position of the molecular mass standards (kDa , $\times 10^{-3}$), are indicated on the right.

[14] as well as by Chang and Hosey [15] using chick heart. The 250-kDa polypeptide expressed in CCAR2823 and CCAR3217 cells was also phosphorylated by PKA and this protein was not detected in non-transformed CHO cells. These results clearly showed that the $\alpha 1$ subunit of cardiac Ca channels can be phosphorylated by PKA and that a large part of the $\alpha 1$ subunit in the cardiac membranes lost the phosphorylation site, probably due to partial proteolysis. The partial cleavage of the $\alpha 1$ subunit similar to that found in the present study occurs in skeletal muscle [28]. The molecular mass of a large portion of the $\alpha 1$ subunit from the T-tubular membrane fraction was 170 kDa [5], which was smaller than that deduced from the amino acid sequence (212,018) [6] and that expressed in L cells (195 kDa) [29]. A minor component of the $\alpha 1$ subunit having a molecular mass of 212 kDa was identified in T-tubular membranes using an antibody against the C-terminal peptide of the skeletal $\alpha 1$ subunit [28, 30].

In order to determine whether endogenous PKA can phosphorylate the $\alpha 1$ subunit incorporated in the cellular membrane, cAMP-dependent phosphorylation of the $\alpha 1$ subunit was studied using CCAR cells and




Fig. 4. Cyclic AMP-dependent phosphorylation of the $\alpha 1$ subunit of CCAR3217 cells in vivo. (A) CCAR3217 cultured on 35 mm plastic dishes were treated with 100 μ M forskolin (lane 5 to 8) or DMSO (lanes 1–4) for 10 min. After solubilizing the cells with CHAPS, the $\alpha 1$ subunit was immunoprecipitated with CR2 (lanes 2–4 and 6–8) or control rabbit IgG (lanes 1 and 5), then phosphorylated with $[\gamma\text{-}^{32}\text{P}]\text{ATP}$ and PKA. The phosphorylated proteins were separated by SDS-PAGE and analyzed with a Fuji Bioimage-analyzer. The migration position of the 250-kDa phosphoprotein is indicated by an arrow. (B) Results shown in panel A are quantified and the radioactivity incorporated into the 250-kDa bands is expressed in arbitrary units. (C) CCAR3217 cells were treated with 5 mM db-cAMP and solubilized with CHAPS in NEHDPF at the indicated times. The $\alpha 1$ subunit was immunoprecipitated with CR2, phosphorylated with $[\gamma\text{-}^{32}\text{P}]\text{ATP}$ and PKA, and the radioactivity incorporated into the $\alpha 1$ subunit was determined as described above.

backphosphorylation. The CCAR3217 cells were treated with or without forskolin and the $\alpha 1$ subunit was solubilized, immunoprecipitated with either CR2 or control rabbit IgG and incubated with PKA and $[\gamma\text{-}^{32}\text{P}]\text{ATP}$. As shown in Fig. 4, the incorporation of ^{32}P into the 250-kDa protein was suppressed by 35% by forskolin. This suppression was also observed after treatment with the membrane permeable cAMP analog, db-cAMP (Fig. 4C). These results indicated that the 250-kDa protein expressed in the CCAR3217 cells was phosphorylated in vivo probably by endogenous PKA in response to an elevated intracellular cAMP concentration. As shown in Fig. 4C, phosphorylation occurred very rapidly and ceased within 5 min after the treatment.

DHP-sensitive Ca channels in cardiac muscles are activated by phosphorylation with PKA in β -adrenergic stimulation [1]. In order to determine whether PKA-mediated phosphorylation of the 250-kDa form of $\alpha 1$

subunit could affect the Ca channel function, we investigated the electrophysiological properties of the CCAR cells. Under our experimental conditions, we did not observe I_{Ca} in CHO cells. In CCAR3217, the depolarizing pulses from the holding potential at -60 mV induced inward currents (Fig. 5). The activation threshold was -20 mV and the peak amplitude of the current was obtained at $+20$ mV. The current was completely inhibited by 10 μ M nifedipine (inset of Fig. 5B). Although these characteristics of the inward current recorded from CCAR were similar to those of the DHP-sensitive Ca channel current recorded in the heart [31], the inactivation time course of the current was very slow as shown in Fig. 5A. The inactivation time constant was 1.11 s at $+20$ mV. However, the voltage dependency of the steady state inactivation of I_{Ca} in CCAR was similar to that in the heart [31]. The apparent inactivation of the current was observed at below -60 mV and 50% inactivation was obtained at -32 mV (data not shown). An




Fig. 5. The Ba current recorded from CCAR3217. (A) The family of currents obtained by depolarization to -50, -30, -10, +10 and +30 mV for 500 ms from a holding potential of -60 mV. (B) The current-voltage relationship of I_{Ba} . The current amplitude measured at 80 ms after the onset of a depolarizing pulse is plotted against the membrane potential. The inset shows superimposed current records at +30 mV obtained in the absence (O) and the presence (●) of 10 μ M nifedipine.

I_{Ba} of similar characteristics was also obtained in CCAR2823, however, the amplitude of the current was significantly lower than that in CCAR3217, possibly due to the lower expression of the $\alpha 1$ subunit in CCAR2823 cells (see Fig. 2). These results are consistent with those previously reported [17, 32-34] indicating that the $\alpha 1$ subunit plays a central role in the DHP-sensitive Ca channel function.

Fig. 6A and B shows the increasing effect of db-cAMP on I_{Ba} . The effect was most remarkably observed in this particular experiment. In the presence of 200 μ M db-cAMP, the amplitude of I_{Ba} elicited by depolarization to +20 mV from a holding potential of -60 mV, was increased from -47.1 pA to -119.0 pA. The inactivation time constants of I_{Ba} at this potential were 1.29 s and 1.53 s, in the absence and the presence of the drug, respectively. The amplitude of I_{Ba} at +20 mV gradually increased after adding db-cAMP, and reached a steady

level about 5 min thereafter (Fig. 6C). This was consistent with the time course of the phosphorylation of the $\alpha 1$ subunit (see Fig. 4C). A similar effect of db-cAMP on I_{Ba} was also observed in CCAR2823. We observed an increase in the amplitude of I_{Ba} by db-cAMP in 8 experiments out of 19 in both types of CCAR cells. Another membrane-permeable cAMP analog, 8-bromo-cAMP, also increased the amplitude of I_{Ba} in 14 experiments out of 25. The degree of the increase of the current varied among cells, and it was usually about 20%. Neither drug increased the I_{Ba} at concentrations below 100 μ M. The decrease in the amplitude of I_{Ba} was not observed in both CCAR cells by adding the membrane-permeable cAMP analogs.

The present studies showed that: (i) the $\alpha 1$ subunit of cardiac Ca channel could be phosphorylated by PKA in vitro and in vivo and that the Ca channel was activated by phosphorylation, and (ii) that the segment containing the phosphorylation site was cleaved from a large part of the $\alpha 1$ subunit in the cardiac membrane. It is not known when the cleavage occurs and at least three possibilities could be considered for the structure of the $\alpha 1$ subunit in cardiac cells: (1) the 250-kDa form of the $\alpha 1$ subunit is predominant in living cardiac cells and the partial degradation resulting in the 200-kDa polypeptide due to the proteolytic cleavage occurs during the isolation of the cardiac membrane; (2) the C-terminal segment of most of the $\alpha 1$ subunit is cleaved but still associated with the 200 kDa polypeptide and functions as a subunit. The $\alpha 2$ and δ subunits of skeletal muscle Ca channels are encoded by the same gene, the protein product of which is proteolytically processed to yield the disulfide-linked $\alpha 2$ and δ polypeptide [7,28]; (3) the C-terminal segment containing the phosphorylation site(s) was lost due to the partial proteolysis in cardiac cells. In the third possibility, the phosphorylation of the C-terminal segment could not be involved in the β -adrenergic-mediated Ca channel regulation. Thus, the identification of the $\alpha 1$ subunit in cardiac cells should be studied for further understanding of the regulatory mechanisms of the cardiac Ca channel function.

The amplitude of I_{Ba} observed in considerable numbers of CCAR cells is much lower than that in cardiac myocytes. Furthermore, the extent of the activation by cAMP-dependent phosphorylation in the CCAR cells was also less than that in the heart. It has been shown that I_{Ba} expressed by injecting the $\alpha 1$ message into *Xenopus* oocytes and L cells was increased remarkably by the coexpression of β and γ subunits of the skeletal muscle Ca channel [17, 32-34]. Northern blots have shown that the β subunit homologous to that of skeletal muscle was expressed in cardiac muscle [34]. Thus, the formation of a multisubunit complex might be necessary for expressing the activation of DHP-sensitive Ca channels. Further studies, including the coexpression of the other subunits are necessary to obtain a final conclusion.




Fig. 6. The effect of db-cAMP on I_h in CCAR3217. (A) Currents in the absence of db-cAMP recorded by depolarization to -40, -20, 0, +20 and +40 mV for 500 ms from a holding potential of -60 mV are superimposed. (B) Currents in the presence of db-cAMP (200 μ M) recorded at same membrane potentials as those in A are superimposed. (C) The time course of the effect of db-cAMP on the amplitude of I_h . The amplitude of I_h measured at 80 ms after the onset of the pulse is plotted against the time.

Acknowledgements: We thank A. Mikami for providing the cDNA clone (pCARD1) and professor S. Numa for his continuous support and encouragement throughout this study. This investigation was supported in part by a research grant from the Institute of Physical and Chemical Research.

REFERENCES

- [1] Reuter, H. (1983) *Nature* 301, 569-574.
- [2] Tsien, R.W. (1977) *Adv. Cyclic Nucleotide Res.* 8, 363-420.
- [3] Kameyama, M., Hofmann, F. and Trautwein, W. (1985) *Pflüger's Arch. Eur. J. Physiol.* 405, 285-293.
- [4] Hescheler, J., Kameyama, M., Trautwein, W., Mieskes, G. and Soling, H.-D. (1987) *Eur. J. Biochem.* 165, 261-266.
- [5] Catterall, W.A., Sengar, M.J. and Takahashi, M. (1988) *J. Biol. Chem.* 263, 3535-3538.
- [6] Tanabe, T., Takeshima, H., Mikami, A., Flockerzi, V., Takahashi, H., Kungawa, K., Kojima, M., Matsuo, H., Hirose, T. and Numa, S. (1987) *Nature* 328, 313-318.
- [7] Ellis, S.B., Williams, M.E., Ways, N.R., Brenner, R., Sharp, A.H., Leung, A.T., Campbell, K.P., McKenna, E., Koch, W.J., Hui, A., Schwartz, A. and Harpold, M.M. (1988) *Science* 241, 1661-1664.
- [8] Ruth, P., Rohrkasten, A., Biel, M., Bosse, E., Regulla, S., Meyer, H.E., Flockerzi, V. and Hofmann, F. (1989) *Science* 245, 1115-1118.
- [9] Jay, S.D., Ellis, S.B., McCue, A.F., Williams, M.E., Vedvick, T.S., Harpold, M.M. and Campbell, K.P. (1990) *Science* 248, 490-492.
- [10] Curtis, B.M. and Catterall, W.A. (1985) *Proc. Natl. Acad. Sci. USA* 82, 2528-2532.
- [11] Flockerzi, V., Ocken, H., Hofmann, F., Pelzer, D., Cavalie, A. and Trautwein, W. (1986) *Nature* 323, 66-68.
- [12] Nunoki, K., Florio, V. and Catterall, W.A. (1989) *Proc. Natl. Acad. Sci. USA* 86, 6816-6820.
- [13] Rohrkasten, A., Meyer, H.E., Nastainczyk, W., Sieber, M. and Hofmann, F. (1988) *J. Biol. Chem.* 263, 15325-15329.
- [14] Yoshida, A., Takahashi, M., Fujimoto, Y., Takisawa, H. and Nakamura, T. (1990) *J. Biochem.* 107, 608-612.
- [15] Chang, F.C. and Hosey, M.M. (1988) *J. Biol. Chem.* 263, 18929-18937.
- [16] Schneider, T. and Hofmann, F. (1988) *Eur. J. Biochem.* 174, 369-375.
- [17] Mikami, A., Imoto, K., Tanabe, T., Niidome, T., Mori, Y., Takeshima, H., Narumiya, S. and Numa, S. (1989) *Nature* 340, 230-233.
- [18] Rosenberg, A., Lade, B.N., Chui, D., Lin, S., Dunn, J.J. and Studier, F.W. (1987) *Gene* 56, 125-135.

- [19] Watt, R.A., Shatzman, A.R. and Rosenberg, M. (1985) *Mol Cell. Biol.* 5, 448-456.
- [20] Lee, C., Levin, A. and Branton, D. (1987) *Anal. Biochem.* 166, 308-312.
- [21] Fukuda, K., Higashida, H., Kubo, T., Maeda, A., Akiba, I., Bujo, H., Mishima, M. and Numa, S. (1988) *Nature* 335, 355-358.
- [22] Jones, L.R., Besch, H.R., Fleming, J.W., McConnaughey, M.M. and Watanabe, A.M. (1979) *J. Biol. Chem.* 254, 530-539.
- [23] Takahashi, M. and Catterall, W.A. (1987) *Biochemistry* 26, 5518-5526.
- [24] Yoshida, A., Ogura, A., Imagawa, T., Shigekawa, M. and Takahashi, M. (1992) *J. Neurosci.* 12, 1094-1100.
- [25] Amemiya, Y. and Miyahara, J. (1988) *Nature* 336, 89-90.
- [26] Nestler, E.J. and Greengard, P. (1984) *Protein Phosphorylation in the Nervous System*, Wiley, New York.
- [27] Hamill, O.P., Marty, A., Neher, E., Sakmann, B. and Sigworth, F.J. (1981) *Pflüger's Arch.* 391, 85-100.
- [28] De Jongh, K.S., Merrick, D.K. and Catterall, W.A. (1989) *Proc. Natl. Acad. Sci. USA* 86, 8585-8589.
- [29] Perez-Reyes, E., Kim, H.S., Lacerda, A.E., Horne, W., Wei, X., Rampe, D., Campbell, K.P., Brown, A.M. and Birnbaumer, L. (1989) *Nature* 340, 233-236.
- [30] Lai, Y., Seagar, M.J., Takahashi, M. and Catterall, W.A. (1991) *J. Biol. Chem.* 265, 20839-20848.
- [31] Hess, P., Lansman, J.B. and Tsien, R.W. (1984) *Nature* 311, 538-544.
- [32] Perez-Reyes, E., Castellano, A., Kim, H.S., Bertrand, P., Bagstrom, E., Lacerda, A.E., Wei, X. and Birnbaumer, L. (1992) *J. Biol. Chem.* 267, 1792-1797.
- [33] Itagaki, K., Koch, W.J., Bodi, I., Klockner, U., Slish, D.F. and Schwartz, A. (1992) *FEBS Lett.* 297, 221-225.
- [34] Hullin, R., Singer-Lahat, D., Freichel, M., Biel, M., Dascal, N., Hofmann, F. and Flockerzi, V. (1992) *EMBO J.* 11, 885-890.